

**PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration (grouped)	Breastfeeding (any) duration (grouped)

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/28/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


**PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	1,700	10.9	9.6 - 12.4	500	3.4	2.7 - 4.4	2,300	14.5	13.1 - 16.1	600	3.5	2.8 - 4.4	10,600	67.6	65.5 - 69.7	15,800
2001	1,600	10.3	9.1 - 11.7	600	4.1	3.3 - 5.0	2,400	15.3	13.7 - 17.0	400	2.7	2.0 - 3.5	10,600	67.7	65.5 - 69.7	15,600
2002	1,700	10.8	9.2 - 12.7	600	3.8	2.9 - 5.1	2,200	13.8	12.0 - 15.8	400	2.3	1.6 - 3.2	11,200	69.3	66.6 - 71.8	16,100
2003	1,800	10.8	9.2 - 12.6	700	4.2	3.2 - 5.4	2,400	14.6	12.7 - 16.7	600	3.9	3.0 - 5.1	11,000	66.5	63.9 - 69.1	16,500
2004	1,600	9.0	7.8 - 10.4	700	3.8	3.1 - 4.8	2,500	14.2	12.7 - 15.8	700	3.8	3.1 - 4.7	12,000	69.2	67.1 - 71.2	17,300
2005	1,700	10.0	8.6 - 11.6	600	3.8	2.9 - 4.8	2,500	14.4	12.8 - 16.2	500	3.1	2.3 - 4.0	11,700	68.8	66.5 - 70.9	17,000
2006	1,400	8.0	6.8 - 9.5	500	2.6	1.9 - 3.6	2,800	16.0	14.2 - 17.9	600	3.2	2.5 - 4.2	12,400	70.1	67.8 - 72.3	17,700
2007	1,400	7.6	6.4 - 9.0	600	3.5	2.7 - 4.6	2,700	14.9	13.3 - 16.7	500	3.1	2.3 - 4.0	12,700	70.9	68.6 - 73.1	17,900
2008	1,400	7.8	6.5 - 9.2	500	2.6	2.0 - 3.5	2,600	14.1	12.5 - 15.9	800	4.2	3.3 - 5.3	12,900	71.3	69.0 - 73.4	18,200
2009	1,200	6.7	5.2 - 8.6	800	4.3	3.2 - 5.9	2,600	14.6	12.5 - 17.0	900	5.0	3.7 - 6.6	12,500	69.4	66.4 - 72.3	17,900
2010	1,100	6.0	4.6 - 7.9	600	3.1	2.1 - 4.5	2,400	13.3	11.3 - 15.7	700	4.0	2.9 - 5.5	13,400	73.6	70.6 - 76.4	18,300
2011	800	4.5	3.2 - 6.1	300	1.6	1.0 - 2.7	2,200	12.2	10.2 - 14.5	700	3.8	2.8 - 5.2	14,200	77.9	75.1 - 80.5	18,300
2012	900	5.1	3.8 - 6.9	500	2.6	1.7 - 4.0	2,200	12.3	10.3 - 14.7	600	3.1	2.1 - 4.4	14,000	76.8	73.9 - 79.5	18,200
2013	700	3.7	2.6 - 5.2	400	2.0	1.2 - 3.3	2,100	11.5	9.5 - 13.9	500	2.9	1.9 - 4.4	14,400	79.9	76.9 - 82.5	18,100

**PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	700	3.7	2.6 - 5.2	400	2.0	1.2 - 3.3	2,100	11.5	9.5 - 13.9	500	2.9	1.9 - 4.4	14,400	79.9	76.9 - 82.5	18,100

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.0	3.7 - 9.6	100	2.4	1.1 - 5.0	200	9.6	6.7 - 13.4	100	2.2	1.0 - 4.7	1,900	79.9	74.9 - 84.1	2,300
Honolulu	400	3.4	2.0 - 5.5	200	1.9	0.9 - 3.8	1,500	11.6	8.9 - 14.9	400	3.2	1.9 - 5.3	10,400	80.0	76.0 - 83.5	12,900
Kauai	<50	4.4	2.7 - 7.1	<50	0.9	0.3 - 2.8	100	12.5	9.3 - 16.5	<50	2.8	1.5 - 5.2	700	79.5	74.8 - 83.4	800
Maui	100	3.0	1.6 - 5.7	100	2.6	1.3 - 5.2	200	12.9	9.5 - 17.1	<50	2.3	1.1 - 4.7	1,500	79.2	74.3 - 83.4	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	1.6	0.5 - 4.6	100	2.2	0.7 - 6.4	500	9.4	5.9 - 14.6	100	2.7	1.2 - 6.1	4,100	84.1	78.1 - 88.7	4,900
Native Hawaiian	300	6.3	3.9 - 10.0	100	2.6	1.3 - 5.2	900	16.1	12.0 - 21.4	200	3.8	2.0 - 7.0	3,800	71.2	65.3 - 76.5	5,300
Chinese	<50	3.3	0.5 - 19.7	<50	0.9	0.1 - 5.5	100	16.4	7.5 - 32.1	0	0.0	-	700	79.4	63.2 - 89.6	900
Filipino	100	1.9	0.6 - 5.9	100	3.3	1.2 - 8.7	300	12.0	8.0 - 17.7	100	4.2	1.6 - 10.4	2,200	78.5	70.9 - 84.6	2,900
Japanese	<50	1.7	0.2 - 10.7	0	0.0	-	100	2.9	0.9 - 9.1	0	0.0	-	1,700	95.4	87.8 - 98.4	1,800
Other	100	6.3	3.1 - 12.4	<50	0.5	0.1 - 3.0	200	9.4	4.9 - 17.2	100	3.4	1.0 - 10.4	1,800	80.4	71.4 - 87.1	2,300

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	1.4	0.2 - 8.4	<50	1.8	0.5 - 6.5	200	21.5	11.8 - 35.9	<50	5.8	1.3 - 22.3	600	69.5	54.2 - 81.4	800
20-24	100	2.1	0.8 - 5.2	100	1.9	0.7 - 5.1	400	9.6	6.2 - 14.5	200	5.1	2.4 - 10.5	3,300	81.3	74.8 - 86.4	4,000
25-34	400	3.9	2.4 - 6.2	200	2.2	1.1 - 4.4	1,200	12.6	9.7 - 16.3	200	1.6	0.9 - 3.0	7,800	79.7	75.5 - 83.3	9,800
35+	200	5.7	3.1 - 10.5	100	1.6	0.5 - 5.0	300	8.2	4.9 - 13.4	100	3.4	1.4 - 8.0	2,800	81.0	74.3 - 86.3	3,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	4.2	2.7 - 6.5	200	2.1	1.0 - 4.2	900	10.1	7.6 - 13.4	200	2.4	1.3 - 4.6	7,200	81.2	77.1 - 84.7	8,900
Female	300	3.3	1.9 - 5.5	200	1.9	0.9 - 3.9	1,200	12.8	9.8 - 16.6	300	3.4	2.0 - 5.9	7,200	78.6	74.2 - 82.4	9,200

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.6	1.5 - 4.3	200	2.1	1.0 - 4.0	1,300	10.9	8.3 - 14.1	200	2.1	1.2 - 3.7	9,500	82.4	78.6 - 85.6	11,500
No	400	5.8	3.6 - 9.0	100	1.9	0.9 - 3.7	800	12.6	9.5 - 16.5	300	4.4	2.4 - 7.9	4,900	75.4	70.4 - 79.8	6,500

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.9	3.1 - 7.5	100	2.3	1.1 - 4.6	900	14.4	10.8 - 18.9	300	4.4	2.5 - 7.7	4,900	74.1	68.8 - 78.7	6,600
131-185%	200	6.8	3.2 - 13.7	100	2.3	0.7 - 7.0	300	11.9	6.9 - 19.7	<50	1.7	0.3 - 8.4	1,900	77.4	68.3 - 84.5	2,500
186+%	200	2.4	1.1 - 5.1	200	2.1	0.9 - 4.9	500	7.4	4.9 - 10.9	100	1.9	0.9 - 4.0	6,300	86.3	81.8 - 89.8	7,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Income (2012+)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	200	4.8	2.7 - 8.7	100	2.0	0.8 - 5.0	600	15.8	11.2 - 21.9	200	4.4	2.0 - 9.2	3,000	72.9	66.0 - 78.8	4,100
\$17,001-\$26,000	100	4.3	2.1 - 8.6	<50	0.6	0.1 - 2.6	300	14.3	8.8 - 22.6	100	2.2	0.6 - 8.0	1,900	78.5	69.8 - 85.2	2,400
\$26,001-\$51,000	200	4.8	2.4 - 9.3	200	4.7	2.3 - 9.6	300	8.8	5.3 - 14.1	100	3.5	1.5 - 7.7	3,100	78.3	71.2 - 84.0	3,900
>=\$51,001	200	2.8	1.3 - 5.8	100	1.2	0.3 - 4.0	500	7.5	4.9 - 11.3	100	1.6	0.7 - 3.7	5,400	87.0	82.3 - 90.5	6,200

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	3.2	1.8 - 5.7	<50	1.1	0.4 - 3.0	200	14.7	11.6 - 18.4	100	4.2	2.6 - 6.7	1,000	76.9	72.4 - 80.7	1,300
Normal birth weight	600	3.8	2.6 - 5.4	300	2.1	1.2 - 3.5	1,900	11.3	9.1 - 13.9	500	2.8	1.8 - 4.5	13,400	80.1	76.9 - 82.9	16,800

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.0	3.3 - 18.1	100	6.6	2.0 - 19.7	200	19.6	10.8 - 33.1	<50	1.5	0.5 - 4.5	600	64.3	50.1 - 76.4	900
No	600	3.6	2.4 - 5.1	300	1.8	1.0 - 3.1	1,900	11.1	9.0 - 13.6	500	3.0	1.9 - 4.6	13,500	80.6	77.5 - 83.3	16,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	4.8	2.5 - 9.0	200	5.3	2.6 - 10.5	500	16.0	10.8 - 23.1	100	3.0	1.1 - 8.1	2,300	70.9	62.9 - 77.8	3,200
No	500	3.6	2.4 - 5.3	200	1.3	0.6 - 2.7	1,500	10.6	8.4 - 13.2	400	2.9	1.8 - 4.6	11,800	81.7	78.4 - 84.5	14,400

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	4.8	3.1 - 7.4	200	2.3	1.2 - 4.5	1,200	15.8	12.3 - 20.1	300	4.1	2.4 - 6.8	5,700	73.0	68.1 - 77.4	7,800
No	300	2.9	1.7 - 5.0	200	1.8	0.8 - 3.7	800	8.0	5.8 - 10.8	200	2.1	1.0 - 4.1	8,700	85.3	81.6 - 88.3	10,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	900	5.1	3.8 - 6.9	500	2.6	1.7 - 4.0	2,200	12.3	10.3 - 14.7	600	3.1	2.1 - 4.4	14,000	76.8	73.9 - 79.5	18,200

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.0	2.4 - 6.7	<50	1.2	0.4 - 3.3	300	14.5	11.0 - 18.7	100	4.9	3.0 - 7.9	1,700	75.5	70.6 - 79.8	2,300
Honolulu	800	5.9	4.1 - 8.4	400	3.0	1.8 - 4.9	1,600	11.8	9.3 - 15.0	400	2.7	1.6 - 4.6	10,100	76.6	72.6 - 80.1	13,200
Kauai	<50	3.4	1.7 - 6.8	<50	3.4	1.7 - 6.8	100	11.3	7.8 - 16.1	<50	2.1	0.9 - 4.9	700	79.8	74.1 - 84.5	800
Maui	<50	1.7	0.8 - 3.5	<50	1.9	0.9 - 4.0	300	13.7	10.5 - 17.7	100	3.7	2.3 - 6.1	1,500	78.9	74.4 - 82.8	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.2	2.7 - 9.6	100	2.9	1.2 - 6.9	300	8.3	5.2 - 13.1	100	1.4	0.4 - 4.2	3,400	82.2	76.1 - 87.0	4,100
Native Hawaiian	300	5.3	3.0 - 9.2	100	2.0	0.9 - 4.4	900	17.8	13.7 - 22.9	300	5.3	3.3 - 8.4	3,600	69.6	63.8 - 74.9	5,200
Chinese	100	8.4	3.0 - 21.1	100	4.7	1.2 - 16.7	100	5.2	1.5 - 16.5	<50	2.9	0.4 - 17.5	900	78.8	64.4 - 88.4	1,100
Filipino	100	2.8	1.1 - 6.9	100	4.2	1.9 - 8.8	600	19.5	14.0 - 26.7	100	4.2	1.9 - 9.3	2,200	69.2	61.4 - 76.0	3,200
Japanese	<50	1.6	0.2 - 10.4	<50	0.3	0.1 - 1.7	100	5.8	2.5 - 12.8	<50	1.0	0.3 - 2.9	1,400	91.2	83.5 - 95.5	1,600
Other	200	7.8	4.2 - 14.1	100	2.2	0.7 - 7.2	200	6.8	3.7 - 12.1	<50	1.6	0.4 - 6.0	2,500	81.6	74.1 - 87.4	3,000

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	3.1	0.5 - 18.6	100	8.9	3.3 - 21.8	200	24.5	14.7 - 37.9	<50	2.6	0.9 - 7.5	600	60.8	46.8 - 73.3	1,000
20-24	200	4.8	2.4 - 9.2	100	3.5	1.5 - 7.7	600	16.7	11.9 - 23.0	100	3.6	1.6 - 7.6	2,700	71.5	64.4 - 77.7	3,700
25-34	500	5.1	3.4 - 7.7	200	1.9	1.0 - 3.6	1,200	11.4	8.8 - 14.5	400	3.5	2.2 - 5.7	8,000	78.0	74.1 - 81.6	10,200
35+	200	6.1	3.2 - 11.4	100	2.1	0.7 - 5.9	200	6.6	3.9 - 10.7	<50	1.2	0.6 - 2.4	2,700	84.1	78.0 - 88.8	3,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	4.2	2.6 - 6.9	300	3.5	2.0 - 5.8	1,100	12.7	9.9 - 16.1	200	2.6	1.5 - 4.7	7,000	77.0	72.7 - 80.8	9,000
Female	600	6.0	4.1 - 8.8	200	1.8	0.9 - 3.6	1,100	12.0	9.3 - 15.3	300	3.5	2.2 - 5.6	7,000	76.6	72.5 - 80.3	9,200

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.3	2.8 - 6.5	300	2.3	1.2 - 4.1	1,200	10.3	8.0 - 13.2	300	2.5	1.5 - 4.2	9,100	80.6	77.0 - 83.8	11,200
No	500	6.5	4.2 - 9.9	200	3.3	1.9 - 5.7	1,100	15.6	12.2 - 19.7	300	3.9	2.4 - 6.5	4,900	70.7	65.7 - 75.3	7,000

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.5	4.1 - 10.1	300	4.2	2.4 - 7.1	1,000	16.2	12.6 - 20.6	200	2.4	1.2 - 4.8	4,500	70.7	65.4 - 75.4	6,400
131-185%	200	6.2	3.1 - 12.0	<50	1.3	0.3 - 6.1	400	13.1	8.5 - 19.8	200	5.2	2.6 - 10.2	2,300	74.1	66.1 - 80.7	3,100
186+%	100	1.8	0.8 - 4.2	100	1.7	0.7 - 4.2	600	8.6	6.0 - 12.1	200	2.8	1.5 - 5.1	5,900	85.1	80.8 - 88.6	6,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Income (2012+)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	200	5.7	3.2 - 10.1	200	4.5	2.3 - 8.4	700	16.2	11.7 - 21.9	100	2.2	0.8 - 5.4	2,900	71.5	64.8 - 77.3	4,100
\$17,001-\$26,000	100	2.9	1.0 - 8.4	100	2.3	0.6 - 8.6	300	14.0	9.0 - 21.2	100	6.3	3.1 - 12.1	1,700	74.5	65.9 - 81.5	2,200
\$26,001-\$51,000	400	8.6	5.2 - 13.9	100	2.4	1.0 - 5.8	700	14.7	10.4 - 20.4	100	2.6	1.1 - 5.7	3,200	71.7	65.0 - 77.5	4,400
>=\$51,001	100	1.7	0.6 - 4.3	100	1.6	0.6 - 4.2	500	7.7	5.2 - 11.4	200	2.9	1.5 - 5.6	5,100	86.2	81.6 - 89.7	5,900

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	3.8	2.2 - 6.6	<50	2.6	1.4 - 5.0	200	14.5	10.7 - 19.2	<50	3.2	1.7 - 5.9	900	75.9	70.6 - 80.5	1,200
Normal birth weight	900	5.2	3.8 - 7.2	500	2.6	1.7 - 4.1	2,100	12.2	10.1 - 14.7	500	3.1	2.1 - 4.5	13,100	76.8	73.7 - 79.7	17,000

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	4.5	2.1 - 9.7	<50	3.4	1.3 - 8.6	200	26.1	14.6 - 42.1	100	12.5	4.7 - 29.3	400	53.4	38.0 - 68.3	700
No	900	5.1	3.7 - 7.1	500	2.7	1.7 - 4.1	2,000	12.0	9.9 - 14.4	500	2.8	1.9 - 4.1	13,100	77.4	74.4 - 80.2	16,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.6	2.7 - 11.1	100	4.4	1.9 - 9.7	500	16.6	11.5 - 23.5	200	7.3	3.9 - 13.2	1,900	66.2	58.0 - 73.5	2,800
No	700	5.0	3.6 - 7.1	400	2.4	1.5 - 3.9	1,700	11.7	9.6 - 14.3	400	2.4	1.5 - 3.7	11,600	78.4	75.2 - 81.4	14,700

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	7.1	4.9 - 10.2	400	4.5	2.8 - 7.2	1,400	17.0	13.6 - 21.0	200	3.0	1.7 - 5.0	5,600	68.5	63.7 - 72.8	8,200
No	300	3.2	1.8 - 5.5	100	1.0	0.4 - 2.6	800	8.6	6.4 - 11.4	300	2.9	1.7 - 4.8	8,300	84.3	80.7 - 87.3	9,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	800	4.5	3.2 - 6.1	300	1.6	1.0 - 2.7	2,200	12.2	10.2 - 14.5	700	3.8	2.8 - 5.2	14,200	77.9	75.1 - 80.5	18,300

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.3	2.7 - 7.0	<50	1.8	0.9 - 3.8	400	15.4	12.0 - 19.6	100	4.0	2.4 - 6.6	1,800	74.4	69.6 - 78.7	2,400
Honolulu	600	4.7	3.1 - 7.0	200	1.5	0.7 - 3.0	1,500	11.7	9.1 - 14.8	500	3.7	2.4 - 5.6	10,300	78.5	74.7 - 81.9	13,100
Kauai	<50	2.1	0.9 - 4.8	<50	1.0	0.3 - 3.5	100	9.5	6.4 - 13.9	<50	4.1	2.2 - 7.4	700	83.3	78.1 - 87.5	900
Maui	100	4.2	2.7 - 6.6	100	2.7	1.5 - 4.7	200	12.8	9.9 - 16.3	100	4.6	3.0 - 6.9	1,500	75.8	71.5 - 79.6	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.5	1.1 - 5.7	100	1.3	0.4 - 4.1	500	11.6	7.7 - 17.0	100	3.2	1.6 - 6.3	3,600	81.4	75.5 - 86.1	4,500
Native Hawaiian	300	5.2	3.0 - 8.8	100	1.9	0.8 - 4.1	800	15.0	11.4 - 19.5	300	6.0	3.7 - 9.6	3,700	71.9	66.4 - 76.8	5,200
Chinese	100	9.4	2.9 - 26.6	<50	4.1	0.6 - 23.2	<50	2.1	0.7 - 5.7	<50	1.6	0.6 - 4.5	600	82.8	66.3 - 92.2	700
Filipino	200	5.1	2.7 - 9.5	<50	0.8	0.4 - 1.8	500	15.3	10.8 - 21.2	100	3.4	1.8 - 6.2	2,500	75.4	68.8 - 81.0	3,300
Japanese	<50	2.0	0.5 - 7.4	<50	0.3	0.1 - 2.1	200	10.1	5.2 - 18.9	<50	1.7	0.3 - 7.6	1,600	85.8	76.8 - 91.7	1,900
Other	200	5.8	2.5 - 12.7	100	3.1	1.2 - 7.9	200	8.1	4.4 - 14.5	100	3.2	1.2 - 8.4	2,200	79.8	71.4 - 86.2	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.5	1.8 - 20.9	<50	1.9	0.6 - 5.4	400	37.6	26.2 - 50.5	<50	3.1	1.2 - 7.8	500	50.9	38.4 - 63.2	1,000
20-24	300	6.9	4.0 - 11.6	100	2.1	0.9 - 4.9	700	15.7	11.3 - 21.3	100	1.2	0.7 - 2.2	3,200	74.0	67.6 - 79.6	4,300
25-34	400	4.0	2.5 - 6.3	200	1.8	0.9 - 3.5	1,000	10.2	7.7 - 13.3	500	4.9	3.2 - 7.2	7,700	79.1	75.2 - 82.6	9,800
35+	100	1.8	1.0 - 3.2	<50	0.4	0.1 - 1.3	200	5.2	3.0 - 8.8	100	4.4	2.3 - 8.4	2,800	88.3	83.5 - 91.8	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	4.5	2.8 - 7.0	200	2.5	1.4 - 4.3	1,100	11.9	9.2 - 15.2	400	3.9	2.5 - 6.0	7,200	77.4	73.3 - 81.0	9,300
Female	400	4.4	2.8 - 6.9	100	0.8	0.3 - 1.9	1,100	12.5	9.8 - 15.8	300	3.8	2.5 - 5.8	7,000	78.4	74.5 - 81.9	9,000

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.0	2.6 - 6.2	200	2.1	1.1 - 3.7	1,000	9.1	7.0 - 11.8	500	4.1	2.8 - 5.9	9,200	80.7	77.2 - 83.8	11,400
No	400	5.2	3.2 - 8.2	100	1.0	0.5 - 1.7	1,200	17.2	13.7 - 21.5	200	3.4	2.0 - 5.8	5,000	73.2	68.3 - 77.5	6,900

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.5	4.2 - 10.1	100	2.3	1.2 - 4.4	1,000	15.7	12.3 - 19.8	300	3.9	2.4 - 6.4	4,700	71.6	66.5 - 76.1	6,500
131-185%	100	4.8	2.1 - 10.5	<50	1.6	0.4 - 6.0	400	13.0	8.2 - 20.0	100	3.3	1.4 - 7.6	2,100	77.4	69.4 - 83.8	2,700
186+%	200	2.9	1.7 - 5.1	100	1.2	0.5 - 3.0	700	8.3	5.8 - 11.6	300	4.1	2.5 - 6.5	6,700	83.5	79.4 - 86.9	8,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	5.9	3.3 - 10.6	100	2.7	1.2 - 6.4	600	17.2	12.5 - 23.1	100	2.3	0.9 - 6.1	2,400	71.8	64.9 - 77.9	3,300
\$10,000-\$24,999	200	5.9	3.0 - 11.4	100	1.6	0.5 - 4.7	500	15.4	10.7 - 21.6	100	4.1	2.3 - 7.1	2,500	73.0	65.9 - 79.1	3,400
\$25,000-\$49,999	300	5.7	3.1 - 9.9	100	1.5	0.5 - 4.8	600	13.2	9.3 - 18.5	200	5.3	2.9 - 9.5	3,300	74.3	67.9 - 79.8	4,500
>=\$50,000	200	2.5	1.2 - 5.0	100	1.2	0.5 - 3.1	400	6.4	4.1 - 9.9	200	3.6	2.1 - 6.1	5,300	86.4	82.1 - 89.8	6,100

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	4.0	2.2 - 7.2	<50	2.4	1.2 - 4.9	200	17.7	13.7 - 22.6	100	5.8	3.6 - 9.1	900	70.1	64.6 - 75.1	1,200
Normal birth weight	800	4.5	3.2 - 6.3	300	1.6	0.9 - 2.7	2,000	11.8	9.7 - 14.2	600	3.7	2.6 - 5.2	13,400	78.5	75.5 - 81.2	17,000

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	9.1	3.4 - 22.3	<50	0.6	0.1 - 3.7	300	30.3	19.7 - 43.6	100	8.4	3.4 - 19.5	500	51.5	38.5 - 64.3	900
No	700	4.2	3.0 - 5.9	300	1.7	1.1 - 2.8	1,900	10.9	9.0 - 13.1	600	3.6	2.6 - 5.0	13,700	79.6	76.8 - 82.2	17,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	6.6	3.7 - 11.5	<50	0.5	0.2 - 1.5	800	21.6	16.4 - 27.8	200	4.5	2.2 - 9.0	2,300	66.9	59.8 - 73.2	3,500
No	600	3.9	2.7 - 5.7	300	1.9	1.2 - 3.2	1,400	9.6	7.6 - 12.0	500	3.7	2.6 - 5.2	11,800	80.9	77.8 - 83.6	14,600

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	5.9	3.9 - 8.8	100	1.7	0.9 - 3.1	1,300	15.9	12.8 - 19.7	300	3.6	2.3 - 5.7	6,000	72.9	68.4 - 76.9	8,200
No	300	3.3	2.0 - 5.3	200	1.6	0.8 - 3.3	900	8.9	6.6 - 11.8	400	4.0	2.6 - 6.1	8,100	82.2	78.5 - 85.4	9,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,100	6.0	4.6 - 7.9	600	3.1	2.1 - 4.5	2,400	13.3	11.3 - 15.7	700	4.0	2.9 - 5.5	13,400	73.6	70.6 - 76.4	18,300

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.0	3.1 - 7.9	100	3.7	2.2 - 6.2	300	15.1	11.8 - 19.2	100	2.7	1.4 - 5.1	1,700	73.5	68.7 - 77.8	2,300
Honolulu	900	6.7	4.8 - 9.3	400	3.2	1.9 - 5.1	1,700	13.0	10.3 - 16.2	600	4.6	3.1 - 6.7	9,600	72.6	68.6 - 76.3	13,200
Kauai	<50	4.0	2.2 - 7.1	0	0.0	-	100	10.6	7.5 - 14.9	<50	2.9	1.5 - 5.7	700	82.5	77.6 - 86.6	800
Maui	100	3.2	1.9 - 5.5	100	3.2	1.9 - 5.5	300	15.0	11.9 - 18.8	<50	2.1	1.1 - 4.0	1,500	76.5	72.1 - 80.3	2,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.3	3.6 - 10.8	100	3.3	1.5 - 7.2	400	8.9	5.6 - 14.0	300	7.4	4.5 - 12.0	3,300	74.1	67.6 - 79.7	4,500
Native Hawaiian	300	6.4	4.0 - 10.0	200	3.6	1.9 - 6.7	900	16.3	12.6 - 20.8	200	3.7	2.0 - 6.7	3,800	70.0	64.6 - 75.0	5,400
Chinese	0	0.0	-	<50	1.5	0.4 - 5.4	100	7.6	2.2 - 22.9	<50	4.4	0.8 - 20.7	600	86.5	70.8 - 94.4	700
Filipino	200	6.7	3.8 - 11.8	100	2.5	1.2 - 5.1	700	20.8	15.2 - 27.7	100	2.9	1.2 - 6.5	2,200	67.1	59.7 - 73.7	3,300
Japanese	<50	0.2	0.0 - 1.3	<50	0.5	0.1 - 2.9	100	4.1	1.8 - 9.2	<50	1.8	0.3 - 8.9	1,600	93.4	87.2 - 96.8	1,700
Other	200	9.2	4.9 - 16.5	100	4.4	1.9 - 10.2	400	13.1	8.1 - 20.5	<50	1.6	0.5 - 5.3	1,900	71.7	62.8 - 79.2	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.2	1.4 - 17.4	<50	1.7	0.6 - 4.8	300	27.0	17.3 - 39.6	100	8.9	3.4 - 21.1	600	57.2	44.6 - 69.0	1,100
20-24	300	7.6	4.6 - 12.3	100	2.5	1.1 - 5.9	800	18.4	13.8 - 24.1	100	3.1	1.4 - 6.8	3,000	68.3	61.7 - 74.2	4,400
25-34	500	5.2	3.5 - 7.7	300	3.1	1.8 - 5.1	1,100	11.3	8.7 - 14.5	400	4.0	2.6 - 6.2	7,200	76.5	72.4 - 80.1	9,400
35+	200	6.3	3.3 - 12.0	100	4.4	2.1 - 8.9	300	7.9	4.8 - 12.8	100	3.5	1.6 - 7.5	2,600	77.8	70.8 - 83.6	3,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	600	6.0	4.0 - 8.8	300	3.1	1.9 - 5.3	1,300	14.2	11.3 - 17.7	400	4.2	2.7 - 6.5	6,700	72.5	68.2 - 76.4	9,200
Female	500	6.0	4.1 - 8.8	300	3.0	1.8 - 5.1	1,100	12.5	9.8 - 15.9	300	3.8	2.3 - 6.2	6,700	74.6	70.3 - 78.5	9,000

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.3	2.8 - 6.5	300	2.8	1.7 - 4.6	1,200	10.7	8.3 - 13.7	500	3.9	2.6 - 6.0	9,000	78.3	74.5 - 81.6	11,400
No	600	8.9	6.2 - 12.6	200	3.6	2.0 - 6.2	1,200	17.7	14.2 - 21.9	300	4.1	2.5 - 6.7	4,500	65.7	60.7 - 70.4	6,800

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.8	4.5 - 10.1	200	3.5	2.0 - 6.2	1,100	17.4	13.8 - 21.7	200	3.9	2.2 - 6.8	4,300	68.4	63.3 - 73.2	6,300
131-185%	100	4.7	2.2 - 9.8	200	5.2	2.3 - 11.2	400	13.8	9.0 - 20.8	100	4.6	2.2 - 9.3	2,100	71.7	63.6 - 78.6	3,000
186+%	300	4.3	2.5 - 7.2	200	2.2	1.1 - 4.1	700	8.7	6.2 - 12.2	300	3.9	2.3 - 6.5	6,300	81.0	76.5 - 84.7	7,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	5.0	2.9 - 8.6	100	4.2	2.0 - 8.3	600	19.3	14.2 - 25.6	100	3.7	1.7 - 7.9	2,200	67.8	60.8 - 74.1	3,300
\$10,000-\$24,999	300	9.4	5.5 - 15.7	100	2.8	1.2 - 6.6	400	13.2	9.1 - 18.6	200	4.7	2.3 - 9.4	2,300	69.9	62.5 - 76.4	3,200
\$25,000-\$49,999	200	5.4	3.0 - 9.3	200	4.0	2.0 - 8.0	700	15.2	10.9 - 20.8	200	4.6	2.5 - 8.4	3,200	70.8	64.3 - 76.6	4,500
>=\$50,000	200	3.5	1.8 - 6.8	100	2.3	1.1 - 4.9	500	8.2	5.5 - 12.1	200	3.5	1.9 - 6.3	5,100	82.5	77.5 - 86.5	6,200

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.2	4.8 - 10.6	<50	3.1	1.6 - 5.9	200	13.5	10.1 - 17.8	100	6.3	4.0 - 9.9	900	69.9	64.6 - 74.8	1,300
Normal birth weight	1,000	5.9	4.4 - 7.9	500	3.1	2.1 - 4.6	2,300	13.3	11.1 - 15.9	700	3.8	2.7 - 5.5	12,600	73.8	70.6 - 76.8	17,000

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	19.8	11.8 - 31.3	100	8.1	3.4 - 18.2	200	15.7	9.2 - 25.6	<50	1.8	0.6 - 5.0	700	54.6	42.7 - 66.0	1,400
No	800	4.8	3.4 - 6.6	500	2.7	1.8 - 4.1	2,200	13.2	11.1 - 15.7	700	4.2	3.0 - 5.9	12,500	75.0	71.9 - 77.9	16,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	8.8	5.5 - 13.8	300	7.2	4.1 - 12.1	700	19.9	14.8 - 26.2	200	5.5	3.1 - 9.7	2,200	58.6	51.6 - 65.3	3,700
No	700	5.2	3.7 - 7.3	300	2.1	1.3 - 3.4	1,700	11.7	9.6 - 14.3	500	3.7	2.5 - 5.4	11,100	77.4	74.1 - 80.3	14,300

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	7.3	5.1 - 10.4	300	3.9	2.4 - 6.3	1,300	16.2	13.0 - 20.0	400	4.5	2.8 - 7.1	5,400	68.1	63.4 - 72.4	8,000
No	500	4.8	3.1 - 7.4	300	2.5	1.4 - 4.4	1,100	11.2	8.7 - 14.5	400	3.6	2.3 - 5.7	7,900	77.8	73.8 - 81.4	10,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,200	6.7	5.2 - 8.6	800	4.3	3.2 - 5.9	2,600	14.6	12.5 - 17.0	900	5.0	3.7 - 6.6	12,500	69.4	66.4 - 72.3	17,900

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	7.6	5.2 - 10.9	100	5.7	3.7 - 8.6	400	15.4	11.9 - 19.6	100	3.7	2.2 - 6.3	1,600	67.6	62.6 - 72.3	2,300
Honolulu	900	7.4	5.4 - 10.0	600	4.3	2.8 - 6.6	1,900	14.6	11.8 - 17.9	600	5.0	3.4 - 7.3	8,700	68.7	64.5 - 72.6	12,700
Kauai	<50	4.7	2.7 - 8.2	<50	2.7	1.3 - 5.8	100	12.9	9.3 - 17.5	<50	4.1	2.2 - 7.4	600	75.6	69.9 - 80.5	800
Maui	100	2.5	1.4 - 4.5	100	3.5	2.1 - 5.7	300	14.2	11.1 - 17.9	100	6.4	4.4 - 9.2	1,500	73.4	69.0 - 77.4	2,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.7	1.2 - 5.9	200	3.6	1.6 - 7.7	400	8.9	5.8 - 13.3	200	5.7	3.3 - 9.9	3,300	79.1	73.2 - 84.0	4,200
Native Hawaiian	700	11.5	8.2 - 15.9	300	5.2	3.2 - 8.6	1,100	19.3	15.4 - 24.1	200	4.3	2.6 - 7.0	3,400	59.6	54.0 - 65.0	5,700
Chinese	100	12.1	4.3 - 29.3	0	0.0	-	100	8.0	2.6 - 22.4	<50	1.2	0.3 - 4.5	600	78.7	61.8 - 89.4	800
Filipino	200	4.8	2.5 - 8.9	100	4.2	2.4 - 7.3	600	18.2	13.2 - 24.7	200	5.9	3.1 - 10.7	2,100	66.9	59.6 - 73.4	3,200
Japanese	<50	0.9	0.2 - 3.3	100	3.9	1.3 - 11.5	100	7.6	3.3 - 16.3	100	3.9	1.4 - 10.5	1,300	83.7	74.0 - 90.2	1,600
Other	200	6.9	3.5 - 13.3	100	5.2	2.2 - 12.1	400	15.1	9.6 - 23.1	100	5.9	2.7 - 12.2	1,600	66.8	57.6 - 74.9	2,400

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.5	3.0 - 13.7	100	6.1	2.7 - 13.3	500	27.8	19.5 - 38.0	100	5.8	2.4 - 12.9	900	53.8	43.3 - 64.0	1,800
20-24	300	7.1	4.3 - 11.5	200	5.6	3.1 - 10.0	700	17.0	12.7 - 22.4	300	6.5	3.9 - 10.7	2,600	63.7	57.1 - 69.9	4,100
25-34	700	7.2	5.1 - 10.2	300	3.4	2.0 - 5.5	1,100	11.8	9.2 - 15.0	400	4.6	3.0 - 7.0	6,600	73.0	68.8 - 76.9	9,100
35+	100	4.8	2.5 - 8.8	100	4.5	2.2 - 9.2	400	11.9	7.5 - 18.5	100	3.4	1.6 - 7.2	2,300	75.3	68.0 - 81.5	3,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	7.4	5.3 - 10.2	300	3.6	2.3 - 5.8	1,300	13.7	10.9 - 17.0	600	5.8	4.0 - 8.3	6,600	69.5	65.3 - 73.5	9,500
Female	500	6.0	4.1 - 8.7	400	5.1	3.3 - 7.8	1,300	15.6	12.5 - 19.2	300	4.0	2.6 - 6.3	5,800	69.3	64.9 - 73.4	8,400

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.2	2.8 - 6.3	300	3.3	2.0 - 5.2	1,200	10.8	8.5 - 13.8	500	4.9	3.4 - 7.0	8,200	76.8	73.0 - 80.1	10,600
No	800	10.3	7.5 - 14.0	400	5.9	3.9 - 8.9	1,500	20.1	16.4 - 24.3	400	5.0	3.2 - 7.9	4,300	58.7	53.7 - 63.5	7,300

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	600	10.2	7.3 - 14.1	400	6.3	4.1 - 9.5	1,200	19.3	15.4 - 23.8	200	3.6	2.0 - 6.2	3,800	60.7	55.3 - 65.9	6,300
131-185%	200	6.1	2.9 - 12.1	100	5.2	2.4 - 11.0	400	15.8	10.8 - 22.5	200	6.1	3.3 - 11.0	1,900	66.9	58.7 - 74.1	2,800
186+%	200	2.8	1.6 - 5.0	200	3.1	1.7 - 5.6	700	9.3	6.7 - 12.7	400	5.8	3.8 - 8.7	5,700	79.0	74.6 - 82.9	7,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	10.9	6.7 - 17.2	200	5.3	2.6 - 10.2	700	22.4	16.4 - 29.8	100	4.7	2.2 - 9.7	1,700	56.8	48.6 - 64.5	3,000
\$10,000-\$24,999	300	9.1	5.8 - 14.1	300	7.4	4.5 - 12.0	600	15.4	11.4 - 20.6	200	4.2	2.2 - 7.7	2,400	63.9	57.2 - 70.0	3,800
\$25,000-\$49,999	200	5.8	3.2 - 10.4	200	5.7	2.8 - 11.0	400	12.3	8.4 - 17.7	200	5.6	3.1 - 9.7	2,600	70.6	63.6 - 76.7	3,600
>=\$50,000	100	2.4	1.2 - 4.7	100	1.9	0.9 - 3.9	600	10.4	7.3 - 14.4	300	5.4	3.3 - 8.5	4,800	80.0	75.1 - 84.1	6,000

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.3	6.6 - 13.0	100	4.8	2.8 - 8.1	200	17.1	13.4 - 21.5	100	4.6	2.8 - 7.4	800	64.2	58.9 - 69.2	1,300
Normal birth weight	1,100	6.5	5.0 - 8.5	700	4.3	3.1 - 6.0	2,400	14.4	12.2 - 17.0	800	5.0	3.7 - 6.7	11,600	69.8	66.6 - 72.9	16,700

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	19.2	12.0 - 29.4	200	9.7	4.9 - 18.3	400	21.4	14.2 - 30.9	100	5.0	1.8 - 13.1	700	44.7	34.7 - 55.2	1,700
No	900	5.5	4.1 - 7.4	600	3.8	2.7 - 5.5	2,200	13.6	11.4 - 16.1	700	4.7	3.5 - 6.4	11,500	72.3	69.1 - 75.3	15,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	14.4	10.1 - 20.2	300	6.4	3.6 - 11.1	800	20.0	15.1 - 25.9	200	5.7	3.2 - 9.8	2,100	53.5	46.8 - 60.1	3,900
No	600	4.7	3.3 - 6.5	500	3.8	2.6 - 5.6	1,700	12.8	10.5 - 15.4	600	4.5	3.2 - 6.3	10,100	74.3	70.9 - 77.4	13,600

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	9.2	6.6 - 12.6	500	7.3	5.0 - 10.5	1,400	19.9	16.4 - 24.1	300	4.5	2.9 - 7.0	4,300	59.1	54.1 - 63.8	7,200
No	500	5.2	3.5 - 7.7	200	2.4	1.4 - 4.2	1,100	10.9	8.5 - 14.0	500	4.9	3.3 - 7.2	7,800	76.5	72.6 - 80.1	10,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,400	7.8	6.5 - 9.2	500	2.6	2.0 - 3.5	2,600	14.1	12.5 - 15.9	800	4.2	3.3 - 5.3	12,900	71.3	69.0 - 73.4	18,200

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	11.0	7.3 - 16.3	100	5.4	3.0 - 9.5	300	13.7	9.6 - 19.3	<50	1.6	0.5 - 4.7	1,600	68.3	61.5 - 74.3	2,400
Honolulu	900	7.4	5.9 - 9.1	300	2.1	1.4 - 3.1	1,900	14.9	13.0 - 17.1	600	4.4	3.4 - 5.8	9,100	71.2	68.6 - 73.8	12,800
Kauai	<50	1.9	0.3 - 11.1	<50	1.7	0.4 - 6.9	100	13.1	7.1 - 22.9	100	10.0	4.9 - 19.1	600	73.4	62.4 - 82.1	900
Maui	200	9.1	5.7 - 14.4	100	3.4	1.6 - 7.1	200	10.0	6.4 - 15.3	100	3.6	1.8 - 7.3	1,600	73.9	67.0 - 79.7	2,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	4.3	2.7 - 6.8	100	1.9	0.9 - 3.8	500	12.5	9.6 - 16.1	300	6.1	4.1 - 8.9	3,300	75.2	70.8 - 79.2	4,300
Native Hawaiian	500	9.6	7.0 - 12.9	200	3.6	2.2 - 6.1	800	16.1	12.8 - 20.1	300	5.8	3.8 - 8.6	3,300	64.9	60.0 - 69.5	5,000
Chinese	<50	2.1	0.8 - 5.4	<50	1.2	0.4 - 3.8	100	9.0	5.8 - 13.7	<50	3.8	1.9 - 7.5	500	84.0	78.3 - 88.4	600
Filipino	300	8.4	5.9 - 11.9	100	3.6	2.0 - 6.1	500	16.5	12.9 - 20.8	100	2.9	1.6 - 5.3	2,300	68.7	63.5 - 73.4	3,300
Japanese	100	3.0	1.2 - 7.3	100	2.5	1.0 - 6.4	200	9.2	5.7 - 14.5	100	2.5	1.1 - 5.6	1,700	82.9	76.6 - 87.8	2,000
Other	400	13.8	9.8 - 19.0	<50	1.4	0.6 - 3.5	400	14.8	10.9 - 19.9	<50	1.6	0.6 - 4.3	2,000	68.4	62.2 - 74.0	2,900

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	10.2	5.8 - 17.2	100	4.8	2.1 - 10.7	300	18.3	12.3 - 26.4	<50	3.1	1.2 - 8.2	900	63.6	54.5 - 71.8	1,400
20-24	400	9.8	7.0 - 13.7	100	2.7	1.4 - 5.0	900	21.6	17.6 - 26.3	300	6.9	4.6 - 10.0	2,300	59.0	53.7 - 64.1	4,000
25-34	700	7.6	5.9 - 9.6	200	2.0	1.2 - 3.2	1,100	11.8	9.8 - 14.1	400	3.7	2.6 - 5.2	7,000	75.0	71.9 - 77.8	9,400
35+	200	5.0	3.0 - 8.0	100	3.6	2.1 - 6.2	300	9.9	7.0 - 13.8	100	2.9	1.6 - 5.4	2,700	78.6	73.8 - 82.8	3,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	7.3	5.7 - 9.3	300	2.8	1.9 - 4.2	1,400	14.8	12.5 - 17.3	500	4.8	3.5 - 6.4	6,700	70.4	67.2 - 73.3	9,500
Female	700	8.3	6.5 - 10.6	200	2.4	1.6 - 3.8	1,200	13.4	11.2 - 16.0	300	3.6	2.5 - 5.2	6,300	72.2	69.0 - 75.3	8,700

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	5.6	4.3 - 7.2	200	2.1	1.4 - 3.1	1,600	13.8	11.8 - 16.0	500	4.1	3.1 - 5.4	8,500	74.5	71.8 - 77.0	11,400
No	800	11.4	9.0 - 14.4	200	3.6	2.3 - 5.5	1,000	14.7	12.0 - 17.8	300	4.4	3.0 - 6.5	4,400	65.8	61.8 - 69.7	6,800

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	800	12.2	9.7 - 15.3	200	3.0	1.9 - 4.8	1,100	16.9	14.1 - 20.3	300	3.9	2.6 - 5.8	4,200	63.9	59.9 - 67.8	6,600
131-185%	300	9.9	6.6 - 14.6	<50	1.4	0.5 - 4.0	300	10.9	7.5 - 15.6	100	2.7	1.3 - 5.8	2,000	75.1	69.1 - 80.2	2,600
186+%	300	3.7	2.5 - 5.4	200	2.2	1.4 - 3.5	900	11.8	9.7 - 14.4	400	5.0	3.7 - 6.9	6,000	77.3	74.1 - 80.2	7,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	400	13.2	9.4 - 18.2	100	3.8	2.0 - 7.1	500	18.3	13.9 - 23.6	100	3.2	1.6 - 6.1	1,900	61.6	55.4 - 67.4	3,000
\$10,000-\$24,999	400	10.4	7.5 - 14.3	100	2.9	1.5 - 5.3	600	15.8	12.3 - 20.1	200	5.1	3.2 - 8.1	2,600	65.8	60.5 - 70.7	3,900
\$25,000-\$49,999	300	7.3	5.0 - 10.5	100	2.4	1.3 - 4.4	600	15.4	12.2 - 19.4	200	3.8	2.3 - 6.3	3,000	71.1	66.3 - 75.4	4,200
>=\$50,000	300	4.2	2.8 - 6.2	100	1.6	0.8 - 2.9	600	9.4	7.3 - 12.1	300	4.4	3.0 - 6.4	4,900	80.5	77.1 - 83.5	6,100

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.4	3.7 - 14.5	<50	4.0	1.5 - 9.9	200	13.8	8.4 - 21.8	<50	3.3	1.2 - 8.8	900	71.5	62.2 - 79.2	1,200
Normal birth weight	1,300	7.8	6.5 - 9.3	400	2.5	1.9 - 3.5	2,400	14.1	12.5 - 16.0	700	4.3	3.4 - 5.4	12,100	71.2	68.9 - 73.5	16,900

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	15.5	10.0 - 23.1	<50	1.5	0.4 - 5.6	300	23.3	16.7 - 31.6	100	8.7	4.8 - 15.1	800	51.0	42.2 - 59.7	1,500
No	1,200	7.2	5.9 - 8.6	400	2.7	1.9 - 3.6	2,200	13.3	11.6 - 15.1	600	3.9	3.0 - 5.0	11,900	73.0	70.7 - 75.2	16,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	12.9	9.4 - 17.5	100	1.9	0.8 - 4.4	600	19.3	15.1 - 24.3	300	8.5	5.8 - 12.3	1,900	57.4	51.6 - 63.1	3,300
No	1,000	6.7	5.4 - 8.2	400	2.7	2.0 - 3.8	1,900	12.9	11.2 - 14.8	500	3.3	2.5 - 4.4	10,800	74.4	71.9 - 76.7	14,500

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	11.6	9.3 - 14.4	200	3.2	2.1 - 4.9	1,400	17.7	15.0 - 20.8	300	3.9	2.6 - 5.7	4,800	63.6	59.8 - 67.2	7,600
No	500	5.1	3.8 - 6.8	200	2.3	1.5 - 3.4	1,200	11.4	9.5 - 13.6	500	4.6	3.4 - 6.1	7,900	76.7	73.9 - 79.2	10,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,400	7.6	6.4 - 9.0	600	3.5	2.7 - 4.6	2,700	14.9	13.3 - 16.7	500	3.1	2.3 - 4.0	12,700	70.9	68.6 - 73.1	17,900

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	10.8	7.2 - 16.0	100	3.9	1.8 - 7.9	200	8.8	5.6 - 13.6	100	2.1	0.8 - 5.4	1,800	74.4	67.9 - 79.9	2,400
Honolulu	800	6.8	5.5 - 8.5	400	3.4	2.5 - 4.6	2,000	16.0	14.0 - 18.3	400	3.3	2.5 - 4.5	8,800	70.5	67.8 - 73.0	12,400
Kauai	100	10.3	5.3 - 18.9	<50	1.2	0.2 - 7.7	100	13.1	7.6 - 21.8	<50	4.2	1.6 - 10.3	700	71.2	61.0 - 79.7	1,000
Maui	200	7.4	4.4 - 12.2	100	5.0	2.6 - 9.5	300	16.3	11.5 - 22.6	<50	1.9	0.6 - 5.8	1,500	69.4	62.2 - 75.8	2,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	400	9.7	7.1 - 13.1	100	3.2	1.8 - 5.5	400	10.4	7.7 - 13.9	100	3.4	2.0 - 5.8	3,000	73.3	68.6 - 77.5	4,100
Native Hawaiian	400	7.7	5.4 - 10.9	300	5.8	3.8 - 8.7	1,100	22.6	18.7 - 27.1	100	2.2	1.1 - 4.2	3,000	61.7	56.7 - 66.4	4,900
Chinese	<50	2.8	1.2 - 6.5	<50	3.6	1.7 - 7.5	<50	6.7	3.9 - 11.2	<50	1.6	0.6 - 4.3	600	85.3	79.6 - 89.6	700
Filipino	200	7.1	4.8 - 10.4	100	1.8	0.8 - 3.8	600	18.1	14.4 - 22.5	200	5.4	3.5 - 8.4	2,200	67.5	62.4 - 72.3	3,200
Japanese	100	2.8	1.1 - 7.1	<50	1.3	0.3 - 5.0	200	7.5	4.5 - 12.3	<50	2.1	0.8 - 5.3	1,800	86.3	80.4 - 90.6	2,100
Other	300	9.6	6.4 - 14.1	100	3.6	1.9 - 6.8	300	11.8	8.3 - 16.5	100	2.5	1.1 - 5.2	2,100	72.6	66.6 - 77.8	2,900

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	10.3	5.9 - 17.5	100	8.8	4.7 - 15.8	300	21.3	14.7 - 29.8	100	4.4	1.9 - 10.0	900	55.1	45.9 - 64.0	1,600
20-24	400	9.7	7.2 - 13.1	100	2.8	1.6 - 5.0	1,000	21.5	17.7 - 25.9	100	2.8	1.6 - 4.9	2,800	63.1	58.2 - 67.7	4,500
25-34	600	6.9	5.3 - 8.9	300	3.3	2.2 - 4.8	1,100	12.9	10.8 - 15.4	200	2.7	1.8 - 4.0	6,500	74.2	71.1 - 77.1	8,700
35+	200	5.0	3.0 - 8.4	100	2.4	1.2 - 4.9	200	7.6	5.1 - 11.1	100	3.7	2.1 - 6.5	2,500	81.2	76.4 - 85.2	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	600	7.0	5.4 - 9.0	200	2.4	1.6 - 3.8	1,500	16.2	13.9 - 18.9	300	3.1	2.2 - 4.5	6,600	71.2	68.0 - 74.2	9,300
Female	700	8.3	6.5 - 10.5	400	4.7	3.4 - 6.5	1,200	13.5	11.3 - 16.1	300	3.0	2.0 - 4.4	6,100	70.6	67.3 - 73.7	8,700

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	6.4	5.1 - 8.1	300	2.7	1.9 - 3.9	1,200	11.0	9.3 - 13.0	300	2.3	1.6 - 3.4	8,700	77.5	74.9 - 79.9	11,200
No	600	9.6	7.4 - 12.4	300	4.8	3.3 - 7.0	1,400	21.5	18.3 - 25.2	300	4.3	2.9 - 6.3	4,000	59.8	55.6 - 63.8	6,700

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	700	10.2	7.9 - 13.1	300	4.8	3.3 - 7.0	1,400	21.5	18.3 - 25.1	200	2.8	1.7 - 4.5	4,000	60.6	56.5 - 64.6	6,700
131-185%	200	7.5	4.8 - 11.5	100	3.0	1.5 - 6.1	400	13.3	9.6 - 18.2	100	3.5	1.8 - 6.7	1,900	72.7	66.8 - 77.9	2,600
186+%	400	5.1	3.6 - 7.0	200	3.1	2.0 - 4.7	700	9.4	7.4 - 11.8	200	2.8	1.9 - 4.3	5,800	79.6	76.5 - 82.4	7,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	10.6	7.3 - 15.1	200	6.3	3.8 - 10.3	700	23.1	18.2 - 28.8	100	3.7	2.0 - 7.0	1,700	56.3	50.0 - 62.4	3,000
\$10,000-\$24,999	300	9.0	6.2 - 12.8	100	4.0	2.3 - 6.9	700	18.2	14.4 - 22.7	100	3.0	1.6 - 5.4	2,400	65.9	60.6 - 70.9	3,700
\$25,000-\$49,999	300	7.6	5.3 - 10.8	100	2.4	1.2 - 4.7	700	16.4	13.0 - 20.5	100	2.5	1.4 - 4.7	2,900	71.0	66.2 - 75.4	4,100
>=\$50,000	300	4.9	3.4 - 7.0	200	3.3	2.1 - 5.1	500	7.7	5.8 - 10.2	200	2.7	1.7 - 4.4	4,800	81.4	78.1 - 84.4	5,900

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	8.8	5.1 - 15.0	100	3.6	1.4 - 8.6	300	19.8	13.8 - 27.6	<50	2.8	1.1 - 6.9	1,000	65.0	56.5 - 72.6	1,500
Normal birth weight	1,200	7.5	6.2 - 9.0	600	3.5	2.7 - 4.6	2,400	14.5	12.8 - 16.4	500	3.1	2.3 - 4.1	11,800	71.4	69.1 - 73.7	16,500

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	16.6	11.2 - 23.9	100	6.9	3.5 - 13.1	400	27.0	19.9 - 35.6	100	5.7	2.8 - 11.2	700	43.8	35.4 - 52.6	1,500
No	1,100	6.8	5.6 - 8.2	500	3.2	2.4 - 4.3	2,300	13.9	12.2 - 15.8	500	2.9	2.1 - 3.8	11,900	73.2	70.9 - 75.4	16,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	12.3	9.0 - 16.5	200	4.5	2.6 - 7.6	1,000	27.2	22.5 - 32.5	100	4.0	2.3 - 6.8	1,800	52.0	46.4 - 57.5	3,500
No	900	6.5	5.2 - 8.0	500	3.3	2.4 - 4.5	1,700	12.0	10.4 - 13.9	400	2.9	2.1 - 3.9	10,700	75.3	72.9 - 77.6	14,200

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	10.8	8.5 - 13.5	300	3.8	2.5 - 5.7	1,400	19.4	16.5 - 22.7	200	3.0	1.9 - 4.6	4,600	63.1	59.2 - 66.8	7,400
No	500	5.2	4.0 - 6.9	300	3.3	2.3 - 4.7	1,200	11.5	9.6 - 13.6	300	3.2	2.3 - 4.5	7,800	76.8	74.0 - 79.3	10,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,400	8.0	6.8 - 9.5	500	2.6	1.9 - 3.6	2,800	16.0	14.2 - 17.9	600	3.2	2.5 - 4.2	12,400	70.1	67.8 - 72.3	17,700

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.8	3.8 - 11.8	100	3.3	1.4 - 7.5	400	18.5	13.4 - 25.0	<50	2.0	0.7 - 5.7	1,500	69.4	62.1 - 75.8	2,100
Honolulu	1,100	8.2	6.7 - 10.0	400	2.9	2.1 - 4.1	2,000	15.5	13.5 - 17.8	400	3.3	2.4 - 4.4	9,100	70.1	67.4 - 72.7	13,000
Kauai	100	6.8	2.7 - 16.4	0	0.0	-	100	15.0	8.2 - 25.8	<50	3.2	0.8 - 11.3	600	75.1	63.2 - 84.1	800
Maui	200	9.1	5.4 - 14.9	<50	1.1	0.2 - 5.0	300	16.8	11.7 - 23.6	100	4.4	2.0 - 9.2	1,300	68.6	60.8 - 75.5	1,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	7.2	5.0 - 10.4	100	2.3	1.2 - 4.4	500	12.9	9.8 - 16.8	200	4.8	3.0 - 7.5	3,100	72.7	68.0 - 77.0	4,200
Native Hawaiian	500	10.8	8.0 - 14.4	200	3.2	1.8 - 5.5	1,100	21.7	17.7 - 26.3	100	2.2	1.1 - 4.3	3,000	62.1	57.0 - 66.9	4,900
Chinese	<50	2.1	0.8 - 5.4	<50	2.1	0.8 - 5.4	100	11.1	7.4 - 16.4	<50	2.8	1.2 - 6.4	500	81.9	75.8 - 86.8	600
Filipino	200	7.1	4.8 - 10.5	100	2.6	1.3 - 5.2	600	18.1	14.2 - 22.7	100	4.0	2.3 - 6.9	2,200	68.2	62.9 - 73.1	3,200
Japanese	100	5.0	2.5 - 9.5	<50	1.8	0.6 - 5.4	200	7.6	4.5 - 12.5	<50	0.5	0.1 - 3.4	1,700	85.1	79.0 - 89.6	2,000
Other	200	9.0	5.8 - 13.6	100	3.0	1.4 - 6.3	400	15.3	11.0 - 20.7	100	3.8	2.0 - 6.9	1,900	68.9	62.5 - 74.6	2,800

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.1	4.9 - 16.2	100	3.7	1.4 - 9.3	400	29.4	21.6 - 38.6	<50	1.4	0.3 - 6.1	800	56.5	47.1 - 65.4	1,500
20-24	400	9.0	6.3 - 12.5	100	3.0	1.6 - 5.4	1,000	23.7	19.5 - 28.5	200	4.3	2.7 - 7.0	2,600	60.0	54.7 - 65.0	4,400
25-34	600	6.9	5.3 - 9.0	200	2.8	1.8 - 4.3	1,100	12.7	10.6 - 15.3	300	3.3	2.2 - 4.7	6,500	74.3	71.0 - 77.2	8,700
35+	300	9.3	6.5 - 13.1	<50	1.2	0.5 - 3.0	300	8.1	5.5 - 11.6	100	2.5	1.3 - 4.9	2,500	78.9	74.0 - 83.1	3,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	7.3	5.6 - 9.4	300	2.9	1.9 - 4.4	1,600	18.0	15.4 - 20.9	400	4.6	3.4 - 6.3	6,100	67.2	63.8 - 70.4	9,000
Female	800	8.8	7.0 - 11.1	200	2.3	1.5 - 3.7	1,200	13.9	11.6 - 16.6	200	1.8	1.1 - 3.0	6,400	73.1	69.8 - 76.2	8,700

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	7.7	6.2 - 9.5	300	2.3	1.6 - 3.4	1,300	11.2	9.4 - 13.2	300	3.0	2.2 - 4.2	8,600	75.8	73.1 - 78.3	11,300
No	600	8.7	6.4 - 11.6	200	3.2	2.0 - 5.2	1,600	24.5	20.9 - 28.5	200	3.6	2.3 - 5.6	3,900	60.0	55.7 - 64.3	6,400

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	700	10.9	8.4 - 14.0	300	4.4	2.9 - 6.6	1,400	22.9	19.4 - 26.9	100	2.2	1.3 - 3.9	3,700	59.5	55.1 - 63.8	6,200
131-185%	200	10.5	7.0 - 15.4	100	2.3	0.9 - 5.7	300	12.5	8.5 - 17.9	100	5.2	2.8 - 9.4	1,600	69.5	62.8 - 75.5	2,400
186+%	400	4.5	3.2 - 6.3	100	1.5	0.9 - 2.7	900	11.5	9.4 - 14.0	300	3.7	2.6 - 5.3	6,300	78.8	75.7 - 81.6	8,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	11.4	7.8 - 16.2	100	4.2	2.3 - 7.7	800	26.8	21.5 - 33.0	100	2.2	0.9 - 5.0	1,700	55.4	48.9 - 61.7	3,000
\$10,000-\$24,999	300	8.4	5.7 - 12.1	200	4.8	2.9 - 8.0	700	19.0	14.9 - 23.8	100	2.1	1.0 - 4.5	2,400	65.7	60.2 - 70.9	3,700
\$25,000-\$49,999	300	7.7	5.3 - 10.9	100	2.0	1.0 - 4.0	700	17.5	13.9 - 21.8	200	4.9	3.1 - 7.7	2,900	67.9	62.9 - 72.5	4,200
>=\$50,000	300	5.7	4.0 - 8.0	100	1.1	0.5 - 2.3	400	7.1	5.2 - 9.6	200	3.6	2.3 - 5.5	4,700	82.6	79.2 - 85.5	5,700

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	11.8	7.1 - 19.1	100	4.3	1.9 - 9.7	200	13.4	8.2 - 20.9	<50	2.6	1.0 - 6.6	1,000	67.9	59.1 - 75.6	1,400
Normal birth weight	1,300	7.7	6.4 - 9.3	400	2.5	1.8 - 3.5	2,600	16.2	14.4 - 18.3	500	3.3	2.5 - 4.3	11,500	70.3	67.8 - 72.6	16,300

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	22.5	16.0 - 30.8	100	5.9	2.9 - 11.5	300	18.0	12.0 - 26.0	100	3.9	1.7 - 8.7	800	49.7	41.0 - 58.5	1,600
No	1,100	6.7	5.5 - 8.2	400	2.3	1.6 - 3.3	2,500	15.6	13.7 - 17.6	500	3.2	2.4 - 4.3	11,500	72.2	69.8 - 74.5	15,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	13.5	10.0 - 18.0	100	4.0	2.3 - 6.9	700	19.3	15.1 - 24.2	200	4.6	2.7 - 7.6	2,200	58.6	52.9 - 64.1	3,700
No	900	6.7	5.4 - 8.3	300	2.3	1.6 - 3.3	2,100	14.9	12.9 - 17.0	400	2.9	2.1 - 4.0	10,100	73.2	70.6 - 75.6	13,800

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	10.1	8.0 - 12.8	300	4.4	3.0 - 6.4	1,600	20.9	17.8 - 24.3	200	3.2	2.1 - 5.0	4,600	61.3	57.4 - 65.2	7,500
No	700	6.7	5.2 - 8.6	100	1.0	0.5 - 1.8	1,300	12.5	10.5 - 14.8	300	3.1	2.2 - 4.4	7,700	76.7	73.9 - 79.4	10,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,700	10.0	8.6 - 11.6	600	3.8	2.9 - 4.8	2,500	14.4	12.8 - 16.2	500	3.1	2.3 - 4.0	11,700	68.8	66.5 - 70.9	17,000

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	13.3	9.1 - 18.9	<50	2.2	0.9 - 5.6	300	13.5	9.4 - 19.0	<50	1.7	0.6 - 4.9	1,500	69.4	62.6 - 75.4	2,200
Honolulu	1,300	10.2	8.6 - 12.0	500	3.9	2.9 - 5.2	1,700	13.7	11.9 - 15.7	400	3.6	2.7 - 4.7	8,600	68.7	66.1 - 71.2	12,500
Kauai	<50	6.6	2.6 - 16.1	0	0.0	-	100	13.0	7.0 - 22.9	<50	1.8	0.3 - 10.7	600	78.6	67.3 - 86.8	700
Maui	100	5.8	3.0 - 10.9	100	6.2	3.3 - 11.3	400	22.3	16.5 - 29.5	<50	1.5	0.5 - 4.7	1,000	64.2	56.4 - 71.2	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.9	3.9 - 9.0	100	3.2	1.8 - 5.7	300	9.0	6.4 - 12.5	100	3.9	2.3 - 6.4	2,900	78.0	73.3 - 82.0	3,700
Native Hawaiian	600	13.3	10.2 - 17.1	300	5.7	3.8 - 8.5	800	16.9	13.5 - 21.0	200	4.4	2.8 - 6.8	2,900	59.7	54.8 - 64.5	4,800
Chinese	100	9.4	6.3 - 13.9	<50	1.2	0.4 - 3.8	100	9.4	6.3 - 13.9	<50	2.9	1.4 - 6.1	500	77.1	71.2 - 82.0	600
Filipino	300	9.8	7.2 - 13.3	100	4.4	2.8 - 7.1	700	20.9	17.1 - 25.3	100	3.3	1.9 - 5.7	2,000	61.5	56.5 - 66.3	3,200
Japanese	100	4.5	2.4 - 8.2	<50	0.4	0.1 - 2.9	200	10.3	7.0 - 15.0	<50	0.9	0.2 - 3.3	1,900	83.8	78.5 - 88.0	2,300
Other	400	15.4	11.3 - 20.6	100	3.5	1.8 - 6.9	400	14.6	10.7 - 19.6	<50	0.9	0.2 - 3.1	1,600	65.6	59.4 - 71.4	2,400

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	12.0	7.4 - 18.8	100	4.9	2.3 - 10.3	400	24.2	17.6 - 32.4	<50	3.0	1.2 - 7.4	800	55.9	47.4 - 64.2	1,500
20-24	500	11.2	8.4 - 14.8	200	5.5	3.6 - 8.3	900	22.2	18.4 - 26.6	100	3.4	2.0 - 5.7	2,400	57.6	52.7 - 62.4	4,200
25-34	800	9.9	8.0 - 12.2	300	3.5	2.4 - 5.0	900	10.8	8.9 - 13.1	300	3.3	2.3 - 4.8	6,100	72.5	69.3 - 75.4	8,400
35+	200	7.5	5.0 - 10.9	<50	1.4	0.6 - 3.6	300	8.8	6.2 - 12.4	100	1.7	0.8 - 3.6	2,400	80.6	75.9 - 84.5	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	10.1	8.3 - 12.4	300	4.0	2.8 - 5.5	1,200	13.7	11.5 - 16.2	300	3.9	2.8 - 5.4	5,900	68.3	65.1 - 71.3	8,700
Female	800	9.8	7.9 - 12.2	300	3.5	2.4 - 5.1	1,300	15.2	12.9 - 17.8	200	2.2	1.4 - 3.4	5,800	69.2	66.0 - 72.3	8,400

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	8.0	6.5 - 9.8	300	3.2	2.3 - 4.4	1,200	11.0	9.3 - 13.0	300	3.0	2.1 - 4.2	8,100	74.8	72.2 - 77.3	10,900
No	800	13.6	10.9 - 16.7	300	4.7	3.2 - 6.9	1,300	20.4	17.3 - 23.9	200	3.2	2.0 - 4.9	3,600	58.2	54.1 - 62.2	6,200

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,000	15.3	12.6 - 18.5	300	5.1	3.6 - 7.2	1,200	18.8	15.9 - 22.2	200	3.2	2.1 - 4.9	3,600	57.6	53.5 - 61.5	6,200
131-185%	200	10.9	7.0 - 16.6	100	4.1	1.9 - 8.7	300	15.6	11.0 - 21.6	<50	1.9	0.7 - 5.6	1,200	67.4	60.1 - 74.0	1,800
186+%	400	5.1	3.7 - 6.9	200	2.2	1.3 - 3.6	700	9.8	7.9 - 12.1	300	3.3	2.3 - 4.9	6,000	79.6	76.6 - 82.3	7,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	400	14.0	10.3 - 18.7	200	6.9	4.4 - 10.5	700	22.3	17.7 - 27.6	100	3.9	2.2 - 6.9	1,600	53.0	47.1 - 58.8	3,000
\$10,000-\$24,999	500	14.7	11.3 - 18.9	200	5.1	3.2 - 8.0	600	17.6	13.9 - 21.9	100	2.7	1.5 - 5.0	2,100	59.9	54.6 - 64.9	3,600
\$25,000-\$49,999	300	8.0	5.6 - 11.3	100	2.9	1.6 - 5.3	500	13.0	10.0 - 16.8	100	2.2	1.1 - 4.2	2,900	73.8	69.1 - 78.0	3,900
>=\$50,000	300	5.3	3.7 - 7.6	100	1.5	0.7 - 3.1	400	8.2	6.2 - 10.7	200	3.7	2.4 - 5.8	4,300	81.3	77.8 - 84.4	5,300

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.2	3.8 - 13.4	100	5.1	2.2 - 11.4	200	15.6	10.1 - 23.3	100	4.2	1.7 - 9.7	800	68.0	59.1 - 75.7	1,200
Normal birth weight	1,600	10.2	8.7 - 11.8	600	3.7	2.8 - 4.7	2,300	14.4	12.7 - 16.2	500	3.0	2.2 - 3.9	10,900	68.8	66.5 - 71.1	15,800

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	23.7	17.0 - 32.1	100	10.0	5.9 - 16.5	300	20.5	14.5 - 28.3	100	4.3	1.9 - 9.6	600	41.5	33.3 - 50.1	1,400
No	1,300	8.7	7.4 - 10.3	500	3.0	2.2 - 4.0	2,100	13.8	12.2 - 15.7	500	3.0	2.3 - 4.0	11,000	71.4	69.1 - 73.7	15,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	15.5	11.9 - 19.9	200	6.2	4.1 - 9.4	700	20.7	16.6 - 25.5	200	5.8	3.8 - 8.9	1,800	51.8	46.4 - 57.1	3,600
No	1,100	8.5	7.1 - 10.2	400	2.9	2.1 - 4.0	1,700	12.6	10.9 - 14.5	300	2.4	1.7 - 3.3	9,700	73.6	71.1 - 75.9	13,200

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	12.9	10.5 - 15.7	400	4.8	3.4 - 6.8	1,300	17.4	14.7 - 20.4	200	2.9	1.9 - 4.4	4,500	61.9	58.2 - 65.5	7,300
No	700	7.7	6.2 - 9.7	200	2.6	1.7 - 3.9	1,200	12.2	10.3 - 14.4	300	3.0	2.1 - 4.3	7,000	74.5	71.6 - 77.1	9,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,600	9.0	7.8 - 10.4	700	3.8	3.1 - 4.8	2,500	14.2	12.7 - 15.8	700	3.8	3.1 - 4.7	12,000	69.2	67.1 - 71.2	17,300

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	12.5	8.8 - 17.6	100	4.0	2.1 - 7.5	300	12.0	8.5 - 16.8	<50	1.4	0.5 - 3.6	1,500	70.1	63.8 - 75.7	2,200
Honolulu	1,200	9.6	8.1 - 11.2	400	3.4	2.6 - 4.5	1,800	14.0	12.3 - 15.9	500	4.1	3.2 - 5.3	8,700	68.9	66.5 - 71.2	12,600
Kauai	<50	2.4	0.6 - 8.3	<50	5.2	2.0 - 12.5	100	14.1	8.3 - 23.0	<50	1.3	0.2 - 8.0	600	77.0	67.2 - 84.6	800
Maui	100	3.7	1.8 - 7.2	100	5.9	3.5 - 9.8	300	18.3	13.5 - 24.2	100	5.5	3.2 - 9.4	1,200	66.7	60.0 - 72.7	1,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.6	3.8 - 8.1	<50	1.2	0.5 - 2.8	500	12.7	9.9 - 16.1	100	3.7	2.3 - 5.8	3,000	76.8	72.7 - 80.4	3,900
Native Hawaiian	600	13.4	10.5 - 16.9	200	4.8	3.2 - 7.3	900	19.4	16.0 - 23.3	200	4.0	2.6 - 6.2	2,800	58.4	53.7 - 62.8	4,700
Chinese	<50	5.9	4.0 - 8.5	<50	2.7	1.5 - 4.7	<50	6.3	4.4 - 9.0	<50	2.6	1.5 - 4.6	500	82.5	78.7 - 85.7	700
Filipino	300	8.0	5.8 - 11.0	200	7.3	5.2 - 10.1	500	16.2	13.1 - 19.9	200	4.8	3.2 - 7.2	2,100	63.7	59.1 - 68.0	3,300
Japanese	100	5.2	3.0 - 9.0	100	2.8	1.3 - 6.0	100	5.6	3.3 - 9.3	100	3.7	1.9 - 7.0	1,800	82.7	77.3 - 87.0	2,200
Other	300	11.2	7.9 - 15.7	100	2.7	1.3 - 5.7	300	13.8	10.1 - 18.7	100	2.6	1.3 - 5.1	1,800	69.6	63.7 - 74.8	2,500

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	13.9	8.8 - 21.4	100	7.0	3.8 - 12.6	300	24.1	17.5 - 32.1	<50	3.4	1.3 - 8.4	600	51.6	43.1 - 60.1	1,300
20-24	400	9.8	7.4 - 12.9	200	4.7	3.1 - 7.0	1,000	22.5	18.9 - 26.6	200	3.5	2.2 - 5.6	2,600	59.4	54.9 - 63.8	4,400
25-34	700	8.5	6.9 - 10.5	300	3.3	2.3 - 4.6	900	10.0	8.3 - 12.1	400	4.3	3.2 - 5.7	6,300	73.9	71.1 - 76.5	8,500
35+	200	7.2	5.0 - 10.2	100	2.9	1.7 - 5.2	300	10.0	7.3 - 13.5	100	3.1	1.8 - 5.3	2,500	76.8	72.3 - 80.7	3,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	9.7	8.0 - 11.7	400	3.9	2.9 - 5.2	1,300	14.3	12.3 - 16.5	400	4.1	3.1 - 5.5	6,400	68.0	65.2 - 70.7	9,400
Female	600	8.2	6.5 - 10.2	300	3.8	2.7 - 5.3	1,100	14.1	11.9 - 16.6	300	3.4	2.4 - 4.7	5,500	70.6	67.5 - 73.5	7,900

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	7.8	6.5 - 9.3	300	2.7	2.0 - 3.7	1,500	12.0	10.4 - 13.8	500	4.0	3.1 - 5.2	8,900	73.4	71.1 - 75.7	12,100
No	600	11.8	9.3 - 14.9	300	6.5	4.7 - 8.8	1,000	19.3	16.2 - 22.9	200	3.2	2.1 - 5.0	3,100	59.2	55.0 - 63.2	5,200

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	800	11.9	9.6 - 14.7	300	5.3	3.8 - 7.2	1,300	20.1	17.3 - 23.3	100	1.7	1.0 - 2.9	3,900	61.0	57.2 - 64.6	6,400
131-185%	200	10.9	7.5 - 15.7	100	2.8	1.3 - 5.7	200	11.9	8.3 - 16.9	100	4.4	2.4 - 7.8	1,500	70.0	63.8 - 75.5	2,100
186+%	400	5.2	3.9 - 6.9	200	2.1	1.4 - 3.3	800	10.2	8.4 - 12.4	400	5.8	4.4 - 7.4	5,800	76.7	73.8 - 79.3	7,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	10.1	7.0 - 14.4	200	7.9	5.2 - 11.8	500	20.8	16.4 - 26.0	100	2.9	1.5 - 5.5	1,500	58.3	52.4 - 64.0	2,600
\$10,000-\$24,999	600	13.3	10.4 - 16.8	200	5.0	3.4 - 7.2	800	18.4	15.1 - 22.2	100	1.9	1.0 - 3.6	2,600	61.4	56.9 - 65.7	4,300
\$25,000-\$49,999	300	6.2	4.4 - 8.6	100	1.9	1.0 - 3.6	600	13.0	10.4 - 16.3	300	6.4	4.6 - 8.7	3,200	72.5	68.4 - 76.2	4,400
>=\$50,000	300	5.7	4.0 - 7.9	100	1.6	0.8 - 3.0	400	8.0	6.1 - 10.5	200	4.1	2.7 - 6.0	3,900	80.6	77.3 - 83.6	4,800

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.9	5.7 - 16.8	100	6.4	3.2 - 12.4	200	14.1	9.0 - 21.3	<50	3.1	1.2 - 7.6	700	66.5	57.8 - 74.3	1,100
Normal birth weight	1,400	8.9	7.7 - 10.4	600	3.7	2.9 - 4.6	2,300	14.2	12.7 - 15.9	600	3.8	3.1 - 4.8	11,200	69.3	67.2 - 71.4	16,200

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	20.1	14.1 - 27.8	100	6.0	3.0 - 11.7	200	16.0	10.8 - 23.0	<50	3.1	1.2 - 7.7	700	54.8	46.4 - 63.0	1,300
No	1,300	8.0	6.8 - 9.4	600	3.6	2.8 - 4.6	2,200	13.9	12.4 - 15.6	600	3.8	3.1 - 4.8	11,100	70.6	68.5 - 72.6	15,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	12.8	9.7 - 16.8	100	3.3	1.9 - 5.7	600	16.8	13.3 - 21.0	200	5.1	3.3 - 7.7	2,200	62.0	56.9 - 66.8	3,500
No	1,100	8.0	6.7 - 9.5	500	3.9	3.1 - 5.0	1,800	13.4	11.8 - 15.2	500	3.4	2.7 - 4.4	9,700	71.3	69.0 - 73.4	13,600

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	10.7	8.8 - 13.1	300	4.2	3.1 - 5.8	1,400	18.1	15.7 - 20.9	200	2.7	1.8 - 4.0	4,900	64.2	60.9 - 67.4	7,700
No	700	7.1	5.6 - 8.8	300	3.5	2.6 - 4.9	1,000	11.1	9.3 - 13.1	400	4.8	3.7 - 6.3	6,800	73.5	70.8 - 76.0	9,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,800	10.8	9.2 - 12.6	700	4.2	3.2 - 5.4	2,400	14.6	12.7 - 16.7	600	3.9	3.0 - 5.1	11,000	66.5	63.9 - 69.1	16,500

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	17.3	14.0 - 21.2	100	3.8	2.3 - 6.3	300	15.1	12.1 - 18.6	<50	2.6	1.5 - 4.5	1,100	61.2	56.5 - 65.6	1,900
Honolulu	1,300	10.3	8.3 - 12.7	500	4.2	3.0 - 5.9	1,800	14.8	12.4 - 17.6	500	3.8	2.6 - 5.4	8,100	66.8	63.4 - 70.1	12,100
Kauai	100	8.9	5.4 - 14.2	<50	4.9	2.5 - 9.5	100	6.5	3.9 - 10.7	100	6.6	3.7 - 11.4	600	73.1	66.1 - 79.1	800
Maui	100	8.1	5.8 - 11.2	100	3.5	2.1 - 6.0	300	16.4	13.0 - 20.5	100	4.9	3.1 - 7.6	1,200	67.1	62.2 - 71.7	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.6	3.3 - 9.3	100	2.1	0.8 - 5.2	400	12.0	8.1 - 17.4	100	4.0	2.1 - 7.6	2,800	76.4	70.2 - 81.6	3,600
Native Hawaiian	600	14.5	12.5 - 16.9	200	5.6	4.2 - 7.3	800	17.6	15.3 - 20.1	200	4.1	3.1 - 5.5	2,600	58.2	55.1 - 61.2	4,500
Chinese	<50	3.3	0.5 - 19.3	0	0.0	-	<50	6.2	1.6 - 21.0	<50	1.9	0.5 - 6.7	600	88.6	74.0 - 95.5	700
Filipino	500	13.6	9.6 - 18.8	200	4.8	2.7 - 8.2	700	18.2	13.7 - 23.8	200	5.1	2.9 - 8.8	2,100	58.4	51.8 - 64.6	3,600
Japanese	100	7.6	4.0 - 14.0	100	5.5	2.4 - 12.3	100	6.3	3.0 - 12.7	<50	2.3	0.7 - 7.4	1,500	78.3	69.6 - 85.0	1,900
Other	300	12.4	7.6 - 19.7	100	3.8	1.6 - 8.9	400	16.7	11.0 - 24.4	100	3.5	1.4 - 8.7	1,400	63.6	54.6 - 71.7	2,200

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	13.9	9.4 - 20.0	200	11.8	6.8 - 19.8	400	29.5	22.2 - 38.0	100	3.5	2.0 - 6.2	600	41.3	33.4 - 49.6	1,500
20-24	600	14.5	11.2 - 18.6	100	3.4	2.3 - 5.0	800	21.9	17.6 - 27.0	100	3.5	1.9 - 6.3	2,200	56.7	51.2 - 62.1	3,800
25-34	800	10.1	7.9 - 12.8	300	4.1	2.8 - 6.0	900	10.4	8.2 - 13.0	400	4.4	3.0 - 6.3	5,900	71.0	67.3 - 74.5	8,300
35+	200	6.2	3.6 - 10.5	<50	1.3	0.4 - 4.4	300	9.1	5.7 - 14.1	100	3.5	1.6 - 7.3	2,300	79.9	73.7 - 85.0	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,000	10.8	8.7 - 13.3	300	3.7	2.6 - 5.4	1,300	15.2	12.6 - 18.2	400	4.1	2.8 - 6.0	5,800	66.1	62.4 - 69.6	8,800
Female	800	10.8	8.5 - 13.5	400	4.6	3.2 - 6.6	1,100	13.9	11.3 - 16.9	300	3.7	2.5 - 5.4	5,200	67.0	63.1 - 70.6	7,800

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	7.1	5.5 - 9.1	200	2.2	1.4 - 3.5	1,300	12.0	9.9 - 14.6	500	4.3	3.0 - 6.0	8,100	74.3	71.1 - 77.3	10,900
No	1,000	18.0	14.9 - 21.7	400	7.9	5.7 - 10.7	1,100	19.5	16.3 - 23.2	200	3.2	2.2 - 4.7	2,900	51.4	47.1 - 55.7	5,600

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,200	16.3	13.5 - 19.5	400	5.1	3.6 - 7.1	1,500	19.7	16.7 - 23.2	300	3.5	2.4 - 5.2	4,100	55.3	51.3 - 59.3	7,400
131-185%	200	9.3	5.7 - 14.9	100	3.9	1.8 - 8.1	300	12.9	8.9 - 18.3	100	4.0	1.8 - 8.5	1,700	70.0	62.8 - 76.3	2,400
186+%	100	2.3	1.3 - 4.0	100	2.1	1.0 - 4.5	400	7.6	5.3 - 11.0	200	3.9	2.3 - 6.4	4,400	84.1	79.9 - 87.5	5,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	800	21.6	17.3 - 26.7	300	6.7	4.5 - 10.0	700	18.7	14.9 - 23.2	100	3.6	2.0 - 6.4	1,900	49.3	43.9 - 54.8	3,900
\$15,001-\$25,000	200	11.8	8.2 - 16.6	100	2.5	1.0 - 6.3	400	19.6	14.1 - 26.6	<50	1.9	1.0 - 3.8	1,300	64.2	56.8 - 70.9	2,100
\$25,001-\$40,000	200	7.0	4.5 - 10.8	200	5.6	3.3 - 9.5	500	16.3	11.9 - 21.8	200	4.9	2.7 - 8.7	2,100	66.1	59.7 - 72.0	3,100
>=\$40,001	300	4.2	2.8 - 6.5	100	1.6	0.8 - 3.4	600	8.9	6.6 - 12.0	300	4.4	2.8 - 6.8	5,100	80.8	76.9 - 84.2	6,300

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.4	5.5 - 15.6	<50	4.0	1.4 - 11.1	200	19.8	12.2 - 30.6	100	5.1	1.8 - 13.5	700	61.8	50.9 - 71.5	1,100
Normal birth weight	1,700	10.9	9.3 - 12.8	600	4.2	3.2 - 5.4	2,200	14.2	12.3 - 16.3	600	3.8	2.9 - 5.1	10,300	66.9	64.2 - 69.5	15,400

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	35.3	27.2 - 44.3	100	5.5	2.8 - 10.3	200	14.3	9.3 - 21.4	<50	2.0	0.9 - 4.5	600	42.9	34.5 - 51.8	1,500
No	1,200	8.3	6.9 - 10.0	600	4.0	3.0 - 5.3	2,200	14.7	12.7 - 16.9	600	4.1	3.1 - 5.4	10,200	68.9	66.1 - 71.5	14,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	21.6	17.0 - 27.0	200	7.2	4.7 - 11.0	600	17.3	13.3 - 22.3	200	5.2	3.1 - 8.4	1,600	48.7	42.8 - 54.5	3,400
No	1,000	7.6	6.1 - 9.4	400	3.4	2.4 - 4.7	1,700	13.7	11.6 - 16.0	500	3.6	2.6 - 5.0	9,100	71.7	68.8 - 74.5	12,700

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	16.4	13.5 - 19.7	400	5.4	3.9 - 7.6	1,400	20.7	17.6 - 24.3	200	3.3	2.2 - 4.9	3,800	54.1	50.1 - 58.2	6,900
No	600	6.8	5.2 - 8.9	300	3.3	2.2 - 4.9	900	10.0	7.9 - 12.6	400	4.3	3.0 - 6.2	7,100	75.5	72.1 - 78.7	9,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,700	10.8	9.2 - 12.7	600	3.8	2.9 - 5.1	2,200	13.8	12.0 - 15.8	400	2.3	1.6 - 3.2	11,200	69.3	66.6 - 71.8	16,100

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	10.5	8.1 - 13.5	100	4.2	2.6 - 6.5	300	17.5	14.3 - 21.1	100	3.2	1.9 - 5.4	1,200	64.7	60.2 - 68.9	1,900
Honolulu	1,300	11.2	9.1 - 13.8	400	3.7	2.5 - 5.5	1,500	12.4	10.1 - 15.1	300	2.3	1.4 - 3.5	8,400	70.4	66.9 - 73.6	11,900
Kauai	100	12.6	8.1 - 19.1	<50	1.1	0.2 - 6.3	100	18.9	13.4 - 25.9	<50	2.5	0.9 - 7.0	400	64.9	56.9 - 72.2	600
Maui	100	7.6	5.3 - 10.8	100	5.0	3.2 - 7.8	300	17.3	13.8 - 21.6	<50	1.4	0.6 - 3.2	1,200	68.6	63.6 - 73.1	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	9.6	6.1 - 14.8	100	2.7	1.3 - 5.7	300	9.5	6.1 - 14.6	200	5.3	3.0 - 9.2	2,500	72.8	66.3 - 78.5	3,400
Native Hawaiian	700	15.5	13.4 - 18.0	200	4.2	3.1 - 5.6	700	16.4	14.3 - 18.8	100	2.9	1.9 - 4.2	2,700	61.0	57.9 - 64.0	4,400
Chinese	100	15.6	6.4 - 33.4	<50	5.5	1.6 - 17.4	100	9.7	3.2 - 25.5	<50	3.2	0.5 - 18.9	400	66.0	48.5 - 80.0	600
Filipino	200	7.1	4.2 - 11.5	200	5.4	2.8 - 9.9	600	19.2	14.4 - 25.1	<50	1.2	0.4 - 3.6	2,100	67.2	60.4 - 73.3	3,200
Japanese	<50	0.5	0.1 - 1.9	<50	0.2	0.0 - 1.4	200	7.8	4.4 - 13.5	<50	0.2	0.0 - 1.4	2,100	91.2	85.5 - 94.8	2,400
Other	400	18.5	12.0 - 27.5	100	6.0	2.7 - 12.6	300	14.8	9.2 - 22.9	0	0.0	-	1,300	60.7	51.1 - 69.6	2,200

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	17.8	11.4 - 26.8	100	6.1	3.3 - 10.8	400	25.4	18.5 - 33.8	<50	0.9	0.3 - 2.7	700	49.8	40.8 - 58.8	1,500
20-24	400	10.2	7.3 - 14.0	300	6.4	3.9 - 10.1	700	17.9	13.9 - 22.6	100	2.5	1.6 - 3.9	2,500	63.1	57.6 - 68.3	4,000
25-34	800	10.3	8.1 - 13.1	200	2.7	1.7 - 4.3	900	11.2	8.9 - 14.0	200	2.6	1.6 - 4.3	5,900	73.2	69.4 - 76.6	8,100
35+	200	9.3	5.8 - 14.6	100	2.1	0.8 - 5.7	200	8.9	5.8 - 13.3	<50	1.9	0.6 - 5.6	2,000	77.8	71.3 - 83.1	2,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,000	12.8	10.2 - 15.8	300	4.3	2.8 - 6.4	1,100	13.7	11.2 - 16.6	200	2.1	1.2 - 3.4	5,400	67.2	63.3 - 70.9	8,100
Female	700	8.9	6.9 - 11.3	300	3.4	2.3 - 4.9	1,100	13.9	11.3 - 16.9	200	2.5	1.6 - 4.0	5,700	71.4	67.7 - 74.8	8,000

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	8.3	6.5 - 10.5	300	3.0	1.9 - 4.6	1,200	11.2	9.1 - 13.7	300	2.4	1.5 - 3.7	8,200	75.2	71.9 - 78.2	10,900
No	800	16.1	12.9 - 20.0	300	5.6	3.9 - 8.1	1,000	19.2	15.9 - 23.0	100	2.2	1.4 - 3.3	2,900	56.8	52.3 - 61.3	5,200

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,000	15.5	12.6 - 19.0	300	5.4	3.6 - 8.1	1,000	16.6	13.6 - 20.0	100	2.0	1.3 - 2.9	3,800	60.5	56.2 - 64.7	6,300
131-185%	300	9.2	5.9 - 14.2	100	4.1	2.3 - 7.4	500	14.4	10.5 - 19.5	100	2.5	1.1 - 5.6	2,300	69.8	63.4 - 75.5	3,300
186+%	200	3.0	1.7 - 5.3	100	1.6	0.7 - 3.3	500	8.9	6.4 - 12.2	200	3.0	1.6 - 5.5	4,300	83.6	79.4 - 87.0	5,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	600	18.6	14.4 - 23.8	200	6.2	3.8 - 9.8	500	17.1	13.0 - 22.3	100	2.6	1.6 - 4.4	1,700	55.4	49.5 - 61.2	3,100
\$15,001-\$25,000	300	13.6	9.1 - 19.9	100	5.3	2.7 - 10.1	300	13.2	9.3 - 18.4	100	2.6	1.2 - 5.3	1,600	65.3	58.0 - 71.9	2,400
\$25,001-\$40,000	400	11.7	8.2 - 16.4	200	4.8	2.6 - 8.8	500	14.5	10.7 - 19.3	100	2.2	0.9 - 5.2	2,300	66.8	60.6 - 72.5	3,400
>=\$40,001	200	3.0	1.7 - 5.1	100	1.7	0.9 - 3.4	700	10.6	8.0 - 13.8	100	2.4	1.3 - 4.3	5,100	82.3	78.5 - 85.6	6,200

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	6.5	3.6 - 11.3	100	5.6	1.8 - 16.0	200	20.8	12.6 - 32.4	<50	2.5	0.5 - 11.1	600	64.6	52.6 - 75.0	1,000
Normal birth weight	1,700	11.1	9.4 - 13.1	600	3.7	2.7 - 5.0	2,000	13.3	11.5 - 15.4	300	2.3	1.6 - 3.2	10,500	69.6	66.8 - 72.2	15,100

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	29.5	21.8 - 38.7	<50	3.9	1.5 - 9.4	300	21.5	14.8 - 30.3	<50	3.7	1.8 - 7.3	500	41.4	32.8 - 50.6	1,200
No	1,400	9.3	7.7 - 11.3	600	3.8	2.8 - 5.2	1,900	13.1	11.2 - 15.2	300	2.2	1.5 - 3.2	10,500	71.6	68.8 - 74.2	14,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	17.1	13.2 - 22.0	200	6.1	3.6 - 10.2	600	18.9	14.6 - 24.1	100	3.7	2.1 - 6.5	1,700	54.1	48.0 - 60.1	3,100
No	1,200	9.3	7.5 - 11.4	300	2.8	1.9 - 4.0	1,600	12.7	10.7 - 15.0	300	2.0	1.3 - 3.1	9,200	73.2	70.2 - 76.0	12,600

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	16.5	13.5 - 20.0	400	5.3	3.6 - 7.9	1,200	18.4	15.3 - 22.0	100	2.1	1.3 - 3.4	3,900	57.7	53.3 - 61.9	6,700
No	600	6.8	5.1 - 9.0	300	2.8	1.8 - 4.1	1,000	10.5	8.5 - 13.0	200	2.4	1.5 - 3.9	7,200	77.6	74.3 - 80.5	9,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,600	10.3	9.1 - 11.7	600	4.1	3.3 - 5.0	2,400	15.3	13.7 - 17.0	400	2.7	2.0 - 3.5	10,600	67.7	65.5 - 69.7	15,600

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	13.6	10.8 - 17.1	100	4.4	2.8 - 6.8	200	13.0	10.3 - 16.3	100	3.0	1.7 - 5.2	1,200	66.0	61.5 - 70.2	1,800
Honolulu	1,200	10.2	8.7 - 12.0	400	3.9	3.0 - 5.0	1,800	15.6	13.6 - 17.8	300	2.5	1.7 - 3.6	7,800	67.8	65.1 - 70.4	11,600
Kauai	<50	4.5	2.1 - 9.4	<50	4.0	1.9 - 8.1	100	14.3	9.4 - 21.3	<50	4.1	1.8 - 8.8	500	73.1	65.3 - 79.7	700
Maui	200	9.9	7.2 - 13.4	100	5.3	3.4 - 8.1	300	15.6	12.2 - 19.7	100	3.1	1.8 - 5.3	1,100	66.2	61.1 - 70.9	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	7.7	5.2 - 11.1	100	2.0	1.0 - 4.0	400	11.7	8.8 - 15.5	100	1.8	0.8 - 3.9	2,600	76.8	72.2 - 80.9	3,400
Native Hawaiian	700	16.1	14.4 - 18.1	300	6.1	5.0 - 7.5	700	17.5	15.7 - 19.5	100	2.7	2.0 - 3.6	2,400	57.5	55.0 - 59.9	4,200
Chinese	<50	4.8	1.3 - 16.6	<50	2.9	1.0 - 8.1	100	15.0	7.1 - 29.0	<50	4.8	1.3 - 16.6	500	72.5	58.1 - 83.4	600
Filipino	300	8.0	5.6 - 11.2	200	5.0	3.1 - 7.8	600	18.8	14.7 - 23.8	100	2.9	1.5 - 5.6	2,100	65.4	59.8 - 70.6	3,200
Japanese	100	4.4	2.1 - 8.9	<50	0.9	0.3 - 2.5	200	11.9	7.9 - 17.5	100	4.1	1.9 - 8.6	1,600	78.8	72.0 - 84.2	2,000
Other	300	13.8	9.4 - 19.7	100	5.4	2.9 - 9.8	300	14.3	9.8 - 20.3	<50	1.7	0.6 - 4.5	1,400	64.9	57.6 - 71.6	2,100

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	16.4	12.8 - 20.7	100	8.3	5.5 - 12.3	400	28.4	23.2 - 34.4	<50	0.7	0.3 - 1.8	700	46.2	40.2 - 52.4	1,500
20-24	500	12.0	9.6 - 14.9	200	5.9	4.0 - 8.4	800	19.9	16.6 - 23.7	100	2.7	1.6 - 4.6	2,400	59.5	55.2 - 63.7	4,100
25-34	700	9.0	7.3 - 11.0	200	2.9	2.1 - 3.8	900	12.2	10.1 - 14.7	200	2.7	1.8 - 4.1	5,500	73.2	70.1 - 76.1	7,500
35+	200	7.9	5.0 - 12.3	100	2.3	1.2 - 4.4	200	8.4	5.5 - 12.6	100	3.7	1.9 - 7.0	1,900	77.7	72.1 - 82.5	2,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	9.2	7.7 - 10.9	400	4.5	3.4 - 5.9	1,200	15.5	13.2 - 18.0	200	2.5	1.7 - 3.9	5,500	68.3	65.3 - 71.2	8,000
Female	900	11.5	9.6 - 13.8	300	3.7	2.8 - 4.9	1,100	15.0	12.9 - 17.5	200	2.8	1.9 - 4.1	5,100	67.0	63.9 - 69.9	7,600

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	7.5	6.1 - 9.1	300	2.9	2.2 - 3.9	1,300	12.7	10.9 - 14.8	300	3.1	2.2 - 4.3	7,700	73.8	71.2 - 76.3	10,400
No	800	16.0	13.6 - 18.7	300	6.4	4.8 - 8.4	1,100	20.3	17.5 - 23.5	100	1.9	1.2 - 2.9	2,900	55.4	51.8 - 59.0	5,200

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	900	14.9	12.8 - 17.4	300	5.4	4.1 - 7.2	1,000	17.5	14.9 - 20.3	100	2.1	1.3 - 3.4	3,500	60.0	56.6 - 63.4	5,900
131-185%	300	9.9	7.4 - 13.2	100	2.7	1.8 - 4.1	500	14.8	11.7 - 18.6	100	2.5	1.5 - 4.1	2,400	70.0	65.5 - 74.2	3,500
186+%	200	4.2	2.7 - 6.3	100	2.5	1.6 - 3.9	500	9.6	7.4 - 12.4	200	3.6	2.3 - 5.8	4,200	80.1	76.4 - 83.3	5,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	500	16.3	13.3 - 19.8	200	6.7	4.7 - 9.5	500	16.8	13.4 - 20.8	100	1.7	0.9 - 3.3	1,800	58.4	53.6 - 63.0	3,200
\$15,001-\$25,000	300	12.0	8.9 - 15.9	100	4.3	2.4 - 7.4	400	20.8	16.3 - 26.2	100	3.8	1.9 - 7.4	1,200	59.2	53.2 - 64.9	2,100
\$25,001-\$40,000	300	9.9	7.4 - 13.0	100	2.6	1.8 - 3.7	600	16.4	13.2 - 20.3	100	2.2	1.3 - 3.8	2,400	68.9	64.4 - 73.1	3,400
>=\$40,001	400	5.8	4.2 - 7.9	200	2.7	1.8 - 3.9	600	10.1	8.0 - 12.8	200	3.4	2.2 - 5.2	4,800	78.0	74.7 - 81.1	6,100

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	15.0	10.0 - 22.0	<50	3.7	1.5 - 8.8	200	19.3	13.0 - 27.6	<50	1.8	0.7 - 4.6	600	60.2	51.1 - 68.6	1,000
Normal birth weight	1,500	10.0	8.7 - 11.4	600	4.1	3.4 - 5.1	2,200	15.0	13.4 - 16.8	400	2.7	2.0 - 3.7	10,000	68.1	65.9 - 70.3	14,600

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	23.6	18.7 - 29.4	100	4.8	3.1 - 7.5	300	21.1	15.4 - 28.2	<50	2.1	1.0 - 4.3	700	48.3	41.4 - 55.3	1,400
No	1,200	8.9	7.7 - 10.4	500	3.9	3.1 - 4.9	2,000	14.6	13.0 - 16.4	400	2.7	2.0 - 3.7	9,800	69.8	67.5 - 72.0	14,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	15.6	12.7 - 18.9	100	4.3	3.1 - 5.9	700	21.9	18.1 - 26.4	100	2.6	1.5 - 4.3	1,700	55.6	50.9 - 60.3	3,100
No	1,100	9.0	7.6 - 10.6	500	3.8	3.0 - 4.9	1,600	13.5	11.8 - 15.4	300	2.6	1.9 - 3.6	8,500	71.1	68.6 - 73.4	12,000

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	13.5	11.6 - 15.8	400	5.9	4.5 - 7.6	1,200	18.7	16.2 - 21.5	200	2.9	1.9 - 4.4	3,900	58.9	55.6 - 62.1	6,600
No	700	7.8	6.3 - 9.7	200	2.8	2.0 - 3.8	1,100	12.7	10.7 - 15.0	200	2.5	1.7 - 3.7	6,600	74.2	71.3 - 76.8	8,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,700	10.9	9.6 - 12.4	500	3.4	2.7 - 4.4	2,300	14.5	13.1 - 16.1	600	3.5	2.8 - 4.4	10,600	67.6	65.5 - 69.7	15,800

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	10.8	8.2 - 14.1	100	4.5	2.8 - 7.2	400	20.7	17.1 - 24.8	100	3.4	2.0 - 5.7	1,100	60.6	55.8 - 65.2	1,800
Honolulu	1,300	11.5	9.8 - 13.4	400	3.3	2.4 - 4.5	1,500	13.2	11.5 - 15.2	400	3.6	2.7 - 4.8	8,000	68.4	65.7 - 71.0	11,700
Kauai	<50	5.8	2.8 - 11.7	<50	3.3	1.3 - 8.0	100	14.7	9.6 - 21.7	<50	1.8	0.5 - 6.1	500	74.4	66.4 - 81.1	600
Maui	100	8.9	6.3 - 12.5	100	3.2	1.8 - 5.6	300	16.7	13.2 - 20.9	100	3.8	2.2 - 6.5	1,100	67.3	62.2 - 72.1	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	400	10.5	7.9 - 13.9	100	3.2	1.8 - 5.6	400	11.7	8.8 - 15.2	200	4.0	2.6 - 6.1	2,700	70.6	66.0 - 74.8	3,800
Native Hawaiian	600	13.2	11.5 - 15.1	200	4.1	3.2 - 5.3	900	21.2	19.1 - 23.4	200	3.8	2.9 - 5.0	2,400	57.6	55.0 - 60.2	4,200
Chinese	100	8.1	3.2 - 18.8	<50	2.3	0.3 - 14.0	100	16.6	8.5 - 29.7	<50	2.3	0.3 - 14.0	400	70.7	56.7 - 81.5	600
Filipino	300	8.7	5.8 - 12.7	100	3.9	2.2 - 7.0	500	15.9	12.4 - 20.1	100	3.1	1.8 - 5.4	2,100	68.4	63.0 - 73.3	3,100
Japanese	100	6.9	3.9 - 12.0	<50	1.0	0.3 - 2.7	100	4.2	2.0 - 8.7	100	4.3	1.9 - 9.4	1,600	83.7	77.1 - 88.6	1,900
Other	300	15.0	10.3 - 21.3	100	4.2	1.9 - 8.9	300	13.3	8.9 - 19.4	100	2.4	1.0 - 6.0	1,400	65.1	57.5 - 71.9	2,100

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	11.2	7.7 - 16.1	100	7.5	5.1 - 10.9	400	25.5	20.6 - 31.0	100	3.3	1.6 - 6.6	900	52.5	46.3 - 58.7	1,800
20-24	500	12.6	10.0 - 15.7	200	5.0	3.2 - 7.7	700	18.7	15.7 - 22.1	100	3.0	2.0 - 4.5	2,300	60.7	56.4 - 64.9	3,800
25-34	800	10.1	8.2 - 12.3	200	2.5	1.6 - 3.8	900	12.0	10.0 - 14.3	300	3.6	2.6 - 5.0	5,400	71.8	68.7 - 74.8	7,500
35+	300	10.6	7.3 - 15.1	<50	1.3	0.4 - 3.7	200	8.6	5.8 - 12.4	100	4.3	2.3 - 7.8	2,000	75.3	69.7 - 80.1	2,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,000	12.6	10.6 - 15.0	300	3.4	2.5 - 4.7	1,200	14.7	12.6 - 17.0	200	2.8	2.0 - 3.9	5,400	66.4	63.4 - 69.3	8,100
Female	700	9.1	7.4 - 11.1	300	3.5	2.4 - 5.0	1,100	14.3	12.3 - 16.6	300	4.3	3.1 - 5.8	5,300	68.8	65.8 - 71.8	7,600

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	10.3	8.7 - 12.3	200	2.1	1.4 - 3.2	1,300	11.8	10.2 - 13.7	400	3.6	2.7 - 4.8	7,800	72.1	69.4 - 74.6	10,800
No	600	12.2	10.0 - 14.7	300	6.3	4.6 - 8.4	1,000	20.4	17.6 - 23.5	200	3.4	2.3 - 4.9	2,800	57.8	54.1 - 61.4	4,900

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	900	13.5	11.4 - 15.8	300	4.4	3.2 - 6.0	1,300	19.2	16.8 - 21.9	200	3.0	2.2 - 4.1	4,000	59.9	56.6 - 63.1	6,700
131-185%	200	10.0	7.1 - 14.0	100	3.3	1.7 - 6.3	300	12.0	9.0 - 15.8	<50	1.7	0.9 - 3.0	1,600	73.1	67.8 - 77.7	2,200
186+%	400	6.4	4.5 - 8.9	100	2.0	1.1 - 3.6	600	9.9	7.8 - 12.5	300	4.9	3.4 - 7.1	4,400	76.8	73.1 - 80.1	5,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	500	13.7	11.1 - 16.8	200	6.7	4.8 - 9.5	800	21.1	17.8 - 24.9	100	2.9	1.8 - 4.7	2,000	55.5	51.0 - 59.9	3,600
\$15,001-\$25,000	400	14.5	11.0 - 18.9	100	2.4	1.2 - 4.8	400	17.3	13.4 - 21.9	100	2.7	1.6 - 4.6	1,600	63.1	57.6 - 68.4	2,500
\$25,001-\$40,000	200	8.1	5.8 - 11.3	100	2.7	1.4 - 5.3	400	14.3	11.4 - 17.9	100	3.0	2.0 - 4.6	2,200	71.8	67.2 - 76.0	3,100
>=\$40,001	400	7.7	5.7 - 10.3	100	1.8	1.1 - 3.2	500	9.1	7.1 - 11.5	300	4.5	3.0 - 6.6	4,400	76.9	73.3 - 80.1	5,800

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	13.5	8.7 - 20.2	<50	4.7	1.8 - 11.7	200	17.9	12.5 - 25.0	<50	3.5	1.6 - 7.3	600	60.4	51.6 - 68.6	1,000
Normal birth weight	1,600	10.7	9.3 - 12.3	500	3.4	2.6 - 4.3	2,100	14.1	12.6 - 15.7	500	3.5	2.7 - 4.4	10,000	68.4	66.2 - 70.5	14,600

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	29.3	22.9 - 36.6	100	7.6	4.4 - 12.7	300	20.8	15.5 - 27.4	<50	2.3	1.1 - 4.6	500	40.0	33.1 - 47.3	1,300
No	1,300	9.0	7.6 - 10.5	400	2.8	2.1 - 3.7	1,900	13.8	12.3 - 15.5	500	3.6	2.8 - 4.6	9,900	70.9	68.7 - 73.0	14,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	17.5	14.0 - 21.6	200	5.3	3.5 - 7.9	700	22.4	18.7 - 26.5	100	3.7	2.2 - 6.3	1,600	51.2	46.3 - 56.0	3,200
No	1,100	8.9	7.5 - 10.5	300	2.7	2.0 - 3.8	1,500	12.3	10.7 - 14.0	400	3.4	2.6 - 4.4	8,800	72.8	70.4 - 75.0	12,000

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	14.5	12.3 - 17.0	300	4.8	3.5 - 6.5	1,200	17.8	15.6 - 20.1	200	2.7	1.9 - 3.8	4,100	60.3	57.1 - 63.4	6,800
No	700	7.9	6.4 - 9.9	200	2.2	1.4 - 3.3	1,000	12.1	10.1 - 14.4	400	4.1	3.0 - 5.5	6,400	73.7	70.8 - 76.5	8,700