

**PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration 8+ weeks	Breastfeeding (any) duration 8+ weeks

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/28/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


**PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	5,000	31.7	29.6 - 33.8	10,800	68.3	66.2 - 70.4	15,800
2001	5,000	31.8	29.7 - 33.9	10,700	68.2	66.1 - 70.3	15,600
2002	4,900	30.4	27.9 - 33.1	11,200	69.6	66.9 - 72.1	16,100
2003	5,400	32.4	29.9 - 35.1	11,200	67.6	64.9 - 70.1	16,500
2004	5,300	30.4	28.4 - 32.5	12,000	69.6	67.5 - 71.6	17,300
2005	5,200	30.6	28.4 - 32.8	11,800	69.4	67.2 - 71.6	17,000
2006	5,200	29.4	27.1 - 31.7	12,500	70.6	68.3 - 72.9	17,700
2007	5,200	29.0	26.8 - 31.2	12,700	71.0	68.8 - 73.2	17,900
2008	5,000	27.5	25.4 - 29.7	13,200	72.5	70.3 - 74.6	18,200
2009	5,300	29.7	26.9 - 32.8	12,600	70.3	67.2 - 73.1	17,900
2010	4,800	26.0	23.2 - 29.0	13,500	74.0	71.0 - 76.8	18,300
2011	4,000	21.8	19.2 - 24.6	14,300	78.2	75.4 - 80.8	18,300
2012	4,000	22.2	19.5 - 25.1	14,200	77.8	74.9 - 80.5	18,200
2013	3,600	19.8	17.2 - 22.8	14,500	80.2	77.2 - 82.8	18,100

**PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,600	19.8	17.2 - 22.8	14,500	80.2	77.2 - 82.8	18,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	19.8	15.6 - 24.8	1,900	80.2	75.2 - 84.4	2,300
Honolulu	2,600	19.7	16.2 - 23.7	10,400	80.3	76.3 - 83.8	12,900
Kauai	200	20.1	16.2 - 24.7	700	79.9	75.3 - 83.8	800
Maui	400	20.4	16.3 - 25.3	1,500	79.6	74.7 - 83.7	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	15.8	11.2 - 21.8	4,100	84.2	78.2 - 88.8	4,900
Native Hawaiian	1,500	28.4	23.2 - 34.3	3,800	71.6	65.7 - 76.8	5,300
Chinese	200	20.6	10.4 - 36.8	700	79.4	63.2 - 89.6	900
Filipino	600	21.5	15.4 - 29.1	2,200	78.5	70.9 - 84.6	2,800
Japanese	100	4.6	1.6 - 12.2	1,700	95.4	87.8 - 98.4	1,800
Other	400	18.0	11.7 - 26.7	1,900	82.0	73.3 - 88.3	2,300

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	30.5	18.6 - 45.8	600	69.5	54.2 - 81.4	800
20-24	700	18.6	13.5 - 25.1	3,300	81.4	74.9 - 86.5	4,000
25-34	1,900	19.9	16.3 - 24.0	7,900	80.1	76.0 - 83.7	9,800
35+	600	18.7	13.4 - 25.4	2,800	81.3	74.6 - 86.6	3,400

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,600	18.4	14.9 - 22.4	7,200	81.6	77.6 - 85.1	8,900
Female	2,000	21.2	17.4 - 25.6	7,300	78.8	74.4 - 82.6	9,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	17.5	14.3 - 21.2	9,500	82.5	78.8 - 85.7	11,500
No	1,600	24.0	19.6 - 29.0	5,000	76.0	71.0 - 80.4	6,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	25.7	21.1 - 31.0	4,900	74.3	69.0 - 78.9	6,600
131-185%	500	21.1	14.3 - 30.1	1,900	78.9	69.9 - 85.7	2,500
186+%	1,000	13.7	10.2 - 18.1	6,300	86.3	81.9 - 89.8	7,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Income (2012+)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	1,100	27.1	21.2 - 34.0	3,000	72.9	66.0 - 78.8	4,100
\$17,001-\$26,000	500	21.0	14.4 - 29.7	1,900	79.0	70.3 - 85.6	2,400
\$26,001-\$51,000	800	20.8	15.3 - 27.7	3,100	79.2	72.3 - 84.7	3,900
>=\$51,001	800	13.0	9.4 - 17.6	5,400	87.0	82.4 - 90.6	6,200

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	22.9	19.0 - 27.3	1,000	77.1	72.7 - 81.0	1,300
Normal birth weight	3,300	19.6	16.8 - 22.8	13,500	80.4	77.2 - 83.2	16,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	35.7	23.6 - 49.9	600	64.3	50.1 - 76.4	900
No	3,200	19.1	16.3 - 22.1	13,500	80.9	77.9 - 83.7	16,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	29.0	22.1 - 37.0	2,300	71.0	63.0 - 77.9	3,200
No	2,600	17.9	15.1 - 21.2	11,800	82.1	78.8 - 84.9	14,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	26.7	22.3 - 31.7	5,700	73.3	68.3 - 77.7	7,800
No	1,500	14.3	11.4 - 17.9	8,800	85.7	82.1 - 88.6	10,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	4,000	22.2	19.5 - 25.1	14,200	77.8	74.9 - 80.5	18,200

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	22.6	18.5 - 27.5	1,800	77.4	72.5 - 81.5	2,300
Honolulu	3,000	22.5	19.1 - 26.4	10,200	77.5	73.6 - 80.9	13,200
Kauai	200	19.7	15.0 - 25.4	700	80.3	74.6 - 85.0	800
Maui	400	20.2	16.4 - 24.7	1,500	79.8	75.3 - 83.6	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	17.5	12.7 - 23.6	3,400	82.5	76.4 - 87.3	4,100
Native Hawaiian	1,500	29.6	24.4 - 35.4	3,600	70.4	64.6 - 75.6	5,200
Chinese	200	18.3	9.6 - 32.2	900	81.7	67.8 - 90.4	1,100
Filipino	900	28.2	21.6 - 35.8	2,300	71.8	64.2 - 78.4	3,200
Japanese	100	8.4	4.2 - 16.2	1,400	91.6	83.8 - 95.8	1,600
Other	500	18.1	12.4 - 25.6	2,500	81.9	74.4 - 87.6	3,000

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	38.3	25.9 - 52.4	600	61.7	47.6 - 74.1	1,000
20-24	1,000	27.3	21.2 - 34.4	2,700	72.7	65.6 - 78.8	3,700
25-34	2,100	20.7	17.3 - 24.6	8,100	79.3	75.4 - 82.7	10,200
35+	500	15.7	11.1 - 21.8	2,700	84.3	78.2 - 88.9	3,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	22.2	18.4 - 26.4	7,000	77.8	73.6 - 81.6	9,000
Female	2,000	22.2	18.6 - 26.3	7,100	77.8	73.7 - 81.4	9,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	18.2	15.1 - 21.7	9,200	81.8	78.3 - 84.9	11,200
No	2,000	28.6	24.1 - 33.6	5,000	71.4	66.4 - 75.9	7,000

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,800	28.6	23.9 - 33.8	4,600	71.4	66.2 - 76.1	6,400
131-185%	800	25.5	18.9 - 33.5	2,300	74.5	66.5 - 81.1	3,100
186+%	1,000	13.8	10.5 - 18.0	6,000	86.2	82.0 - 89.5	6,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Income (2012+)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	1,100	27.8	22.0 - 34.4	2,900	72.2	65.6 - 78.0	4,100
\$17,001-\$26,000	600	24.7	17.8 - 33.3	1,700	75.3	66.7 - 82.2	2,200
\$26,001-\$51,000	1,200	28.1	22.2 - 34.8	3,200	71.9	65.2 - 77.8	4,400
>=\$51,001	700	12.6	9.2 - 17.0	5,200	87.4	83.0 - 90.8	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	23.2	18.7 - 28.5	900	76.8	71.5 - 81.3	1,200
Normal birth weight	3,800	22.1	19.3 - 25.2	13,200	77.9	74.8 - 80.7	17,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	46.6	31.7 - 62.0	400	53.4	38.0 - 68.3	700
No	3,600	21.5	18.8 - 24.5	13,200	78.5	75.5 - 81.2	16,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	32.2	25.0 - 40.3	1,900	67.8	59.7 - 75.0	2,800
No	3,000	20.6	17.8 - 23.8	11,700	79.4	76.2 - 82.2	14,700

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	30.8	26.5 - 35.6	5,700	69.2	64.4 - 73.5	8,200
No	1,400	14.4	11.5 - 18.0	8,400	85.6	82.0 - 88.5	9,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	4,000	21.8	19.2 - 24.6	14,300	78.2	75.4 - 80.8	18,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	25.3	21.1 - 30.1	1,800	74.7	69.9 - 78.9	2,400
Honolulu	2,800	21.3	17.9 - 25.1	10,300	78.7	74.9 - 82.1	13,100
Kauai	100	15.8	11.7 - 20.9	700	84.2	79.1 - 88.3	900
Maui	500	23.7	20.0 - 28.0	1,500	76.3	72.0 - 80.0	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	18.5	13.8 - 24.4	3,600	81.5	75.6 - 86.2	4,500
Native Hawaiian	1,400	27.5	22.6 - 33.0	3,800	72.5	67.0 - 77.4	5,200
Chinese	100	16.6	7.4 - 33.3	600	83.4	66.7 - 92.6	700
Filipino	800	24.0	18.5 - 30.6	2,500	76.0	69.4 - 81.5	3,300
Japanese	300	14.2	8.3 - 23.2	1,600	85.8	76.8 - 91.7	1,900
Other	600	20.1	13.7 - 28.4	2,200	79.9	71.6 - 86.3	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	49.1	36.8 - 61.6	500	50.9	38.4 - 63.2	1,000
20-24	1,100	26.0	20.4 - 32.4	3,200	74.0	67.6 - 79.6	4,300
25-34	2,000	20.4	17.0 - 24.3	7,800	79.6	75.7 - 83.0	9,800
35+	400	11.4	7.9 - 16.2	2,800	88.6	83.8 - 92.1	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,100	22.5	18.9 - 26.6	7,200	77.5	73.4 - 81.1	9,300
Female	1,900	21.0	17.6 - 25.0	7,100	79.0	75.0 - 82.4	9,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	19.0	16.0 - 22.5	9,200	81.0	77.5 - 84.0	11,400
No	1,800	26.4	22.1 - 31.2	5,100	73.6	68.8 - 77.9	6,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,800	27.9	23.4 - 32.9	4,700	72.1	67.1 - 76.6	6,500
131-185%	600	22.4	16.0 - 30.4	2,100	77.6	69.6 - 84.0	2,700
186+%	1,300	16.3	12.9 - 20.4	6,700	83.7	79.6 - 87.1	8,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	28.2	22.1 - 35.1	2,400	71.8	64.9 - 77.9	3,300
\$10,000-\$24,999	900	26.6	20.5 - 33.7	2,500	73.4	66.3 - 79.5	3,400
\$25,000-\$49,999	1,100	25.1	19.6 - 31.4	3,400	74.9	68.6 - 80.4	4,500
>=\$50,000	800	13.4	10.0 - 17.7	5,300	86.6	82.3 - 90.0	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	29.2	24.3 - 34.7	900	70.8	65.3 - 75.7	1,200
Normal birth weight	3,600	21.3	18.5 - 24.2	13,400	78.7	75.8 - 81.5	17,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	48.1	35.3 - 61.1	500	51.9	38.9 - 64.7	900
No	3,500	20.1	17.5 - 22.9	13,700	79.9	77.1 - 82.5	17,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	32.3	26.0 - 39.3	2,400	67.7	60.7 - 74.0	3,500
No	2,800	18.9	16.2 - 22.0	11,800	81.1	78.0 - 83.8	14,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	26.6	22.6 - 31.0	6,000	73.4	69.0 - 77.4	8,200
No	1,700	17.7	14.5 - 21.4	8,100	82.3	78.6 - 85.5	9,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	4,800	26.0	23.2 - 29.0	13,500	74.0	71.0 - 76.8	18,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	25.9	21.6 - 30.7	1,700	74.1	69.3 - 78.4	2,300
Honolulu	3,600	27.0	23.3 - 31.0	9,700	73.0	69.0 - 76.7	13,200
Kauai	100	17.5	13.4 - 22.4	700	82.5	77.6 - 86.6	800
Maui	500	23.2	19.4 - 27.6	1,500	76.8	72.4 - 80.6	2,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,100	25.1	19.6 - 31.6	3,300	74.9	68.4 - 80.4	4,500
Native Hawaiian	1,600	29.2	24.3 - 34.7	3,800	70.8	65.3 - 75.7	5,400
Chinese	100	13.0	5.2 - 28.8	600	87.0	71.2 - 94.8	700
Filipino	1,100	32.9	26.3 - 40.3	2,200	67.1	59.7 - 73.7	3,300
Japanese	100	6.6	3.2 - 12.8	1,600	93.4	87.2 - 96.8	1,700
Other	800	28.3	20.8 - 37.2	1,900	71.7	62.8 - 79.2	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	42.8	31.0 - 55.4	600	57.2	44.6 - 69.0	1,100
20-24	1,400	31.6	25.6 - 38.1	3,000	68.4	61.9 - 74.4	4,400
25-34	2,200	23.0	19.4 - 27.1	7,200	77.0	72.9 - 80.6	9,400
35+	700	21.4	15.8 - 28.4	2,600	78.6	71.6 - 84.2	3,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,500	26.6	22.8 - 30.9	6,800	73.4	69.1 - 77.2	9,200
Female	2,300	25.4	21.5 - 29.7	6,700	74.6	70.3 - 78.5	9,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	21.2	17.9 - 24.9	9,000	78.8	75.1 - 82.1	11,400
No	2,300	34.1	29.4 - 39.1	4,500	65.9	60.9 - 70.6	6,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,000	31.1	26.4 - 36.2	4,300	68.9	63.8 - 73.6	6,300
131-185%	800	27.7	20.8 - 35.9	2,100	72.3	64.1 - 79.2	3,000
186+%	1,500	18.7	14.9 - 23.1	6,300	81.3	76.9 - 85.1	7,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,100	32.0	25.7 - 39.0	2,300	68.0	61.0 - 74.3	3,300
\$10,000-\$24,999	1,000	30.0	23.5 - 37.4	2,300	70.0	62.6 - 76.5	3,200
\$25,000-\$49,999	1,300	28.3	22.6 - 34.8	3,200	71.7	65.2 - 77.4	4,500
>=\$50,000	1,100	17.1	13.1 - 22.0	5,100	82.9	78.0 - 86.9	6,200

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	29.8	24.9 - 35.1	900	70.2	64.9 - 75.1	1,300
Normal birth weight	4,400	25.7	22.8 - 28.9	12,600	74.3	71.1 - 77.2	17,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	44.8	33.5 - 56.7	800	55.2	43.3 - 66.5	1,400
No	4,100	24.6	21.7 - 27.6	12,600	75.4	72.4 - 78.3	16,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	40.9	34.2 - 48.0	2,200	59.1	52.0 - 65.8	3,700
No	3,200	22.2	19.3 - 25.5	11,100	77.8	74.5 - 80.7	14,300

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	31.4	27.1 - 36.0	5,500	68.6	64.0 - 72.9	8,000
No	2,200	21.8	18.3 - 25.8	7,900	78.2	74.2 - 81.7	10,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,300	29.7	26.9 - 32.8	12,600	70.3	67.2 - 73.1	17,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	31.8	27.2 - 36.9	1,600	68.2	63.1 - 72.8	2,300
Honolulu	3,900	30.5	26.7 - 34.6	8,800	69.5	65.4 - 73.3	12,700
Kauai	200	23.5	18.7 - 29.1	600	76.5	70.9 - 81.3	800
Maui	500	25.3	21.3 - 29.6	1,500	74.7	70.4 - 78.7	2,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	20.1	15.2 - 26.0	3,400	79.9	74.0 - 84.8	4,200
Native Hawaiian	2,200	39.1	33.8 - 44.7	3,500	60.9	55.3 - 66.2	5,700
Chinese	200	21.3	10.6 - 38.2	600	78.7	61.8 - 89.4	800
Filipino	1,000	32.0	25.6 - 39.2	2,200	68.0	60.8 - 74.4	3,200
Japanese	300	16.3	9.8 - 26.0	1,300	83.7	74.0 - 90.2	1,600
Other	800	32.9	24.8 - 42.1	1,600	67.1	57.9 - 75.2	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	46.2	36.0 - 56.7	900	53.8	43.3 - 64.0	1,800
20-24	1,500	35.7	29.6 - 42.4	2,600	64.3	57.6 - 70.4	4,100
25-34	2,300	25.9	22.1 - 30.1	6,700	74.1	69.9 - 77.9	9,100
35+	700	23.6	17.6 - 30.9	2,300	76.4	69.1 - 82.4	3,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,800	29.4	25.5 - 33.7	6,700	70.6	66.3 - 74.5	9,500
Female	2,500	30.1	26.0 - 34.5	5,900	69.9	65.5 - 74.0	8,400

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	21.9	18.6 - 25.6	8,300	78.1	74.4 - 81.4	10,600
No	3,000	41.2	36.3 - 46.2	4,300	58.8	53.8 - 63.7	7,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,400	39.1	34.0 - 44.5	3,800	60.9	55.5 - 66.0	6,300
131-185%	900	32.9	25.7 - 41.1	1,900	67.1	58.9 - 74.3	2,800
186+%	1,400	19.1	15.4 - 23.4	5,900	80.9	76.6 - 84.6	7,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,300	42.9	35.2 - 51.1	1,700	57.1	48.9 - 64.8	3,000
\$10,000-\$24,999	1,400	36.1	30.0 - 42.8	2,400	63.9	57.2 - 70.0	3,800
\$25,000-\$49,999	1,000	28.2	22.1 - 35.1	2,600	71.8	64.9 - 77.9	3,600
>=\$50,000	1,100	18.4	14.4 - 23.1	4,900	81.6	76.9 - 85.6	6,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	34.7	29.7 - 40.0	800	65.3	60.0 - 70.3	1,300
Normal birth weight	4,900	29.4	26.3 - 32.6	11,800	70.6	67.4 - 73.7	16,700

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	54.9	44.4 - 65.0	800	45.1	35.0 - 55.6	1,700
No	4,200	26.8	23.9 - 29.9	11,600	73.2	70.1 - 76.1	15,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	46.1	39.5 - 52.8	2,100	53.9	47.2 - 60.5	3,900
No	3,400	24.7	21.7 - 28.0	10,200	75.3	72.0 - 78.3	13,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	40.5	35.7 - 45.4	4,300	59.5	54.6 - 64.3	7,200
No	2,300	22.3	18.8 - 26.2	7,900	77.7	73.8 - 81.2	10,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,000	27.5	25.4 - 29.7	13,200	72.5	70.3 - 74.6	18,200

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	31.2	25.2 - 38.0	1,600	68.8	62.0 - 74.8	2,400
Honolulu	3,500	27.5	25.0 - 30.1	9,300	72.5	69.9 - 75.0	12,800
Kauai	200	24.9	16.5 - 35.8	600	75.1	64.2 - 83.5	900
Maui	500	24.4	18.7 - 31.1	1,600	75.6	68.9 - 81.3	2,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,000	22.3	18.5 - 26.6	3,400	77.7	73.4 - 81.5	4,300
Native Hawaiian	1,700	32.9	28.4 - 37.7	3,400	67.1	62.3 - 71.6	5,000
Chinese	100	15.4	11.1 - 21.0	500	84.6	79.0 - 88.9	600
Filipino	1,000	31.3	26.6 - 36.5	2,300	68.7	63.5 - 73.4	3,300
Japanese	300	17.1	12.2 - 23.4	1,700	82.9	76.6 - 87.8	2,000
Other	900	31.5	25.9 - 37.6	2,000	68.5	62.4 - 74.1	2,900

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	34.6	26.6 - 43.7	900	65.4	56.3 - 73.4	1,500
20-24	1,500	39.0	33.9 - 44.3	2,400	61.0	55.7 - 66.1	4,000
25-34	2,200	23.9	21.2 - 26.9	7,100	76.1	73.1 - 78.8	9,400
35+	700	21.0	16.9 - 25.8	2,700	79.0	74.2 - 83.1	3,400

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,700	28.2	25.2 - 31.3	6,800	71.8	68.7 - 74.8	9,500
Female	2,300	26.8	23.8 - 30.0	6,300	73.2	70.0 - 76.2	8,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	24.4	21.9 - 27.1	8,600	75.6	72.9 - 78.1	11,400
No	2,200	32.7	29.0 - 36.8	4,500	67.3	63.2 - 71.0	6,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,300	34.4	30.6 - 38.5	4,400	65.6	61.5 - 69.4	6,600
131-185%	600	24.1	19.0 - 30.0	2,000	75.9	70.0 - 81.0	2,600
186+%	1,700	21.5	18.7 - 24.6	6,100	78.5	75.4 - 81.3	7,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,100	37.5	31.7 - 43.7	1,900	62.5	56.3 - 68.3	3,000
\$10,000-\$24,999	1,200	32.1	27.3 - 37.4	2,600	67.9	62.6 - 72.7	3,900
\$25,000-\$49,999	1,200	27.9	23.6 - 32.6	3,000	72.1	67.4 - 76.4	4,200
>=\$50,000	1,100	18.2	15.3 - 21.6	5,000	81.8	78.4 - 84.7	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	27.3	19.7 - 36.5	900	72.7	63.5 - 80.3	1,200
Normal birth weight	4,700	27.5	25.3 - 29.8	12,300	72.5	70.2 - 74.7	16,900

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	45.6	37.0 - 54.4	800	54.4	45.6 - 63.0	1,500
No	4,200	25.9	23.7 - 28.2	12,100	74.1	71.8 - 76.3	16,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	40.7	35.1 - 46.5	2,000	59.3	53.5 - 64.9	3,300
No	3,600	24.5	22.3 - 26.9	10,900	75.5	73.1 - 77.7	14,500

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	35.1	31.5 - 38.8	5,000	64.9	61.2 - 68.5	7,600
No	2,300	22.1	19.6 - 24.9	8,000	77.9	75.1 - 80.4	10,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,200	29.0	26.8 - 31.2	12,700	71.0	68.8 - 73.2	17,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	25.6	20.1 - 32.1	1,800	74.4	67.9 - 79.9	2,400
Honolulu	3,700	29.4	26.9 - 32.1	8,800	70.6	67.9 - 73.1	12,400
Kauai	300	27.9	19.5 - 38.1	700	72.1	61.9 - 80.5	1,000
Maui	600	30.6	24.2 - 37.8	1,500	69.4	62.2 - 75.8	2,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,100	26.7	22.5 - 31.4	3,000	73.3	68.6 - 77.5	4,100
Native Hawaiian	1,900	38.3	33.6 - 43.3	3,000	61.7	56.7 - 66.4	4,900
Chinese	100	14.7	10.4 - 20.4	600	85.3	79.6 - 89.6	700
Filipino	1,000	32.2	27.5 - 37.3	2,200	67.8	62.7 - 72.5	3,200
Japanese	300	13.3	9.0 - 19.1	1,800	86.7	80.9 - 91.0	2,100
Other	800	27.3	22.0 - 33.2	2,100	72.7	66.8 - 78.0	2,900

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	44.9	36.0 - 54.1	900	55.1	45.9 - 64.0	1,600
20-24	1,700	36.9	32.3 - 41.8	2,800	63.1	58.2 - 67.7	4,500
25-34	2,200	25.6	22.7 - 28.7	6,500	74.4	71.3 - 77.3	8,700
35+	600	18.5	14.5 - 23.3	2,500	81.5	76.7 - 85.5	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,700	28.8	25.8 - 31.9	6,600	71.2	68.1 - 74.2	9,300
Female	2,500	29.2	26.1 - 32.5	6,100	70.8	67.5 - 73.9	8,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	22.3	19.9 - 24.9	8,700	77.7	75.1 - 80.1	11,200
No	2,700	40.2	36.2 - 44.4	4,000	59.8	55.6 - 63.8	6,700

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,600	39.4	35.4 - 43.5	4,000	60.6	56.5 - 64.6	6,700
131-185%	700	27.0	21.8 - 32.9	1,900	73.0	67.1 - 78.2	2,600
186+%	1,500	20.3	17.5 - 23.4	5,800	79.7	76.6 - 82.5	7,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,300	43.7	37.6 - 50.0	1,700	56.3	50.0 - 62.4	3,000
\$10,000-\$24,999	1,200	34.1	29.1 - 39.4	2,400	65.9	60.6 - 70.9	3,700
\$25,000-\$49,999	1,200	28.9	24.6 - 33.7	2,900	71.1	66.3 - 75.4	4,100
>=\$50,000	1,100	18.4	15.5 - 21.8	4,800	81.6	78.2 - 84.5	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	35.0	27.4 - 43.5	1,000	65.0	56.5 - 72.6	1,500
Normal birth weight	4,700	28.4	26.2 - 30.8	11,800	71.6	69.2 - 73.8	16,500

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	56.2	47.4 - 64.6	700	43.8	35.4 - 52.6	1,500
No	4,300	26.7	24.5 - 29.0	11,900	73.3	71.0 - 75.5	16,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	48.0	42.5 - 53.6	1,800	52.0	46.4 - 57.5	3,500
No	3,500	24.5	22.3 - 26.9	10,700	75.5	73.1 - 77.7	14,200

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	36.9	33.2 - 40.8	4,600	63.1	59.2 - 66.8	7,400
No	2,300	23.0	20.4 - 25.8	7,900	77.0	74.2 - 79.6	10,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,200	29.4	27.1 - 31.7	12,500	70.6	68.3 - 72.9	17,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	29.9	23.5 - 37.1	1,500	70.1	62.9 - 76.5	2,100
Honolulu	3,800	29.3	26.7 - 32.0	9,200	70.7	68.0 - 73.3	13,000
Kauai	200	24.9	15.9 - 36.8	600	75.1	63.2 - 84.1	800
Maui	600	31.4	24.5 - 39.2	1,300	68.6	60.8 - 75.5	1,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,100	26.7	22.4 - 31.4	3,100	73.3	68.6 - 77.6	4,200
Native Hawaiian	1,800	37.6	32.8 - 42.7	3,100	62.4	57.3 - 67.2	4,900
Chinese	100	17.4	12.6 - 23.4	500	82.6	76.6 - 87.4	600
Filipino	1,000	30.7	25.8 - 36.0	2,200	69.3	64.0 - 74.2	3,200
Japanese	300	14.9	10.4 - 21.0	1,700	85.1	79.0 - 89.6	2,000
Other	800	30.6	24.8 - 36.9	1,900	69.4	63.1 - 75.2	2,800

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	43.5	34.6 - 52.9	800	56.5	47.1 - 65.4	1,500
20-24	1,700	39.2	34.2 - 44.4	2,700	60.8	55.6 - 65.8	4,400
25-34	2,200	25.1	22.1 - 28.3	6,500	74.9	71.7 - 77.9	8,700
35+	700	21.1	16.9 - 26.0	2,500	78.9	74.0 - 83.1	3,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,900	32.2	29.0 - 35.6	6,100	67.8	64.4 - 71.0	9,000
Female	2,300	26.5	23.4 - 29.7	6,400	73.5	70.3 - 76.6	8,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	23.8	21.3 - 26.5	8,600	76.2	73.5 - 78.7	11,300
No	2,500	39.1	34.9 - 43.5	3,900	60.9	56.5 - 65.1	6,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,500	40.2	36.0 - 44.6	3,700	59.8	55.4 - 64.0	6,200
131-185%	700	29.2	23.3 - 35.9	1,700	70.8	64.1 - 76.7	2,400
186+%	1,600	20.6	17.8 - 23.7	6,300	79.4	76.3 - 82.2	8,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,300	44.5	38.2 - 50.9	1,700	55.5	49.1 - 61.8	3,000
\$10,000-\$24,999	1,300	34.0	28.9 - 39.5	2,400	66.0	60.5 - 71.1	3,700
\$25,000-\$49,999	1,300	31.2	26.7 - 36.1	2,900	68.8	63.9 - 73.3	4,200
>=\$50,000	1,000	16.7	13.8 - 20.0	4,800	83.3	80.0 - 86.2	5,700

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	30.9	23.3 - 39.7	1,000	69.1	60.3 - 76.7	1,400
Normal birth weight	4,800	29.2	26.9 - 31.7	11,500	70.8	68.3 - 73.1	16,300

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	50.3	41.5 - 59.0	800	49.7	41.0 - 58.5	1,600
No	4,300	27.2	24.9 - 29.6	11,600	72.8	70.4 - 75.1	15,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	40.6	35.1 - 46.3	2,200	59.4	53.7 - 64.9	3,700
No	3,600	26.3	23.9 - 28.9	10,200	73.7	71.1 - 76.1	13,800

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	37.9	34.1 - 41.9	4,600	62.1	58.1 - 65.9	7,500
No	2,300	23.0	20.4 - 25.9	7,700	77.0	74.1 - 79.6	10,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,200	30.6	28.4 - 32.8	11,800	69.4	67.2 - 71.6	17,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	30.6	24.6 - 37.4	1,500	69.4	62.6 - 75.4	2,200
Honolulu	3,800	30.5	28.0 - 33.1	8,700	69.5	66.9 - 72.0	12,500
Kauai	100	19.6	11.8 - 30.7	600	80.4	69.3 - 88.2	700
Maui	600	35.8	28.8 - 43.6	1,000	64.2	56.4 - 71.2	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	21.2	17.2 - 25.7	2,900	78.8	74.3 - 82.8	3,700
Native Hawaiian	1,900	39.3	34.5 - 44.2	2,900	60.7	55.8 - 65.5	4,800
Chinese	100	22.9	18.0 - 28.8	500	77.1	71.2 - 82.0	600
Filipino	1,200	37.6	32.8 - 42.6	2,000	62.4	57.4 - 67.2	3,200
Japanese	400	16.2	12.0 - 21.5	1,900	83.8	78.5 - 88.0	2,300
Other	800	34.4	28.6 - 40.6	1,600	65.6	59.4 - 71.4	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	44.1	35.8 - 52.6	800	55.9	47.4 - 64.2	1,500
20-24	1,800	42.4	37.6 - 47.3	2,400	57.6	52.7 - 62.4	4,200
25-34	2,200	26.5	23.6 - 29.6	6,200	73.5	70.4 - 76.4	8,400
35+	600	18.8	14.9 - 23.4	2,400	81.2	76.6 - 85.1	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,700	30.9	27.9 - 34.1	6,000	69.1	65.9 - 72.1	8,700
Female	2,500	30.2	27.2 - 33.5	5,800	69.8	66.5 - 72.8	8,400

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	24.4	21.9 - 27.0	8,200	75.6	73.0 - 78.1	10,900
No	2,600	41.5	37.5 - 45.5	3,600	58.5	54.5 - 62.5	6,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,600	42.1	38.2 - 46.1	3,600	57.9	53.9 - 61.8	6,200
131-185%	600	32.6	26.0 - 39.9	1,200	67.4	60.1 - 74.0	1,800
186+%	1,500	19.2	16.6 - 22.1	6,100	80.8	77.9 - 83.4	7,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,400	46.7	40.9 - 52.6	1,600	53.3	47.4 - 59.1	3,000
\$10,000-\$24,999	1,400	39.8	34.8 - 45.0	2,100	60.2	55.0 - 65.2	3,600
\$25,000-\$49,999	1,000	25.9	21.7 - 30.6	2,900	74.1	69.4 - 78.3	3,900
>=\$50,000	900	17.2	14.3 - 20.7	4,400	82.8	79.3 - 85.7	5,300

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	31.2	23.5 - 40.1	800	68.8	59.9 - 76.5	1,200
Normal birth weight	4,800	30.5	28.3 - 32.9	11,000	69.5	67.1 - 71.7	15,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	57.7	49.0 - 65.9	600	42.3	34.1 - 51.0	1,400
No	4,300	27.9	25.7 - 30.2	11,100	72.1	69.8 - 74.3	15,400

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	47.9	42.6 - 53.3	1,900	52.1	46.7 - 57.4	3,600
No	3,400	25.7	23.4 - 28.1	9,800	74.3	71.9 - 76.6	13,200

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	37.9	34.3 - 41.6	4,500	62.1	58.4 - 65.7	7,300
No	2,300	24.5	21.9 - 27.3	7,100	75.5	72.7 - 78.1	9,400

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,300	30.4	28.4 - 32.5	12,000	69.6	67.5 - 71.6	17,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	29.8	24.2 - 36.1	1,500	70.2	63.9 - 75.8	2,200
Honolulu	3,800	30.6	28.3 - 33.0	8,700	69.4	67.0 - 71.7	12,600
Kauai	200	23.0	15.4 - 32.8	600	77.0	67.2 - 84.6	800
Maui	600	32.9	26.9 - 39.5	1,200	67.1	60.5 - 73.1	1,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	23.0	19.3 - 27.0	3,000	77.0	73.0 - 80.7	3,900
Native Hawaiian	2,000	41.2	36.7 - 45.8	2,800	58.8	54.2 - 63.3	4,700
Chinese	100	17.2	14.0 - 21.0	500	82.8	79.0 - 86.0	700
Filipino	1,200	35.6	31.3 - 40.1	2,100	64.4	59.9 - 68.7	3,300
Japanese	400	16.9	12.6 - 22.3	1,900	83.1	77.7 - 87.4	2,200
Other	800	30.1	24.9 - 36.0	1,800	69.9	64.0 - 75.1	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	48.4	39.9 - 56.9	600	51.6	43.1 - 60.1	1,300
20-24	1,800	40.5	36.1 - 45.0	2,600	59.5	55.0 - 63.9	4,400
25-34	2,100	25.3	22.7 - 28.1	6,300	74.7	71.9 - 77.3	8,500
35+	700	23.1	19.2 - 27.6	2,500	76.9	72.4 - 80.8	3,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,000	31.3	28.6 - 34.1	6,500	68.7	65.9 - 71.4	9,400
Female	2,300	29.4	26.4 - 32.5	5,600	70.6	67.5 - 73.6	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,200	26.0	23.8 - 28.4	9,000	74.0	71.6 - 76.2	12,100
No	2,100	40.6	36.6 - 44.8	3,100	59.4	55.2 - 63.4	5,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,500	38.9	35.3 - 42.7	3,900	61.1	57.3 - 64.7	6,400
131-185%	600	30.0	24.5 - 36.2	1,500	70.0	63.8 - 75.5	2,100
186+%	1,700	22.5	19.9 - 25.3	5,900	77.5	74.7 - 80.1	7,500

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,100	41.4	35.7 - 47.3	1,500	58.6	52.7 - 64.3	2,600
\$10,000-\$24,999	1,700	38.6	34.3 - 43.1	2,600	61.4	56.9 - 65.7	4,300
\$25,000-\$49,999	1,200	26.4	22.7 - 30.4	3,200	73.6	69.6 - 77.3	4,400
>=\$50,000	900	19.0	16.1 - 22.4	3,900	81.0	77.6 - 83.9	4,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	33.5	25.7 - 42.2	700	66.5	57.8 - 74.3	1,100
Normal birth weight	4,900	30.2	28.2 - 32.3	11,300	69.8	67.7 - 71.8	16,200

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	45.2	37.0 - 53.6	700	54.8	46.4 - 63.0	1,300
No	4,600	28.9	26.9 - 31.1	11,200	71.1	68.9 - 73.1	15,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	37.8	32.9 - 42.8	2,200	62.2	57.2 - 67.1	3,500
No	3,800	28.3	26.1 - 30.5	9,800	71.7	69.5 - 73.9	13,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	35.4	32.2 - 38.7	5,000	64.6	61.3 - 67.8	7,700
No	2,400	26.1	23.5 - 28.7	6,800	73.9	71.3 - 76.5	9,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,400	32.4	29.9 - 35.1	11,200	67.6	64.9 - 70.1	16,500

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	38.1	33.7 - 42.8	1,200	61.9	57.2 - 66.3	1,900
Honolulu	3,900	32.2	29.0 - 35.6	8,200	67.8	64.4 - 71.0	12,100
Kauai	200	24.1	18.4 - 30.9	600	75.9	69.1 - 81.6	800
Maui	600	32.0	27.5 - 36.8	1,200	68.0	63.2 - 72.5	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	22.9	17.8 - 29.0	2,800	77.1	71.0 - 82.2	3,600
Native Hawaiian	1,800	41.2	38.2 - 44.3	2,600	58.8	55.7 - 61.8	4,500
Chinese	100	10.5	3.9 - 25.5	600	89.5	74.5 - 96.1	700
Filipino	1,400	39.8	33.6 - 46.3	2,200	60.2	53.7 - 66.4	3,600
Japanese	400	20.4	13.9 - 29.1	1,500	79.6	70.9 - 86.1	1,900
Other	800	35.5	27.5 - 44.5	1,400	64.5	55.5 - 72.5	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	58.5	50.2 - 66.3	600	41.5	33.7 - 49.8	1,500
20-24	1,600	42.5	37.2 - 48.0	2,200	57.5	52.0 - 62.8	3,800
25-34	2,300	28.2	24.7 - 31.9	6,000	71.8	68.1 - 75.3	8,300
35+	500	17.7	12.9 - 23.7	2,400	82.3	76.3 - 87.1	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,900	33.1	29.6 - 36.8	5,900	66.9	63.2 - 70.4	8,800
Female	2,500	31.7	28.1 - 35.5	5,300	68.3	64.5 - 71.9	7,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	24.4	21.5 - 27.6	8,300	75.6	72.4 - 78.5	10,900
No	2,700	48.0	43.7 - 52.4	2,900	52.0	47.6 - 56.3	5,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,200	43.4	39.5 - 47.5	4,200	56.6	52.5 - 60.5	7,400
131-185%	700	29.7	23.5 - 36.9	1,700	70.3	63.1 - 76.5	2,400
186+%	800	14.7	11.4 - 18.8	4,500	85.3	81.2 - 88.6	5,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,000	49.9	44.5 - 55.4	2,000	50.1	44.6 - 55.5	3,900
\$15,001-\$25,000	700	34.7	28.0 - 42.0	1,400	65.3	58.0 - 72.0	2,100
\$25,001-\$40,000	1,000	32.4	26.6 - 38.8	2,100	67.6	61.2 - 73.4	3,100
>=\$40,001	1,100	18.1	14.8 - 21.9	5,200	81.9	78.1 - 85.2	6,300

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	35.5	26.0 - 46.3	700	64.5	53.7 - 74.0	1,100
Normal birth weight	5,000	32.2	29.6 - 34.9	10,500	67.8	65.1 - 70.4	15,400

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	56.3	47.4 - 64.7	700	43.7	35.3 - 52.6	1,500
No	4,500	30.1	27.5 - 32.9	10,300	69.9	67.1 - 72.5	14,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	50.1	44.2 - 55.9	1,700	49.9	44.1 - 55.8	3,400
No	3,500	27.3	24.6 - 30.3	9,300	72.7	69.7 - 75.4	12,700

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,100	45.1	41.1 - 49.2	3,800	54.9	50.8 - 58.9	6,900
No	2,200	23.3	20.2 - 26.6	7,200	76.7	73.4 - 79.8	9,400

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	4,900	30.4	27.9 - 33.1	11,200	69.6	66.9 - 72.1	16,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	34.8	30.6 - 39.2	1,200	65.2	60.8 - 69.4	1,900
Honolulu	3,500	29.4	26.2 - 32.9	8,400	70.6	67.1 - 73.8	11,900
Kauai	200	33.5	26.3 - 41.4	400	66.5	58.6 - 73.7	600
Maui	500	31.1	26.5 - 36.0	1,200	68.9	64.0 - 73.5	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	26.0	20.4 - 32.5	2,500	74.0	67.5 - 79.6	3,400
Native Hawaiian	1,700	38.7	35.7 - 41.8	2,700	61.3	58.2 - 64.3	4,400
Chinese	200	34.0	20.0 - 51.5	400	66.0	48.5 - 80.0	600
Filipino	1,000	32.8	26.7 - 39.6	2,100	67.2	60.4 - 73.3	3,200
Japanese	200	8.8	5.2 - 14.5	2,100	91.2	85.5 - 94.8	2,300
Other	900	39.3	30.4 - 48.9	1,300	60.7	51.1 - 69.6	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	50.0	41.0 - 58.9	700	50.0	41.1 - 59.0	1,500
20-24	1,500	36.5	31.3 - 42.0	2,500	63.5	58.0 - 68.7	4,000
25-34	2,100	26.4	23.0 - 30.2	5,900	73.6	69.8 - 77.0	8,100
35+	600	22.2	16.9 - 28.7	2,000	77.8	71.3 - 83.1	2,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,600	32.6	29.0 - 36.5	5,400	67.4	63.5 - 71.0	8,100
Female	2,300	28.2	24.7 - 31.8	5,800	71.8	68.2 - 75.3	8,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	24.4	21.4 - 27.7	8,200	75.6	72.3 - 78.6	10,900
No	2,200	43.0	38.5 - 47.5	3,000	57.0	52.5 - 61.5	5,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,500	39.2	35.0 - 43.5	3,800	60.8	56.5 - 65.0	6,300
131-185%	1,000	30.0	24.3 - 36.4	2,300	70.0	63.6 - 75.7	3,300
186+%	800	15.9	12.5 - 20.0	4,300	84.1	80.0 - 87.5	5,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,400	44.0	38.3 - 49.9	1,700	56.0	50.1 - 61.7	3,100
\$15,001-\$25,000	800	34.6	28.0 - 41.9	1,600	65.4	58.1 - 72.0	2,400
\$25,001-\$40,000	1,100	33.0	27.3 - 39.2	2,300	67.0	60.8 - 72.7	3,400
>=\$40,001	1,100	17.3	14.0 - 21.1	5,100	82.7	78.9 - 86.0	6,200

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	35.4	25.0 - 47.4	600	64.6	52.6 - 75.0	1,000
Normal birth weight	4,500	30.1	27.5 - 32.8	10,600	69.9	67.2 - 72.5	15,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	58.0	48.9 - 66.7	500	42.0	33.3 - 51.1	1,200
No	4,100	28.1	25.5 - 30.9	10,500	71.9	69.1 - 74.5	14,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	45.4	39.4 - 51.5	1,700	54.6	48.5 - 60.6	3,100
No	3,300	26.5	23.7 - 29.5	9,300	73.5	70.5 - 76.3	12,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	42.1	37.9 - 46.4	3,900	57.9	53.6 - 62.1	6,700
No	2,000	22.1	19.1 - 25.3	7,200	77.9	74.7 - 80.9	9,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,000	31.8	29.7 - 33.9	10,700	68.2	66.1 - 70.3	15,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	33.4	29.2 - 38.0	1,200	66.6	62.0 - 70.8	1,800
Honolulu	3,700	31.6	29.0 - 34.3	7,900	68.4	65.7 - 71.0	11,600
Kauai	200	26.9	20.3 - 34.7	500	73.1	65.3 - 79.7	700
Maui	500	33.2	28.6 - 38.2	1,100	66.8	61.8 - 71.4	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	22.5	18.5 - 27.2	2,600	77.5	72.8 - 81.5	3,400
Native Hawaiian	1,800	41.8	39.3 - 44.2	2,500	58.2	55.8 - 60.7	4,200
Chinese	200	27.5	16.6 - 41.9	500	72.5	58.1 - 83.4	600
Filipino	1,100	34.1	29.0 - 39.7	2,100	65.9	60.3 - 71.0	3,200
Japanese	400	20.5	15.1 - 27.2	1,600	79.5	72.8 - 84.9	2,000
Other	700	34.7	28.1 - 42.0	1,400	65.3	58.0 - 71.9	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	53.8	47.6 - 59.8	700	46.2	40.2 - 52.4	1,500
20-24	1,700	40.3	36.1 - 44.6	2,500	59.7	55.4 - 63.9	4,100
25-34	1,900	26.0	23.1 - 29.0	5,600	74.0	71.0 - 76.9	7,500
35+	500	21.4	16.7 - 27.0	2,000	78.6	73.0 - 83.3	2,500

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,500	31.1	28.2 - 34.1	5,500	68.9	65.9 - 71.8	8,000
Female	2,500	32.4	29.5 - 35.5	5,100	67.6	64.5 - 70.5	7,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	25.5	23.0 - 28.1	7,800	74.5	71.9 - 77.0	10,400
No	2,300	44.3	40.7 - 48.0	2,900	55.7	52.0 - 59.3	5,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,300	39.5	36.2 - 42.9	3,600	60.5	57.1 - 63.8	5,900
131-185%	1,000	29.0	24.9 - 33.5	2,500	71.0	66.5 - 75.1	3,500
186+%	1,000	19.3	16.2 - 22.9	4,200	80.7	77.1 - 83.8	5,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,300	41.4	36.8 - 46.2	1,800	58.6	53.8 - 63.2	3,200
\$15,001-\$25,000	800	39.7	34.1 - 45.7	1,300	60.3	54.3 - 65.9	2,100
\$25,001-\$40,000	1,100	30.9	26.7 - 35.4	2,400	69.1	64.6 - 73.3	3,400
>=\$40,001	1,300	21.0	18.1 - 24.3	4,800	79.0	75.7 - 81.9	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	39.8	31.4 - 48.9	600	60.2	51.1 - 68.6	1,000
Normal birth weight	4,600	31.2	29.1 - 33.4	10,000	68.8	66.6 - 70.9	14,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	51.2	44.3 - 58.2	700	48.8	41.8 - 55.7	1,400
No	4,100	29.6	27.4 - 31.9	9,900	70.4	68.1 - 72.6	14,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	44.1	39.4 - 48.8	1,800	55.9	51.2 - 60.6	3,100
No	3,400	28.2	25.9 - 30.7	8,600	71.8	69.3 - 74.1	12,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	40.5	37.3 - 43.8	3,900	59.5	56.2 - 62.7	6,600
No	2,200	25.2	22.6 - 28.0	6,600	74.8	72.0 - 77.4	8,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,000	31.7	29.6 - 33.8	10,800	68.3	66.2 - 70.4	15,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	39.0	34.4 - 43.8	1,100	61.0	56.2 - 65.6	1,800
Honolulu	3,600	30.7	28.2 - 33.4	8,100	69.3	66.6 - 71.8	11,700
Kauai	200	25.6	18.9 - 33.6	500	74.4	66.4 - 81.1	600
Maui	500	32.7	27.9 - 37.8	1,100	67.3	62.2 - 72.1	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,100	28.7	24.5 - 33.3	2,700	71.3	66.7 - 75.5	3,800
Native Hawaiian	1,700	41.6	39.0 - 44.2	2,400	58.4	55.8 - 61.0	4,200
Chinese	200	27.0	16.6 - 40.8	500	73.0	59.2 - 83.4	600
Filipino	1,000	31.4	26.5 - 36.8	2,200	68.6	63.2 - 73.5	3,100
Japanese	300	15.5	10.7 - 21.9	1,600	84.5	78.1 - 89.3	1,900
Other	700	34.2	27.4 - 41.8	1,400	65.8	58.2 - 72.6	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	47.3	41.1 - 53.5	900	52.7	46.5 - 58.9	1,800
20-24	1,500	38.7	34.6 - 43.0	2,300	61.3	57.0 - 65.4	3,800
25-34	2,100	27.3	24.4 - 30.5	5,500	72.7	69.5 - 75.6	7,500
35+	600	23.8	19.0 - 29.3	2,100	76.2	70.7 - 81.0	2,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,700	33.1	30.2 - 36.1	5,400	66.9	63.9 - 69.8	8,100
Female	2,300	30.2	27.3 - 33.3	5,300	69.8	66.7 - 72.7	7,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	27.0	24.5 - 29.6	7,900	73.0	70.4 - 75.5	10,800
No	2,100	42.0	38.4 - 45.7	2,900	58.0	54.3 - 61.6	4,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,700	39.7	36.5 - 43.0	4,000	60.3	57.0 - 63.5	6,700
131-185%	600	26.6	21.9 - 31.8	1,600	73.4	68.2 - 78.1	2,200
186+%	1,300	21.9	18.7 - 25.5	4,400	78.1	74.5 - 81.3	5,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,600	44.2	39.8 - 48.7	2,000	55.8	51.3 - 60.2	3,600
\$15,001-\$25,000	900	36.9	31.6 - 42.4	1,600	63.1	57.6 - 68.4	2,500
\$25,001-\$40,000	800	27.0	22.9 - 31.5	2,200	73.0	68.5 - 77.1	3,100
>=\$40,001	1,300	22.0	18.9 - 25.6	4,500	78.0	74.4 - 81.1	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	38.9	30.8 - 47.8	600	61.1	52.2 - 69.2	1,000
Normal birth weight	4,500	30.9	28.8 - 33.1	10,100	69.1	66.9 - 71.2	14,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	59.7	52.4 - 66.6	500	40.3	33.4 - 47.6	1,300
No	4,000	28.4	26.3 - 30.6	10,000	71.6	69.4 - 73.7	14,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	47.3	42.5 - 52.1	1,700	52.7	47.9 - 57.5	3,200
No	3,200	26.7	24.5 - 29.1	8,800	73.3	70.9 - 75.5	12,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	39.3	36.2 - 42.5	4,100	60.7	57.5 - 63.8	6,800
No	2,200	25.3	22.6 - 28.2	6,500	74.7	71.8 - 77.4	8,700