

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (exclusive) duration (grouped)	Breastfeeding (exclusive) duration (grouped)

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/28/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	1,700	12.6	11.1 - 14.4	2,900	21.5	19.5 - 23.6	3,000	22.2	20.3 - 24.3	800	5.5	4.5 - 6.7	5,200	38.1	35.8 - 40.6	13,600
2001	1,600	10.3	9.1 - 11.7	4,600	29.7	27.6 - 31.9	3,000	19.5	17.7 - 21.4	700	4.2	3.4 - 5.3	5,700	36.2	34.0 - 38.5	15,600
2002	1,700	10.9	9.2 - 12.8	5,000	31.3	28.7 - 34.1	2,800	17.7	15.7 - 20.0	700	4.2	3.2 - 5.5	5,800	35.9	33.2 - 38.7	16,000
2003	1,800	10.9	9.3 - 12.7	5,100	31.4	28.8 - 34.1	3,200	19.4	17.3 - 21.7	600	3.4	2.5 - 4.6	5,700	34.9	32.3 - 37.5	16,400
2004	1,600	9.1	7.9 - 10.5	5,700	33.6	31.6 - 35.7	3,300	19.2	17.5 - 21.0	600	3.7	3.0 - 4.6	5,900	34.4	32.4 - 36.5	17,100
2005	1,700	10.1	8.7 - 11.7	5,500	32.4	30.3 - 34.7	3,300	19.3	17.5 - 21.3	500	3.1	2.3 - 4.0	5,900	35.1	32.9 - 37.4	16,900
2006	1,400	8.2	6.9 - 9.7	5,600	32.0	29.7 - 34.3	3,400	19.4	17.4 - 21.4	800	4.4	3.5 - 5.5	6,300	36.1	33.7 - 38.5	17,400
2007	1,400	7.7	6.4 - 9.1	6,300	35.6	33.3 - 37.9	3,100	17.5	15.7 - 19.4	500	2.9	2.2 - 3.9	6,400	36.3	34.0 - 38.6	17,800
2008	1,400	7.8	6.6 - 9.3	6,500	36.2	34.0 - 38.5	3,100	17.0	15.2 - 18.8	600	3.1	2.4 - 4.1	6,500	35.9	33.6 - 38.2	18,000
2009	1,200	6.8	5.3 - 8.7	6,100	34.6	31.6 - 37.7	3,000	17.1	14.8 - 19.7	900	5.2	4.0 - 6.8	6,400	36.3	33.3 - 39.4	17,700
2010	1,100	6.1	4.6 - 8.0	4,900	27.0	24.2 - 29.9	3,200	18.0	15.6 - 20.7	800	4.6	3.4 - 6.2	8,000	44.3	41.1 - 47.5	18,000
2011	800	4.5	3.3 - 6.2	5,500	30.6	27.7 - 33.6	3,100	17.0	14.6 - 19.6	700	3.9	2.8 - 5.4	7,900	44.0	40.8 - 47.3	18,000
2012	900	5.2	3.9 - 7.1	6,000	33.5	30.4 - 36.7	2,800	16.0	13.7 - 18.5	600	3.2	2.1 - 4.7	7,500	42.2	38.9 - 45.5	17,800
2013	700	3.9	2.8 - 5.5	4,700	27.2	24.1 - 30.4	2,900	16.7	14.1 - 19.5	400	2.5	1.6 - 3.9	8,600	49.8	46.2 - 53.4	17,300

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	700	3.9	2.8 - 5.5	4,700	27.2	24.1 - 30.4	2,900	16.7	14.1 - 19.5	400	2.5	1.6 - 3.9	8,600	49.8	46.2 - 53.4	17,300

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.4	4.0 - 10.2	700	30.3	25.2 - 35.8	300	13.3	9.8 - 17.7	<50	1.6	0.6 - 3.9	1,100	48.5	42.7 - 54.2	2,200
Honolulu	400	3.5	2.1 - 5.7	3,300	26.1	22.2 - 30.5	2,200	17.6	14.2 - 21.5	300	2.6	1.5 - 4.7	6,300	50.2	45.4 - 55.0	12,500
Kauai	<50	4.6	2.8 - 7.4	200	18.6	14.8 - 23.2	100	18.5	14.6 - 23.1	<50	2.1	1.0 - 4.3	500	56.2	50.8 - 61.6	800
Maui	100	3.1	1.7 - 5.9	600	34.2	29.0 - 39.8	300	13.8	10.3 - 18.2	100	2.8	1.4 - 5.4	900	46.1	40.5 - 51.9	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	1.7	0.6 - 4.8	900	19.1	14.1 - 25.4	700	14.3	10.0 - 20.0	100	2.2	0.8 - 5.4	2,900	62.8	55.7 - 69.4	4,600
Native Hawaiian	300	6.5	4.0 - 10.3	1,300	24.3	19.7 - 29.6	1,200	22.5	17.5 - 28.5	100	1.6	0.7 - 4.0	2,300	45.1	39.0 - 51.3	5,200
Chinese	<50	3.7	0.5 - 21.4	400	48.2	31.7 - 65.1	200	18.9	8.8 - 36.2	<50	0.9	0.3 - 2.8	200	28.4	15.5 - 46.0	800
Filipino	100	2.0	0.7 - 6.2	1,100	40.7	32.2 - 49.7	400	13.6	8.8 - 20.4	100	4.4	1.7 - 11.0	1,100	39.3	30.7 - 48.6	2,700
Japanese	<50	1.7	0.2 - 10.8	400	23.7	15.5 - 34.5	300	16.2	9.2 - 26.9	<50	2.1	0.4 - 9.8	1,000	56.2	44.5 - 67.3	1,800
Other	100	6.5	3.2 - 12.8	600	29.1	20.6 - 39.4	300	11.3	6.3 - 19.5	100	3.6	1.2 - 10.6	1,100	49.4	39.2 - 59.7	2,200

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	1.5	0.2 - 8.8	300	41.9	27.3 - 58.1	100	9.9	3.8 - 23.7	<50	1.9	0.4 - 8.0	300	44.8	29.8 - 60.7	800
20-24	100	2.1	0.8 - 5.3	800	19.6	14.7 - 25.6	600	15.5	10.5 - 22.1	200	3.9	1.8 - 8.4	2,300	59.0	51.3 - 66.2	3,900
25-34	400	4.0	2.5 - 6.4	2,600	27.0	22.9 - 31.6	1,700	17.7	14.2 - 21.8	200	2.0	1.0 - 3.9	4,700	49.3	44.4 - 54.2	9,500
35+	200	6.3	3.4 - 11.4	1,100	33.6	26.5 - 41.5	500	16.6	11.5 - 23.4	100	2.3	0.8 - 7.0	1,300	41.2	33.4 - 49.3	3,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	4.4	2.8 - 6.8	2,100	25.3	21.3 - 29.8	1,200	13.9	10.8 - 17.8	200	2.1	1.1 - 4.1	4,600	54.2	49.2 - 59.2	8,500
Female	300	3.4	2.0 - 5.7	2,600	28.9	24.5 - 33.7	1,700	19.3	15.5 - 23.7	300	2.8	1.5 - 5.3	4,000	45.6	40.5 - 50.7	8,900

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.7	1.6 - 4.5	2,700	24.6	20.9 - 28.7	2,000	17.7	14.4 - 21.6	300	2.3	1.2 - 4.2	5,800	52.8	48.1 - 57.4	11,000
No	400	6.0	3.8 - 9.4	2,000	31.7	26.8 - 37.2	900	14.9	11.4 - 19.2	200	2.8	1.4 - 5.6	2,800	44.6	39.0 - 50.3	6,300

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.1	3.3 - 7.9	1,800	28.6	23.7 - 34.0	900	14.3	10.7 - 18.8	200	2.7	1.2 - 5.8	3,100	49.3	43.5 - 55.1	6,300
131-185%	200	6.9	3.3 - 13.9	800	32.5	24.1 - 42.3	400	15.4	9.7 - 23.5	100	3.7	1.2 - 10.6	1,000	41.4	32.2 - 51.3	2,400
186+%	200	2.5	1.2 - 5.4	1,700	24.7	20.2 - 29.9	1,300	18.4	14.2 - 23.4	100	1.9	0.8 - 4.1	3,600	52.5	46.8 - 58.2	6,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Income (2012+)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	200	5.1	2.8 - 9.1	1,100	29.4	23.3 - 36.3	600	15.6	11.0 - 21.7	100	1.9	0.7 - 5.1	1,900	48.0	40.8 - 55.3	3,900
\$17,001-\$26,000	100	4.5	2.2 - 9.0	600	28.0	20.0 - 37.6	300	13.2	8.0 - 21.1	100	3.6	1.1 - 11.0	1,200	50.7	40.9 - 60.5	2,300
\$26,001-\$51,000	200	5.0	2.5 - 9.8	1,100	30.2	23.5 - 37.8	500	14.6	9.8 - 21.0	100	3.4	1.4 - 8.1	1,700	46.9	39.2 - 54.7	3,700
>=\$51,001	200	2.9	1.4 - 6.1	1,400	23.9	19.1 - 29.4	1,100	18.9	14.5 - 24.4	100	2.1	0.9 - 4.8	3,100	52.2	45.9 - 58.3	5,900

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	3.6	2.0 - 6.3	500	41.6	36.4 - 46.9	200	20.1	16.3 - 24.5	<50	2.4	1.3 - 4.3	400	32.4	27.6 - 37.6	1,200
Normal birth weight	600	3.9	2.7 - 5.6	4,200	26.1	22.9 - 29.6	2,700	16.4	13.7 - 19.5	400	2.5	1.5 - 4.0	8,300	51.1	47.2 - 54.9	16,200

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.2	3.4 - 18.5	300	28.4	17.1 - 43.2	100	13.6	6.7 - 25.8	<50	1.2	0.3 - 4.5	400	48.6	34.7 - 62.8	900
No	600	3.7	2.6 - 5.4	4,400	27.2	24.0 - 30.6	2,700	16.9	14.2 - 19.9	400	2.6	1.6 - 4.1	8,000	49.7	45.9 - 53.4	16,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.1	2.7 - 9.6	800	26.2	19.5 - 34.3	400	14.7	9.7 - 21.5	100	3.0	1.0 - 8.6	1,500	51.0	42.5 - 59.5	3,000
No	500	3.7	2.5 - 5.5	3,800	27.5	24.1 - 31.1	2,400	17.1	14.3 - 20.4	300	2.4	1.4 - 4.0	6,900	49.3	45.3 - 53.3	13,900

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.0	3.2 - 7.7	2,400	32.7	27.9 - 37.9	1,200	15.7	12.2 - 20.1	200	2.9	1.5 - 5.6	3,300	43.7	38.4 - 49.1	7,500
No	300	3.1	1.8 - 5.2	2,200	23.0	19.3 - 27.1	1,700	17.4	14.0 - 21.4	200	2.2	1.1 - 4.0	5,300	54.4	49.6 - 59.2	9,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	900	5.2	3.9 - 7.1	6,000	33.5	30.4 - 36.7	2,800	16.0	13.7 - 18.5	600	3.2	2.1 - 4.7	7,500	42.2	38.9 - 45.5	17,800

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.1	2.4 - 6.9	700	31.9	27.0 - 37.1	400	16.7	13.0 - 21.2	100	4.4	2.6 - 7.3	900	42.9	37.7 - 48.4	2,200
Honolulu	800	6.0	4.2 - 8.6	4,300	33.5	29.5 - 37.8	2,000	15.9	12.9 - 19.4	400	3.2	1.9 - 5.3	5,300	41.4	37.2 - 45.8	12,900
Kauai	<50	3.5	1.8 - 7.0	200	30.0	24.3 - 36.3	100	12.2	8.6 - 17.2	<50	2.7	1.2 - 5.7	400	51.6	45.0 - 58.1	800
Maui	<50	1.8	0.8 - 3.6	700	36.7	31.9 - 41.7	300	17.2	13.6 - 21.5	<50	2.1	1.0 - 4.0	800	42.3	37.4 - 47.4	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.3	2.8 - 9.7	800	19.6	14.9 - 25.4	700	16.8	12.3 - 22.5	100	1.7	0.5 - 5.3	2,300	56.7	49.9 - 63.2	4,100
Native Hawaiian	300	5.4	3.1 - 9.5	1,600	31.7	26.4 - 37.5	1,000	20.9	16.3 - 26.4	200	4.8	2.7 - 8.4	1,900	37.1	31.5 - 43.1	5,000
Chinese	100	8.5	3.1 - 21.2	400	39.9	27.2 - 54.2	100	6.3	2.7 - 14.2	<50	3.0	0.4 - 17.6	500	42.4	29.5 - 56.4	1,100
Filipino	100	3.0	1.2 - 7.1	1,300	41.1	33.5 - 49.0	600	19.9	14.2 - 27.0	200	4.9	2.3 - 10.4	1,000	31.2	24.3 - 39.1	3,100
Japanese	<50	1.7	0.2 - 10.8	700	48.5	37.2 - 59.9	100	9.7	4.7 - 19.0	<50	0.3	0.0 - 1.5	600	39.9	29.3 - 51.4	1,500
Other	200	7.9	4.2 - 14.3	1,100	37.6	29.6 - 46.2	300	9.1	5.3 - 15.1	100	2.2	0.8 - 6.0	1,300	43.3	35.0 - 51.9	3,000

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	3.2	0.5 - 18.9	500	50.0	36.2 - 63.8	200	18.0	9.1 - 32.5	<50	2.7	0.9 - 7.7	300	26.0	15.9 - 39.6	1,000
20-24	200	4.9	2.5 - 9.4	1,100	29.3	23.2 - 36.3	700	17.7	12.8 - 24.1	200	5.5	2.8 - 10.5	1,600	42.6	35.4 - 50.0	3,700
25-34	500	5.2	3.4 - 7.9	3,100	30.9	26.9 - 35.1	1,700	16.6	13.5 - 20.1	300	3.2	1.8 - 5.4	4,400	44.2	39.8 - 48.6	10,000
35+	200	6.3	3.3 - 11.8	1,300	41.3	33.9 - 49.2	400	11.2	7.5 - 16.6	<50	0.6	0.2 - 1.6	1,300	40.5	33.2 - 48.4	3,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	4.3	2.6 - 7.1	3,000	34.3	29.9 - 38.9	1,400	16.4	13.2 - 20.2	300	3.1	1.7 - 5.4	3,700	41.9	37.3 - 46.7	8,800
Female	600	6.1	4.2 - 9.0	2,900	32.7	28.5 - 37.2	1,400	15.5	12.4 - 19.2	300	3.3	1.9 - 5.5	3,800	42.4	37.9 - 47.0	9,000

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.4	2.8 - 6.7	3,500	32.0	28.1 - 36.1	1,800	16.7	13.8 - 20.2	400	3.3	2.0 - 5.4	4,800	43.7	39.5 - 48.0	11,000
No	500	6.7	4.3 - 10.2	2,400	35.9	31.0 - 41.1	1,000	14.7	11.4 - 18.8	200	3.0	1.7 - 5.4	2,700	39.7	34.7 - 45.0	6,800

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.6	4.2 - 10.3	2,300	37.1	31.9 - 42.6	900	14.7	11.3 - 18.9	200	2.8	1.4 - 5.4	2,400	38.8	33.6 - 44.3	6,300
131-185%	200	6.3	3.1 - 12.2	900	30.7	23.8 - 38.6	700	22.8	16.5 - 30.7	200	6.8	3.5 - 12.7	1,000	33.4	26.2 - 41.5	3,000
186+%	100	1.9	0.8 - 4.3	2,200	33.0	28.1 - 38.2	900	13.4	10.1 - 17.4	100	1.5	0.6 - 3.7	3,400	50.3	44.9 - 55.6	6,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Income (2012+)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	200	5.8	3.2 - 10.3	1,600	40.7	33.9 - 47.8	600	14.1	10.0 - 19.4	<50	1.0	0.3 - 3.6	1,500	38.4	31.7 - 45.5	4,000
\$17,001-\$26,000	100	3.0	1.0 - 8.7	700	31.9	23.9 - 41.1	400	16.5	10.7 - 24.8	100	6.1	3.0 - 12.2	900	42.4	33.7 - 51.6	2,200
\$26,001-\$51,000	400	8.7	5.3 - 14.1	1,300	29.3	23.7 - 35.7	900	19.5	14.6 - 25.6	200	5.0	2.6 - 9.4	1,600	37.4	31.1 - 44.1	4,400
>=\$51,001	100	1.7	0.7 - 4.4	1,900	33.8	28.5 - 39.6	800	14.2	10.6 - 18.7	100	1.6	0.6 - 4.2	2,800	48.7	42.9 - 54.5	5,700

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	3.9	2.3 - 6.8	500	43.9	38.4 - 49.6	200	18.5	14.3 - 23.7	<50	1.3	0.5 - 3.7	400	32.3	27.2 - 37.7	1,200
Normal birth weight	900	5.3	3.9 - 7.3	5,400	32.8	29.5 - 36.2	2,600	15.6	13.2 - 18.4	500	3.3	2.2 - 4.9	7,100	42.9	39.5 - 46.4	16,600

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	4.6	2.1 - 9.9	200	28.4	16.6 - 44.2	300	35.9	21.7 - 53.2	0	0.0	-	200	31.1	18.9 - 46.6	700
No	900	5.3	3.8 - 7.2	5,600	33.8	30.6 - 37.1	2,500	15.2	12.9 - 17.9	600	3.4	2.3 - 5.0	7,000	42.4	39.0 - 45.8	16,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.8	2.8 - 11.5	700	26.9	20.4 - 34.6	600	20.8	14.6 - 28.6	100	3.3	1.3 - 8.1	1,200	43.2	35.2 - 51.5	2,700
No	700	5.1	3.6 - 7.2	5,000	34.8	31.4 - 38.4	2,200	15.2	12.8 - 18.0	500	3.2	2.1 - 5.0	6,000	41.6	38.0 - 45.3	14,400

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	7.2	5.0 - 10.4	3,000	36.7	32.1 - 41.6	1,300	15.9	12.7 - 19.8	300	3.6	2.2 - 5.9	2,900	36.5	31.9 - 41.3	8,100
No	300	3.3	1.9 - 5.7	3,000	31.0	27.0 - 35.3	1,500	15.8	12.7 - 19.4	300	2.8	1.6 - 5.2	4,500	47.1	42.6 - 51.7	9,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	800	4.5	3.3 - 6.2	5,500	30.6	27.7 - 33.6	3,100	17.0	14.6 - 19.6	700	3.9	2.8 - 5.4	7,900	44.0	40.8 - 47.3	18,000

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.4	2.7 - 7.0	800	32.3	27.7 - 37.3	400	16.4	12.8 - 20.6	100	3.6	2.1 - 6.2	1,000	43.4	38.3 - 48.5	2,300
Honolulu	600	4.7	3.1 - 7.1	4,000	30.6	26.8 - 34.8	2,300	17.7	14.6 - 21.4	500	4.0	2.6 - 6.1	5,600	42.9	38.7 - 47.3	13,000
Kauai	<50	2.1	0.9 - 5.0	200	25.7	20.5 - 31.6	100	17.4	13.1 - 22.8	<50	3.3	1.6 - 6.5	400	51.5	45.1 - 57.9	800
Maui	100	4.3	2.7 - 6.7	600	30.1	26.0 - 34.6	200	12.4	9.6 - 15.8	100	4.1	2.6 - 6.4	900	49.1	44.5 - 53.8	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.5	1.1 - 5.7	900	20.0	15.4 - 25.7	800	18.5	13.7 - 24.5	200	3.7	1.8 - 7.3	2,500	55.2	48.7 - 61.6	4,500
Native Hawaiian	300	5.3	3.1 - 8.9	1,600	31.7	26.6 - 37.3	1,000	20.1	15.7 - 25.3	300	5.6	3.2 - 9.4	1,900	37.4	32.0 - 43.2	5,200
Chinese	100	9.2	2.8 - 26.0	300	35.6	21.3 - 53.1	100	10.1	3.9 - 24.1	<50	1.6	0.6 - 4.4	300	43.4	28.0 - 60.3	700
Filipino	200	5.3	2.8 - 9.8	1,200	39.1	31.9 - 46.9	600	18.0	13.0 - 24.5	100	4.6	2.5 - 8.3	1,000	33.0	26.2 - 40.5	3,200
Japanese	<50	2.0	0.5 - 7.4	800	41.7	31.8 - 52.3	300	13.9	7.6 - 24.1	<50	1.7	0.4 - 7.7	700	40.7	31.0 - 51.1	1,800
Other	200	5.9	2.6 - 12.9	700	26.8	19.7 - 35.3	300	11.1	6.4 - 18.5	100	2.6	0.9 - 7.3	1,400	53.7	44.5 - 62.5	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.9	1.9 - 21.9	300	34.2	23.5 - 46.8	200	22.1	13.3 - 34.6	<50	2.4	0.8 - 7.4	300	34.3	23.3 - 47.3	1,000
20-24	300	7.0	4.1 - 11.7	1,300	30.7	24.7 - 37.3	900	21.8	16.5 - 28.3	200	5.1	2.7 - 9.5	1,500	35.4	29.2 - 42.2	4,300
25-34	400	4.0	2.5 - 6.3	2,800	28.9	25.0 - 33.1	1,500	15.4	12.4 - 19.0	400	3.9	2.5 - 5.9	4,600	47.8	43.4 - 52.3	9,700
35+	100	1.8	1.0 - 3.3	1,100	34.6	27.6 - 42.3	400	13.4	8.8 - 19.8	100	2.9	1.1 - 7.0	1,400	47.3	39.7 - 55.1	3,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	4.5	2.9 - 7.0	2,700	29.0	25.0 - 33.2	1,700	17.9	14.5 - 21.9	300	3.1	1.9 - 4.9	4,200	45.5	41.0 - 50.1	9,200
Female	400	4.5	2.9 - 7.0	2,800	32.2	28.0 - 36.8	1,400	16.0	12.8 - 19.7	400	4.8	3.1 - 7.4	3,700	42.5	38.0 - 47.1	8,800

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.1	2.6 - 6.2	3,200	28.4	24.9 - 32.3	1,800	15.8	13.0 - 19.1	400	3.5	2.2 - 5.4	5,500	48.2	44.0 - 52.4	11,300
No	400	5.3	3.3 - 8.3	2,300	34.1	29.4 - 39.3	1,300	18.9	14.9 - 23.6	300	4.6	2.9 - 7.3	2,500	37.1	32.2 - 42.2	6,700

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.7	4.3 - 10.4	2,000	31.9	27.2 - 37.1	1,300	20.0	15.9 - 24.7	200	3.1	1.8 - 5.4	2,400	38.3	33.2 - 43.6	6,300
131-185%	100	4.7	2.1 - 10.5	700	25.6	18.8 - 33.7	500	17.2	11.4 - 25.0	100	4.5	2.2 - 9.0	1,300	48.0	39.5 - 56.6	2,700
186+%	200	3.0	1.7 - 5.2	2,500	31.5	27.1 - 36.3	1,200	14.6	11.4 - 18.7	300	4.2	2.5 - 7.0	3,700	46.7	41.8 - 51.6	8,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	6.2	3.4 - 11.1	1,000	33.2	26.3 - 40.8	700	23.6	17.7 - 30.8	100	2.5	1.0 - 6.1	1,100	34.4	27.6 - 42.0	3,100
\$10,000-\$24,999	200	6.0	3.0 - 11.6	900	28.0	22.0 - 34.8	700	21.0	15.3 - 28.0	100	4.0	2.1 - 7.6	1,400	41.1	34.2 - 48.4	3,400
\$25,000-\$49,999	300	5.6	3.1 - 9.9	1,300	28.1	22.5 - 34.4	700	16.5	12.1 - 22.2	100	3.2	1.7 - 6.3	2,100	46.5	40.0 - 53.2	4,500
>=\$50,000	200	2.5	1.2 - 5.1	2,000	32.8	27.7 - 38.4	700	11.9	8.7 - 16.3	300	4.9	2.8 - 8.4	2,900	47.8	42.2 - 53.5	6,100

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	4.1	2.2 - 7.4	600	45.8	40.2 - 51.4	200	14.2	10.6 - 18.7	<50	3.7	2.0 - 6.7	400	32.3	27.3 - 37.7	1,200
Normal birth weight	800	4.5	3.2 - 6.3	5,000	29.5	26.4 - 32.8	2,900	17.2	14.6 - 20.0	700	3.9	2.8 - 5.5	7,600	44.9	41.5 - 48.4	16,800

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	9.1	3.4 - 22.3	300	37.4	25.6 - 51.0	200	17.7	10.1 - 29.3	100	13.9	6.4 - 27.6	200	21.8	13.8 - 32.7	900
No	700	4.2	3.0 - 5.9	5,200	30.4	27.4 - 33.6	2,800	16.6	14.2 - 19.3	600	3.4	2.4 - 4.9	7,700	45.3	42.0 - 48.7	17,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	6.6	3.7 - 11.5	1,100	32.7	26.3 - 39.9	600	17.5	12.6 - 23.8	200	6.8	3.8 - 12.0	1,300	36.3	29.6 - 43.6	3,500
No	600	4.0	2.7 - 5.8	4,400	30.3	27.0 - 33.7	2,400	16.5	13.8 - 19.5	500	3.3	2.2 - 4.8	6,600	46.0	42.4 - 49.7	14,400

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	6.0	4.0 - 9.0	2,600	31.7	27.4 - 36.3	1,500	18.9	15.4 - 23.1	300	3.8	2.4 - 5.9	3,200	39.6	35.0 - 44.4	8,100
No	300	3.3	2.0 - 5.3	2,900	29.9	26.0 - 34.1	1,500	15.3	12.3 - 19.0	400	3.8	2.4 - 6.0	4,700	47.7	43.3 - 52.2	9,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,100	6.1	4.6 - 8.0	4,900	27.0	24.2 - 29.9	3,200	18.0	15.6 - 20.7	800	4.6	3.4 - 6.2	8,000	44.3	41.1 - 47.5	18,000

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.0	3.2 - 7.9	600	26.8	22.5 - 31.6	400	17.2	13.6 - 21.4	100	5.2	3.3 - 8.0	1,000	45.8	40.8 - 51.0	2,300
Honolulu	900	6.8	4.9 - 9.4	3,500	27.0	23.4 - 30.9	2,400	18.2	15.1 - 21.8	600	4.8	3.3 - 7.0	5,600	43.2	39.0 - 47.5	13,000
Kauai	<50	4.1	2.3 - 7.3	200	23.6	18.9 - 29.1	100	15.7	11.8 - 20.6	<50	4.5	2.6 - 7.8	400	52.0	46.0 - 58.0	800
Maui	100	3.3	1.9 - 5.6	600	28.7	24.5 - 33.3	400	18.3	14.8 - 22.3	100	2.7	1.5 - 4.8	900	47.0	42.2 - 51.9	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.4	3.6 - 10.9	600	13.9	10.1 - 18.8	800	18.8	14.1 - 24.8	200	5.4	3.0 - 9.4	2,500	55.6	48.9 - 62.0	4,400
Native Hawaiian	300	6.4	4.0 - 10.1	1,600	29.2	24.2 - 34.7	900	16.0	12.3 - 20.5	300	6.3	4.1 - 9.6	2,300	42.1	36.6 - 47.8	5,400
Chinese	0	0.0	-	200	21.9	11.8 - 37.0	100	12.7	5.0 - 28.6	100	7.8	2.2 - 24.1	400	57.6	41.1 - 72.6	700
Filipino	200	7.0	3.9 - 12.2	1,200	38.4	31.4 - 46.0	700	23.1	17.1 - 30.3	100	3.9	1.6 - 8.8	900	27.7	21.4 - 35.0	3,200
Japanese	<50	0.2	0.0 - 1.3	700	39.9	30.0 - 50.8	200	14.6	8.7 - 23.4	<50	0.9	0.3 - 2.6	700	44.4	34.0 - 55.2	1,700
Other	200	9.4	5.0 - 16.9	600	24.2	17.4 - 32.5	500	18.0	11.8 - 26.7	100	2.4	0.8 - 7.1	1,200	46.0	37.3 - 55.1	2,600

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.2	1.4 - 17.4	300	26.1	16.8 - 38.2	300	27.7	17.6 - 40.7	<50	4.0	1.2 - 12.2	400	37.0	26.5 - 48.9	1,100
20-24	300	7.8	4.8 - 12.6	1,100	24.9	19.7 - 31.0	800	19.0	14.1 - 25.1	200	3.9	2.0 - 7.5	1,900	44.3	37.8 - 51.1	4,300
25-34	500	5.3	3.5 - 7.8	2,500	26.6	22.8 - 30.7	1,600	17.2	14.0 - 21.0	500	5.2	3.5 - 7.7	4,200	45.7	41.3 - 50.2	9,300
35+	200	6.5	3.3 - 12.2	1,000	31.2	24.6 - 38.6	500	15.3	10.7 - 21.5	100	4.2	2.1 - 7.9	1,400	42.9	35.6 - 50.5	3,300

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	600	6.1	4.1 - 8.9	2,600	28.2	24.3 - 32.4	1,500	16.3	13.2 - 20.0	500	5.8	4.0 - 8.5	4,000	43.6	39.2 - 48.1	9,100
Female	500	6.1	4.1 - 8.9	2,300	25.8	22.0 - 30.0	1,800	19.6	16.1 - 23.7	300	3.4	2.1 - 5.4	4,000	45.1	40.5 - 49.7	8,900

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.4	2.8 - 6.6	3,000	26.8	23.3 - 30.6	2,000	17.5	14.5 - 21.0	500	4.1	2.7 - 6.0	5,300	47.3	43.1 - 51.5	11,200
No	600	9.0	6.3 - 12.7	1,800	27.3	23.1 - 32.1	1,300	18.7	15.0 - 23.1	400	5.6	3.6 - 8.6	2,700	39.4	34.6 - 44.4	6,800

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.8	4.5 - 10.2	1,800	28.6	24.1 - 33.7	1,300	20.3	16.2 - 25.2	200	3.3	1.9 - 5.6	2,600	40.9	35.8 - 46.3	6,200
131-185%	100	4.8	2.2 - 10.1	900	31.0	23.8 - 39.3	400	15.3	10.4 - 22.0	200	6.3	3.4 - 11.5	1,200	42.5	34.7 - 50.8	2,900
186+%	300	4.3	2.5 - 7.2	1,900	25.2	21.2 - 29.7	1,300	17.0	13.5 - 21.3	400	5.2	3.3 - 8.1	3,700	48.3	43.3 - 53.3	7,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	5.1	2.9 - 8.6	900	28.1	22.1 - 35.1	800	23.7	17.8 - 30.9	100	3.6	1.6 - 7.6	1,300	39.5	32.7 - 46.7	3,300
\$10,000-\$24,999	300	9.6	5.6 - 16.0	900	29.2	22.8 - 36.4	500	16.7	11.8 - 23.0	100	3.5	1.6 - 7.3	1,300	41.1	33.8 - 48.8	3,200
\$25,000-\$49,999	200	5.5	3.1 - 9.5	1,200	26.4	20.9 - 32.6	800	17.4	12.9 - 23.0	200	5.5	3.2 - 9.3	2,000	45.3	38.9 - 51.9	4,400
>=\$50,000	200	3.6	1.8 - 6.9	1,600	27.1	22.4 - 32.3	1,000	16.1	12.2 - 20.8	300	5.3	3.2 - 8.6	2,900	48.0	42.4 - 53.7	6,100

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.5	5.0 - 11.1	500	37.7	32.5 - 43.1	200	15.0	11.3 - 19.4	100	5.0	3.1 - 8.1	400	34.9	29.9 - 40.3	1,200
Normal birth weight	1,000	6.0	4.4 - 8.0	4,400	26.2	23.3 - 29.4	3,100	18.2	15.6 - 21.1	800	4.6	3.3 - 6.3	7,500	45.0	41.6 - 48.4	16,800

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	19.3	11.5 - 30.6	400	27.5	18.0 - 39.6	200	12.4	6.9 - 21.3	<50	1.7	0.6 - 4.9	500	39.0	28.1 - 51.2	1,400
No	800	4.9	3.5 - 6.7	4,500	27.2	24.3 - 30.2	3,100	18.6	16.1 - 21.5	800	4.9	3.6 - 6.7	7,300	44.4	41.1 - 47.8	16,400

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	8.8	5.5 - 13.8	1,200	31.0	24.9 - 37.8	700	19.8	14.6 - 26.3	100	3.4	1.8 - 6.3	1,400	37.0	30.6 - 43.9	3,800
No	700	5.3	3.7 - 7.4	3,700	26.2	23.1 - 29.5	2,500	17.7	15.0 - 20.7	700	5.0	3.6 - 7.0	6,400	45.9	42.2 - 49.5	14,000

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	7.4	5.2 - 10.6	2,300	29.8	25.6 - 34.4	1,300	16.5	13.2 - 20.4	400	4.9	3.2 - 7.5	3,200	41.3	36.7 - 46.1	7,800
No	500	4.9	3.1 - 7.5	2,500	25.2	21.7 - 29.2	1,900	19.4	16.0 - 23.2	400	4.4	2.9 - 6.7	4,600	46.1	41.7 - 50.5	10,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,200	6.8	5.3 - 8.7	6,100	34.6	31.6 - 37.7	3,000	17.1	14.8 - 19.7	900	5.2	4.0 - 6.8	6,400	36.3	33.3 - 39.4	17,700

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	7.6	5.3 - 11.0	700	31.6	27.0 - 36.6	400	18.1	14.4 - 22.5	100	6.2	4.1 - 9.1	800	36.5	31.7 - 41.6	2,300
Honolulu	900	7.4	5.4 - 10.1	4,500	35.7	31.7 - 40.0	2,100	16.7	13.7 - 20.2	600	4.9	3.3 - 7.1	4,400	35.3	31.3 - 39.5	12,600
Kauai	<50	4.8	2.7 - 8.4	200	21.9	17.2 - 27.5	200	20.8	16.2 - 26.3	100	8.4	5.5 - 12.6	400	44.0	38.1 - 50.2	800
Maui	100	2.6	1.4 - 4.7	700	35.8	31.4 - 40.5	300	17.1	13.8 - 21.0	100	4.9	3.2 - 7.4	800	39.6	35.1 - 44.2	2,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.8	1.3 - 6.1	1,000	23.9	18.8 - 29.8	600	14.4	10.4 - 19.5	300	8.0	5.0 - 12.5	2,100	51.0	44.6 - 57.3	4,100
Native Hawaiian	700	11.6	8.3 - 16.0	1,900	32.8	27.8 - 38.3	1,200	20.7	16.4 - 25.7	300	5.2	3.2 - 8.1	1,700	29.7	24.8 - 35.1	5,700
Chinese	100	11.8	4.2 - 28.8	300	42.2	27.2 - 58.7	100	11.8	4.6 - 27.2	<50	1.6	0.6 - 4.7	300	32.6	19.4 - 49.4	800
Filipino	200	4.8	2.5 - 8.8	1,400	44.3	37.1 - 51.8	600	18.2	13.2 - 24.6	100	2.1	1.2 - 3.6	1,000	30.6	24.0 - 38.1	3,200
Japanese	<50	0.9	0.2 - 3.3	800	48.8	38.4 - 59.4	200	11.2	6.1 - 19.6	<50	2.4	0.7 - 7.9	600	36.7	27.2 - 47.3	1,600
Other	200	7.1	3.5 - 13.6	800	31.9	24.0 - 41.1	400	17.6	11.7 - 25.6	200	7.7	3.8 - 14.8	800	35.7	27.7 - 44.7	2,400

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.4	2.9 - 13.5	600	35.3	26.2 - 45.6	400	22.2	14.6 - 32.3	100	3.0	0.9 - 9.1	600	33.1	23.8 - 43.9	1,800
20-24	300	7.3	4.4 - 11.8	1,300	32.9	26.9 - 39.5	900	22.6	17.4 - 28.9	300	6.5	3.8 - 10.7	1,200	30.8	25.0 - 37.2	4,000
25-34	700	7.3	5.1 - 10.3	2,800	30.8	26.8 - 35.1	1,400	15.9	12.9 - 19.4	500	5.4	3.8 - 7.7	3,600	40.6	36.3 - 45.1	9,000
35+	100	4.8	2.5 - 9.0	1,400	47.9	40.3 - 55.6	300	10.3	6.5 - 15.9	100	4.1	1.9 - 8.6	1,000	32.8	26.1 - 40.3	3,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	7.4	5.3 - 10.3	3,200	33.9	29.8 - 38.3	1,800	19.2	15.9 - 22.9	400	4.6	3.1 - 6.8	3,300	34.9	30.7 - 39.2	9,400
Female	500	6.1	4.1 - 8.8	2,900	35.3	31.0 - 39.8	1,200	14.8	11.8 - 18.4	500	5.8	4.0 - 8.4	3,200	38.0	33.7 - 42.5	8,300

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.3	2.9 - 6.4	3,300	32.0	28.2 - 36.0	1,500	14.2	11.5 - 17.3	600	6.0	4.3 - 8.3	4,600	43.6	39.5 - 47.7	10,500
No	800	10.4	7.6 - 14.1	2,800	38.3	33.6 - 43.4	1,500	21.4	17.5 - 25.8	300	4.0	2.5 - 6.4	1,900	25.9	21.8 - 30.5	7,300

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	600	10.2	7.3 - 14.1	2,400	39.1	33.8 - 44.6	1,000	15.3	11.9 - 19.5	300	4.3	2.7 - 6.8	1,900	31.1	26.3 - 36.3	6,200
131-185%	200	6.1	2.9 - 12.2	800	29.6	22.7 - 37.5	600	21.8	15.8 - 29.3	200	8.2	4.6 - 14.2	1,000	34.3	27.2 - 42.3	2,800
186+%	200	2.9	1.6 - 5.0	2,400	33.4	28.8 - 38.3	1,200	16.2	12.9 - 20.3	400	5.3	3.5 - 8.1	3,000	42.2	37.4 - 47.1	7,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	11.0	6.7 - 17.4	1,200	39.2	31.4 - 47.4	500	16.1	11.0 - 22.9	100	2.6	1.1 - 6.1	900	31.2	24.4 - 39.1	2,900
\$10,000-\$24,999	300	9.2	5.8 - 14.1	1,500	39.3	32.9 - 46.1	600	16.7	12.4 - 22.2	300	6.6	4.0 - 10.6	1,100	28.2	22.4 - 34.8	3,800
\$25,000-\$49,999	200	5.9	3.2 - 10.5	1,100	29.4	23.3 - 36.4	600	17.2	12.6 - 23.0	200	6.7	3.9 - 11.4	1,500	40.8	34.1 - 47.8	3,600
>=\$50,000	100	2.4	1.2 - 4.7	2,000	33.4	28.4 - 38.8	1,000	16.9	13.1 - 21.6	300	5.4	3.4 - 8.5	2,500	41.9	36.5 - 47.4	5,900

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.6	6.8 - 13.4	600	45.1	39.7 - 50.5	200	13.8	10.5 - 18.0	100	4.5	2.8 - 7.3	300	27.0	22.4 - 32.2	1,200
Normal birth weight	1,100	6.6	5.0 - 8.6	5,600	33.8	30.6 - 37.1	2,900	17.4	14.9 - 20.1	900	5.2	3.9 - 6.9	6,100	37.0	33.8 - 40.3	16,500

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	19.5	12.1 - 29.8	700	42.5	32.4 - 53.3	200	10.2	5.6 - 17.6	<50	1.3	0.4 - 3.7	400	26.5	18.3 - 36.9	1,600
No	900	5.6	4.2 - 7.5	5,300	33.6	30.4 - 36.8	2,800	17.8	15.3 - 20.6	900	5.7	4.3 - 7.5	5,800	37.4	34.1 - 40.7	15,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	14.6	10.2 - 20.4	1,500	37.8	31.4 - 44.6	500	11.9	8.6 - 16.2	200	5.2	3.1 - 8.8	1,200	30.5	24.7 - 37.1	3,900
No	600	4.7	3.3 - 6.6	4,500	33.5	30.1 - 37.0	2,500	18.5	15.8 - 21.6	700	5.3	3.9 - 7.2	5,100	38.0	34.5 - 41.6	13,400

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	9.2	6.6 - 12.6	2,600	35.7	31.1 - 40.6	1,300	17.4	14.0 - 21.5	400	5.3	3.6 - 8.0	2,300	32.4	27.9 - 37.2	7,200
No	500	5.3	3.6 - 7.8	3,300	33.4	29.5 - 37.6	1,700	16.9	13.9 - 20.5	500	5.3	3.7 - 7.6	3,900	39.0	35.0 - 43.3	10,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,400	7.8	6.6 - 9.3	6,500	36.2	34.0 - 38.5	3,100	17.0	15.2 - 18.8	600	3.1	2.4 - 4.1	6,500	35.9	33.6 - 38.2	18,000

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	11.2	7.4 - 16.5	700	31.3	25.2 - 38.0	400	15.7	11.2 - 21.4	<50	1.3	0.4 - 3.8	1,000	40.6	34.0 - 47.5	2,400
Honolulu	900	7.4	6.0 - 9.1	4,800	38.0	35.3 - 40.8	2,100	16.3	14.4 - 18.5	400	3.4	2.5 - 4.6	4,400	34.8	32.2 - 37.5	12,700
Kauai	<50	1.9	0.3 - 11.6	300	30.7	21.3 - 41.9	200	24.8	16.3 - 35.9	<50	3.0	0.8 - 10.9	300	39.6	29.3 - 50.8	800
Maui	200	9.1	5.7 - 14.3	700	32.8	26.5 - 39.8	400	19.1	14.1 - 25.3	100	3.7	1.7 - 7.7	700	35.3	28.8 - 42.4	2,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	4.4	2.7 - 6.9	900	22.1	18.3 - 26.5	800	18.3	14.9 - 22.4	200	5.5	3.6 - 8.2	2,100	49.7	44.8 - 54.6	4,300
Native Hawaiian	500	9.6	7.0 - 13.0	1,900	37.2	32.5 - 42.1	900	17.1	13.7 - 21.2	200	3.8	2.3 - 6.3	1,600	32.3	27.9 - 37.1	5,000
Chinese	<50	2.1	0.8 - 5.5	200	39.9	33.5 - 46.6	200	24.5	19.1 - 30.9	<50	1.8	0.7 - 4.6	200	31.7	25.8 - 38.3	600
Filipino	300	8.5	6.0 - 12.0	1,600	47.8	42.5 - 53.1	500	15.4	11.9 - 19.6	<50	1.0	0.3 - 2.8	900	27.3	22.8 - 32.3	3,200
Japanese	100	3.0	1.2 - 7.4	1,000	47.2	40.3 - 54.2	300	16.3	11.7 - 22.3	100	2.9	1.4 - 6.2	600	30.6	24.6 - 37.3	2,000
Other	400	13.9	9.9 - 19.1	1,000	33.9	28.4 - 40.0	400	15.3	11.3 - 20.4	<50	1.3	0.5 - 3.7	1,000	35.6	29.9 - 41.7	2,900

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	10.3	5.9 - 17.3	500	33.8	25.9 - 42.8	200	15.7	10.2 - 23.4	<50	3.0	1.0 - 8.6	500	37.2	28.9 - 46.3	1,400
20-24	400	9.8	7.0 - 13.7	1,400	34.3	29.5 - 39.5	700	18.6	14.8 - 23.2	200	4.4	2.7 - 7.1	1,300	32.8	28.0 - 37.9	4,000
25-34	700	7.6	6.0 - 9.7	3,300	35.6	32.5 - 38.8	1,500	15.7	13.4 - 18.2	300	2.9	1.9 - 4.2	3,600	38.3	35.1 - 41.5	9,300
35+	200	5.0	3.1 - 8.1	1,400	41.2	36.1 - 46.5	600	19.2	15.3 - 23.7	100	2.5	1.2 - 4.8	1,100	32.2	27.5 - 37.2	3,300

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	7.3	5.7 - 9.4	3,300	35.0	32.0 - 38.2	1,700	18.2	15.8 - 20.9	300	3.3	2.3 - 4.8	3,400	36.1	33.0 - 39.3	9,400
Female	700	8.4	6.6 - 10.7	3,200	37.5	34.2 - 40.9	1,300	15.6	13.2 - 18.3	300	3.0	2.0 - 4.4	3,100	35.6	32.3 - 39.0	8,600

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	5.7	4.4 - 7.3	4,100	36.5	33.8 - 39.4	1,800	16.3	14.2 - 18.6	300	2.8	2.0 - 4.0	4,400	38.7	35.9 - 41.5	11,300
No	800	11.5	9.1 - 14.5	2,400	35.6	31.8 - 39.6	1,200	18.1	15.1 - 21.5	200	3.7	2.4 - 5.6	2,100	31.1	27.5 - 35.1	6,700

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	800	12.2	9.7 - 15.3	2,400	36.0	32.2 - 40.1	1,100	16.7	13.9 - 20.0	200	3.0	1.8 - 4.7	2,100	32.0	28.3 - 36.0	6,600
131-185%	300	10.0	6.7 - 14.6	800	31.5	25.9 - 37.7	500	17.3	13.0 - 22.6	100	2.7	1.3 - 5.8	1,000	38.6	32.7 - 44.9	2,600
186+%	300	3.7	2.5 - 5.4	2,800	36.7	33.4 - 40.1	1,300	17.0	14.5 - 19.9	300	3.3	2.3 - 4.9	3,000	39.2	35.9 - 42.7	7,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	400	13.3	9.5 - 18.3	1,200	38.6	32.8 - 44.8	500	16.8	12.7 - 21.9	<50	1.6	0.6 - 4.2	900	29.7	24.4 - 35.7	3,000
\$10,000-\$24,999	400	10.4	7.5 - 14.3	1,300	32.8	28.0 - 38.1	700	17.0	13.3 - 21.3	200	4.5	2.7 - 7.3	1,400	35.3	30.4 - 40.6	3,900
\$25,000-\$49,999	300	7.3	5.0 - 10.5	1,500	35.4	30.8 - 40.2	800	18.8	15.2 - 23.0	200	3.8	2.3 - 6.1	1,400	34.8	30.3 - 39.6	4,100
>=\$50,000	300	4.2	2.8 - 6.2	2,200	36.5	32.8 - 40.3	1,000	16.1	13.4 - 19.3	100	2.4	1.4 - 4.1	2,500	40.8	37.0 - 44.7	6,100

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.4	3.6 - 14.3	600	49.3	40.2 - 58.5	200	15.7	10.1 - 23.5	<50	3.4	1.3 - 8.3	300	24.3	17.2 - 33.0	1,200
Normal birth weight	1,300	7.9	6.6 - 9.4	5,900	35.2	32.9 - 37.6	2,900	17.1	15.3 - 19.0	500	3.1	2.4 - 4.1	6,200	36.7	34.4 - 39.1	16,800

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	15.5	10.0 - 23.1	500	34.8	26.9 - 43.6	300	21.9	15.4 - 30.0	100	7.0	3.6 - 13.1	300	20.8	14.6 - 28.9	1,500
No	1,200	7.2	6.0 - 8.7	5,900	36.3	34.0 - 38.8	2,700	16.6	14.8 - 18.6	500	2.9	2.1 - 3.8	6,000	37.0	34.6 - 39.4	16,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	12.9	9.4 - 17.5	1,100	34.4	29.2 - 40.1	700	19.7	15.5 - 24.8	200	6.0	3.8 - 9.4	900	26.9	22.1 - 32.4	3,300
No	1,000	6.7	5.5 - 8.3	5,300	36.6	34.1 - 39.2	2,400	16.4	14.5 - 18.5	400	2.6	1.8 - 3.6	5,400	37.7	35.2 - 40.3	14,400

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	11.7	9.4 - 14.5	2,700	36.2	32.6 - 40.0	1,300	17.4	14.6 - 20.4	200	3.1	2.0 - 4.7	2,400	31.7	28.2 - 35.4	7,600
No	500	5.1	3.9 - 6.8	3,700	36.5	33.6 - 39.5	1,700	16.6	14.4 - 19.1	300	3.3	2.3 - 4.6	3,900	38.4	35.5 - 41.5	10,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,400	7.7	6.4 - 9.1	6,300	35.6	33.3 - 37.9	3,100	17.5	15.7 - 19.4	500	2.9	2.2 - 3.9	6,400	36.3	34.0 - 38.6	17,800

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	10.9	7.2 - 16.1	700	29.1	23.3 - 35.6	300	11.8	8.0 - 16.9	100	2.5	1.1 - 5.6	1,100	45.8	39.1 - 52.7	2,300
Honolulu	800	6.9	5.5 - 8.5	4,700	37.9	35.2 - 40.7	2,300	18.6	16.5 - 21.0	400	2.9	2.1 - 4.0	4,200	33.7	31.1 - 36.5	12,300
Kauai	100	10.4	5.4 - 19.1	200	15.3	9.2 - 24.2	300	24.8	16.9 - 34.7	100	5.6	2.4 - 12.4	400	44.0	34.1 - 54.5	1,000
Maui	200	7.5	4.4 - 12.4	800	39.0	32.1 - 46.4	300	14.1	9.7 - 19.9	100	2.4	0.9 - 6.3	800	37.0	30.3 - 44.3	2,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	400	9.8	7.2 - 13.3	900	22.7	18.8 - 27.2	600	14.5	11.3 - 18.4	200	4.0	2.4 - 6.5	2,000	49.0	44.0 - 54.0	4,100
Native Hawaiian	400	7.7	5.4 - 10.9	1,700	33.6	29.1 - 38.5	1,100	22.7	18.8 - 27.1	100	2.8	1.6 - 5.1	1,600	33.1	28.5 - 37.9	4,900
Chinese	<50	2.8	1.2 - 6.5	300	39.3	32.9 - 46.0	100	17.5	12.9 - 23.3	<50	2.4	1.0 - 5.6	300	38.0	31.7 - 44.7	700
Filipino	200	7.3	4.9 - 10.6	1,400	45.0	39.8 - 50.3	600	18.1	14.4 - 22.5	100	3.2	1.8 - 5.7	800	26.4	22.1 - 31.3	3,200
Japanese	100	2.8	1.1 - 7.1	1,100	55.0	48.0 - 61.8	300	13.7	9.6 - 19.2	<50	1.9	0.7 - 4.7	600	26.5	20.8 - 33.2	2,100
Other	300	9.6	6.4 - 14.2	900	31.8	26.4 - 37.8	400	15.0	11.0 - 20.1	100	2.1	0.9 - 4.9	1,200	41.4	35.4 - 47.7	2,900

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	10.7	6.1 - 18.1	500	30.4	22.5 - 39.7	400	25.4	18.2 - 34.1	<50	2.9	0.9 - 8.5	500	30.6	22.7 - 39.9	1,600
20-24	400	9.8	7.3 - 13.2	1,400	31.8	27.4 - 36.6	900	19.9	16.2 - 24.2	100	2.5	1.3 - 4.6	1,600	36.0	31.4 - 40.9	4,500
25-34	600	6.9	5.3 - 8.9	3,200	37.1	33.9 - 40.4	1,300	15.3	13.0 - 17.8	300	3.1	2.1 - 4.5	3,300	37.7	34.5 - 40.9	8,700
35+	200	5.1	3.0 - 8.6	1,200	39.5	34.3 - 45.0	500	16.4	12.8 - 20.8	100	3.2	1.7 - 5.9	1,100	35.8	30.7 - 41.2	3,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	600	7.0	5.4 - 9.0	3,200	35.2	32.1 - 38.4	1,700	18.3	15.8 - 21.0	300	2.7	1.9 - 4.0	3,400	36.8	33.6 - 40.0	9,200
Female	700	8.4	6.6 - 10.6	3,100	36.0	32.8 - 39.4	1,400	16.7	14.2 - 19.5	300	3.1	2.1 - 4.7	3,100	35.8	32.5 - 39.2	8,600

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	6.5	5.1 - 8.2	4,000	36.1	33.4 - 38.9	1,900	16.8	14.7 - 19.1	300	2.7	1.9 - 3.9	4,200	37.9	35.1 - 40.8	11,100
No	600	9.7	7.5 - 12.5	2,300	34.8	30.9 - 38.8	1,200	18.7	15.6 - 22.2	200	3.3	2.1 - 5.1	2,200	33.5	29.7 - 37.6	6,600

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	700	10.4	8.0 - 13.3	2,300	34.6	30.8 - 38.7	1,200	19.0	15.9 - 22.5	200	2.6	1.5 - 4.3	2,200	33.4	29.6 - 37.5	6,600
131-185%	200	7.6	4.9 - 11.7	900	33.6	28.0 - 39.6	500	19.7	15.2 - 25.2	100	2.3	1.0 - 5.3	1,000	36.7	31.0 - 42.9	2,600
186+%	400	5.1	3.7 - 7.0	2,800	38.3	34.9 - 41.8	1,100	14.9	12.5 - 17.6	300	3.7	2.5 - 5.3	2,800	38.1	34.7 - 41.7	7,300

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	10.9	7.5 - 15.5	1,000	33.4	27.7 - 39.6	500	17.0	12.7 - 22.4	100	3.1	1.5 - 6.4	1,100	35.6	29.8 - 41.9	3,000
\$10,000-\$24,999	300	9.1	6.3 - 13.0	1,300	34.6	29.7 - 40.0	800	22.4	18.2 - 27.3	100	1.4	0.6 - 3.3	1,200	32.5	27.6 - 37.8	3,600
\$25,000-\$49,999	300	7.6	5.3 - 10.8	1,400	35.3	30.7 - 40.2	700	17.9	14.4 - 22.1	100	3.6	2.1 - 6.0	1,500	35.6	31.0 - 40.6	4,100
>=\$50,000	300	4.9	3.4 - 7.0	2,300	38.8	35.0 - 42.7	800	14.3	11.7 - 17.4	200	3.5	2.3 - 5.4	2,300	38.4	34.6 - 42.4	5,900

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.2	5.3 - 15.5	600	40.1	32.2 - 48.6	300	22.4	15.9 - 30.5	100	4.7	2.2 - 9.8	300	23.7	17.1 - 31.8	1,400
Normal birth weight	1,200	7.6	6.3 - 9.1	5,800	35.2	32.9 - 37.6	2,800	17.1	15.3 - 19.1	500	2.8	2.1 - 3.7	6,100	37.4	35.0 - 39.8	16,300

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	16.6	11.2 - 23.9	500	34.3	26.5 - 43.0	300	18.5	12.6 - 26.3	100	4.4	1.9 - 10.0	400	26.2	19.3 - 34.6	1,500
No	1,100	6.9	5.6 - 8.3	5,700	35.8	33.5 - 38.2	2,800	17.5	15.7 - 19.5	400	2.7	2.0 - 3.7	5,900	37.1	34.7 - 39.5	16,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	12.3	9.0 - 16.5	1,200	33.0	28.0 - 38.5	800	24.1	19.6 - 29.3	100	3.2	1.7 - 5.9	1,000	27.4	22.7 - 32.6	3,500
No	900	6.6	5.3 - 8.1	5,100	36.3	33.8 - 38.9	2,200	16.0	14.1 - 18.0	400	2.8	2.1 - 3.9	5,400	38.3	35.8 - 41.0	14,000

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	10.9	8.7 - 13.7	2,300	31.8	28.3 - 35.6	1,300	17.9	15.1 - 21.1	200	2.7	1.6 - 4.3	2,700	36.7	33.0 - 40.6	7,300
No	500	5.3	4.0 - 6.9	3,800	38.0	35.1 - 41.0	1,700	17.2	14.9 - 19.7	300	3.2	2.3 - 4.5	3,700	36.3	33.4 - 39.3	10,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,400	8.2	6.9 - 9.7	5,600	32.0	29.7 - 34.3	3,400	19.4	17.4 - 21.4	800	4.4	3.5 - 5.5	6,300	36.1	33.7 - 38.5	17,400

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	7.0	3.9 - 12.1	600	26.6	20.6 - 33.7	400	20.3	14.9 - 27.0	100	4.4	2.3 - 8.5	900	41.7	34.5 - 49.2	2,100
Honolulu	1,100	8.3	6.8 - 10.1	4,500	35.2	32.5 - 38.0	2,400	18.5	16.4 - 20.9	500	4.2	3.2 - 5.5	4,300	33.8	31.1 - 36.5	12,700
Kauai	100	6.9	2.7 - 16.6	100	12.9	7.0 - 22.7	200	22.5	14.0 - 34.0	100	6.3	2.5 - 15.1	400	51.4	39.6 - 63.0	800
Maui	200	9.2	5.5 - 15.1	400	23.9	17.9 - 31.3	400	22.7	16.7 - 30.1	100	4.7	2.3 - 9.4	700	39.4	31.9 - 47.3	1,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	7.3	5.0 - 10.5	900	21.8	17.9 - 26.4	600	15.1	11.8 - 19.1	300	7.5	5.2 - 10.6	2,000	48.3	43.2 - 53.4	4,200
Native Hawaiian	500	11.2	8.3 - 14.9	1,500	30.9	26.3 - 35.9	1,100	24.0	19.8 - 28.7	100	2.7	1.4 - 4.9	1,500	31.3	26.7 - 36.3	4,700
Chinese	<50	2.2	0.8 - 5.6	200	42.4	35.5 - 49.7	100	20.9	15.6 - 27.3	<50	3.6	1.7 - 7.5	200	30.9	24.7 - 38.0	600
Filipino	200	7.2	4.8 - 10.7	1,200	37.6	32.4 - 43.1	600	18.3	14.4 - 23.0	100	4.5	2.7 - 7.4	1,000	32.4	27.5 - 37.8	3,200
Japanese	100	5.1	2.6 - 9.7	900	46.8	39.8 - 54.0	300	16.8	12.0 - 23.1	<50	2.1	0.8 - 5.4	600	29.1	23.0 - 36.1	2,000
Other	200	9.1	5.9 - 13.8	800	29.9	24.3 - 36.1	600	20.7	16.0 - 26.5	100	4.2	2.2 - 8.0	1,000	36.0	30.1 - 42.5	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.0	4.8 - 16.0	400	30.1	22.4 - 39.1	400	26.9	19.4 - 35.9	0	0.0	-	500	34.0	25.9 - 43.3	1,500
20-24	400	9.2	6.5 - 12.9	1,400	33.0	28.2 - 38.1	900	21.3	17.3 - 26.0	200	5.2	3.3 - 8.3	1,300	31.3	26.6 - 36.4	4,200
25-34	600	7.1	5.4 - 9.2	2,600	30.5	27.4 - 33.8	1,600	18.5	15.9 - 21.4	400	4.8	3.5 - 6.5	3,300	39.1	35.7 - 42.6	8,500
35+	300	9.4	6.6 - 13.4	1,100	35.4	30.3 - 40.7	500	15.6	12.0 - 20.0	100	4.1	2.4 - 7.0	1,100	35.5	30.5 - 40.9	3,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	7.4	5.7 - 9.5	2,900	32.8	29.7 - 36.2	1,900	21.4	18.6 - 24.5	400	4.9	3.6 - 6.6	3,000	33.5	30.3 - 36.9	8,900
Female	800	9.1	7.1 - 11.4	2,600	31.1	27.9 - 34.4	1,500	17.2	14.7 - 20.1	300	3.9	2.7 - 5.5	3,300	38.8	35.4 - 42.3	8,500

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	7.9	6.3 - 9.7	3,600	32.7	29.9 - 35.5	1,900	16.7	14.6 - 19.1	500	4.6	3.5 - 6.1	4,200	38.2	35.3 - 41.1	11,100
No	600	8.8	6.5 - 11.7	1,900	30.7	26.8 - 35.0	1,500	24.0	20.4 - 28.0	300	4.0	2.6 - 6.1	2,100	32.5	28.5 - 36.7	6,300

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	700	11.1	8.6 - 14.3	1,900	30.9	27.0 - 35.1	1,400	22.2	18.7 - 26.1	200	3.6	2.3 - 5.6	2,000	32.2	28.2 - 36.4	6,100
131-185%	200	10.8	7.2 - 15.8	700	29.0	23.2 - 35.7	400	17.9	13.1 - 24.1	100	5.6	3.1 - 9.9	800	36.7	30.2 - 43.7	2,300
186+%	400	4.6	3.3 - 6.4	2,700	34.0	30.7 - 37.4	1,400	17.5	14.9 - 20.3	400	4.5	3.2 - 6.2	3,100	39.5	36.1 - 43.0	7,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	11.6	8.0 - 16.5	800	27.4	22.0 - 33.5	700	24.1	19.0 - 30.2	100	3.5	1.8 - 6.8	1,000	33.4	27.6 - 39.7	2,900
\$10,000-\$24,999	300	8.5	5.8 - 12.3	1,300	35.3	30.1 - 40.8	800	20.7	16.5 - 25.8	100	3.8	2.2 - 6.7	1,200	31.6	26.7 - 37.1	3,600
\$25,000-\$49,999	300	7.8	5.5 - 11.1	1,300	31.4	26.9 - 36.3	800	19.8	15.9 - 24.3	200	5.4	3.5 - 8.3	1,500	35.6	30.8 - 40.7	4,100
>=\$50,000	300	5.8	4.0 - 8.2	1,900	32.9	29.2 - 36.9	900	15.7	12.9 - 18.9	200	4.1	2.8 - 6.1	2,300	41.6	37.5 - 45.7	5,600

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	12.2	7.3 - 19.6	500	36.3	28.4 - 45.1	200	16.9	11.1 - 24.8	<50	2.0	0.7 - 5.3	500	32.6	24.8 - 41.4	1,400
Normal birth weight	1,300	7.9	6.5 - 9.4	5,100	31.6	29.3 - 34.1	3,100	19.6	17.6 - 21.8	700	4.6	3.6 - 5.8	5,800	36.4	33.9 - 38.9	16,000

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	22.3	15.8 - 30.5	400	27.9	20.8 - 36.3	300	19.0	13.0 - 27.1	100	5.6	2.7 - 11.1	400	25.2	18.3 - 33.5	1,600
No	1,100	6.9	5.6 - 8.4	5,000	32.1	29.7 - 34.5	3,000	19.4	17.4 - 21.6	700	4.3	3.4 - 5.5	5,800	37.3	34.8 - 39.9	15,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	13.7	10.2 - 18.3	1,000	26.9	22.1 - 32.2	800	23.1	18.5 - 28.3	300	7.3	4.9 - 10.9	1,100	29.0	24.1 - 34.4	3,600
No	900	6.9	5.5 - 8.5	4,500	33.0	30.4 - 35.6	2,500	18.4	16.3 - 20.7	500	3.7	2.8 - 4.8	5,200	38.1	35.4 - 40.9	13,600

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	10.3	8.1 - 13.1	2,300	31.9	28.3 - 35.8	1,500	20.6	17.5 - 24.1	300	4.6	3.2 - 6.6	2,400	32.5	28.9 - 36.4	7,300
No	700	6.8	5.3 - 8.7	3,100	31.8	29.0 - 34.9	1,800	18.1	15.7 - 20.7	400	4.1	3.0 - 5.5	3,900	39.2	36.1 - 42.4	9,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,700	10.1	8.7 - 11.7	5,500	32.4	30.3 - 34.7	3,300	19.3	17.5 - 21.3	500	3.1	2.3 - 4.0	5,900	35.1	32.9 - 37.4	16,900

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	13.3	9.1 - 18.9	500	23.4	18.2 - 29.7	500	23.2	17.8 - 29.5	100	2.3	1.0 - 5.5	800	37.8	31.4 - 44.7	2,200
Honolulu	1,300	10.3	8.7 - 12.1	4,400	35.2	32.6 - 37.9	2,400	19.1	17.0 - 21.4	400	2.9	2.1 - 4.0	4,000	32.4	29.9 - 35.1	12,400
Kauai	<50	6.6	2.6 - 16.1	100	13.5	7.5 - 23.1	200	22.5	14.2 - 33.8	<50	3.5	0.9 - 12.3	400	53.9	42.1 - 65.2	700
Maui	100	5.8	3.0 - 11.0	500	31.9	25.2 - 39.5	200	13.8	9.2 - 20.0	100	4.7	2.3 - 9.3	700	43.8	36.3 - 51.6	1,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	6.0	3.9 - 9.0	800	20.5	16.6 - 25.1	700	19.0	15.3 - 23.5	100	3.1	1.7 - 5.5	1,900	51.4	46.2 - 56.5	3,700
Native Hawaiian	600	13.3	10.2 - 17.2	1,500	31.9	27.4 - 36.6	800	17.8	14.3 - 21.9	200	3.8	2.3 - 6.2	1,600	33.3	28.8 - 38.1	4,800
Chinese	100	9.6	6.4 - 14.1	200	37.7	31.7 - 44.2	100	18.0	13.5 - 23.5	<50	3.6	1.8 - 7.0	200	31.1	25.5 - 37.4	600
Filipino	300	10.0	7.3 - 13.5	1,300	40.4	35.5 - 45.5	700	21.9	18.0 - 26.5	100	3.6	2.1 - 6.0	800	24.1	20.0 - 28.8	3,100
Japanese	100	4.6	2.4 - 8.3	1,100	47.3	41.1 - 53.7	300	15.3	11.3 - 20.5	100	2.3	1.0 - 5.2	700	30.5	25.0 - 36.7	2,300
Other	400	15.5	11.4 - 20.7	600	26.4	21.3 - 32.1	600	23.2	18.3 - 29.0	<50	1.4	0.6 - 3.7	800	33.5	27.8 - 39.7	2,400

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	12.1	7.4 - 19.0	400	30.2	23.0 - 38.6	300	22.3	16.0 - 30.3	100	3.8	1.6 - 9.0	500	31.6	24.2 - 40.0	1,500
20-24	500	11.3	8.5 - 14.9	1,200	29.3	25.1 - 34.0	1,000	24.0	20.0 - 28.4	100	3.5	2.2 - 5.8	1,300	31.8	27.4 - 36.7	4,200
25-34	800	10.0	8.1 - 12.3	2,800	33.1	30.1 - 36.3	1,400	17.2	14.8 - 19.9	300	3.4	2.3 - 4.9	3,000	36.3	33.1 - 39.6	8,400
35+	200	7.6	5.1 - 11.1	1,000	36.1	31.1 - 41.4	500	17.1	13.3 - 21.7	<50	1.0	0.4 - 2.2	1,100	38.3	33.1 - 43.7	2,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	10.2	8.3 - 12.5	2,800	32.9	29.9 - 36.1	1,600	19.0	16.4 - 21.8	300	3.0	2.0 - 4.4	3,000	34.9	31.8 - 38.1	8,600
Female	800	9.9	8.0 - 12.2	2,700	32.0	28.9 - 35.2	1,600	19.6	17.1 - 22.5	300	3.1	2.1 - 4.5	2,900	35.4	32.2 - 38.7	8,300

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	8.0	6.5 - 9.8	3,500	32.8	30.1 - 35.5	1,900	17.9	15.8 - 20.3	300	2.9	2.1 - 4.1	4,100	38.3	35.5 - 41.2	10,800
No	800	13.7	11.0 - 16.8	2,000	31.9	28.2 - 35.8	1,300	21.6	18.5 - 25.2	200	3.3	2.2 - 5.0	1,800	29.5	25.9 - 33.4	6,100

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,000	15.4	12.6 - 18.6	2,000	31.5	27.9 - 35.3	1,300	20.6	17.5 - 24.0	100	2.2	1.3 - 3.6	1,900	30.4	26.8 - 34.2	6,200
131-185%	200	11.0	7.1 - 16.7	500	29.5	23.2 - 36.7	400	20.2	14.8 - 26.9	100	3.6	1.7 - 7.5	600	35.8	29.0 - 43.2	1,800
186+%	400	5.1	3.7 - 6.9	2,600	34.7	31.5 - 38.0	1,300	16.8	14.3 - 19.6	300	3.7	2.6 - 5.3	3,000	39.7	36.4 - 43.2	7,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	400	14.1	10.4 - 18.9	1,000	33.5	28.2 - 39.2	600	20.2	15.9 - 25.3	100	3.5	1.8 - 6.4	900	28.8	23.7 - 34.4	3,000
\$10,000-\$24,999	500	14.7	11.3 - 18.9	1,100	32.2	27.5 - 37.3	800	21.2	17.2 - 25.8	<50	1.2	0.5 - 2.8	1,100	30.7	26.0 - 35.7	3,600
\$25,000-\$49,999	300	8.0	5.6 - 11.3	1,100	28.2	23.9 - 32.9	700	16.9	13.5 - 21.0	200	4.4	2.8 - 6.9	1,700	42.5	37.6 - 47.5	3,900
>=\$50,000	300	5.3	3.7 - 7.7	2,000	37.2	33.4 - 41.2	900	17.4	14.5 - 20.9	200	3.3	2.0 - 5.2	1,900	36.8	32.9 - 40.8	5,300

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.3	3.8 - 13.6	600	47.2	38.4 - 56.2	100	12.3	7.5 - 19.6	100	4.3	1.9 - 9.5	300	28.9	21.4 - 37.8	1,200
Normal birth weight	1,600	10.2	8.8 - 11.9	4,900	31.3	29.1 - 33.6	3,100	19.8	17.9 - 21.9	500	3.0	2.2 - 3.9	5,600	35.6	33.3 - 38.0	15,700

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	23.7	17.0 - 32.1	500	34.8	27.1 - 43.4	200	12.9	8.2 - 19.8	<50	0.9	0.1 - 5.4	400	27.7	20.6 - 36.1	1,400
No	1,300	8.8	7.4 - 10.4	4,900	32.1	29.9 - 34.5	3,000	19.6	17.7 - 21.7	500	3.3	2.5 - 4.3	5,500	36.1	33.8 - 38.5	15,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	15.4	11.8 - 19.8	1,200	32.9	28.1 - 38.1	700	20.0	16.0 - 24.6	100	4.2	2.5 - 7.0	1,000	27.5	23.0 - 32.6	3,600
No	1,100	8.6	7.2 - 10.3	4,200	32.2	29.8 - 34.7	2,500	18.9	16.8 - 21.1	400	2.8	2.1 - 3.9	4,900	37.5	34.9 - 40.1	13,000

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	13.0	10.6 - 15.8	2,100	29.6	26.3 - 33.1	1,500	20.5	17.6 - 23.7	200	2.9	2.0 - 4.4	2,500	34.0	30.5 - 37.6	7,200
No	700	7.8	6.2 - 9.7	3,200	34.4	31.5 - 37.3	1,700	18.1	15.8 - 20.7	300	3.2	2.3 - 4.6	3,400	36.5	33.6 - 39.5	9,400

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,600	9.1	7.9 - 10.5	5,700	33.6	31.6 - 35.7	3,300	19.2	17.5 - 21.0	600	3.7	3.0 - 4.6	5,900	34.4	32.4 - 36.5	17,100

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	12.6	8.8 - 17.8	600	27.0	21.6 - 33.2	400	17.8	13.3 - 23.4	100	2.9	1.4 - 6.3	800	39.6	33.5 - 46.1	2,100
Honolulu	1,200	9.7	8.2 - 11.4	4,500	36.3	33.9 - 38.8	2,300	18.5	16.5 - 20.6	400	3.2	2.4 - 4.2	4,000	32.3	30.0 - 34.7	12,400
Kauai	<50	2.3	0.6 - 8.2	100	15.6	9.5 - 24.6	200	24.5	16.8 - 34.3	<50	4.9	1.9 - 11.8	400	52.7	42.5 - 62.7	800
Maui	100	3.7	1.8 - 7.3	500	30.4	24.5 - 36.9	400	23.6	18.3 - 29.9	100	7.5	4.5 - 12.2	600	34.9	28.8 - 41.6	1,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.7	3.9 - 8.3	800	19.8	16.4 - 23.8	700	17.3	14.1 - 21.1	200	5.2	3.5 - 7.7	2,000	51.9	47.3 - 56.5	3,800
Native Hawaiian	600	13.4	10.6 - 17.0	1,700	35.1	30.9 - 39.6	1,000	21.8	18.2 - 25.8	200	4.2	2.7 - 6.5	1,200	25.5	21.7 - 29.7	4,700
Chinese	<50	5.9	4.1 - 8.6	300	41.7	37.3 - 46.3	100	16.9	13.8 - 20.6	<50	2.0	1.0 - 3.8	200	33.5	29.3 - 37.9	700
Filipino	300	8.2	5.9 - 11.2	1,200	36.9	32.6 - 41.5	600	20.0	16.5 - 23.9	100	4.1	2.6 - 6.4	1,000	30.9	26.8 - 35.3	3,200
Japanese	100	5.2	3.0 - 9.0	1,000	45.5	39.3 - 51.8	400	18.8	14.4 - 24.2	<50	1.8	0.7 - 4.5	600	28.7	23.3 - 34.7	2,200
Other	300	11.4	8.1 - 16.0	900	34.6	29.1 - 40.5	400	17.1	13.0 - 22.2	100	2.2	1.0 - 4.6	900	34.7	29.2 - 40.6	2,500

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	14.2	8.9 - 21.7	400	28.7	21.6 - 37.1	300	24.5	17.7 - 32.8	100	4.7	2.2 - 9.8	300	27.9	20.9 - 36.2	1,200
20-24	400	9.8	7.4 - 12.9	1,300	30.3	26.3 - 34.6	1,000	23.4	19.8 - 27.4	200	4.6	3.0 - 6.9	1,400	31.9	27.9 - 36.2	4,400
25-34	700	8.7	7.0 - 10.7	2,800	34.3	31.4 - 37.4	1,400	16.8	14.6 - 19.3	300	3.4	2.4 - 4.7	3,000	36.8	33.8 - 39.8	8,300
35+	200	7.2	5.0 - 10.2	1,200	37.9	33.3 - 42.8	600	17.6	14.1 - 21.7	100	3.0	1.7 - 5.2	1,100	34.3	29.8 - 39.1	3,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	9.8	8.1 - 11.8	3,200	34.7	31.9 - 37.6	1,800	18.9	16.6 - 21.3	300	3.4	2.4 - 4.7	3,100	33.2	30.5 - 36.0	9,300
Female	600	8.2	6.6 - 10.3	2,500	32.3	29.3 - 35.4	1,500	19.6	17.1 - 22.3	300	4.1	3.0 - 5.6	2,800	35.9	32.8 - 39.0	7,800

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	7.9	6.5 - 9.4	4,000	33.1	30.7 - 35.5	2,200	18.3	16.4 - 20.4	500	3.9	3.0 - 5.0	4,400	36.9	34.4 - 39.4	12,000
No	600	11.9	9.4 - 15.0	1,800	34.8	30.9 - 38.9	1,100	21.3	18.1 - 25.0	200	3.3	2.1 - 5.1	1,500	28.7	25.1 - 32.6	5,100

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	800	12.1	9.7 - 14.8	2,000	31.6	28.1 - 35.2	1,400	22.0	19.0 - 25.3	200	2.8	1.8 - 4.5	2,000	31.6	28.2 - 35.2	6,300
131-185%	200	11.1	7.7 - 15.9	500	25.0	20.0 - 30.9	400	18.7	14.2 - 24.2	100	5.8	3.4 - 9.5	800	39.4	33.4 - 45.7	2,100
186+%	400	5.2	4.0 - 6.9	2,800	37.3	34.3 - 40.5	1,300	17.2	14.9 - 19.7	300	4.1	3.0 - 5.6	2,700	36.2	33.1 - 39.3	7,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	10.2	7.0 - 14.6	1,000	37.6	32.0 - 43.5	600	22.1	17.6 - 27.4	100	2.3	1.1 - 4.8	700	27.9	22.9 - 33.4	2,600
\$10,000-\$24,999	600	13.5	10.6 - 17.0	1,200	27.3	23.4 - 31.5	900	20.1	16.7 - 23.9	200	4.6	3.0 - 7.0	1,500	34.5	30.4 - 39.0	4,200
\$25,000-\$49,999	300	6.2	4.4 - 8.6	1,300	29.6	25.8 - 33.7	900	21.2	17.9 - 25.0	200	4.5	3.1 - 6.7	1,700	38.5	34.4 - 42.7	4,400
>=\$50,000	300	5.7	4.1 - 8.0	1,900	40.1	36.3 - 44.1	700	15.1	12.4 - 18.1	200	3.3	2.2 - 5.1	1,700	35.8	32.0 - 39.7	4,700

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	10.3	5.9 - 17.4	500	50.9	42.0 - 59.7	100	12.7	7.9 - 19.8	100	5.0	2.2 - 11.0	200	21.1	14.8 - 29.2	1,000
Normal birth weight	1,400	9.0	7.7 - 10.5	5,200	32.5	30.4 - 34.6	3,200	19.6	17.9 - 21.5	600	3.6	2.9 - 4.6	5,700	35.3	33.2 - 37.4	16,100

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	19.9	13.9 - 27.6	400	28.4	21.5 - 36.6	300	21.1	15.2 - 28.6	100	5.4	2.7 - 10.6	300	25.1	18.7 - 32.9	1,400
No	1,300	8.1	6.9 - 9.5	5,300	34.0	31.8 - 36.1	3,000	19.1	17.4 - 21.0	600	3.6	2.8 - 4.6	5,500	35.2	33.1 - 37.4	15,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	12.9	9.7 - 16.8	1,000	30.2	25.7 - 35.1	800	21.7	17.8 - 26.2	200	6.1	4.1 - 9.1	1,000	29.1	24.8 - 34.0	3,500
No	1,100	8.1	6.8 - 9.6	4,600	34.4	32.1 - 36.7	2,500	18.7	16.8 - 20.7	400	3.1	2.4 - 4.1	4,800	35.8	33.5 - 38.1	13,400

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	10.9	8.9 - 13.2	2,400	31.4	28.3 - 34.6	1,600	21.3	18.7 - 24.3	200	2.9	2.0 - 4.3	2,600	33.5	30.4 - 36.8	7,600
No	700	7.1	5.7 - 8.9	3,300	35.4	32.7 - 38.3	1,600	17.7	15.6 - 20.1	400	4.5	3.4 - 5.9	3,200	35.3	32.5 - 38.1	9,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,800	10.9	9.3 - 12.7	5,100	31.4	28.8 - 34.1	3,200	19.4	17.3 - 21.7	600	3.4	2.5 - 4.6	5,700	34.9	32.3 - 37.5	16,400

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	17.4	14.1 - 21.3	400	21.0	17.4 - 25.1	300	16.6	13.4 - 20.3	100	3.5	2.0 - 5.8	800	41.6	37.0 - 46.3	1,900
Honolulu	1,300	10.4	8.4 - 12.8	4,100	34.2	30.8 - 37.7	2,400	19.7	17.0 - 22.8	400	3.3	2.2 - 5.0	3,900	32.4	29.1 - 35.8	12,000
Kauai	100	9.1	5.6 - 14.6	200	20.3	14.8 - 27.2	200	20.0	14.6 - 26.7	<50	3.4	1.5 - 7.6	400	47.1	39.7 - 54.7	800
Maui	100	8.2	5.8 - 11.4	500	28.4	24.1 - 33.1	300	19.7	15.9 - 24.1	100	4.2	2.6 - 6.8	700	39.6	34.7 - 44.6	1,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.6	3.3 - 9.5	800	22.4	17.3 - 28.6	600	18.0	13.5 - 23.6	300	7.5	4.8 - 11.7	1,700	46.4	40.1 - 52.8	3,600
Native Hawaiian	600	14.6	12.5 - 16.9	1,300	28.5	25.8 - 31.4	900	19.6	17.3 - 22.2	100	2.6	1.8 - 3.7	1,500	34.7	31.9 - 37.7	4,500
Chinese	<50	3.3	0.5 - 19.3	300	45.1	30.1 - 61.0	100	14.7	6.4 - 30.3	<50	0.8	0.1 - 5.1	200	36.1	22.7 - 52.0	700
Filipino	500	13.9	9.9 - 19.2	1,200	33.2	27.4 - 39.7	800	21.8	16.9 - 27.8	100	1.8	0.7 - 4.8	1,000	29.2	23.6 - 35.5	3,600
Japanese	100	7.7	4.0 - 14.2	900	44.9	35.8 - 54.3	300	13.9	8.6 - 21.7	<50	1.0	0.1 - 6.6	600	32.5	24.5 - 41.7	1,900
Other	300	12.4	7.6 - 19.7	700	32.5	24.7 - 41.4	500	23.5	16.8 - 31.8	100	4.1	1.7 - 9.7	600	27.5	20.3 - 36.0	2,200

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	14.1	9.6 - 20.3	500	34.2	26.4 - 43.0	300	22.8	16.5 - 30.7	<50	1.2	0.4 - 3.5	400	27.7	21.1 - 35.4	1,500
20-24	600	14.5	11.2 - 18.6	1,000	26.2	21.7 - 31.4	900	22.8	18.4 - 27.8	100	3.1	1.6 - 5.9	1,300	33.3	28.4 - 38.7	3,800
25-34	800	10.2	8.0 - 13.0	2,600	31.8	28.1 - 35.7	1,500	18.4	15.5 - 21.8	400	4.4	3.0 - 6.5	2,900	35.1	31.4 - 39.0	8,200
35+	200	6.3	3.6 - 10.7	1,000	35.5	29.0 - 42.7	500	16.0	11.5 - 21.7	100	2.2	0.9 - 5.3	1,100	40.1	33.5 - 47.0	2,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,000	11.0	8.8 - 13.5	2,700	30.7	27.2 - 34.4	1,800	20.7	17.7 - 24.0	300	3.8	2.6 - 5.7	2,900	33.9	30.4 - 37.5	8,700
Female	800	10.9	8.6 - 13.6	2,500	32.1	28.4 - 36.1	1,400	18.0	15.1 - 21.3	200	3.0	1.9 - 4.7	2,800	36.0	32.2 - 40.0	7,700

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	7.2	5.6 - 9.2	3,400	31.9	28.6 - 35.4	2,000	18.9	16.2 - 21.9	400	3.6	2.5 - 5.2	4,200	38.5	35.1 - 42.0	10,800
No	1,000	18.1	14.9 - 21.8	1,700	30.3	26.4 - 34.5	1,100	20.4	17.2 - 24.2	200	3.1	1.9 - 5.1	1,600	28.0	24.4 - 31.8	5,600

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,200	16.5	13.7 - 19.7	2,200	30.3	26.7 - 34.3	1,500	20.7	17.6 - 24.3	200	2.8	1.7 - 4.5	2,200	29.7	26.2 - 33.5	7,300
131-185%	200	9.3	5.7 - 14.9	700	30.7	24.3 - 38.1	500	19.5	14.4 - 25.9	100	3.9	1.7 - 8.4	900	36.6	29.8 - 43.8	2,400
186+%	100	2.3	1.3 - 4.1	1,700	32.6	27.8 - 37.8	1,000	18.8	15.0 - 23.3	200	4.1	2.5 - 6.8	2,200	42.2	37.2 - 47.3	5,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	800	21.6	17.3 - 26.7	1,200	29.7	25.0 - 35.0	800	20.7	16.6 - 25.5	100	2.8	1.4 - 5.7	1,000	25.1	20.9 - 29.9	3,900
\$15,001-\$25,000	200	12.0	8.3 - 16.9	500	24.9	18.9 - 32.0	500	23.1	17.0 - 30.5	100	4.4	2.2 - 8.5	700	35.7	28.9 - 43.1	2,100
\$25,001-\$40,000	200	7.1	4.6 - 10.9	1,000	32.9	26.9 - 39.6	600	18.7	14.2 - 24.3	100	3.0	1.5 - 5.9	1,200	38.3	32.1 - 44.9	3,100
>=\$40,001	300	4.3	2.8 - 6.6	2,100	33.4	29.0 - 38.1	1,100	18.4	15.0 - 22.3	300	4.1	2.6 - 6.5	2,500	39.7	35.3 - 44.4	6,200

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.7	5.7 - 16.1	400	34.6	25.0 - 45.7	300	24.6	16.1 - 35.6	<50	3.9	1.1 - 12.7	300	27.2	19.4 - 36.8	1,100
Normal birth weight	1,700	11.0	9.3 - 12.9	4,800	31.1	28.5 - 33.9	2,900	19.0	16.9 - 21.4	500	3.4	2.5 - 4.6	5,400	35.4	32.7 - 38.2	15,300

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	35.4	27.3 - 44.5	400	27.4	20.3 - 36.0	200	13.0	8.1 - 20.1	<50	1.6	0.6 - 3.9	300	22.6	16.2 - 30.6	1,500
No	1,200	8.4	7.0 - 10.1	4,600	31.8	29.0 - 34.7	3,000	20.2	17.9 - 22.7	500	3.6	2.6 - 5.0	5,300	36.0	33.2 - 38.8	14,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	21.6	17.1 - 27.1	1,100	32.1	26.8 - 37.9	600	16.8	12.8 - 21.7	100	3.6	2.0 - 6.4	900	25.9	21.2 - 31.2	3,400
No	1,000	7.7	6.2 - 9.5	3,900	31.1	28.2 - 34.3	2,600	20.3	17.8 - 23.0	400	3.5	2.4 - 4.9	4,700	37.4	34.4 - 40.6	12,600

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	16.5	13.6 - 19.8	2,100	30.7	27.0 - 34.6	1,300	19.5	16.5 - 23.0	200	3.1	1.9 - 5.0	2,100	30.2	26.7 - 34.0	6,900
No	600	6.9	5.3 - 9.0	3,000	31.9	28.3 - 35.6	1,800	19.1	16.3 - 22.3	300	3.6	2.4 - 5.3	3,600	38.5	34.9 - 42.2	9,300

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,700	10.9	9.2 - 12.8	5,000	31.3	28.7 - 34.1	2,800	17.7	15.7 - 20.0	700	4.2	3.2 - 5.5	5,800	35.9	33.2 - 38.7	16,000

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	10.6	8.2 - 13.7	600	30.8	26.7 - 35.3	400	19.0	15.8 - 22.7	100	3.0	1.7 - 5.1	700	36.6	32.2 - 41.1	1,900
Honolulu	1,300	11.3	9.1 - 13.8	3,800	32.4	29.0 - 36.0	2,000	17.2	14.6 - 20.2	500	4.6	3.3 - 6.4	4,100	34.5	31.1 - 38.1	11,800
Kauai	100	12.5	8.0 - 18.9	100	14.7	9.9 - 21.4	100	20.2	14.5 - 27.3	<50	3.4	1.5 - 7.3	300	49.3	41.4 - 57.2	700
Maui	100	7.7	5.3 - 10.9	500	30.5	26.0 - 35.5	300	18.6	15.0 - 23.0	100	3.2	1.8 - 5.5	700	40.0	35.1 - 45.2	1,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	9.7	6.2 - 14.8	600	17.5	13.0 - 23.2	400	12.8	8.8 - 18.2	300	9.7	6.2 - 14.7	1,700	50.4	43.7 - 57.0	3,400
Native Hawaiian	700	15.6	13.4 - 18.1	1,300	29.2	26.4 - 32.1	900	20.1	17.8 - 22.7	200	3.6	2.6 - 5.0	1,400	31.5	28.7 - 34.4	4,400
Chinese	100	15.6	6.4 - 33.4	200	38.6	24.0 - 55.7	100	23.5	12.1 - 40.5	<50	3.2	0.5 - 18.9	100	19.1	9.3 - 35.1	600
Filipino	200	7.0	4.2 - 11.4	1,300	40.5	33.7 - 47.7	600	18.3	13.5 - 24.2	100	2.4	1.1 - 5.1	1,000	31.8	25.8 - 38.6	3,200
Japanese	<50	0.5	0.1 - 1.9	1,000	43.0	34.8 - 51.5	500	22.5	16.2 - 30.4	100	2.5	0.8 - 7.4	700	31.4	24.2 - 39.7	2,300
Other	400	18.7	12.1 - 27.7	600	29.1	21.3 - 38.2	300	12.9	7.8 - 20.7	<50	1.5	0.4 - 5.2	800	37.9	29.2 - 47.4	2,100

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	17.8	11.4 - 26.8	600	38.5	30.2 - 47.6	300	21.3	14.8 - 29.7	<50	3.3	0.9 - 10.8	300	19.1	13.7 - 25.9	1,500
20-24	400	10.2	7.3 - 14.1	1,200	30.2	25.3 - 35.6	800	20.5	16.4 - 25.3	100	2.7	1.5 - 4.9	1,400	36.4	31.3 - 41.8	4,000
25-34	800	10.4	8.1 - 13.1	2,400	30.1	26.4 - 34.1	1,200	15.5	12.7 - 18.7	400	5.1	3.6 - 7.2	3,100	39.0	35.0 - 43.1	8,100
35+	200	9.5	5.9 - 14.8	800	32.8	26.2 - 40.1	500	18.4	13.5 - 24.5	100	4.2	2.0 - 8.5	900	35.2	28.5 - 42.5	2,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,000	12.8	10.2 - 15.8	2,500	31.0	27.4 - 34.9	1,300	16.4	13.7 - 19.5	400	4.4	3.0 - 6.3	2,900	35.4	31.7 - 39.3	8,100
Female	700	8.9	6.9 - 11.4	2,500	31.6	27.9 - 35.6	1,500	19.0	16.0 - 22.4	300	4.0	2.6 - 6.0	2,900	36.4	32.6 - 40.4	8,000

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	8.3	6.5 - 10.6	3,400	30.8	27.5 - 34.3	1,800	16.6	14.1 - 19.4	500	4.8	3.5 - 6.7	4,300	39.4	35.9 - 43.0	10,900
No	800	16.2	13.0 - 20.1	1,700	32.3	28.1 - 36.8	1,000	20.0	16.6 - 24.0	100	2.8	1.8 - 4.3	1,500	28.7	24.9 - 32.7	5,200

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,000	15.6	12.7 - 19.0	1,900	30.8	26.8 - 35.0	1,100	17.4	14.5 - 20.8	100	2.2	1.3 - 3.8	2,100	34.0	30.1 - 38.2	6,300
131-185%	300	9.1	5.8 - 14.0	1,000	29.7	23.9 - 36.1	500	16.3	12.1 - 21.7	200	5.8	3.5 - 9.5	1,300	39.1	32.9 - 45.7	3,300
186+%	200	3.0	1.7 - 5.4	1,700	33.3	28.5 - 38.4	900	17.8	14.2 - 22.1	300	5.7	3.6 - 8.8	2,000	40.2	35.2 - 45.4	5,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	600	18.7	14.5 - 23.9	900	30.7	25.5 - 36.5	500	17.1	13.3 - 21.7	100	2.8	1.2 - 6.0	900	30.7	25.6 - 36.2	3,100
\$15,001-\$25,000	300	13.6	9.1 - 19.9	600	25.8	19.9 - 32.7	400	17.5	12.5 - 23.8	100	2.4	1.1 - 5.3	1,000	40.7	33.7 - 48.1	2,500
\$25,001-\$40,000	400	11.7	8.2 - 16.4	1,000	28.4	22.7 - 34.8	700	20.1	15.5 - 25.7	200	5.3	3.1 - 8.9	1,200	34.5	28.8 - 40.7	3,400
>=\$40,001	200	3.0	1.7 - 5.2	2,200	36.1	31.6 - 40.8	1,000	16.1	13.0 - 19.8	300	4.8	3.1 - 7.4	2,400	40.0	35.4 - 44.7	6,100

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	6.4	3.6 - 11.1	400	39.9	29.0 - 52.0	200	23.3	14.5 - 35.2	<50	2.3	0.5 - 11.1	300	28.0	18.2 - 40.5	1,000
Normal birth weight	1,700	11.2	9.4 - 13.2	4,600	30.7	28.0 - 33.6	2,600	17.3	15.3 - 19.6	600	4.3	3.2 - 5.7	5,500	36.5	33.7 - 39.3	15,000

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	29.4	21.7 - 38.5	300	27.1	19.7 - 35.9	200	17.7	11.8 - 25.8	<50	1.9	0.8 - 4.8	300	23.9	17.1 - 32.4	1,200
No	1,400	9.4	7.7 - 11.3	4,600	31.8	29.0 - 34.7	2,600	17.9	15.7 - 20.3	600	4.4	3.3 - 5.9	5,300	36.5	33.7 - 39.5	14,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	17.2	13.2 - 22.0	900	30.0	24.6 - 36.0	600	19.3	14.8 - 24.7	100	3.0	1.7 - 5.4	900	30.5	25.1 - 36.4	3,100
No	1,200	9.3	7.5 - 11.5	4,000	31.6	28.6 - 34.8	2,200	17.4	15.1 - 20.0	600	4.6	3.4 - 6.2	4,700	37.1	34.0 - 40.3	12,600

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	16.5	13.5 - 20.0	2,100	30.7	26.8 - 34.9	1,200	18.6	15.5 - 22.1	200	3.1	1.9 - 4.8	2,100	31.2	27.4 - 35.3	6,700
No	600	6.8	5.1 - 9.0	2,900	31.7	28.2 - 35.4	1,600	17.2	14.6 - 20.3	500	5.0	3.5 - 7.0	3,600	39.3	35.6 - 43.1	9,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,600	10.3	9.1 - 11.7	4,600	29.7	27.6 - 31.9	3,000	19.5	17.7 - 21.4	700	4.2	3.4 - 5.3	5,700	36.2	34.0 - 38.5	15,600

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	13.7	10.8 - 17.1	400	23.1	19.3 - 27.3	300	18.8	15.3 - 22.9	100	3.7	2.2 - 5.9	700	40.8	36.2 - 45.6	1,700
Honolulu	1,200	10.2	8.7 - 12.0	3,700	31.8	29.1 - 34.5	2,200	19.3	17.1 - 21.7	500	4.5	3.4 - 5.9	4,000	34.2	31.5 - 37.1	11,600
Kauai	<50	4.5	2.1 - 9.4	100	17.0	11.8 - 23.8	200	23.4	17.4 - 30.8	<50	6.2	3.3 - 11.5	300	48.9	41.0 - 56.8	700
Maui	200	10.0	7.3 - 13.6	400	27.6	23.2 - 32.5	300	20.2	16.3 - 24.6	<50	1.9	0.9 - 4.1	600	40.2	35.2 - 45.4	1,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	7.7	5.3 - 11.1	700	20.0	16.1 - 24.5	600	16.7	13.3 - 20.6	200	5.2	3.3 - 8.3	1,700	50.4	45.3 - 55.5	3,400
Native Hawaiian	700	16.1	14.3 - 18.0	1,300	30.0	27.8 - 32.4	800	18.1	16.3 - 20.1	200	3.6	2.8 - 4.6	1,400	32.2	29.9 - 34.6	4,300
Chinese	<50	4.6	1.2 - 16.1	200	35.6	23.6 - 49.8	200	24.1	14.0 - 38.1	<50	4.6	1.2 - 16.1	200	31.0	19.5 - 45.5	600
Filipino	300	8.0	5.6 - 11.3	1,100	32.6	27.3 - 38.3	700	22.3	18.0 - 27.4	100	3.7	2.0 - 6.9	1,100	33.4	28.1 - 39.0	3,200
Japanese	100	4.4	2.1 - 8.9	900	42.4	35.1 - 50.0	400	22.0	16.3 - 29.0	100	5.7	3.0 - 10.5	500	25.5	19.5 - 32.6	2,000
Other	300	13.8	9.5 - 19.8	600	26.4	20.4 - 33.5	400	18.6	13.7 - 24.9	100	3.2	1.4 - 6.9	800	37.9	31.1 - 45.2	2,100

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	16.4	12.9 - 20.8	500	32.8	27.2 - 38.9	400	24.9	19.7 - 31.0	<50	2.2	1.2 - 4.2	400	23.6	18.8 - 29.1	1,500
20-24	500	12.0	9.6 - 14.9	1,100	27.1	23.4 - 31.2	900	23.2	19.7 - 27.2	200	4.0	2.5 - 6.4	1,400	33.6	29.7 - 37.8	4,100
25-34	700	8.9	7.3 - 10.9	2,200	29.6	26.5 - 32.9	1,200	16.3	14.0 - 18.9	400	4.7	3.3 - 6.5	3,000	40.5	37.2 - 43.9	7,500
35+	200	7.9	5.0 - 12.3	800	32.7	27.1 - 38.8	500	19.4	14.9 - 24.9	100	4.5	2.8 - 7.3	900	35.4	29.8 - 41.5	2,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	9.2	7.6 - 10.9	2,400	30.3	27.3 - 33.4	1,700	20.7	18.2 - 23.5	300	3.8	2.7 - 5.3	2,900	36.1	33.0 - 39.3	8,000
Female	900	11.6	9.7 - 13.8	2,200	29.1	26.2 - 32.2	1,400	18.2	15.8 - 20.9	400	4.7	3.4 - 6.4	2,800	36.4	33.3 - 39.6	7,600

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	7.5	6.1 - 9.1	3,100	29.9	27.2 - 32.7	2,000	18.9	16.7 - 21.3	500	4.9	3.7 - 6.5	4,000	38.8	35.9 - 41.7	10,400
No	800	16.0	13.6 - 18.8	1,500	29.4	26.1 - 32.8	1,100	20.6	17.7 - 23.9	100	2.8	1.9 - 4.0	1,600	31.2	27.9 - 34.7	5,200

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	900	14.9	12.8 - 17.3	1,700	28.1	25.0 - 31.4	1,300	21.2	18.4 - 24.3	200	2.8	1.9 - 4.0	1,900	33.0	29.7 - 36.5	5,900
131-185%	300	9.9	7.4 - 13.2	900	26.8	22.7 - 31.4	700	20.3	16.7 - 24.4	200	5.5	3.5 - 8.6	1,300	37.5	32.9 - 42.4	3,500
186+%	200	4.2	2.7 - 6.4	1,700	33.0	29.0 - 37.2	900	16.4	13.6 - 19.8	300	5.6	3.8 - 8.0	2,100	40.9	36.7 - 45.1	5,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	500	16.5	13.4 - 20.0	900	27.3	23.1 - 32.0	800	24.0	20.0 - 28.6	100	3.3	2.1 - 5.1	900	28.9	24.7 - 33.5	3,100
\$15,001-\$25,000	300	11.8	8.8 - 15.7	600	27.6	22.6 - 33.2	400	20.3	16.0 - 25.5	100	2.6	1.1 - 6.3	800	37.6	31.9 - 43.6	2,100
\$25,001-\$40,000	300	9.8	7.4 - 12.9	900	25.3	21.3 - 29.7	700	21.6	17.9 - 25.7	200	4.5	2.8 - 7.3	1,300	38.8	34.2 - 43.6	3,400
>=\$40,001	400	5.8	4.2 - 7.9	2,000	33.2	29.6 - 37.1	1,000	16.1	13.4 - 19.1	300	5.6	4.0 - 7.8	2,400	39.3	35.6 - 43.1	6,100

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	15.0	10.0 - 22.0	400	34.7	26.4 - 44.1	200	19.6	13.3 - 28.0	<50	1.9	0.7 - 4.7	300	28.8	21.2 - 37.8	1,000
Normal birth weight	1,500	10.0	8.8 - 11.4	4,300	29.3	27.1 - 31.6	2,800	19.5	17.7 - 21.5	600	4.4	3.5 - 5.6	5,400	36.8	34.5 - 39.1	14,600

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	23.5	18.6 - 29.2	400	28.1	22.1 - 35.2	300	20.8	15.7 - 27.1	100	4.3	2.3 - 7.9	300	23.2	17.8 - 29.7	1,400
No	1,200	8.9	7.7 - 10.4	4,200	29.8	27.5 - 32.1	2,700	19.3	17.4 - 21.3	600	4.3	3.4 - 5.5	5,300	37.7	35.3 - 40.1	14,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	15.5	12.7 - 18.8	800	26.1	22.2 - 30.3	700	22.1	18.4 - 26.3	100	4.0	2.4 - 6.5	1,000	32.4	27.9 - 37.2	3,100
No	1,100	9.0	7.6 - 10.6	3,600	30.3	27.8 - 32.9	2,300	18.9	16.9 - 21.2	500	4.3	3.3 - 5.6	4,500	37.5	35.0 - 40.1	12,000

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	13.5	11.6 - 15.7	2,000	29.8	26.8 - 33.0	1,300	19.2	16.6 - 21.9	300	3.8	2.6 - 5.5	2,200	33.8	30.7 - 37.0	6,700
No	700	7.9	6.3 - 9.7	2,600	29.8	26.9 - 32.8	1,700	19.6	17.2 - 22.3	400	4.6	3.4 - 6.2	3,400	38.1	35.1 - 41.3	8,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,700	12.6	11.1 - 14.4	2,900	21.5	19.5 - 23.6	3,000	22.2	20.3 - 24.3	800	5.5	4.5 - 6.7	5,200	38.1	35.8 - 40.6	13,600

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	12.7	9.6 - 16.5	400	23.3	19.2 - 28.1	300	19.6	15.8 - 24.0	100	4.8	3.0 - 7.6	600	39.6	34.6 - 44.8	1,500
Honolulu	1,300	13.2	11.3 - 15.5	2,200	21.8	19.3 - 24.5	2,200	21.7	19.4 - 24.3	600	5.7	4.5 - 7.3	3,800	37.5	34.5 - 40.5	10,200
Kauai	<50	6.9	3.3 - 13.7	100	13.1	8.1 - 20.6	100	21.6	14.9 - 30.1	<50	4.9	2.2 - 10.5	300	53.5	44.4 - 62.4	500
Maui	100	10.4	7.3 - 14.5	300	20.5	16.2 - 25.6	400	28.5	23.7 - 33.9	100	5.1	3.0 - 8.4	500	35.5	30.2 - 41.2	1,400

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	400	12.4	9.3 - 16.3	400	12.1	9.0 - 16.0	700	21.3	17.2 - 25.9	200	7.2	5.1 - 10.1	1,500	47.0	41.9 - 52.2	3,200
Native Hawaiian	600	14.7	12.8 - 16.8	900	23.1	20.9 - 25.6	900	25.3	22.9 - 27.7	200	4.9	3.8 - 6.2	1,200	32.0	29.5 - 34.7	3,700
Chinese	100	9.5	3.8 - 21.6	200	28.8	17.2 - 44.2	100	15.6	7.8 - 28.8	<50	2.7	0.4 - 16.1	200	43.4	29.9 - 57.9	500
Filipino	300	9.9	6.7 - 14.4	800	28.5	23.3 - 34.4	700	24.4	19.6 - 29.8	200	5.6	3.3 - 9.2	900	31.7	26.3 - 37.6	2,800
Japanese	100	9.1	5.1 - 15.6	400	25.3	18.4 - 33.8	300	18.4	12.5 - 26.3	100	5.7	2.8 - 11.3	600	41.5	33.2 - 50.3	1,500
Other	300	16.7	11.5 - 23.6	400	18.9	13.4 - 26.1	400	19.4	14.1 - 26.1	100	4.5	2.2 - 9.1	800	40.5	33.0 - 48.4	1,900

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	12.1	8.4 - 17.3	400	26.0	20.4 - 32.5	400	23.5	19.1 - 28.6	100	3.8	1.9 - 7.5	600	34.5	28.6 - 41.0	1,600
20-24	500	14.0	11.2 - 17.5	700	19.4	16.0 - 23.2	900	26.5	22.7 - 30.8	100	3.2	2.0 - 5.3	1,200	36.8	32.4 - 41.4	3,400
25-34	800	11.9	9.7 - 14.4	1,300	20.4	17.5 - 23.6	1,300	20.7	17.9 - 23.9	400	6.9	5.3 - 9.0	2,600	40.1	36.5 - 43.8	6,400
35+	300	13.1	9.1 - 18.5	500	24.6	19.3 - 30.7	400	18.8	14.3 - 24.5	100	6.3	4.0 - 9.7	800	37.2	31.3 - 43.5	2,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,000	14.4	12.1 - 17.0	1,500	20.9	18.2 - 23.9	1,500	21.1	18.6 - 24.0	400	5.4	4.0 - 7.1	2,700	38.2	35.0 - 41.6	7,100
Female	700	10.7	8.7 - 13.1	1,400	22.1	19.3 - 25.3	1,500	23.4	20.5 - 26.5	400	5.7	4.3 - 7.5	2,500	38.0	34.6 - 41.6	6,500

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	12.3	10.3 - 14.5	1,800	19.7	17.3 - 22.4	1,900	21.3	18.9 - 23.9	500	5.7	4.5 - 7.3	3,700	41.0	38.0 - 44.1	9,100
No	600	13.4	11.0 - 16.2	1,100	25.2	21.9 - 28.8	1,100	24.0	20.9 - 27.4	200	5.1	3.7 - 7.2	1,400	32.3	28.7 - 36.1	4,500

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	900	14.9	12.7 - 17.5	1,300	21.1	18.3 - 24.2	1,400	23.5	20.7 - 26.5	300	4.3	3.1 - 6.0	2,200	36.2	32.8 - 39.7	6,000
131-185%	200	12.1	8.6 - 16.9	400	19.9	15.0 - 26.0	400	24.3	19.2 - 30.2	100	8.0	5.1 - 12.3	600	35.6	29.7 - 41.9	1,800
186+%	400	7.7	5.5 - 10.6	1,100	22.2	18.7 - 26.1	900	19.6	16.4 - 23.3	300	6.8	4.9 - 9.2	2,100	43.8	39.6 - 48.2	4,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	500	15.2	12.4 - 18.6	700	22.6	18.9 - 26.8	800	23.6	19.9 - 27.7	100	4.5	2.9 - 6.8	1,100	34.1	29.7 - 38.9	3,300
\$15,001-\$25,000	400	16.1	12.2 - 20.9	400	18.0	13.8 - 23.2	500	23.9	19.1 - 29.5	100	4.0	2.2 - 7.0	900	37.9	32.2 - 44.0	2,200
\$25,001-\$40,000	200	9.7	6.9 - 13.4	500	21.0	16.6 - 26.1	600	23.4	19.3 - 28.1	200	7.1	4.8 - 10.6	1,000	38.8	33.5 - 44.3	2,600
>=\$40,001	400	9.2	6.9 - 12.3	1,100	22.0	18.6 - 25.8	1,000	20.6	17.3 - 24.4	300	6.5	4.7 - 8.8	2,000	41.7	37.5 - 46.0	4,800

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	14.3	9.3 - 21.4	300	27.8	19.8 - 37.7	200	23.2	16.5 - 31.5	<50	4.8	2.4 - 9.3	300	29.9	21.9 - 39.3	900
Normal birth weight	1,600	12.4	10.8 - 14.2	2,600	21.0	19.0 - 23.2	2,800	22.2	20.1 - 24.3	700	5.6	4.6 - 6.9	4,900	38.8	36.3 - 41.3	12,600

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	30.5	23.9 - 38.0	300	22.5	16.9 - 29.4	300	24.3	18.4 - 31.2	<50	2.8	1.0 - 7.5	200	20.0	14.9 - 26.3	1,200
No	1,300	10.4	8.9 - 12.2	2,600	21.5	19.3 - 23.7	2,700	22.2	20.1 - 24.4	700	5.8	4.7 - 7.1	4,800	40.1	37.5 - 42.7	12,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	19.1	15.4 - 23.5	600	19.9	16.2 - 24.0	800	25.9	21.8 - 30.6	100	4.3	2.6 - 6.9	900	30.8	26.3 - 35.7	2,900
No	1,100	10.4	8.8 - 12.3	2,200	21.9	19.5 - 24.4	2,200	21.3	19.1 - 23.7	600	5.8	4.6 - 7.2	4,200	40.6	37.8 - 43.5	10,300

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	16.1	13.7 - 18.9	1,300	21.8	19.0 - 24.9	1,300	21.3	18.7 - 24.0	300	5.4	4.0 - 7.1	2,200	35.4	32.2 - 38.9	6,100
No	700	9.5	7.6 - 11.7	1,600	21.4	18.6 - 24.4	1,700	23.0	20.2 - 26.1	400	5.6	4.2 - 7.3	2,900	40.6	37.2 - 44.0	7,300