

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (exclusive) duration 8+ weeks	Breastfeeding (exclusive) duration 8+ weeks

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/28/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	8,200	60.6	58.1 - 63.0	5,400	39.4	37.0 - 41.9	13,600
2001	9,800	62.8	60.5 - 65.0	5,800	37.2	35.0 - 39.5	15,600
2002	10,100	62.9	60.1 - 65.6	5,900	37.1	34.4 - 39.9	16,000
2003	10,600	64.5	61.8 - 67.1	5,800	35.5	32.9 - 38.2	16,400
2004	11,100	64.8	62.8 - 66.9	6,000	35.2	33.1 - 37.2	17,100
2005	10,900	64.5	62.2 - 66.7	6,000	35.5	33.3 - 37.8	16,900
2006	11,000	63.0	60.6 - 65.4	6,400	37.0	34.6 - 39.4	17,400
2007	11,300	63.4	61.0 - 65.7	6,500	36.6	34.3 - 39.0	17,800
2008	11,500	63.7	61.4 - 65.9	6,500	36.3	34.1 - 38.6	18,000
2009	11,100	62.5	59.3 - 65.5	6,700	37.5	34.5 - 40.7	17,700
2010	9,800	54.7	51.4 - 57.9	8,200	45.3	42.1 - 48.6	18,000
2011	10,000	55.3	52.0 - 58.5	8,100	44.7	41.5 - 48.0	18,000
2012	10,100	56.8	53.5 - 60.1	7,700	43.2	39.9 - 46.5	17,800
2013	8,600	49.6	46.0 - 53.2	8,700	50.4	46.8 - 54.0	17,300

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,600	49.6	46.0 - 53.2	8,700	50.4	46.8 - 54.0	17,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	50.7	45.0 - 56.5	1,100	49.3	43.5 - 55.0	2,200
Honolulu	6,100	49.2	44.4 - 54.0	6,300	50.8	46.0 - 55.6	12,500
Kauai	300	43.3	38.0 - 48.8	500	56.7	51.2 - 62.0	800
Maui	1,000	53.1	47.3 - 58.7	900	46.9	41.3 - 52.7	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,700	36.9	30.3 - 44.0	2,900	63.1	56.0 - 69.7	4,600
Native Hawaiian	2,800	54.6	48.4 - 60.8	2,300	45.4	39.2 - 51.6	5,200
Chinese	600	71.2	53.6 - 84.1	200	28.8	15.9 - 46.4	800
Filipino	1,600	59.6	50.3 - 68.3	1,100	40.4	31.7 - 49.7	2,700
Japanese	800	43.3	32.3 - 55.1	1,000	56.7	44.9 - 67.7	1,800
Other	1,100	48.8	38.7 - 59.1	1,100	51.2	40.9 - 61.3	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	53.8	37.9 - 68.9	400	46.2	31.1 - 62.1	800
20-24	1,600	40.8	33.6 - 48.4	2,300	59.2	51.6 - 66.4	3,900
25-34	4,700	49.8	44.9 - 54.7	4,800	50.2	45.3 - 55.1	9,500
35+	1,900	58.8	50.7 - 66.6	1,300	41.2	33.4 - 49.3	3,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,800	45.1	40.1 - 50.1	4,600	54.9	49.9 - 59.9	8,500
Female	4,800	53.8	48.7 - 58.9	4,100	46.2	41.1 - 51.3	8,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,200	46.9	42.3 - 51.5	5,900	53.1	48.5 - 57.7	11,000
No	3,400	54.2	48.5 - 59.8	2,900	45.8	40.2 - 51.5	6,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,200	50.4	44.6 - 56.2	3,100	49.6	43.8 - 55.4	6,300
131-185%	1,400	56.9	47.1 - 66.3	1,000	43.1	33.7 - 52.9	2,400
186+%	3,200	46.9	41.2 - 52.7	3,700	53.1	47.3 - 58.8	6,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Income (2012+)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	2,000	51.5	44.2 - 58.8	1,900	48.5	41.2 - 55.8	3,900
\$17,001-\$26,000	1,100	49.1	39.4 - 59.0	1,200	50.9	41.0 - 60.6	2,300
\$26,001-\$51,000	1,900	51.9	44.0 - 59.6	1,800	48.1	40.4 - 56.0	3,700
>=\$51,001	2,800	47.2	41.0 - 53.4	3,100	52.8	46.6 - 59.0	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	67.0	61.8 - 71.9	400	33.0	28.1 - 38.2	1,200
Normal birth weight	7,800	48.3	44.5 - 52.2	8,400	51.7	47.8 - 55.5	16,200

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	50.6	36.5 - 64.6	500	49.4	35.4 - 63.5	900
No	8,000	49.7	46.0 - 53.5	8,100	50.3	46.5 - 54.0	16,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	48.6	40.2 - 57.2	1,500	51.4	42.8 - 59.8	3,000
No	7,000	50.0	46.0 - 54.0	7,000	50.0	46.0 - 54.0	13,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,200	55.6	50.2 - 60.9	3,300	44.4	39.1 - 49.8	7,500
No	4,400	45.0	40.2 - 49.8	5,400	55.0	50.2 - 59.8	9,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,100	56.8	53.5 - 60.1	7,700	43.2	39.9 - 46.5	17,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	55.2	49.8 - 60.5	1,000	44.8	39.5 - 50.2	2,200
Honolulu	7,400	57.6	53.2 - 61.9	5,500	42.4	38.1 - 46.8	12,900
Kauai	400	47.4	40.9 - 53.9	400	52.6	46.1 - 59.1	800
Maui	1,100	57.1	52.0 - 62.1	800	42.9	37.9 - 48.0	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	43.2	36.6 - 50.0	2,300	56.8	50.0 - 63.4	4,100
Native Hawaiian	3,100	61.8	55.7 - 67.5	1,900	38.2	32.5 - 44.3	5,000
Chinese	600	54.7	40.7 - 67.9	500	45.3	32.1 - 59.3	1,100
Filipino	2,100	66.5	58.5 - 73.6	1,000	33.5	26.4 - 41.5	3,100
Japanese	900	60.1	48.6 - 70.7	600	39.9	29.3 - 51.4	1,500
Other	1,700	56.2	47.6 - 64.5	1,300	43.8	35.5 - 52.4	3,000

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	73.1	59.5 - 83.4	300	26.9	16.6 - 40.5	1,000
20-24	2,100	55.9	48.5 - 63.1	1,600	44.1	36.9 - 51.5	3,700
25-34	5,500	54.8	50.4 - 59.2	4,500	45.2	40.8 - 49.6	10,000
35+	1,800	59.1	51.3 - 66.4	1,300	40.9	33.6 - 48.7	3,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,100	57.3	52.6 - 61.9	3,800	42.7	38.1 - 47.4	8,800
Female	5,000	56.3	51.7 - 60.9	3,900	43.7	39.1 - 48.3	9,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,000	54.9	50.6 - 59.1	4,900	45.1	40.9 - 49.4	11,000
No	4,100	59.8	54.6 - 64.9	2,700	40.2	35.1 - 45.4	6,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,800	60.6	55.1 - 65.8	2,500	39.4	34.2 - 44.9	6,300
131-185%	2,000	64.8	56.6 - 72.2	1,100	35.2	27.8 - 43.4	3,000
186+%	3,300	49.1	43.8 - 54.5	3,400	50.9	45.5 - 56.2	6,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Income (2012+)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	2,500	61.6	54.5 - 68.3	1,500	38.4	31.7 - 45.5	4,000
\$17,001-\$26,000	1,200	56.7	47.5 - 65.4	900	43.3	34.6 - 52.5	2,200
\$26,001-\$51,000	2,700	60.8	54.1 - 67.2	1,700	39.2	32.8 - 45.9	4,400
>=\$51,001	2,900	50.7	44.9 - 56.4	2,800	49.3	43.6 - 55.1	5,700

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	67.2	61.6 - 72.2	400	32.8	27.8 - 38.4	1,200
Normal birth weight	9,300	56.0	52.5 - 59.5	7,300	44.0	40.5 - 47.5	16,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	68.9	53.4 - 81.1	200	31.1	18.9 - 46.6	700
No	9,300	56.6	53.1 - 60.0	7,200	43.4	40.0 - 46.9	16,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	56.0	47.7 - 64.0	1,200	44.0	36.0 - 52.3	2,700
No	8,300	57.3	53.6 - 60.9	6,200	42.7	39.1 - 46.4	14,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,000	62.2	57.3 - 66.8	3,100	37.8	33.2 - 42.7	8,100
No	5,000	52.1	47.5 - 56.6	4,600	47.9	43.4 - 52.5	9,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,000	55.3	52.0 - 58.5	8,100	44.7	41.5 - 48.0	18,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,300	56.6	51.5 - 61.7	1,000	43.4	38.3 - 48.5	2,300
Honolulu	7,300	56.4	52.0 - 60.6	5,700	43.6	39.4 - 48.0	13,000
Kauai	400	48.5	42.1 - 54.9	400	51.5	45.1 - 57.9	800
Maui	1,000	49.5	44.9 - 54.2	1,000	50.5	45.8 - 55.1	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,000	44.5	38.1 - 51.1	2,500	55.5	48.9 - 61.9	4,500
Native Hawaiian	3,200	61.3	55.5 - 66.9	2,000	38.7	33.1 - 44.5	5,200
Chinese	400	56.6	39.7 - 72.0	300	43.4	28.0 - 60.3	700
Filipino	2,100	65.8	58.2 - 72.7	1,100	34.2	27.3 - 41.8	3,200
Japanese	1,100	59.3	48.9 - 69.0	700	40.7	31.0 - 51.1	1,800
Other	1,200	46.1	37.3 - 55.3	1,500	53.9	44.7 - 62.7	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	65.7	52.7 - 76.7	300	34.3	23.3 - 47.3	1,000
20-24	2,700	63.7	56.9 - 70.0	1,600	36.3	30.0 - 43.1	4,300
25-34	5,000	51.4	47.0 - 55.9	4,700	48.6	44.1 - 53.0	9,700
35+	1,600	52.5	44.8 - 60.1	1,500	47.5	39.9 - 55.2	3,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,000	54.3	49.7 - 58.8	4,200	45.7	41.2 - 50.3	9,200
Female	5,000	56.4	51.8 - 61.0	3,800	43.6	39.0 - 48.2	8,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,800	51.4	47.3 - 55.6	5,500	48.6	44.4 - 52.7	11,300
No	4,200	61.8	56.6 - 66.8	2,600	38.2	33.2 - 43.4	6,700

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,900	61.2	55.9 - 66.3	2,500	38.8	33.7 - 44.1	6,300
131-185%	1,400	52.0	43.4 - 60.5	1,300	48.0	39.5 - 56.6	2,700
186+%	4,200	52.3	47.3 - 57.2	3,800	47.7	42.8 - 52.7	8,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	65.4	57.8 - 72.2	1,100	34.6	27.8 - 42.2	3,100
\$10,000-\$24,999	2,000	58.0	50.7 - 65.1	1,400	42.0	34.9 - 49.3	3,400
\$25,000-\$49,999	2,400	52.7	46.0 - 59.3	2,100	47.3	40.7 - 54.0	4,500
>=\$50,000	3,100	51.4	45.7 - 57.0	3,000	48.6	43.0 - 54.3	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	67.7	62.3 - 72.7	400	32.3	27.3 - 37.7	1,200
Normal birth weight	9,200	54.4	50.9 - 57.8	7,700	45.6	42.2 - 49.1	16,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	75.5	63.7 - 84.4	200	24.5	15.6 - 36.3	900
No	9,200	54.1	50.7 - 57.5	7,800	45.9	42.5 - 49.3	17,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	61.7	54.4 - 68.5	1,300	38.3	31.5 - 45.6	3,500
No	7,700	53.6	49.9 - 57.3	6,700	46.4	42.7 - 50.1	14,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,800	60.0	55.2 - 64.6	3,200	40.0	35.4 - 44.8	8,100
No	5,100	51.7	47.2 - 56.1	4,800	48.3	43.9 - 52.8	9,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,800	54.7	51.4 - 57.9	8,200	45.3	42.1 - 48.6	18,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	53.0	47.9 - 58.1	1,100	47.0	41.9 - 52.1	2,300
Honolulu	7,300	55.7	51.4 - 59.9	5,800	44.3	40.1 - 48.6	13,000
Kauai	400	47.4	41.5 - 53.5	400	52.6	46.5 - 58.5	800
Maui	1,000	52.7	47.8 - 57.5	900	47.3	42.5 - 52.2	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	43.1	36.7 - 49.7	2,500	56.9	50.3 - 63.3	4,400
Native Hawaiian	3,100	56.9	51.2 - 62.4	2,300	43.1	37.6 - 48.8	5,400
Chinese	300	38.4	24.1 - 55.1	500	61.6	44.9 - 75.9	700
Filipino	2,300	71.2	63.9 - 77.6	900	28.8	22.4 - 36.1	3,200
Japanese	900	55.4	44.6 - 65.8	700	44.6	34.2 - 55.4	1,700
Other	1,400	54.0	44.9 - 62.7	1,200	46.0	37.3 - 55.1	2,600

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	63.0	51.1 - 73.5	400	37.0	26.5 - 48.9	1,100
20-24	2,400	54.8	48.1 - 61.4	2,000	45.2	38.6 - 51.9	4,300
25-34	4,900	53.2	48.7 - 57.6	4,300	46.8	42.4 - 51.3	9,300
35+	1,800	56.0	48.3 - 63.3	1,400	44.0	36.7 - 51.7	3,300

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,000	54.9	50.4 - 59.3	4,100	45.1	40.7 - 49.6	9,100
Female	4,900	54.5	49.8 - 59.1	4,100	45.5	40.9 - 50.2	8,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,800	51.5	47.3 - 55.7	5,400	48.5	44.3 - 52.7	11,200
No	4,100	60.0	54.9 - 64.8	2,700	40.0	35.2 - 45.1	6,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,700	58.7	53.4 - 63.9	2,600	41.3	36.1 - 46.6	6,200
131-185%	1,600	56.3	48.0 - 64.2	1,200	43.7	35.8 - 52.0	2,900
186+%	3,900	50.1	45.1 - 55.1	3,800	49.9	44.9 - 54.9	7,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	60.3	53.1 - 67.1	1,300	39.7	32.9 - 46.9	3,300
\$10,000-\$24,999	1,800	57.7	50.0 - 65.1	1,300	42.3	34.9 - 50.0	3,200
\$25,000-\$49,999	2,400	54.4	47.8 - 60.8	2,000	45.6	39.2 - 52.2	4,400
>=\$50,000	3,000	50.0	44.3 - 55.6	3,000	50.0	44.4 - 55.7	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	64.0	58.5 - 69.1	400	36.0	30.9 - 41.5	1,200
Normal birth weight	9,100	54.0	50.6 - 57.4	7,700	46.0	42.6 - 49.4	16,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	60.4	48.3 - 71.3	600	39.6	28.7 - 51.7	1,400
No	8,900	54.5	51.2 - 57.9	7,500	45.5	42.1 - 48.8	16,400

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	62.5	55.6 - 69.0	1,400	37.5	31.0 - 44.4	3,800
No	7,400	53.0	49.3 - 56.6	6,600	47.0	43.4 - 50.7	14,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,500	57.7	52.9 - 62.3	3,300	42.3	37.7 - 47.1	7,800
No	5,300	52.9	48.4 - 57.3	4,700	47.1	42.7 - 51.6	10,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,100	62.5	59.3 - 65.5	6,700	37.5	34.5 - 40.7	17,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,400	61.3	56.2 - 66.2	900	38.7	33.8 - 43.8	2,300
Honolulu	8,000	63.9	59.6 - 67.9	4,500	36.1	32.1 - 40.4	12,600
Kauai	400	52.2	46.1 - 58.3	400	47.8	41.7 - 53.9	800
Maui	1,200	59.1	54.4 - 63.6	800	40.9	36.4 - 45.6	2,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	47.5	41.2 - 53.9	2,100	52.5	46.1 - 58.8	4,100
Native Hawaiian	3,900	68.9	63.4 - 73.9	1,800	31.1	26.1 - 36.6	5,700
Chinese	500	67.4	50.6 - 80.6	300	32.6	19.4 - 49.4	800
Filipino	2,200	68.6	61.2 - 75.2	1,000	31.4	24.8 - 38.8	3,200
Japanese	1,000	63.3	52.7 - 72.8	600	36.7	27.2 - 47.3	1,600
Other	1,500	62.3	53.3 - 70.5	900	37.7	29.5 - 46.7	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	66.4	55.6 - 75.7	600	33.6	24.3 - 44.4	1,800
20-24	2,700	67.7	61.2 - 73.6	1,300	32.3	26.4 - 38.8	4,000
25-34	5,200	58.2	53.7 - 62.5	3,700	41.8	37.5 - 46.3	9,000
35+	1,900	65.8	58.3 - 72.6	1,000	34.2	27.4 - 41.7	2,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,000	63.9	59.5 - 68.1	3,400	36.1	31.9 - 40.5	9,400
Female	5,100	60.8	56.3 - 65.2	3,300	39.2	34.8 - 43.7	8,300

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,800	55.2	51.0 - 59.3	4,700	44.8	40.7 - 49.0	10,500
No	5,300	73.0	68.4 - 77.1	2,000	27.0	22.9 - 31.6	7,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,200	67.6	62.3 - 72.5	2,000	32.4	27.5 - 37.7	6,200
131-185%	1,800	65.1	57.2 - 72.3	1,000	34.9	27.7 - 42.8	2,800
186+%	4,000	56.3	51.3 - 61.1	3,100	43.7	38.9 - 48.7	7,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	68.4	60.5 - 75.3	900	31.6	24.7 - 39.5	2,900
\$10,000-\$24,999	2,700	69.9	63.3 - 75.8	1,100	30.1	24.2 - 36.7	3,800
\$25,000-\$49,999	2,100	58.3	51.3 - 65.0	1,500	41.7	35.0 - 48.7	3,600
>=\$50,000	3,400	56.6	51.1 - 62.0	2,600	43.4	38.0 - 48.9	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	71.4	66.2 - 76.0	300	28.6	24.0 - 33.8	1,200
Normal birth weight	10,200	61.8	58.5 - 65.0	6,300	38.2	35.0 - 41.5	16,500

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	73.0	62.7 - 81.3	400	27.0	18.7 - 37.3	1,600
No	9,600	61.3	58.0 - 64.6	6,100	38.7	35.4 - 42.0	15,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	68.7	62.2 - 74.6	1,200	31.3	25.4 - 37.8	3,900
No	8,200	60.6	57.0 - 64.2	5,300	39.4	35.8 - 43.0	13,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,800	66.2	61.4 - 70.7	2,400	33.8	29.3 - 38.6	7,200
No	6,000	59.9	55.6 - 64.0	4,000	40.1	36.0 - 44.4	10,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,500	63.7	61.4 - 65.9	6,500	36.3	34.1 - 38.6	18,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,400	59.4	52.5 - 66.0	1,000	40.6	34.0 - 47.5	2,400
Honolulu	8,200	64.5	61.8 - 67.2	4,500	35.5	32.8 - 38.2	12,700
Kauai	500	60.4	49.2 - 70.7	300	39.6	29.3 - 50.8	800
Maui	1,400	64.7	57.6 - 71.2	700	35.3	28.8 - 42.4	2,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,100	49.2	44.3 - 54.0	2,200	50.8	46.0 - 55.7	4,300
Native Hawaiian	3,400	67.1	62.3 - 71.6	1,600	32.9	28.4 - 37.7	5,000
Chinese	400	67.7	61.1 - 73.7	200	32.3	26.3 - 38.9	600
Filipino	2,400	72.7	67.7 - 77.2	900	27.3	22.8 - 32.3	3,200
Japanese	1,400	69.4	62.7 - 75.4	600	30.6	24.6 - 37.3	2,000
Other	1,800	64.3	58.2 - 70.0	1,000	35.7	30.0 - 41.8	2,900

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	61.8	52.7 - 70.1	500	38.2	29.9 - 47.3	1,400
20-24	2,600	66.9	61.8 - 71.7	1,300	33.1	28.3 - 38.2	4,000
25-34	5,700	61.3	58.0 - 64.4	3,600	38.7	35.6 - 42.0	9,300
35+	2,300	67.4	62.4 - 72.1	1,100	32.6	27.9 - 37.6	3,300

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,000	63.5	60.3 - 66.6	3,400	36.5	33.4 - 39.7	9,400
Female	5,500	63.9	60.5 - 67.2	3,100	36.1	32.8 - 39.5	8,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,900	60.8	57.9 - 63.6	4,400	39.2	36.4 - 42.1	11,300
No	4,600	68.6	64.6 - 72.3	2,100	31.4	27.7 - 35.4	6,700

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,500	67.7	63.8 - 71.5	2,100	32.3	28.5 - 36.2	6,600
131-185%	1,600	61.0	54.7 - 66.9	1,000	39.0	33.1 - 45.3	2,600
186+%	4,600	60.0	56.5 - 63.4	3,100	40.0	36.6 - 43.5	7,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,100	70.3	64.3 - 75.6	900	29.7	24.4 - 35.7	3,000
\$10,000-\$24,999	2,500	64.3	59.0 - 69.3	1,400	35.7	30.7 - 41.0	3,900
\$25,000-\$49,999	2,700	64.4	59.6 - 69.0	1,500	35.6	31.0 - 40.4	4,100
>=\$50,000	3,600	58.6	54.7 - 62.4	2,500	41.4	37.6 - 45.3	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	75.5	66.7 - 82.5	300	24.5	17.5 - 33.3	1,200
Normal birth weight	10,600	62.8	60.4 - 65.2	6,200	37.2	34.8 - 39.6	16,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	78.3	70.2 - 84.7	300	21.7	15.3 - 29.8	1,500
No	10,100	62.6	60.1 - 64.9	6,100	37.4	35.1 - 39.9	16,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	72.7	67.3 - 77.6	900	27.3	22.4 - 32.7	3,300
No	8,900	61.8	59.3 - 64.4	5,500	38.2	35.6 - 40.7	14,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,100	67.9	64.2 - 71.4	2,400	32.1	28.6 - 35.8	7,600
No	6,300	61.0	58.0 - 64.0	4,000	39.0	36.0 - 42.0	10,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,300	63.4	61.0 - 65.7	6,500	36.6	34.3 - 39.0	17,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,300	53.7	46.9 - 60.4	1,100	46.3	39.6 - 53.1	2,300
Honolulu	8,100	66.0	63.2 - 68.6	4,200	34.0	31.4 - 36.8	12,300
Kauai	600	55.0	44.6 - 65.0	500	45.0	35.0 - 55.4	1,000
Maui	1,300	63.0	55.7 - 69.7	800	37.0	30.3 - 44.3	2,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,100	50.7	45.7 - 55.7	2,000	49.3	44.3 - 54.3	4,100
Native Hawaiian	3,300	66.7	61.8 - 71.2	1,600	33.3	28.8 - 38.2	4,900
Chinese	400	62.0	55.3 - 68.3	300	38.0	31.7 - 44.7	700
Filipino	2,300	73.0	68.0 - 77.4	900	27.0	22.6 - 32.0	3,200
Japanese	1,500	73.0	66.3 - 78.7	600	27.0	21.3 - 33.7	2,100
Other	1,700	58.4	52.1 - 64.4	1,200	41.6	35.6 - 47.9	2,900

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	68.8	59.5 - 76.7	500	31.2	23.3 - 40.5	1,600
20-24	2,900	64.0	59.1 - 68.6	1,600	36.0	31.4 - 40.9	4,500
25-34	5,400	61.8	58.5 - 65.0	3,300	38.2	35.0 - 41.5	8,700
35+	1,900	64.2	58.8 - 69.3	1,100	35.8	30.7 - 41.2	3,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,800	63.1	59.8 - 66.2	3,400	36.9	33.8 - 40.2	9,200
Female	5,500	63.7	60.3 - 67.0	3,100	36.3	33.0 - 39.7	8,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,900	61.8	58.9 - 64.6	4,300	38.2	35.4 - 41.1	11,100
No	4,400	66.0	62.0 - 69.9	2,200	34.0	30.1 - 38.0	6,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,300	66.1	62.1 - 69.9	2,200	33.9	30.1 - 37.9	6,600
131-185%	1,700	63.3	57.1 - 69.0	1,000	36.7	31.0 - 42.9	2,600
186+%	4,500	61.6	58.0 - 65.0	2,800	38.4	35.0 - 42.0	7,300

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,900	63.7	57.4 - 69.6	1,100	36.3	30.4 - 42.6	3,000
\$10,000-\$24,999	2,400	67.2	61.9 - 72.1	1,200	32.8	27.9 - 38.1	3,600
\$25,000-\$49,999	2,600	64.4	59.4 - 69.0	1,500	35.6	31.0 - 40.6	4,100
>=\$50,000	3,600	61.2	57.3 - 65.0	2,300	38.8	35.0 - 42.7	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	74.3	66.1 - 81.0	400	25.7	19.0 - 33.9	1,400
Normal birth weight	10,200	62.4	60.0 - 64.8	6,100	37.6	35.2 - 40.0	16,300

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	72.9	64.4 - 79.9	400	27.1	20.1 - 35.6	1,500
No	10,000	62.6	60.2 - 65.0	6,000	37.4	35.0 - 39.8	16,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	72.1	66.8 - 76.8	1,000	27.9	23.2 - 33.2	3,500
No	8,600	61.4	58.7 - 63.9	5,400	38.6	36.1 - 41.3	14,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,600	62.7	58.9 - 66.4	2,700	37.3	33.6 - 41.1	7,300
No	6,400	63.5	60.5 - 66.4	3,700	36.5	33.6 - 39.5	10,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,000	63.0	60.6 - 65.4	6,400	37.0	34.6 - 39.4	17,400

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	57.9	50.4 - 65.0	900	42.1	35.0 - 49.6	2,100
Honolulu	8,300	65.3	62.5 - 68.0	4,400	34.7	32.0 - 37.5	12,700
Kauai	400	48.6	37.0 - 60.4	400	51.4	39.6 - 63.0	800
Maui	1,100	59.5	51.5 - 67.0	700	40.5	33.0 - 48.5	1,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,100	50.3	45.3 - 55.4	2,100	49.7	44.6 - 54.7	4,200
Native Hawaiian	3,200	68.4	63.4 - 73.0	1,500	31.6	27.0 - 36.6	4,700
Chinese	400	68.3	61.3 - 74.7	200	31.7	25.3 - 38.7	600
Filipino	2,100	66.3	60.9 - 71.3	1,100	33.7	28.7 - 39.1	3,200
Japanese	1,400	70.3	63.3 - 76.5	600	29.7	23.5 - 36.7	2,000
Other	1,700	63.2	56.7 - 69.2	1,000	36.8	30.8 - 43.3	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	66.0	56.7 - 74.1	500	34.0	25.9 - 43.3	1,500
20-24	2,900	68.2	63.1 - 73.0	1,300	31.8	27.0 - 36.9	4,200
25-34	5,100	59.8	56.3 - 63.2	3,400	40.2	36.8 - 43.7	8,500
35+	2,000	63.5	58.1 - 68.6	1,200	36.5	31.4 - 41.9	3,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,800	65.5	62.2 - 68.8	3,100	34.5	31.2 - 37.8	8,900
Female	5,100	60.4	56.9 - 63.9	3,400	39.6	36.1 - 43.1	8,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,700	60.6	57.6 - 63.5	4,400	39.4	36.5 - 42.4	11,100
No	4,300	67.3	63.1 - 71.3	2,100	32.7	28.7 - 36.9	6,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,100	67.4	63.2 - 71.4	2,000	32.6	28.6 - 36.8	6,100
131-185%	1,400	62.3	55.3 - 68.9	900	37.7	31.1 - 44.7	2,300
186+%	4,700	59.2	55.7 - 62.6	3,200	40.8	37.4 - 44.3	7,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	66.6	60.3 - 72.4	1,000	33.4	27.6 - 39.7	2,900
\$10,000-\$24,999	2,500	67.7	62.3 - 72.7	1,200	32.3	27.3 - 37.7	3,600
\$25,000-\$49,999	2,600	63.6	58.5 - 68.4	1,500	36.4	31.6 - 41.5	4,100
>=\$50,000	3,200	56.8	52.7 - 60.9	2,400	43.2	39.1 - 47.3	5,600

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	66.7	57.9 - 74.5	500	33.3	25.5 - 42.1	1,400
Normal birth weight	10,100	62.8	60.2 - 65.2	6,000	37.2	34.8 - 39.8	16,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	73.4	65.0 - 80.3	400	26.6	19.7 - 35.0	1,600
No	9,600	61.9	59.3 - 64.4	5,900	38.1	35.6 - 40.7	15,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	70.0	64.5 - 75.0	1,100	30.0	25.0 - 35.5	3,600
No	8,300	61.1	58.3 - 63.7	5,300	38.9	36.3 - 41.7	13,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,900	66.7	62.8 - 70.3	2,400	33.3	29.7 - 37.2	7,300
No	5,900	59.9	56.7 - 63.0	3,900	40.1	37.0 - 43.3	9,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,900	64.5	62.2 - 66.7	6,000	35.5	33.3 - 37.8	16,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,400	62.2	55.3 - 68.6	800	37.8	31.4 - 44.7	2,200
Honolulu	8,300	67.1	64.5 - 69.7	4,100	32.9	30.3 - 35.5	12,400
Kauai	300	46.1	34.8 - 57.9	400	53.9	42.1 - 65.2	700
Maui	900	55.2	47.4 - 62.7	700	44.8	37.3 - 52.6	1,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	48.0	42.8 - 53.2	1,900	52.0	46.8 - 57.2	3,700
Native Hawaiian	3,200	66.5	61.6 - 71.0	1,600	33.5	29.0 - 38.4	4,800
Chinese	400	67.1	60.7 - 72.8	200	32.9	27.2 - 39.3	600
Filipino	2,400	75.6	70.9 - 79.7	800	24.4	20.3 - 29.1	3,100
Japanese	1,600	69.0	62.9 - 74.6	700	31.0	25.4 - 37.1	2,300
Other	1,600	66.5	60.3 - 72.2	800	33.5	27.8 - 39.7	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	68.4	60.0 - 75.8	500	31.6	24.2 - 40.0	1,500
20-24	2,800	67.6	62.8 - 72.1	1,400	32.4	27.9 - 37.2	4,200
25-34	5,300	63.3	60.0 - 66.5	3,100	36.7	33.5 - 40.0	8,400
35+	1,800	61.3	55.9 - 66.5	1,100	38.7	33.5 - 44.1	2,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,600	64.7	61.4 - 67.8	3,000	35.3	32.2 - 38.6	8,600
Female	5,300	64.3	60.9 - 67.4	3,000	35.7	32.6 - 39.1	8,300

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,600	61.2	58.3 - 64.0	4,200	38.8	36.0 - 41.7	10,800
No	4,300	70.2	66.4 - 73.9	1,800	29.8	26.1 - 33.6	6,100

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,300	69.4	65.6 - 73.0	1,900	30.6	27.0 - 34.4	6,200
131-185%	1,200	64.2	56.8 - 71.0	600	35.8	29.0 - 43.2	1,800
186+%	4,500	59.7	56.3 - 63.1	3,000	40.3	36.9 - 43.7	7,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,100	71.2	65.6 - 76.3	900	28.8	23.7 - 34.4	3,000
\$10,000-\$24,999	2,500	69.0	64.0 - 73.6	1,100	31.0	26.4 - 36.0	3,600
\$25,000-\$49,999	2,200	57.3	52.3 - 62.2	1,700	42.7	37.8 - 47.7	3,900
>=\$50,000	3,300	62.6	58.6 - 66.5	2,000	37.4	33.5 - 41.4	5,300

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	71.1	62.2 - 78.6	300	28.9	21.4 - 37.8	1,200
Normal birth weight	10,000	63.9	61.6 - 66.3	5,700	36.1	33.7 - 38.4	15,700

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	71.4	63.0 - 78.6	400	28.6	21.4 - 37.0	1,400
No	9,700	63.5	61.1 - 65.9	5,600	36.5	34.1 - 38.9	15,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	71.5	66.4 - 76.1	1,000	28.5	23.9 - 33.6	3,600
No	8,100	62.3	59.6 - 64.8	4,900	37.7	35.2 - 40.4	13,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,800	65.7	62.1 - 69.2	2,500	34.3	30.8 - 37.9	7,200
No	5,900	63.0	60.0 - 65.9	3,500	37.0	34.1 - 40.0	9,400

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,100	64.8	62.8 - 66.9	6,000	35.2	33.1 - 37.2	17,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,300	60.0	53.5 - 66.1	900	40.0	33.9 - 46.5	2,100
Honolulu	8,300	67.0	64.5 - 69.3	4,100	33.0	30.7 - 35.5	12,400
Kauai	400	45.8	35.9 - 56.1	400	54.2	43.9 - 64.1	800
Maui	1,200	64.1	57.4 - 70.3	600	35.9	29.7 - 42.6	1,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	46.3	41.8 - 51.0	2,000	53.7	49.0 - 58.2	3,800
Native Hawaiian	3,500	73.8	69.6 - 77.6	1,200	26.2	22.4 - 30.4	4,700
Chinese	400	66.1	61.7 - 70.3	200	33.9	29.7 - 38.3	700
Filipino	2,200	68.9	64.4 - 73.0	1,000	31.1	27.0 - 35.6	3,200
Japanese	1,600	71.3	65.3 - 76.7	600	28.7	23.3 - 34.7	2,200
Other	1,600	64.7	58.7 - 70.2	900	35.3	29.8 - 41.3	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	70.6	62.2 - 77.7	400	29.4	22.3 - 37.8	1,200
20-24	3,000	67.5	63.2 - 71.5	1,400	32.5	28.5 - 36.8	4,400
25-34	5,200	62.5	59.5 - 65.5	3,100	37.5	34.5 - 40.5	8,300
35+	2,100	65.0	60.2 - 69.5	1,100	35.0	30.5 - 39.8	3,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,200	66.4	63.5 - 69.1	3,100	33.6	30.9 - 36.5	9,300
Female	4,900	63.0	59.8 - 66.1	2,900	37.0	33.9 - 40.2	7,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,500	62.5	60.0 - 64.9	4,500	37.5	35.1 - 40.0	12,000
No	3,600	70.3	66.3 - 73.9	1,500	29.7	26.1 - 33.7	5,100

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,300	68.1	64.5 - 71.5	2,000	31.9	28.5 - 35.5	6,300
131-185%	1,200	60.2	53.8 - 66.2	800	39.8	33.8 - 46.2	2,100
186+%	4,700	62.7	59.5 - 65.7	2,800	37.3	34.3 - 40.5	7,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,900	71.7	66.2 - 76.7	700	28.3	23.3 - 33.8	2,600
\$10,000-\$24,999	2,800	65.3	60.9 - 69.4	1,500	34.7	30.6 - 39.1	4,200
\$25,000-\$49,999	2,700	60.3	56.0 - 64.4	1,700	39.7	35.6 - 44.0	4,400
>=\$50,000	3,000	63.3	59.4 - 67.1	1,700	36.7	32.9 - 40.6	4,700

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	77.8	69.6 - 84.3	200	22.2	15.7 - 30.4	1,000
Normal birth weight	10,300	64.0	61.8 - 66.1	5,800	36.0	33.9 - 38.2	16,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	74.0	66.2 - 80.6	400	26.0	19.4 - 33.8	1,400
No	10,000	64.1	61.9 - 66.2	5,600	35.9	33.8 - 38.1	15,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	69.7	64.8 - 74.1	1,100	30.3	25.9 - 35.2	3,500
No	8,600	63.6	61.3 - 65.9	4,900	36.4	34.1 - 38.7	13,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,000	66.1	62.8 - 69.2	2,600	33.9	30.8 - 37.2	7,600
No	5,900	63.7	60.9 - 66.4	3,300	36.3	33.6 - 39.1	9,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,600	64.5	61.8 - 67.1	5,800	35.5	32.9 - 38.2	16,400

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	57.7	53.0 - 62.3	800	42.3	37.7 - 47.0	1,900
Honolulu	8,100	67.1	63.6 - 70.4	4,000	32.9	29.6 - 36.4	12,000
Kauai	400	50.4	42.9 - 57.9	400	49.6	42.1 - 57.1	800
Maui	1,000	59.9	54.8 - 64.8	700	40.1	35.2 - 45.2	1,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	52.1	45.7 - 58.4	1,700	47.9	41.6 - 54.3	3,600
Native Hawaiian	2,900	65.0	62.0 - 67.8	1,600	35.0	32.2 - 38.0	4,500
Chinese	400	63.1	47.2 - 76.6	300	36.9	23.4 - 52.8	700
Filipino	2,500	70.4	64.1 - 76.0	1,100	29.6	24.0 - 35.9	3,600
Japanese	1,300	67.5	58.3 - 75.5	600	32.5	24.5 - 41.7	1,900
Other	1,600	71.7	63.0 - 78.9	600	28.3	21.1 - 37.0	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	72.0	64.3 - 78.7	400	28.0	21.3 - 35.7	1,500
20-24	2,500	66.4	61.0 - 71.3	1,300	33.6	28.7 - 39.0	3,800
25-34	5,200	63.9	60.0 - 67.6	3,000	36.1	32.4 - 40.0	8,200
35+	1,700	59.7	52.8 - 66.3	1,100	40.3	33.7 - 47.2	2,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,700	65.3	61.7 - 68.8	3,000	34.7	31.2 - 38.3	8,700
Female	4,900	63.5	59.5 - 67.3	2,800	36.5	32.7 - 40.5	7,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,600	60.8	57.3 - 64.3	4,200	39.2	35.7 - 42.7	10,800
No	4,000	71.5	67.6 - 75.0	1,600	28.5	25.0 - 32.4	5,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,100	69.6	65.8 - 73.1	2,200	30.4	26.9 - 34.2	7,300
131-185%	1,500	63.1	55.9 - 69.9	900	36.9	30.1 - 44.1	2,400
186+%	2,900	57.1	51.9 - 62.1	2,200	42.9	37.9 - 48.1	5,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,900	74.9	70.1 - 79.1	1,000	25.1	20.9 - 29.9	3,900
\$15,001-\$25,000	1,300	62.7	55.2 - 69.6	800	37.3	30.4 - 44.8	2,100
\$25,001-\$40,000	1,900	60.5	53.9 - 66.7	1,200	39.5	33.3 - 46.1	3,100
>=\$40,001	3,700	59.8	55.1 - 64.2	2,500	40.2	35.8 - 44.9	6,200

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	72.8	63.2 - 80.6	300	27.2	19.4 - 36.8	1,100
Normal birth weight	9,800	63.9	61.1 - 66.6	5,500	36.1	33.4 - 38.9	15,300

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	77.4	69.4 - 83.8	300	22.6	16.2 - 30.6	1,500
No	9,300	63.4	60.5 - 66.2	5,300	36.6	33.8 - 39.5	14,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	73.6	68.3 - 78.3	900	26.4	21.7 - 31.7	3,400
No	7,800	61.9	58.8 - 65.0	4,800	38.1	35.0 - 41.2	12,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,800	69.1	65.3 - 72.7	2,100	30.9	27.3 - 34.7	6,900
No	5,700	60.9	57.2 - 64.6	3,600	39.1	35.4 - 42.8	9,300

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,100	62.9	60.1 - 65.6	5,900	37.1	34.4 - 39.9	16,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	62.9	58.3 - 67.2	700	37.1	32.8 - 41.7	1,900
Honolulu	7,600	64.1	60.4 - 67.5	4,300	35.9	32.5 - 39.6	11,800
Kauai	300	50.7	42.8 - 58.6	300	49.3	41.4 - 57.2	600
Maui	1,000	59.6	54.5 - 64.6	700	40.4	35.4 - 45.5	1,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	46.0	39.4 - 52.8	1,800	54.0	47.2 - 60.6	3,400
Native Hawaiian	3,000	67.9	65.0 - 70.8	1,400	32.1	29.2 - 35.0	4,400
Chinese	500	80.9	64.9 - 90.7	100	19.1	9.3 - 35.1	600
Filipino	2,100	66.9	60.1 - 73.1	1,100	33.1	26.9 - 39.9	3,200
Japanese	1,600	68.6	60.3 - 75.8	700	31.4	24.2 - 39.7	2,300
Other	1,300	62.1	52.6 - 70.8	800	37.9	29.2 - 47.4	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	78.6	70.8 - 84.7	300	21.4	15.3 - 29.2	1,500
20-24	2,500	63.2	57.8 - 68.3	1,500	36.8	31.7 - 42.2	4,000
25-34	4,900	60.2	56.1 - 64.2	3,200	39.8	35.8 - 43.9	8,100
35+	1,600	62.1	54.7 - 68.9	1,000	37.9	31.1 - 45.3	2,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,100	63.8	59.9 - 67.6	2,900	36.2	32.4 - 40.1	8,100
Female	5,000	62.0	57.9 - 65.9	3,000	38.0	34.1 - 42.1	8,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,400	59.2	55.5 - 62.7	4,400	40.8	37.3 - 44.5	10,900
No	3,700	70.8	66.7 - 74.6	1,500	29.2	25.4 - 33.3	5,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,100	65.3	61.1 - 69.3	2,200	34.7	30.7 - 38.9	6,300
131-185%	1,900	58.7	52.0 - 65.0	1,400	41.3	35.0 - 48.0	3,300
186+%	3,000	58.4	53.2 - 63.5	2,100	41.6	36.5 - 46.8	5,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,100	68.0	62.2 - 73.2	1,000	32.0	26.8 - 37.8	3,100
\$15,001-\$25,000	1,500	59.3	51.9 - 66.3	1,000	40.7	33.7 - 48.1	2,500
\$25,001-\$40,000	2,200	64.1	57.9 - 69.9	1,200	35.9	30.1 - 42.1	3,400
>=\$40,001	3,600	58.4	53.7 - 63.1	2,500	41.6	36.9 - 46.3	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	72.0	59.5 - 81.8	300	28.0	18.2 - 40.5	1,000
Normal birth weight	9,400	62.3	59.4 - 65.1	5,700	37.7	34.9 - 40.6	15,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	76.1	67.6 - 82.9	300	23.9	17.1 - 32.4	1,200
No	9,100	62.2	59.2 - 65.1	5,500	37.8	34.9 - 40.8	14,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	68.8	62.8 - 74.2	1,000	31.2	25.8 - 37.2	3,100
No	7,700	61.6	58.4 - 64.8	4,800	38.4	35.2 - 41.6	12,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,600	68.0	63.8 - 71.8	2,200	32.0	28.2 - 36.2	6,700
No	5,500	59.3	55.5 - 63.0	3,800	40.7	37.0 - 44.5	9,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,800	62.8	60.5 - 65.0	5,800	37.2	35.0 - 39.5	15,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	58.8	54.0 - 63.5	700	41.2	36.5 - 46.0	1,700
Honolulu	7,500	64.6	61.8 - 67.4	4,100	35.4	32.6 - 38.2	11,600
Kauai	300	49.2	41.3 - 57.1	400	50.8	42.9 - 58.7	700
Maui	1,000	59.6	54.3 - 64.6	600	40.4	35.4 - 45.7	1,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	48.6	43.5 - 53.6	1,700	51.4	46.4 - 56.5	3,400
Native Hawaiian	2,900	67.2	64.9 - 69.5	1,400	32.8	30.5 - 35.1	4,300
Chinese	400	69.0	54.5 - 80.5	200	31.0	19.5 - 45.5	600
Filipino	2,100	64.6	58.8 - 69.9	1,100	35.4	30.1 - 41.2	3,200
Japanese	1,500	73.8	66.6 - 79.8	500	26.2	20.2 - 33.4	2,000
Other	1,300	61.4	54.0 - 68.3	800	38.6	31.7 - 46.0	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	76.4	70.9 - 81.2	400	23.6	18.8 - 29.1	1,500
20-24	2,700	65.7	61.5 - 69.7	1,400	34.3	30.3 - 38.5	4,100
25-34	4,400	58.3	54.8 - 61.6	3,100	41.7	38.4 - 45.2	7,500
35+	1,600	63.1	57.1 - 68.8	900	36.9	31.2 - 42.9	2,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,100	62.8	59.6 - 65.9	3,000	37.2	34.1 - 40.4	8,000
Female	4,800	62.7	59.5 - 65.8	2,800	37.3	34.2 - 40.5	7,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,200	59.9	57.0 - 62.7	4,200	40.1	37.3 - 43.0	10,400
No	3,600	68.5	65.0 - 71.9	1,600	31.5	28.1 - 35.0	5,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,900	66.4	62.9 - 69.7	2,000	33.6	30.3 - 37.1	5,900
131-185%	2,100	60.3	55.3 - 65.0	1,400	39.7	35.0 - 44.7	3,500
186+%	3,000	58.3	54.1 - 62.4	2,200	41.7	37.6 - 45.9	5,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,200	70.8	66.2 - 75.0	900	29.2	25.0 - 33.8	3,100
\$15,001-\$25,000	1,300	62.1	56.1 - 67.8	800	37.9	32.2 - 43.9	2,100
\$25,001-\$40,000	2,100	60.4	55.6 - 65.1	1,400	39.6	34.9 - 44.4	3,400
>=\$40,001	3,600	58.9	55.0 - 62.7	2,500	41.1	37.3 - 45.0	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	70.1	61.1 - 77.8	300	29.9	22.2 - 38.9	1,000
Normal birth weight	9,100	62.2	59.9 - 64.5	5,500	37.8	35.5 - 40.1	14,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	76.1	69.6 - 81.6	300	23.9	18.4 - 30.4	1,400
No	8,600	61.2	58.8 - 63.6	5,400	38.8	36.4 - 41.2	14,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	66.4	61.6 - 70.9	1,100	33.6	29.1 - 38.4	3,100
No	7,400	61.5	58.9 - 64.1	4,600	38.5	35.9 - 41.1	12,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,300	65.4	62.1 - 68.5	2,300	34.6	31.5 - 37.9	6,700
No	5,400	60.8	57.6 - 63.8	3,500	39.2	36.2 - 42.4	8,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,200	60.6	58.1 - 63.0	5,400	39.4	37.0 - 41.9	13,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	900	59.1	53.9 - 64.2	600	40.9	35.8 - 46.1	1,500
Honolulu	6,200	61.2	58.1 - 64.1	3,900	38.8	35.9 - 41.9	10,200
Kauai	300	46.5	37.6 - 55.6	300	53.5	44.4 - 62.4	500
Maui	900	63.1	57.3 - 68.5	500	36.9	31.5 - 42.7	1,400

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,700	51.8	46.6 - 57.0	1,600	48.2	43.0 - 53.4	3,200
Native Hawaiian	2,500	66.5	63.9 - 69.1	1,300	33.5	30.9 - 36.1	3,700
Chinese	300	56.6	42.1 - 70.1	200	43.4	29.9 - 57.9	500
Filipino	1,900	67.3	61.3 - 72.7	900	32.7	27.3 - 38.7	2,800
Japanese	800	55.2	46.4 - 63.7	700	44.8	36.3 - 53.6	1,500
Other	1,100	59.2	51.3 - 66.7	800	40.8	33.3 - 48.7	1,900

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	65.5	59.0 - 71.4	600	34.5	28.6 - 41.0	1,600
20-24	2,100	62.2	57.6 - 66.6	1,300	37.8	33.4 - 42.4	3,400
25-34	3,700	58.5	54.8 - 62.2	2,700	41.5	37.8 - 45.2	6,400
35+	1,300	60.4	54.1 - 66.4	900	39.6	33.6 - 45.9	2,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,300	60.3	56.9 - 63.6	2,800	39.7	36.4 - 43.1	7,100
Female	3,900	60.9	57.4 - 64.4	2,500	39.1	35.6 - 42.6	6,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,200	57.5	54.4 - 60.5	3,900	42.5	39.5 - 45.6	9,100
No	3,000	66.9	63.1 - 70.5	1,500	33.1	29.5 - 36.9	4,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,800	62.8	59.3 - 66.2	2,200	37.2	33.8 - 40.7	6,000
131-185%	1,100	61.9	55.5 - 68.0	700	38.1	32.0 - 44.5	1,800
186+%	2,600	54.7	50.4 - 59.1	2,100	45.3	40.9 - 49.6	4,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,100	64.7	60.0 - 69.2	1,100	35.3	30.8 - 40.0	3,300
\$15,001-\$25,000	1,400	61.6	55.6 - 67.3	900	38.4	32.7 - 44.4	2,200
\$25,001-\$40,000	1,500	59.2	53.6 - 64.5	1,100	40.8	35.5 - 46.4	2,600
>=\$40,001	2,700	56.8	52.5 - 61.0	2,100	43.2	39.0 - 47.5	4,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	67.9	58.5 - 76.1	300	32.1	23.9 - 41.5	900
Normal birth weight	7,600	60.0	57.5 - 62.5	5,000	40.0	37.5 - 42.5	12,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	79.3	72.9 - 84.4	300	20.7	15.6 - 27.1	1,200
No	7,000	58.6	55.9 - 61.2	5,000	41.4	38.8 - 44.1	12,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	68.3	63.4 - 72.9	900	31.7	27.1 - 36.6	2,900
No	6,000	58.0	55.2 - 60.8	4,300	42.0	39.2 - 44.8	10,300

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,900	63.2	59.8 - 66.5	2,300	36.8	33.5 - 40.2	6,100
No	4,200	58.3	54.8 - 61.7	3,000	41.7	38.3 - 45.2	7,300