

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	2.8	1.8 - 4.2	2,200	13.0	10.9 - 15.4	600	3.2	2.3 - 4.7	14,000	81.0	78.2 - 83.5	17,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	<50	1.2	0.4 - 3.5	300	15.1	11.5 - 19.4	100	5.1	3.1 - 8.2	1,700	78.6	73.8 - 82.8	2,200
Honolulu	400	3.1	1.9 - 5.2	1,600	12.6	9.9 - 15.9	400	2.9	1.7 - 4.9	10,100	81.4	77.6 - 84.7	12,400
Kauai	<50	3.5	1.8 - 7.0	100	11.7	8.0 - 16.6	<50	2.2	0.9 - 5.1	700	82.6	77.0 - 87.1	800
Maui	<50	2.0	0.9 - 4.1	300	13.9	10.7 - 18.0	100	3.8	2.3 - 6.2	1,500	80.3	75.8 - 84.1	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.1	1.3 - 7.2	300	8.8	5.5 - 13.8	100	1.4	0.5 - 4.4	3,400	86.7	81.0 - 90.9	3,900
Native Hawaiian	100	2.1	0.9 - 4.7	900	18.8	14.5 - 24.1	300	5.6	3.5 - 8.9	3,600	73.5	67.8 - 78.5	4,900
Chinese	100	5.2	1.3 - 18.0	100	5.7	1.6 - 17.9	<50	3.2	0.5 - 18.9	900	86.0	71.5 - 93.7	1,000
Filipino	100	4.3	2.0 - 9.1	600	20.1	14.4 - 27.4	100	4.3	1.9 - 9.6	2,200	71.2	63.4 - 78.0	3,100
Japanese	<50	0.3	0.1 - 1.7	100	5.9	2.6 - 13.0	<50	1.0	0.4 - 3.0	1,400	92.7	85.9 - 96.4	1,500
Other	100	2.4	0.7 - 7.8	200	7.3	4.0 - 13.1	<50	1.7	0.4 - 6.5	2,500	88.5	81.8 - 93.0	2,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.2	3.5 - 22.4	200	25.3	15.2 - 39.0	<50	2.7	0.9 - 7.7	600	62.8	48.7 - 75.0	1,000
20-24	100	3.6	1.6 - 8.1	600	17.5	12.5 - 24.1	100	3.7	1.7 - 8.0	2,700	75.1	68.0 - 81.1	3,600
25-34	200	2.0	1.1 - 3.8	1,200	12.0	9.3 - 15.3	400	3.7	2.3 - 6.0	8,000	82.3	78.5 - 85.5	9,700
35+	100	2.2	0.7 - 6.3	200	7.0	4.2 - 11.4	<50	1.3	0.6 - 2.6	2,700	89.5	84.5 - 93.1	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.6	2.1 - 6.0	1,100	13.3	10.4 - 16.8	200	2.7	1.5 - 4.9	7,000	80.4	76.3 - 83.9	8,700
Female	200	2.0	1.0 - 3.9	1,100	12.7	9.9 - 16.3	300	3.8	2.4 - 6.0	7,000	81.5	77.6 - 84.9	8,600

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.4	1.3 - 4.3	1,200	10.8	8.4 - 13.8	300	2.7	1.6 - 4.4	9,100	84.2	80.7 - 87.1	10,700
No	200	3.5	2.0 - 6.1	1,100	16.7	13.0 - 21.1	300	4.2	2.5 - 7.0	4,900	75.6	70.7 - 79.9	6,500

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.5	2.6 - 7.6	1,000	17.3	13.5 - 22.0	200	2.6	1.3 - 5.1	4,500	75.6	70.4 - 80.1	6,000
131-185%	<50	1.4	0.3 - 6.4	400	14.0	9.1 - 21.1	200	5.6	2.8 - 10.9	2,300	79.0	71.2 - 85.1	2,900
186+%	100	1.8	0.7 - 4.2	600	8.7	6.1 - 12.3	200	2.8	1.5 - 5.2	5,900	86.7	82.5 - 89.9	6,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Income (2012+)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	200	4.7	2.5 - 8.9	700	17.2	12.5 - 23.1	100	2.3	0.9 - 5.7	2,900	75.8	69.2 - 81.4	3,800
\$17,001-\$26,000	100	2.4	0.6 - 8.8	300	14.5	9.3 - 21.8	100	6.5	3.2 - 12.5	1,700	76.7	68.2 - 83.5	2,200
\$26,001-\$51,000	100	2.7	1.1 - 6.3	700	16.1	11.4 - 22.2	100	2.8	1.2 - 6.2	3,200	78.4	71.9 - 83.8	4,000
>=\$51,001	100	1.6	0.6 - 4.3	500	7.9	5.3 - 11.6	200	2.9	1.5 - 5.7	5,100	87.6	83.2 - 91.0	5,800

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	2.7	1.4 - 5.2	200	15.0	11.1 - 20.0	<50	3.3	1.8 - 6.1	900	78.9	73.6 - 83.4	1,100
Normal birth weight	500	2.8	1.8 - 4.3	2,100	12.9	10.7 - 15.5	500	3.2	2.2 - 4.8	13,100	81.1	78.1 - 83.7	16,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	3.6	1.4 - 9.1	200	27.3	15.3 - 44.0	100	13.1	4.9 - 30.5	400	56.0	39.6 - 71.1	700
No	500	2.8	1.8 - 4.4	2,000	12.6	10.5 - 15.2	500	2.9	2.0 - 4.3	13,100	81.6	78.7 - 84.2	16,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.6	2.0 - 10.2	500	17.6	12.2 - 24.8	200	7.7	4.1 - 14.0	1,900	70.1	61.8 - 77.2	2,700
No	400	2.5	1.5 - 4.1	1,700	12.4	10.1 - 15.1	400	2.5	1.6 - 3.9	11,600	82.6	79.6 - 85.3	14,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	4.9	3.0 - 7.7	1,400	18.3	14.7 - 22.5	200	3.2	1.9 - 5.3	5,600	73.7	69.0 - 77.9	7,600
No	100	1.1	0.4 - 2.7	800	8.9	6.6 - 11.8	300	3.0	1.8 - 5.0	8,300	87.1	83.7 - 89.8	9,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	300	1.7	1.1 - 2.8	2,200	12.7	10.7 - 15.1	700	4.0	2.9 - 5.4	14,200	81.5	78.8 - 83.9	17,500

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	<50	1.9	0.9 - 4.0	400	16.1	12.5 - 20.5	100	4.2	2.5 - 6.9	1,800	77.7	73.0 - 81.9	2,300
Honolulu	200	1.6	0.8 - 3.1	1,500	12.2	9.6 - 15.5	500	3.8	2.5 - 5.9	10,300	82.3	78.7 - 85.5	12,500
Kauai	<50	1.0	0.3 - 3.5	100	9.7	6.5 - 14.2	<50	4.2	2.3 - 7.6	700	85.0	79.9 - 89.0	800
Maui	100	2.8	1.5 - 4.9	200	13.3	10.4 - 17.0	100	4.8	3.1 - 7.2	1,500	79.1	75.0 - 82.7	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	1.3	0.4 - 4.2	500	11.9	7.9 - 17.4	100	3.3	1.7 - 6.4	3,600	83.5	77.7 - 88.1	4,400
Native Hawaiian	100	2.0	0.9 - 4.4	800	15.8	12.0 - 20.5	300	6.4	3.9 - 10.1	3,700	75.9	70.4 - 80.6	4,900
Chinese	<50	4.5	0.7 - 25.2	<50	2.3	0.8 - 6.3	<50	1.8	0.6 - 5.0	600	91.4	77.4 - 97.1	600
Filipino	<50	0.9	0.4 - 1.9	500	16.1	11.4 - 22.3	100	3.5	1.9 - 6.5	2,500	79.5	73.1 - 84.7	3,100
Japanese	<50	0.4	0.1 - 2.2	200	10.3	5.3 - 19.3	<50	1.7	0.4 - 7.8	1,600	87.6	78.5 - 93.2	1,800
Other	100	3.3	1.3 - 8.3	200	8.6	4.6 - 15.3	100	3.4	1.3 - 8.9	2,200	84.7	76.9 - 90.2	2,600

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	2.0	0.7 - 5.8	400	40.2	28.3 - 53.3	<50	3.4	1.3 - 8.3	500	54.4	41.7 - 66.6	1,000
20-24	100	2.3	1.0 - 5.3	700	16.8	12.2 - 22.8	100	1.3	0.7 - 2.3	3,200	79.5	73.4 - 84.6	4,000
25-34	200	1.9	1.0 - 3.7	1,000	10.6	8.1 - 13.8	500	5.1	3.4 - 7.5	7,700	82.4	78.6 - 85.7	9,400
35+	<50	0.4	0.1 - 1.3	200	5.3	3.0 - 9.0	100	4.5	2.3 - 8.6	2,800	89.9	85.0 - 93.3	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	200	2.6	1.4 - 4.5	1,100	12.4	9.6 - 15.9	400	4.0	2.6 - 6.3	7,200	81.0	77.0 - 84.4	8,900
Female	100	0.9	0.4 - 2.0	1,100	13.1	10.3 - 16.5	300	4.0	2.6 - 6.1	7,000	82.1	78.3 - 85.3	8,600

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.2	1.2 - 3.9	1,000	9.5	7.3 - 12.3	500	4.2	2.9 - 6.2	9,200	84.1	80.8 - 87.0	10,900
No	100	1.0	0.6 - 1.8	1,200	18.2	14.4 - 22.7	200	3.6	2.1 - 6.2	5,000	77.2	72.4 - 81.3	6,500

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	100	2.5	1.3 - 4.7	1,000	16.8	13.2 - 21.2	300	4.2	2.5 - 6.9	4,700	76.6	71.8 - 80.7	6,100
131-185%	<50	1.6	0.4 - 6.3	400	13.7	8.7 - 20.9	100	3.5	1.5 - 8.0	2,100	81.2	73.5 - 87.1	2,600
186+%	100	1.3	0.5 - 3.1	700	8.5	6.0 - 12.0	300	4.2	2.6 - 6.7	6,700	86.0	82.1 - 89.2	7,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	2.9	1.2 - 6.7	600	18.2	13.3 - 24.5	100	2.5	0.9 - 6.5	2,400	76.4	69.5 - 82.1	3,100
\$10,000-\$24,999	100	1.7	0.6 - 5.0	500	16.4	11.4 - 22.9	100	4.3	2.4 - 7.6	2,500	77.6	70.8 - 83.2	3,200
\$25,000-\$49,999	100	1.6	0.5 - 5.0	600	14.0	9.9 - 19.5	200	5.6	3.0 - 10.0	3,300	78.8	72.5 - 83.9	4,200
>=\$50,000	100	1.2	0.5 - 3.1	400	6.5	4.2 - 10.1	200	3.6	2.1 - 6.2	5,300	88.6	84.5 - 91.7	6,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	2.5	1.2 - 5.1	200	18.4	14.3 - 23.4	100	6.0	3.8 - 9.4	900	73.0	67.6 - 77.9	1,200
Normal birth weight	300	1.7	1.0 - 2.8	2,000	12.3	10.2 - 14.9	600	3.9	2.7 - 5.4	13,400	82.1	79.3 - 84.7	16,300

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	0.7	0.1 - 4.1	300	33.4	21.9 - 47.2	100	9.3	3.7 - 21.3	500	56.6	43.0 - 69.4	800
No	300	1.8	1.1 - 2.9	1,900	11.4	9.4 - 13.7	600	3.8	2.7 - 5.2	13,700	83.1	80.4 - 85.5	16,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	0.6	0.2 - 1.6	800	23.1	17.6 - 29.6	200	4.8	2.3 - 9.6	2,300	71.6	64.6 - 77.7	3,300
No	300	2.0	1.2 - 3.3	1,400	10.0	7.9 - 12.5	500	3.8	2.7 - 5.4	11,800	84.2	81.3 - 86.7	14,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	1.8	0.9 - 3.3	1,300	16.9	13.6 - 20.9	300	3.9	2.5 - 6.0	6,000	77.4	73.2 - 81.2	7,800
No	200	1.7	0.8 - 3.4	900	9.2	6.8 - 12.2	400	4.2	2.7 - 6.3	8,100	85.0	81.4 - 88.0	9,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.3	2.3 - 4.7	2,400	14.2	12.0 - 16.7	700	4.3	3.1 - 5.9	13,400	78.3	75.4 - 80.9	17,200

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.9	2.3 - 6.5	300	15.9	12.4 - 20.2	100	2.9	1.5 - 5.4	1,700	77.4	72.6 - 81.5	2,200
Honolulu	400	3.4	2.1 - 5.4	1,700	13.9	11.1 - 17.3	600	4.9	3.3 - 7.2	9,600	77.8	73.9 - 81.3	12,300
Kauai	0	0.0	-	100	11.1	7.8 - 15.5	<50	3.0	1.5 - 5.9	700	85.9	81.2 - 89.6	800
Maui	100	3.3	2.0 - 5.6	300	15.5	12.3 - 19.4	<50	2.1	1.1 - 4.1	1,500	79.0	74.8 - 82.8	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.5	1.6 - 7.6	400	9.5	6.0 - 14.8	300	7.9	4.8 - 12.8	3,300	79.1	72.6 - 84.4	4,200
Native Hawaiian	200	3.9	2.0 - 7.1	900	17.4	13.5 - 22.1	200	4.0	2.2 - 7.2	3,800	74.8	69.4 - 79.5	5,100
Chinese	<50	1.5	0.4 - 5.4	100	7.6	2.2 - 22.9	<50	4.4	0.8 - 20.7	600	86.5	70.8 - 94.4	700
Filipino	100	2.7	1.3 - 5.5	700	22.3	16.4 - 29.5	100	3.1	1.3 - 7.0	2,200	71.9	64.5 - 78.4	3,000
Japanese	<50	0.5	0.1 - 2.9	100	4.1	1.8 - 9.2	<50	1.8	0.3 - 9.0	1,600	93.7	87.3 - 96.9	1,700
Other	100	4.9	2.0 - 11.2	400	14.4	8.9 - 22.4	<50	1.7	0.5 - 5.8	1,900	79.0	70.3 - 85.6	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	1.8	0.6 - 5.1	300	28.5	18.3 - 41.4	100	9.4	3.6 - 22.1	600	60.4	47.4 - 72.0	1,100
20-24	100	2.7	1.2 - 6.4	800	19.9	15.0 - 26.0	100	3.4	1.5 - 7.3	3,000	74.0	67.4 - 79.6	4,100
25-34	300	3.2	1.9 - 5.4	1,100	11.9	9.2 - 15.3	400	4.2	2.7 - 6.6	7,200	80.7	76.7 - 84.1	8,900
35+	100	4.7	2.3 - 9.5	300	8.5	5.2 - 13.6	100	3.7	1.7 - 8.0	2,600	83.1	76.6 - 88.1	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.3	2.0 - 5.6	1,300	15.1	12.0 - 18.7	400	4.4	2.9 - 6.9	6,700	77.1	73.0 - 80.9	8,700
Female	300	3.2	1.9 - 5.4	1,100	13.3	10.4 - 16.9	300	4.1	2.5 - 6.6	6,700	79.4	75.3 - 83.0	8,500

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	1.8 - 4.8	1,200	11.2	8.7 - 14.3	500	4.1	2.7 - 6.3	9,000	81.8	78.2 - 84.9	10,900
No	200	3.9	2.2 - 6.8	1,200	19.5	15.6 - 24.0	300	4.5	2.7 - 7.4	4,500	72.1	67.1 - 76.6	6,200

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	200	3.8	2.1 - 6.6	1,100	18.6	14.8 - 23.2	200	4.2	2.3 - 7.3	4,300	73.4	68.3 - 78.0	5,900
131-185%	200	5.5	2.5 - 11.8	400	14.5	9.4 - 21.8	100	4.8	2.3 - 9.8	2,100	75.2	67.0 - 81.9	2,800
186+%	200	2.3	1.2 - 4.3	700	9.1	6.4 - 12.7	300	4.1	2.4 - 6.7	6,300	84.6	80.4 - 88.0	7,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	4.4	2.2 - 8.7	600	20.3	15.0 - 26.9	100	3.9	1.8 - 8.3	2,200	71.4	64.2 - 77.7	3,100
\$10,000-\$24,999	100	3.1	1.3 - 7.2	400	14.5	10.1 - 20.4	200	5.2	2.5 - 10.4	2,300	77.2	70.2 - 82.9	2,900
\$25,000-\$49,999	200	4.2	2.1 - 8.4	700	16.1	11.5 - 22.0	200	4.8	2.6 - 8.9	3,200	74.8	68.3 - 80.4	4,300
>=\$50,000	100	2.4	1.2 - 5.0	500	8.5	5.7 - 12.5	200	3.6	2.0 - 6.5	5,100	85.5	80.8 - 89.2	6,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	3.3	1.7 - 6.4	200	14.6	10.9 - 19.1	100	6.8	4.3 - 10.7	900	75.3	69.9 - 80.0	1,200
Normal birth weight	500	3.3	2.2 - 4.9	2,300	14.2	11.9 - 16.8	700	4.1	2.8 - 5.8	12,600	78.5	75.4 - 81.3	16,000

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	10.1	4.3 - 22.2	200	19.6	11.5 - 31.3	<50	2.2	0.8 - 6.2	700	68.1	55.1 - 78.8	1,100
No	500	2.9	1.9 - 4.3	2,200	13.9	11.7 - 16.5	700	4.4	3.2 - 6.2	12,500	78.8	75.8 - 81.5	15,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	7.8	4.5 - 13.2	700	21.8	16.3 - 28.6	200	6.1	3.4 - 10.6	2,200	64.3	56.9 - 71.0	3,400
No	300	2.2	1.3 - 3.6	1,700	12.4	10.1 - 15.1	500	3.9	2.6 - 5.7	11,100	81.6	78.5 - 84.3	13,600

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.2	2.6 - 6.8	1,300	17.5	14.1 - 21.5	400	4.9	3.1 - 7.7	5,400	73.4	68.8 - 77.6	7,400
No	300	2.6	1.5 - 4.6	1,100	11.8	9.1 - 15.2	400	3.8	2.4 - 6.0	7,900	81.7	77.9 - 85.1	9,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	800	4.6	3.4 - 6.3	2,600	15.6	13.4 - 18.2	900	5.3	4.0 - 7.0	12,500	74.4	71.4 - 77.2	16,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.2	4.0 - 9.3	400	16.6	12.9 - 21.1	100	4.1	2.4 - 6.8	1,600	73.1	68.1 - 77.7	2,200
Honolulu	600	4.7	3.1 - 7.1	1,900	15.8	12.8 - 19.3	600	5.4	3.7 - 7.8	8,700	74.1	70.0 - 77.9	11,800
Kauai	<50	2.9	1.3 - 6.1	100	13.5	9.8 - 18.4	<50	4.3	2.4 - 7.7	600	79.3	73.8 - 83.9	800
Maui	100	3.6	2.1 - 5.9	300	14.5	11.4 - 18.3	100	6.6	4.6 - 9.4	1,500	75.3	70.9 - 79.2	2,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	3.7	1.7 - 7.9	400	9.1	6.0 - 13.6	200	5.9	3.4 - 10.2	3,300	81.3	75.4 - 86.0	4,100
Native Hawaiian	300	5.9	3.6 - 9.6	1,100	21.9	17.4 - 27.1	200	4.9	2.9 - 7.9	3,400	67.4	61.6 - 72.6	5,100
Chinese	0	0.0	-	100	9.1	2.9 - 25.0	<50	1.4	0.4 - 5.1	600	89.5	74.4 - 96.1	700
Filipino	100	4.5	2.6 - 7.6	600	19.2	13.8 - 25.9	200	6.2	3.3 - 11.2	2,100	70.2	62.9 - 76.6	3,000
Japanese	100	4.0	1.3 - 11.6	100	7.6	3.4 - 16.4	100	3.9	1.4 - 10.6	1,300	84.4	74.6 - 90.9	1,600
Other	100	5.6	2.3 - 13.0	400	16.2	10.3 - 24.7	100	6.3	2.9 - 13.1	1,600	71.8	62.3 - 79.7	2,300

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.5	2.8 - 14.2	500	29.8	21.0 - 40.4	100	6.2	2.6 - 13.8	900	57.6	46.6 - 67.8	1,600
20-24	200	6.0	3.3 - 10.8	700	18.3	13.7 - 24.0	300	7.0	4.2 - 11.5	2,600	68.6	61.9 - 74.6	3,800
25-34	300	3.6	2.2 - 5.9	1,100	12.7	9.9 - 16.1	400	5.0	3.3 - 7.5	6,600	78.7	74.6 - 82.3	8,400
35+	100	4.8	2.3 - 9.7	400	12.5	7.9 - 19.4	100	3.6	1.7 - 7.5	2,300	79.1	71.6 - 85.0	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.9	2.5 - 6.2	1,300	14.8	11.8 - 18.3	600	6.2	4.3 - 8.9	6,600	75.0	70.8 - 78.8	8,800
Female	400	5.4	3.6 - 8.2	1,300	16.6	13.3 - 20.4	300	4.3	2.7 - 6.6	5,800	73.7	69.3 - 77.7	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.4	2.1 - 5.4	1,200	11.3	8.8 - 14.4	500	5.1	3.6 - 7.3	8,200	80.2	76.5 - 83.4	10,200
No	400	6.6	4.3 - 9.9	1,500	22.4	18.4 - 27.0	400	5.6	3.6 - 8.7	4,300	65.4	60.3 - 70.3	6,500

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	7.0	4.5 - 10.5	1,200	21.5	17.2 - 26.4	200	4.0	2.3 - 6.9	3,800	67.6	62.1 - 72.7	5,600
131-185%	100	5.6	2.6 - 11.7	400	16.8	11.5 - 23.9	200	6.5	3.5 - 11.7	1,900	71.2	63.1 - 78.1	2,600
186+%	200	3.2	1.7 - 5.7	700	9.6	6.9 - 13.1	400	5.9	3.9 - 9.0	5,700	81.3	76.9 - 85.0	7,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	5.9	3.0 - 11.4	700	25.1	18.5 - 33.2	100	5.3	2.5 - 10.9	1,700	63.7	55.2 - 71.4	2,700
\$10,000-\$24,999	300	8.1	4.9 - 13.2	600	17.0	12.5 - 22.6	200	4.6	2.4 - 8.5	2,400	70.3	63.6 - 76.2	3,500
\$25,000-\$49,999	200	6.0	3.0 - 11.7	400	13.1	9.0 - 18.7	200	5.9	3.3 - 10.3	2,600	74.9	68.0 - 80.8	3,400
>=\$50,000	100	2.0	0.9 - 4.0	600	10.6	7.5 - 14.8	300	5.5	3.4 - 8.8	4,800	81.9	77.1 - 85.9	5,900

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.3	3.1 - 8.9	200	18.8	14.8 - 23.6	100	5.1	3.1 - 8.1	800	70.8	65.3 - 75.7	1,100
Normal birth weight	700	4.6	3.3 - 6.4	2,400	15.4	13.0 - 18.1	800	5.3	4.0 - 7.2	11,600	74.7	71.5 - 77.6	15,600

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	12.0	6.1 - 22.3	400	26.5	17.8 - 37.5	100	6.2	2.2 - 16.0	700	55.4	43.8 - 66.4	1,300
No	600	4.1	2.9 - 5.8	2,200	14.4	12.1 - 17.0	700	5.0	3.7 - 6.7	11,500	76.5	73.4 - 79.4	15,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	7.5	4.3 - 12.9	800	23.3	17.8 - 30.0	200	6.6	3.8 - 11.4	2,100	62.5	55.2 - 69.2	3,400
No	500	4.0	2.7 - 5.8	1,700	13.4	11.0 - 16.1	600	4.7	3.3 - 6.6	10,100	77.9	74.6 - 80.9	13,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	8.0	5.5 - 11.6	1,400	22.0	18.1 - 26.4	300	5.0	3.2 - 7.7	4,300	65.0	60.0 - 69.8	6,500
No	200	2.5	1.4 - 4.5	1,100	11.5	8.9 - 14.8	500	5.2	3.5 - 7.6	7,800	80.7	76.9 - 84.1	9,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	2.9	2.1 - 3.8	2,600	15.3	13.6 - 17.2	800	4.6	3.6 - 5.7	12,900	77.3	75.1 - 79.3	16,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.1	3.4 - 10.7	300	15.4	10.8 - 21.5	<50	1.8	0.6 - 5.3	1,600	76.7	69.9 - 82.3	2,100
Honolulu	300	2.2	1.5 - 3.3	1,900	16.1	14.0 - 18.4	600	4.8	3.6 - 6.2	9,100	76.9	74.3 - 79.3	11,900
Kauai	<50	1.7	0.4 - 7.0	100	13.3	7.3 - 23.3	100	10.2	5.0 - 19.5	600	74.8	63.8 - 83.3	800
Maui	100	3.7	1.7 - 7.8	200	11.0	7.1 - 16.8	100	4.0	1.9 - 8.1	1,600	81.3	74.7 - 86.5	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.0	1.0 - 3.9	500	13.1	10.1 - 16.8	300	6.3	4.3 - 9.3	3,300	78.6	74.3 - 82.4	4,100
Native Hawaiian	200	4.0	2.4 - 6.7	800	17.8	14.2 - 22.2	300	6.4	4.3 - 9.4	3,300	71.8	66.8 - 76.2	4,500
Chinese	<50	1.2	0.4 - 3.8	100	9.2	5.9 - 14.0	<50	3.9	2.0 - 7.6	500	85.8	80.3 - 89.9	600
Filipino	100	3.9	2.2 - 6.7	500	18.0	14.1 - 22.6	100	3.2	1.7 - 5.8	2,300	75.0	69.9 - 79.5	3,000
Japanese	100	2.6	1.0 - 6.6	200	9.5	5.8 - 14.9	100	2.5	1.1 - 5.8	1,700	85.4	79.3 - 89.9	2,000
Other	<50	1.6	0.7 - 4.1	400	17.2	12.6 - 22.9	<50	1.8	0.6 - 4.9	2,000	79.4	73.4 - 84.3	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.4	2.3 - 11.9	300	20.4	13.7 - 29.1	<50	3.5	1.3 - 9.1	900	70.8	61.4 - 78.6	1,300
20-24	100	2.9	1.6 - 5.5	900	24.0	19.5 - 29.1	300	7.6	5.1 - 11.1	2,300	65.5	60.0 - 70.6	3,600
25-34	200	2.1	1.3 - 3.4	1,100	12.8	10.6 - 15.3	400	4.0	2.8 - 5.7	7,000	81.1	78.2 - 83.7	8,700
35+	100	3.8	2.2 - 6.5	300	10.4	7.4 - 14.5	100	3.1	1.7 - 5.7	2,700	82.7	78.1 - 86.6	3,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.1	2.0 - 4.5	1,400	15.9	13.5 - 18.7	500	5.1	3.8 - 6.9	6,700	75.9	72.8 - 78.7	8,800
Female	200	2.7	1.7 - 4.1	1,200	14.6	12.2 - 17.4	300	4.0	2.7 - 5.7	6,300	78.8	75.7 - 81.6	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.2	1.5 - 3.3	1,600	14.6	12.5 - 16.9	500	4.3	3.2 - 5.8	8,500	78.9	76.3 - 81.3	10,800
No	200	4.1	2.6 - 6.2	1,000	16.6	13.6 - 20.1	300	5.0	3.4 - 7.3	4,400	74.4	70.3 - 78.0	6,000

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	200	3.4	2.1 - 5.5	1,100	19.3	16.0 - 23.0	300	4.5	3.0 - 6.6	4,200	72.8	68.7 - 76.6	5,800
131-185%	<50	1.5	0.5 - 4.5	300	12.1	8.3 - 17.3	100	3.0	1.4 - 6.4	2,000	83.3	77.7 - 87.8	2,400
186+%	200	2.3	1.4 - 3.6	900	12.3	10.0 - 14.9	400	5.2	3.8 - 7.2	6,000	80.2	77.1 - 83.0	7,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	4.4	2.3 - 8.2	500	21.0	16.1 - 27.0	100	3.7	1.9 - 7.0	1,900	70.9	64.6 - 76.6	2,600
\$10,000-\$24,999	100	3.2	1.7 - 5.9	600	17.6	13.7 - 22.4	200	5.7	3.6 - 9.0	2,600	73.4	68.1 - 78.1	3,500
\$25,000-\$49,999	100	2.5	1.4 - 4.7	600	16.6	13.1 - 20.9	200	4.1	2.5 - 6.8	3,000	76.7	72.0 - 80.8	3,800
>=\$50,000	100	1.6	0.9 - 3.1	600	9.8	7.6 - 12.6	300	4.6	3.1 - 6.7	4,900	84.0	80.7 - 86.8	5,800

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	4.3	1.7 - 10.6	200	14.9	9.1 - 23.4	<50	3.6	1.3 - 9.4	900	77.2	68.0 - 84.4	1,100
Normal birth weight	400	2.8	2.0 - 3.8	2,400	15.3	13.5 - 17.3	700	4.6	3.7 - 5.9	12,100	77.3	75.0 - 79.4	15,600

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	1.8	0.5 - 6.6	300	27.6	19.9 - 36.9	100	10.3	5.7 - 17.8	800	60.3	50.7 - 69.2	1,300
No	400	2.9	2.1 - 3.9	2,200	14.3	12.5 - 16.2	600	4.2	3.3 - 5.4	11,900	78.7	76.4 - 80.7	15,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	2.2	0.9 - 5.0	600	22.1	17.4 - 27.7	300	9.7	6.6 - 14.1	1,900	66.0	59.9 - 71.6	2,900
No	400	2.9	2.1 - 4.0	1,900	13.8	12.0 - 15.9	500	3.6	2.7 - 4.8	10,800	79.7	77.4 - 81.8	13,500

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	3.7	2.4 - 5.6	1,400	20.0	17.0 - 23.5	300	4.4	3.0 - 6.4	4,800	71.9	68.1 - 75.5	6,700
No	200	2.4	1.6 - 3.6	1,200	12.0	10.0 - 14.3	500	4.8	3.6 - 6.4	7,900	80.8	78.1 - 83.2	9,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.8	2.9 - 4.9	2,700	16.2	14.4 - 18.1	500	3.3	2.5 - 4.3	12,700	76.7	74.5 - 78.8	16,600

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.3	2.1 - 8.8	200	9.9	6.3 - 15.2	100	2.4	0.9 - 6.0	1,800	83.4	77.1 - 88.2	2,100
Honolulu	400	3.6	2.6 - 5.0	2,000	17.2	15.0 - 19.6	400	3.6	2.6 - 4.8	8,800	75.6	73.0 - 78.1	11,600
Kauai	<50	1.4	0.2 - 8.5	100	14.6	8.4 - 24.1	<50	4.7	1.8 - 11.4	700	79.3	69.3 - 86.7	900
Maui	100	5.4	2.8 - 10.2	300	17.6	12.4 - 24.3	<50	2.1	0.7 - 6.2	1,500	75.0	67.7 - 81.0	2,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.5	2.0 - 6.1	400	11.5	8.6 - 15.4	100	3.8	2.2 - 6.4	3,000	81.2	76.7 - 85.0	3,700
Native Hawaiian	300	6.3	4.2 - 9.5	1,100	24.5	20.3 - 29.3	100	2.4	1.2 - 4.5	3,000	66.8	61.7 - 71.5	4,600
Chinese	<50	3.7	1.7 - 7.7	<50	6.9	4.0 - 11.5	<50	1.7	0.6 - 4.4	600	87.8	82.3 - 91.7	700
Filipino	100	1.9	0.9 - 4.0	600	19.5	15.5 - 24.2	200	5.8	3.7 - 9.0	2,200	72.7	67.6 - 77.3	3,000
Japanese	<50	1.3	0.3 - 5.2	200	7.7	4.6 - 12.7	<50	2.2	0.8 - 5.5	1,800	88.7	83.2 - 92.6	2,000
Other	100	4.0	2.1 - 7.5	300	13.0	9.2 - 18.2	100	2.7	1.3 - 5.7	2,100	80.3	74.5 - 85.0	2,600

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.8	5.3 - 17.6	300	23.7	16.5 - 33.0	100	4.9	2.1 - 11.1	900	61.5	51.7 - 70.4	1,500
20-24	100	3.1	1.7 - 5.5	1,000	23.8	19.7 - 28.6	100	3.2	1.8 - 5.4	2,800	69.9	64.9 - 74.4	4,100
25-34	300	3.5	2.4 - 5.1	1,100	13.9	11.6 - 16.5	200	2.9	1.9 - 4.3	6,500	79.7	76.8 - 82.4	8,100
35+	100	2.6	1.3 - 5.1	200	8.0	5.4 - 11.7	100	3.9	2.2 - 6.8	2,500	85.5	81.0 - 89.1	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	200	2.6	1.7 - 4.1	1,500	17.5	14.9 - 20.3	300	3.4	2.3 - 4.9	6,600	76.5	73.4 - 79.4	8,600
Female	400	5.1	3.7 - 7.0	1,200	14.7	12.3 - 17.5	300	3.2	2.2 - 4.7	6,100	76.9	73.7 - 79.9	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	2.0 - 4.2	1,200	11.8	9.9 - 13.9	300	2.5	1.7 - 3.6	8,700	82.8	80.4 - 85.0	10,500
No	300	5.3	3.6 - 7.8	1,400	23.8	20.3 - 27.8	300	4.8	3.3 - 6.9	4,000	66.1	61.8 - 70.1	6,000

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.4	3.7 - 7.8	1,400	24.0	20.4 - 27.9	200	3.1	1.9 - 5.1	4,000	67.5	63.3 - 71.5	6,000
131-185%	100	3.2	1.6 - 6.6	400	14.4	10.4 - 19.6	100	3.8	2.0 - 7.3	1,900	78.6	72.7 - 83.4	2,400
186+%	200	3.3	2.2 - 5.0	700	9.9	7.8 - 12.4	200	3.0	1.9 - 4.5	5,800	83.9	80.9 - 86.4	7,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	7.1	4.2 - 11.5	700	25.8	20.4 - 32.0	100	4.2	2.2 - 7.8	1,700	63.0	56.4 - 69.2	2,700
\$10,000-\$24,999	100	4.4	2.5 - 7.5	700	20.0	15.8 - 24.9	100	3.2	1.8 - 5.9	2,400	72.4	67.1 - 77.2	3,300
\$25,000-\$49,999	100	2.6	1.4 - 5.0	700	17.8	14.1 - 22.1	100	2.7	1.5 - 5.0	2,900	76.9	72.2 - 81.0	3,800
>=\$50,000	200	3.4	2.2 - 5.3	500	8.1	6.1 - 10.7	200	2.9	1.8 - 4.6	4,800	85.6	82.4 - 88.3	5,600

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	3.9	1.6 - 9.4	300	21.8	15.2 - 30.1	<50	3.1	1.2 - 7.6	1,000	71.3	62.6 - 78.7	1,300
Normal birth weight	600	3.8	2.9 - 5.0	2,400	15.7	13.8 - 17.7	500	3.3	2.5 - 4.4	11,800	77.2	74.9 - 79.3	15,200

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.3	4.2 - 15.5	400	32.4	24.0 - 42.1	100	6.9	3.4 - 13.4	700	52.5	42.9 - 61.9	1,300
No	500	3.5	2.6 - 4.6	2,300	14.9	13.1 - 16.9	500	3.1	2.3 - 4.1	11,900	78.5	76.3 - 80.6	15,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.1	3.0 - 8.6	1,000	31.0	25.7 - 36.9	100	4.6	2.7 - 7.8	1,800	59.3	53.3 - 65.0	3,100
No	500	3.5	2.6 - 4.8	1,700	12.9	11.1 - 14.9	400	3.1	2.3 - 4.2	10,700	80.5	78.2 - 82.6	13,300

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.2	2.8 - 6.4	1,400	21.7	18.5 - 25.3	200	3.3	2.2 - 5.1	4,600	70.7	66.8 - 74.3	6,600
No	300	3.5	2.5 - 4.9	1,200	12.1	10.1 - 14.4	300	3.4	2.4 - 4.8	7,800	81.0	78.4 - 83.4	9,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	2.9	2.1 - 3.9	2,800	17.4	15.5 - 19.5	600	3.5	2.7 - 4.6	12,400	76.2	73.9 - 78.4	16,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.6	1.5 - 8.0	400	19.9	14.4 - 26.7	<50	2.1	0.7 - 6.1	1,500	74.5	67.2 - 80.6	2,000
Honolulu	400	3.2	2.3 - 4.4	2,000	16.9	14.7 - 19.4	400	3.6	2.6 - 4.8	9,100	76.4	73.7 - 78.8	11,900
Kauai	0	0.0	-	100	16.0	8.8 - 27.5	<50	3.4	0.9 - 12.1	600	80.5	68.7 - 88.6	800
Maui	<50	1.2	0.3 - 5.5	300	18.5	12.9 - 25.8	100	4.8	2.2 - 10.1	1,300	75.5	67.6 - 81.9	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.5	1.3 - 4.8	500	13.9	10.6 - 18.1	200	5.2	3.3 - 8.0	3,100	78.4	73.7 - 82.5	3,900
Native Hawaiian	200	3.5	2.0 - 6.2	1,100	24.4	20.0 - 29.4	100	2.5	1.3 - 4.8	3,000	69.6	64.4 - 74.4	4,400
Chinese	<50	2.1	0.8 - 5.6	100	11.3	7.5 - 16.8	<50	2.8	1.2 - 6.5	500	83.7	77.7 - 88.3	600
Filipino	100	2.8	1.4 - 5.6	600	19.5	15.3 - 24.4	100	4.3	2.5 - 7.4	2,200	73.5	68.1 - 78.2	3,000
Japanese	<50	1.9	0.6 - 5.7	200	8.0	4.8 - 13.1	<50	0.6	0.1 - 3.5	1,700	89.5	84.0 - 93.3	1,900
Other	100	3.3	1.6 - 6.9	400	16.8	12.2 - 22.7	100	4.2	2.2 - 7.6	1,900	75.7	69.3 - 81.1	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	4.1	1.6 - 10.2	400	32.3	23.9 - 42.1	<50	1.5	0.3 - 6.7	800	62.1	52.3 - 71.0	1,300
20-24	100	3.3	1.8 - 5.9	1,000	26.1	21.5 - 31.2	200	4.8	2.9 - 7.7	2,600	65.9	60.5 - 70.9	4,000
25-34	200	3.0	2.0 - 4.6	1,100	13.7	11.4 - 16.4	300	3.5	2.4 - 5.0	6,500	79.8	76.7 - 82.6	8,100
35+	<50	1.4	0.6 - 3.3	300	8.9	6.1 - 12.8	100	2.8	1.4 - 5.4	2,500	87.0	82.6 - 90.4	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.2	2.1 - 4.7	1,600	19.4	16.6 - 22.5	400	5.0	3.6 - 6.8	6,100	72.4	69.0 - 75.6	8,400
Female	200	2.6	1.6 - 4.1	1,200	15.2	12.7 - 18.1	200	2.0	1.2 - 3.3	6,400	80.2	77.1 - 83.0	8,000

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.5	1.7 - 3.7	1,300	12.1	10.2 - 14.3	300	3.3	2.4 - 4.6	8,600	82.1	79.6 - 84.4	10,500
No	200	3.5	2.2 - 5.7	1,600	26.8	22.9 - 31.1	200	3.9	2.5 - 6.1	3,900	65.7	61.2 - 70.0	5,900

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.9	3.3 - 7.4	1,400	25.7	21.8 - 30.1	100	2.5	1.4 - 4.4	3,700	66.8	62.3 - 71.1	5,500
131-185%	100	2.5	1.0 - 6.3	300	13.9	9.5 - 20.0	100	5.8	3.2 - 10.5	1,600	77.7	70.9 - 83.3	2,100
186+%	100	1.6	0.9 - 2.9	900	12.0	9.8 - 14.7	300	3.9	2.7 - 5.5	6,300	82.5	79.5 - 85.1	7,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	4.8	2.6 - 8.7	800	30.3	24.3 - 36.9	100	2.5	1.1 - 5.6	1,700	62.5	55.7 - 68.9	2,700
\$10,000-\$24,999	200	5.3	3.2 - 8.7	700	20.7	16.3 - 25.9	100	2.3	1.1 - 4.9	2,400	71.7	66.1 - 76.7	3,400
\$25,000-\$49,999	100	2.1	1.1 - 4.3	700	19.0	15.1 - 23.6	200	5.4	3.4 - 8.3	2,900	73.5	68.5 - 78.0	3,900
>=\$50,000	100	1.1	0.5 - 2.5	400	7.5	5.6 - 10.1	200	3.8	2.5 - 5.8	4,700	87.5	84.4 - 90.1	5,400

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	4.9	2.1 - 10.9	200	15.1	9.4 - 23.5	<50	3.0	1.1 - 7.4	1,000	77.0	68.1 - 84.0	1,300
Normal birth weight	400	2.7	1.9 - 3.8	2,600	17.6	15.6 - 19.8	500	3.6	2.7 - 4.7	11,500	76.1	73.7 - 78.4	15,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.6	3.7 - 14.7	300	23.2	15.7 - 33.0	100	5.0	2.2 - 11.1	800	64.2	54.1 - 73.2	1,200
No	400	2.5	1.7 - 3.5	2,500	16.7	14.7 - 18.9	500	3.4	2.6 - 4.6	11,500	77.4	75.0 - 79.6	14,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.6	2.6 - 8.0	700	22.3	17.5 - 27.9	200	5.3	3.2 - 8.8	2,200	67.8	61.8 - 73.3	3,200
No	300	2.4	1.7 - 3.5	2,100	15.9	13.9 - 18.2	400	3.1	2.3 - 4.3	10,100	78.5	76.0 - 80.8	12,900

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.9	3.4 - 7.1	1,600	23.2	19.8 - 27.0	200	3.6	2.3 - 5.5	4,600	68.3	64.2 - 72.1	6,700
No	100	1.1	0.6 - 1.9	1,300	13.4	11.3 - 15.9	300	3.3	2.3 - 4.7	7,700	82.2	79.5 - 84.6	9,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.2	3.2 - 5.3	2,500	16.0	14.3 - 18.0	500	3.4	2.6 - 4.4	11,700	76.4	74.2 - 78.5	15,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	<50	2.6	1.0 - 6.5	300	15.5	10.8 - 21.7	<50	1.9	0.7 - 5.6	1,500	80.0	73.3 - 85.3	1,900
Honolulu	500	4.4	3.3 - 5.8	1,700	15.2	13.2 - 17.4	400	4.0	3.0 - 5.3	8,600	76.5	73.9 - 78.9	11,200
Kauai	0	0.0	-	100	13.9	7.5 - 24.4	<50	1.9	0.3 - 11.4	600	84.2	73.3 - 91.2	700
Maui	100	6.6	3.5 - 12.0	400	23.7	17.6 - 31.2	<50	1.6	0.5 - 5.0	1,000	68.1	60.3 - 75.0	1,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.4	1.9 - 6.1	300	9.6	6.8 - 13.2	100	4.1	2.5 - 6.8	2,900	82.9	78.5 - 86.6	3,500
Native Hawaiian	300	6.6	4.4 - 9.8	800	19.5	15.6 - 24.1	200	5.1	3.2 - 7.9	2,900	68.8	63.7 - 73.5	4,100
Chinese	<50	1.3	0.4 - 4.2	100	10.4	6.9 - 15.3	<50	3.2	1.5 - 6.7	500	85.1	79.6 - 89.3	600
Filipino	100	4.9	3.1 - 7.8	700	23.2	19.0 - 28.0	100	3.7	2.1 - 6.3	2,000	68.2	63.0 - 73.0	2,900
Japanese	<50	0.5	0.1 - 3.0	200	10.8	7.4 - 15.6	<50	0.9	0.3 - 3.5	1,900	87.8	82.8 - 91.5	2,200
Other	100	4.2	2.1 - 8.1	400	17.2	12.7 - 23.0	<50	1.0	0.3 - 3.6	1,600	77.6	71.4 - 82.7	2,100

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.6	2.6 - 11.6	400	27.5	20.1 - 36.4	<50	3.4	1.3 - 8.4	800	63.6	54.4 - 71.8	1,300
20-24	200	6.2	4.1 - 9.3	900	25.0	20.8 - 29.8	100	3.9	2.3 - 6.5	2,400	64.9	59.8 - 69.7	3,700
25-34	300	3.9	2.7 - 5.6	900	12.0	9.9 - 14.5	300	3.7	2.6 - 5.3	6,100	80.4	77.5 - 83.1	7,600
35+	<50	1.5	0.6 - 3.9	300	9.6	6.7 - 13.4	100	1.9	0.9 - 3.9	2,400	87.1	82.8 - 90.4	2,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.4	3.2 - 6.2	1,200	15.3	12.9 - 18.0	300	4.3	3.1 - 6.0	5,900	76.0	72.9 - 78.9	7,800
Female	300	3.9	2.7 - 5.6	1,300	16.8	14.3 - 19.7	200	2.4	1.6 - 3.8	5,800	76.8	73.6 - 79.7	7,500

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.5	2.5 - 4.8	1,200	12.0	10.1 - 14.1	300	3.3	2.3 - 4.5	8,100	81.3	78.8 - 83.6	10,000
No	300	5.5	3.7 - 7.9	1,300	23.6	20.1 - 27.5	200	3.6	2.3 - 5.6	3,600	67.3	63.1 - 71.3	5,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.0	4.3 - 8.5	1,200	22.3	18.8 - 26.1	200	3.8	2.4 - 5.8	3,600	68.0	63.8 - 71.9	5,300
131-185%	100	4.6	2.2 - 9.7	300	17.5	12.4 - 24.1	<50	2.2	0.7 - 6.3	1,200	75.7	68.3 - 81.8	1,600
186+%	200	2.3	1.4 - 3.8	700	10.3	8.3 - 12.7	300	3.5	2.4 - 5.1	6,000	83.9	81.0 - 86.4	7,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	8.0	5.2 - 12.1	700	25.9	20.7 - 31.8	100	4.6	2.6 - 8.0	1,600	61.6	55.3 - 67.5	2,600
\$10,000-\$24,999	200	6.0	3.8 - 9.3	600	20.6	16.4 - 25.6	100	3.2	1.7 - 5.9	2,100	70.2	64.8 - 75.1	3,000
\$25,000-\$49,999	100	3.2	1.7 - 5.8	500	14.2	10.9 - 18.2	100	2.4	1.2 - 4.6	2,900	80.2	75.8 - 84.1	3,600
>=\$50,000	100	1.6	0.8 - 3.3	400	8.6	6.5 - 11.3	200	4.0	2.6 - 6.1	4,300	85.8	82.5 - 88.6	5,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.5	2.3 - 12.2	200	16.8	10.9 - 25.0	100	4.5	1.9 - 10.4	800	73.2	64.2 - 80.7	1,100
Normal birth weight	600	4.1	3.1 - 5.3	2,300	16.0	14.2 - 18.0	500	3.3	2.5 - 4.4	10,900	76.6	74.4 - 78.8	14,200

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	13.1	7.8 - 21.3	300	26.9	19.2 - 36.4	100	5.6	2.4 - 12.4	600	54.3	44.6 - 63.8	1,100
No	500	3.3	2.5 - 4.4	2,100	15.1	13.4 - 17.1	500	3.3	2.5 - 4.3	11,000	78.3	76.0 - 80.4	14,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	7.4	4.8 - 11.1	700	24.5	19.8 - 29.9	200	6.9	4.5 - 10.5	1,800	61.2	55.4 - 66.7	3,000
No	400	3.2	2.3 - 4.4	1,700	13.8	12.0 - 15.8	300	2.6	1.8 - 3.6	9,700	80.4	78.1 - 82.6	12,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.6	4.0 - 7.7	1,300	20.0	17.0 - 23.4	200	3.4	2.2 - 5.1	4,500	71.1	67.4 - 74.6	6,300
No	200	2.8	1.9 - 4.2	1,200	13.2	11.1 - 15.6	300	3.2	2.3 - 4.6	7,000	80.7	78.0 - 83.2	8,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	700	4.2	3.4 - 5.3	2,500	15.6	14.0 - 17.4	700	4.2	3.4 - 5.2	12,000	76.0	74.0 - 77.9	15,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.6	2.4 - 8.5	300	13.8	9.7 - 19.1	<50	1.6	0.6 - 4.2	1,500	80.1	74.1 - 85.0	1,900
Honolulu	400	3.8	2.9 - 5.0	1,800	15.5	13.6 - 17.6	500	4.6	3.6 - 5.8	8,700	76.2	73.8 - 78.4	11,400
Kauai	<50	5.3	2.1 - 12.8	100	14.5	8.5 - 23.5	<50	1.3	0.2 - 8.1	600	78.9	69.0 - 86.3	700
Maui	100	6.1	3.6 - 10.2	300	19.0	14.0 - 25.1	100	5.7	3.3 - 9.8	1,200	69.2	62.5 - 75.2	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	<50	1.3	0.6 - 2.9	500	13.5	10.5 - 17.0	100	3.9	2.4 - 6.2	3,000	81.4	77.4 - 84.8	3,600
Native Hawaiian	200	5.6	3.7 - 8.4	900	22.4	18.5 - 26.8	200	4.7	3.0 - 7.1	2,800	67.4	62.6 - 71.8	4,100
Chinese	<50	2.9	1.6 - 5.0	<50	6.7	4.7 - 9.5	<50	2.8	1.6 - 4.9	500	87.6	84.1 - 90.4	600
Filipino	200	7.9	5.7 - 10.9	500	17.6	14.3 - 21.6	200	5.3	3.5 - 7.8	2,100	69.2	64.7 - 73.4	3,000
Japanese	100	3.0	1.4 - 6.4	100	5.9	3.4 - 9.8	100	3.9	2.0 - 7.3	1,800	87.3	82.2 - 91.0	2,100
Other	100	3.1	1.4 - 6.4	300	15.6	11.4 - 20.9	100	3.0	1.5 - 5.7	1,800	78.4	72.6 - 83.2	2,200

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.1	4.4 - 14.6	300	27.9	20.5 - 36.8	<50	4.0	1.6 - 9.7	600	60.0	50.8 - 68.5	1,100
20-24	200	5.2	3.4 - 7.8	1,000	25.0	21.0 - 29.4	200	3.9	2.5 - 6.1	2,600	65.9	61.2 - 70.3	3,900
25-34	300	3.6	2.6 - 5.0	900	11.0	9.1 - 13.2	400	4.7	3.5 - 6.2	6,300	80.8	78.1 - 83.2	7,700
35+	100	3.2	1.8 - 5.6	300	10.8	7.9 - 14.5	100	3.3	1.9 - 5.7	2,500	82.8	78.4 - 86.4	3,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	4.3	3.2 - 5.8	1,300	15.8	13.6 - 18.2	400	4.6	3.5 - 6.1	6,400	75.3	72.5 - 77.9	8,500
Female	300	4.1	2.9 - 5.8	1,100	15.4	13.0 - 18.1	300	3.7	2.6 - 5.2	5,500	76.8	73.8 - 79.6	7,200

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.0	2.2 - 4.0	1,500	13.0	11.3 - 15.0	500	4.4	3.4 - 5.6	8,900	79.6	77.4 - 81.7	11,200
No	300	7.3	5.3 - 10.0	1,000	21.9	18.5 - 25.8	200	3.7	2.4 - 5.7	3,100	67.1	62.8 - 71.1	4,600

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.0	4.4 - 8.2	1,300	22.8	19.6 - 26.4	100	1.9	1.1 - 3.3	3,900	69.2	65.4 - 72.8	5,600
131-185%	100	3.1	1.5 - 6.4	200	13.4	9.3 - 18.9	100	4.9	2.7 - 8.8	1,500	78.6	72.5 - 83.7	1,900
186+%	200	2.3	1.4 - 3.5	800	10.8	8.9 - 13.1	400	6.1	4.7 - 7.8	5,800	80.9	78.1 - 83.3	7,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	8.8	5.8 - 13.1	500	23.1	18.3 - 28.7	100	3.2	1.6 - 6.1	1,500	64.9	58.8 - 70.6	2,400
\$10,000-\$24,999	200	5.7	3.9 - 8.3	800	21.2	17.5 - 25.5	100	2.2	1.2 - 4.2	2,600	70.8	66.2 - 75.0	3,700
\$25,000-\$49,999	100	2.1	1.1 - 3.9	600	13.9	11.0 - 17.3	300	6.8	4.9 - 9.3	3,200	77.2	73.3 - 80.8	4,100
>=\$50,000	100	1.7	0.9 - 3.1	400	8.5	6.5 - 11.1	200	4.3	2.9 - 6.4	3,900	85.5	82.3 - 88.2	4,500

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.1	3.6 - 13.7	200	15.6	10.0 - 23.5	<50	3.4	1.3 - 8.5	700	73.9	65.0 - 81.1	1,000
Normal birth weight	600	4.0	3.2 - 5.1	2,300	15.6	13.9 - 17.4	600	4.2	3.4 - 5.3	11,200	76.1	74.0 - 78.1	14,800

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.5	3.8 - 14.5	200	20.0	13.6 - 28.4	<50	3.9	1.5 - 9.5	700	68.6	59.4 - 76.5	1,100
No	600	3.9	3.1 - 5.0	2,200	15.1	13.5 - 17.0	600	4.2	3.3 - 5.2	11,100	76.8	74.7 - 78.7	14,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	3.8	2.2 - 6.5	600	19.3	15.3 - 24.0	200	5.8	3.8 - 8.9	2,200	71.1	65.9 - 75.8	3,000
No	500	4.3	3.3 - 5.4	1,800	14.5	12.8 - 16.5	500	3.7	2.9 - 4.8	9,700	77.5	75.2 - 79.5	12,500

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.7	3.4 - 6.5	1,400	20.3	17.6 - 23.4	200	3.0	2.0 - 4.4	4,900	71.9	68.6 - 75.0	6,900
No	300	3.8	2.8 - 5.2	1,000	11.9	10.0 - 14.1	400	5.2	4.0 - 6.7	6,800	79.1	76.5 - 81.4	8,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	700	4.7	3.6 - 6.0	2,400	16.3	14.3 - 18.6	600	4.4	3.4 - 5.8	11,000	74.6	72.0 - 77.1	14,800

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.6	2.8 - 7.6	300	18.2	14.7 - 22.4	<50	3.2	1.8 - 5.4	1,100	74.0	69.3 - 78.2	1,500
Honolulu	500	4.7	3.4 - 6.6	1,800	16.5	13.9 - 19.5	500	4.2	3.0 - 6.1	8,100	74.5	71.1 - 77.7	10,900
Kauai	<50	5.4	2.7 - 10.4	100	7.2	4.3 - 11.7	100	7.2	4.1 - 12.5	600	80.2	73.4 - 85.6	800
Maui	100	3.9	2.3 - 6.5	300	17.8	14.1 - 22.2	100	5.3	3.4 - 8.2	1,200	73.0	68.1 - 77.4	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.2	0.9 - 5.5	400	12.7	8.6 - 18.3	100	4.3	2.2 - 8.1	2,800	80.9	74.7 - 85.8	3,400
Native Hawaiian	200	6.5	5.0 - 8.5	800	20.6	18.0 - 23.4	200	4.8	3.6 - 6.4	2,600	68.1	64.9 - 71.2	3,800
Chinese	0	0.0	-	<50	6.4	1.7 - 21.6	<50	1.9	0.5 - 6.9	600	91.7	78.0 - 97.2	700
Filipino	200	5.5	3.2 - 9.4	700	21.1	15.9 - 27.3	200	5.9	3.4 - 10.1	2,100	67.5	60.7 - 73.7	3,100
Japanese	100	6.0	2.6 - 13.3	100	6.8	3.3 - 13.7	<50	2.5	0.8 - 8.0	1,500	84.7	76.1 - 90.6	1,800
Other	100	4.4	1.8 - 10.2	400	19.0	12.6 - 27.7	100	4.0	1.6 - 9.9	1,400	72.6	63.3 - 80.2	1,900

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	13.8	7.9 - 22.7	400	34.3	26.0 - 43.6	100	4.1	2.3 - 7.2	600	47.9	39.0 - 57.0	1,300
20-24	100	3.9	2.7 - 5.8	800	25.7	20.7 - 31.4	100	4.0	2.2 - 7.4	2,200	66.4	60.5 - 71.8	3,300
25-34	300	4.6	3.1 - 6.6	900	11.5	9.1 - 14.5	400	4.9	3.4 - 7.0	5,900	79.0	75.4 - 82.2	7,500
35+	<50	1.4	0.4 - 4.6	300	9.7	6.1 - 15.0	100	3.7	1.7 - 7.8	2,300	85.2	79.2 - 89.7	2,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.2	2.9 - 6.1	1,300	17.0	14.2 - 20.3	400	4.6	3.1 - 6.7	5,800	74.2	70.4 - 77.6	7,800
Female	400	5.2	3.6 - 7.4	1,100	15.6	12.7 - 18.9	300	4.2	2.9 - 6.0	5,200	75.1	71.3 - 78.5	6,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.4	1.6 - 3.7	1,300	13.0	10.6 - 15.7	500	4.6	3.3 - 6.5	8,100	80.0	76.8 - 82.8	10,200
No	400	9.6	7.0 - 13.0	1,100	23.8	19.9 - 28.2	200	3.9	2.7 - 5.7	2,900	62.7	58.0 - 67.2	4,600

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.1	4.3 - 8.5	1,500	23.6	20.0 - 27.6	300	4.2	2.9 - 6.2	4,100	66.1	61.8 - 70.2	6,200
131-185%	100	4.3	2.0 - 8.9	300	14.2	9.8 - 20.1	100	4.4	2.0 - 9.3	1,700	77.1	70.1 - 82.9	2,200
186+%	100	2.2	1.0 - 4.6	400	7.8	5.4 - 11.2	200	4.0	2.4 - 6.6	4,400	86.1	81.9 - 89.4	5,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	300	8.6	5.7 - 12.6	700	23.9	19.1 - 29.4	100	4.6	2.6 - 8.1	1,900	62.9	56.9 - 68.6	3,100
\$15,001-\$25,000	100	2.9	1.1 - 7.1	400	22.2	16.1 - 30.0	<50	2.2	1.1 - 4.3	1,300	72.7	64.9 - 79.4	1,900
\$25,001-\$40,000	200	6.0	3.5 - 10.2	500	17.5	12.9 - 23.4	200	5.3	3.0 - 9.4	2,100	71.1	64.5 - 76.9	2,900
>=\$40,001	100	1.7	0.8 - 3.5	600	9.3	6.9 - 12.5	300	4.6	3.0 - 7.0	5,100	84.4	80.6 - 87.6	6,100

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	4.4	1.5 - 12.1	200	21.9	13.5 - 33.5	100	5.6	2.0 - 14.8	700	68.1	56.3 - 78.1	1,000
Normal birth weight	600	4.7	3.6 - 6.1	2,200	15.9	13.8 - 18.3	600	4.3	3.3 - 5.7	10,300	75.1	72.4 - 77.6	13,700

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.4	4.4 - 15.7	200	22.1	14.6 - 32.1	<50	3.1	1.4 - 7.0	600	66.3	55.9 - 75.3	1,000
No	600	4.4	3.3 - 5.8	2,200	16.0	13.9 - 18.4	600	4.4	3.3 - 5.9	10,200	75.1	72.4 - 77.7	13,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	9.2	6.0 - 13.9	600	22.1	17.1 - 28.1	200	6.6	4.0 - 10.7	1,600	62.1	55.5 - 68.2	2,600
No	400	3.7	2.6 - 5.1	1,700	14.8	12.6 - 17.3	500	3.9	2.8 - 5.5	9,100	77.6	74.7 - 80.2	11,800

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	6.5	4.6 - 9.1	1,400	24.8	21.1 - 28.9	200	3.9	2.6 - 5.8	3,800	64.8	60.4 - 68.9	5,800
No	300	3.5	2.4 - 5.2	900	10.8	8.5 - 13.6	400	4.6	3.2 - 6.7	7,100	81.1	77.7 - 84.0	8,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.3	3.2 - 5.7	2,200	15.4	13.4 - 17.7	400	2.6	1.8 - 3.6	11,200	77.7	75.1 - 80.1	14,400

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.7	3.0 - 7.3	300	19.5	16.0 - 23.5	100	3.6	2.1 - 6.0	1,200	72.3	67.7 - 76.4	1,700
Honolulu	400	4.2	2.9 - 6.1	1,500	14.0	11.4 - 17.0	300	2.6	1.6 - 4.0	8,400	79.3	75.9 - 82.3	10,500
Kauai	<50	1.2	0.2 - 7.2	100	21.6	15.4 - 29.4	<50	2.9	1.0 - 8.0	400	74.3	66.1 - 81.1	600
Maui	100	5.5	3.5 - 8.5	300	18.8	14.9 - 23.3	<50	1.5	0.7 - 3.4	1,200	74.2	69.3 - 78.6	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.0	1.4 - 6.3	300	10.5	6.7 - 16.1	200	5.9	3.3 - 10.1	2,500	80.6	74.3 - 85.6	3,100
Native Hawaiian	200	4.9	3.6 - 6.7	700	19.4	17.0 - 22.2	100	3.4	2.3 - 4.9	2,700	72.2	69.1 - 75.1	3,700
Chinese	<50	6.5	1.9 - 20.3	100	11.4	3.9 - 29.4	<50	3.8	0.6 - 21.8	400	78.2	60.1 - 89.5	500
Filipino	200	5.8	3.1 - 10.7	600	20.6	15.5 - 26.9	<50	1.3	0.4 - 3.9	2,100	72.3	65.4 - 78.2	3,000
Japanese	<50	0.2	0.0 - 1.4	200	7.8	4.4 - 13.6	<50	0.2	0.0 - 1.4	2,100	91.7	86.0 - 95.2	2,300
Other	100	7.3	3.3 - 15.2	300	18.2	11.4 - 27.7	0	0.0	-	1,300	74.5	64.4 - 82.6	1,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	7.4	4.1 - 13.1	400	30.9	22.7 - 40.4	<50	1.1	0.4 - 3.2	700	60.6	51.0 - 69.4	1,200
20-24	300	7.1	4.4 - 11.3	700	19.9	15.5 - 25.1	100	2.8	1.7 - 4.4	2,500	70.3	64.6 - 75.4	3,600
25-34	200	3.0	1.9 - 4.8	900	12.5	10.0 - 15.6	200	2.9	1.8 - 4.7	5,900	81.6	78.1 - 84.7	7,200
35+	100	2.3	0.9 - 6.3	200	9.8	6.4 - 14.7	<50	2.1	0.7 - 6.2	2,000	85.8	79.9 - 90.1	2,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.9	3.2 - 7.4	1,100	15.7	12.9 - 19.0	200	2.4	1.4 - 3.9	5,400	77.0	73.3 - 80.4	7,000
Female	300	3.7	2.5 - 5.4	1,100	15.2	12.4 - 18.5	200	2.8	1.7 - 4.4	5,700	78.3	74.7 - 81.5	7,300

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.2	2.1 - 5.0	1,200	12.2	10.0 - 14.9	300	2.6	1.6 - 4.0	8,200	82.0	78.9 - 84.7	10,000
No	300	6.7	4.6 - 9.7	1,000	22.9	19.0 - 27.3	100	2.6	1.7 - 4.0	2,900	67.8	63.1 - 72.1	4,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.4	4.3 - 9.5	1,000	19.6	16.2 - 23.6	100	2.3	1.5 - 3.5	3,800	71.6	67.2 - 75.7	5,300
131-185%	100	4.5	2.5 - 8.2	500	15.9	11.6 - 21.4	100	2.7	1.2 - 6.1	2,300	76.9	70.7 - 82.0	3,000
186+%	100	1.6	0.8 - 3.4	500	9.1	6.6 - 12.6	200	3.1	1.7 - 5.7	4,300	86.2	82.2 - 89.4	5,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	200	7.6	4.7 - 12.0	500	21.1	16.1 - 27.1	100	3.2	1.9 - 5.3	1,700	68.2	61.8 - 73.9	2,500
\$15,001-\$25,000	100	6.2	3.2 - 11.6	300	15.3	10.8 - 21.2	100	3.0	1.4 - 6.2	1,600	75.6	68.6 - 81.4	2,100
\$25,001-\$40,000	200	5.5	2.9 - 9.9	500	16.4	12.2 - 21.8	100	2.5	1.0 - 5.9	2,300	75.7	69.4 - 81.0	3,000
>=\$40,001	100	1.8	0.9 - 3.5	700	10.9	8.3 - 14.3	100	2.4	1.3 - 4.4	5,100	84.9	81.2 - 88.0	6,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	6.0	1.9 - 17.0	200	22.2	13.4 - 34.5	<50	2.7	0.6 - 11.9	600	69.0	56.1 - 79.6	900
Normal birth weight	600	4.2	3.1 - 5.6	2,000	15.0	12.9 - 17.3	300	2.6	1.8 - 3.6	10,500	78.3	75.7 - 80.7	13,400

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	5.5	2.2 - 13.1	300	30.5	21.3 - 41.6	<50	5.2	2.6 - 10.3	500	58.7	47.8 - 68.9	900
No	600	4.2	3.1 - 5.7	1,900	14.4	12.4 - 16.7	300	2.4	1.7 - 3.6	10,500	78.9	76.3 - 81.4	13,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	7.4	4.3 - 12.2	600	22.8	17.7 - 28.9	100	4.5	2.6 - 7.9	1,700	65.3	58.6 - 71.5	2,600
No	300	3.0	2.1 - 4.4	1,600	14.0	11.9 - 16.5	300	2.2	1.4 - 3.4	9,200	80.7	77.9 - 83.2	11,400

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	6.4	4.3 - 9.4	1,200	22.1	18.4 - 26.2	100	2.5	1.5 - 4.1	3,900	69.0	64.5 - 73.2	5,600
No	300	3.0	2.0 - 4.4	1,000	11.3	9.1 - 13.9	200	2.6	1.6 - 4.1	7,200	83.2	80.1 - 85.8	8,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.6	3.7 - 5.6	2,400	17.0	15.3 - 18.9	400	3.0	2.2 - 3.9	10,600	75.4	73.3 - 77.5	14,000

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.1	3.3 - 7.8	200	15.0	11.9 - 18.9	100	3.5	2.0 - 6.0	1,200	76.4	71.9 - 80.4	1,500
Honolulu	400	4.3	3.3 - 5.6	1,800	17.4	15.1 - 19.8	300	2.8	1.9 - 4.0	7,800	75.6	72.8 - 78.1	10,400
Kauai	<50	4.2	2.0 - 8.4	100	15.0	9.9 - 22.2	<50	4.2	1.9 - 9.2	500	76.6	68.8 - 82.9	700
Maui	100	5.9	3.8 - 9.0	300	17.3	13.5 - 21.8	100	3.4	2.0 - 5.8	1,100	73.4	68.4 - 77.9	1,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.2	1.1 - 4.3	400	12.7	9.5 - 16.7	100	1.9	0.9 - 4.2	2,600	83.2	78.8 - 86.8	3,100
Native Hawaiian	300	7.3	6.0 - 8.9	700	20.9	18.7 - 23.2	100	3.2	2.4 - 4.3	2,400	68.6	66.0 - 71.0	3,600
Chinese	<50	3.1	1.1 - 8.5	100	15.8	7.5 - 30.3	<50	5.0	1.3 - 17.3	500	76.1	61.6 - 86.4	600
Filipino	200	5.4	3.4 - 8.5	600	20.4	15.9 - 25.7	100	3.1	1.6 - 6.1	2,100	71.1	65.3 - 76.2	3,000
Japanese	<50	0.9	0.3 - 2.6	200	12.4	8.3 - 18.3	100	4.2	2.0 - 9.0	1,600	82.4	75.9 - 87.4	1,900
Other	100	6.2	3.3 - 11.3	300	16.5	11.4 - 23.4	<50	2.0	0.7 - 5.2	1,400	75.3	67.8 - 81.5	1,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.9	6.6 - 14.7	400	34.0	27.8 - 40.8	<50	0.8	0.3 - 2.1	700	55.3	48.4 - 62.0	1,300
20-24	200	6.6	4.6 - 9.6	800	22.6	18.9 - 26.8	100	3.1	1.8 - 5.2	2,400	67.6	63.1 - 71.9	3,600
25-34	200	3.2	2.3 - 4.2	900	13.4	11.1 - 16.2	200	3.0	2.0 - 4.5	5,500	80.4	77.4 - 83.1	6,800
35+	100	2.5	1.3 - 4.8	200	9.1	5.9 - 13.6	100	4.0	2.1 - 7.5	1,900	84.4	79.2 - 88.5	2,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	5.0	3.7 - 6.5	1,200	17.0	14.6 - 19.8	200	2.8	1.8 - 4.3	5,500	75.2	72.1 - 78.1	7,300
Female	300	4.1	3.1 - 5.5	1,100	17.0	14.6 - 19.7	200	3.2	2.1 - 4.6	5,100	75.7	72.7 - 78.5	6,700

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.2	2.4 - 4.2	1,300	13.8	11.8 - 16.0	300	3.3	2.3 - 4.7	7,700	79.8	77.2 - 82.1	9,600
No	300	7.6	5.8 - 10.0	1,100	24.2	20.9 - 27.8	100	2.2	1.4 - 3.5	2,900	66.0	62.1 - 69.7	4,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.4	4.8 - 8.4	1,000	20.5	17.6 - 23.8	100	2.5	1.6 - 4.0	3,500	70.6	66.9 - 73.9	5,000
131-185%	100	3.0	2.0 - 4.5	500	16.5	13.0 - 20.6	100	2.8	1.7 - 4.6	2,400	77.8	73.4 - 81.6	3,100
186+%	100	2.7	1.7 - 4.1	500	10.0	7.7 - 12.9	200	3.8	2.4 - 6.0	4,200	83.5	80.1 - 86.5	5,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	200	8.0	5.7 - 11.3	500	20.1	16.1 - 24.7	100	2.1	1.1 - 4.0	1,800	69.8	64.7 - 74.5	2,600
\$15,001-\$25,000	100	4.9	2.8 - 8.4	400	23.6	18.5 - 29.6	100	4.3	2.2 - 8.4	1,200	67.2	60.8 - 73.0	1,800
\$25,001-\$40,000	100	2.8	2.0 - 4.1	600	18.2	14.6 - 22.4	100	2.5	1.4 - 4.3	2,400	76.5	72.1 - 80.4	3,100
>=\$40,001	200	2.8	2.0 - 4.1	600	10.7	8.5 - 13.5	200	3.6	2.3 - 5.5	4,800	82.8	79.6 - 85.6	5,800

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	4.4	1.8 - 10.3	200	22.7	15.4 - 32.2	<50	2.1	0.8 - 5.4	600	70.8	61.1 - 78.9	900
Normal birth weight	600	4.6	3.7 - 5.6	2,200	16.7	14.9 - 18.6	400	3.0	2.3 - 4.1	10,000	75.7	73.5 - 77.8	13,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	6.3	4.0 - 9.9	300	27.7	20.4 - 36.3	<50	2.8	1.3 - 5.7	700	63.2	54.8 - 70.9	1,100
No	500	4.3	3.4 - 5.3	2,000	16.0	14.3 - 18.0	400	3.0	2.2 - 4.1	9,800	76.6	74.4 - 78.7	12,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	5.1	3.6 - 7.0	700	26.0	21.5 - 31.1	100	3.0	1.8 - 5.1	1,700	65.9	60.7 - 70.7	2,600
No	500	4.2	3.3 - 5.4	1,600	14.8	12.9 - 16.9	300	2.9	2.1 - 4.0	8,500	78.1	75.7 - 80.3	10,900

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	6.8	5.3 - 8.7	1,200	21.7	18.8 - 24.8	200	3.4	2.3 - 5.1	3,900	68.2	64.7 - 71.4	5,700
No	200	3.0	2.2 - 4.1	1,100	13.8	11.6 - 16.3	200	2.7	1.8 - 4.0	6,600	80.5	77.8 - 83.0	8,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	3.8	3.0 - 4.9	2,300	16.3	14.7 - 18.1	600	4.0	3.2 - 5.0	10,600	75.9	73.8 - 77.8	14,000

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.0	3.1 - 8.1	400	23.2	19.3 - 27.7	100	3.8	2.3 - 6.4	1,100	67.9	63.0 - 72.5	1,600
Honolulu	400	3.7	2.7 - 5.1	1,500	14.9	13.0 - 17.1	400	4.1	3.1 - 5.4	8,000	77.3	74.7 - 79.7	10,400
Kauai	<50	3.5	1.4 - 8.5	100	15.6	10.2 - 23.0	<50	1.9	0.5 - 6.4	500	79.0	71.1 - 85.2	600
Maui	100	3.5	2.0 - 6.2	300	18.3	14.5 - 22.9	100	4.2	2.4 - 7.1	1,100	73.9	68.9 - 78.4	1,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.6	2.1 - 6.3	400	13.0	9.9 - 17.0	200	4.4	2.9 - 6.8	2,700	78.9	74.4 - 82.8	3,400
Native Hawaiian	200	4.8	3.7 - 6.1	900	24.4	22.0 - 26.9	200	4.4	3.4 - 5.7	2,400	66.4	63.7 - 69.0	3,600
Chinese	<50	2.5	0.4 - 15.1	100	18.0	9.3 - 32.0	<50	2.5	0.4 - 15.1	400	76.9	62.4 - 87.0	600
Filipino	100	4.3	2.4 - 7.6	500	17.4	13.6 - 22.0	100	3.4	2.0 - 5.9	2,100	74.9	69.7 - 79.5	2,900
Japanese	<50	1.0	0.4 - 2.9	100	4.5	2.1 - 9.4	100	4.6	2.0 - 10.1	1,600	89.8	83.7 - 93.8	1,800
Other	100	5.0	2.3 - 10.4	300	15.7	10.5 - 22.7	100	2.9	1.1 - 7.0	1,400	76.5	68.8 - 82.8	1,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.4	5.7 - 12.3	400	28.7	23.3 - 34.8	100	3.7	1.8 - 7.4	900	59.2	52.7 - 65.3	1,600
20-24	200	5.7	3.7 - 8.7	700	21.4	18.0 - 25.3	100	3.4	2.3 - 5.1	2,300	69.5	65.1 - 73.5	3,300
25-34	200	2.8	1.8 - 4.3	900	13.3	11.1 - 15.9	300	4.0	2.9 - 5.5	5,400	79.9	76.9 - 82.6	6,800
35+	<50	1.4	0.5 - 4.1	200	9.6	6.6 - 13.9	100	4.8	2.6 - 8.7	2,000	84.2	79.0 - 88.3	2,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.9	2.8 - 5.4	1,200	16.8	14.5 - 19.4	200	3.3	2.3 - 4.5	5,400	76.0	73.1 - 78.7	7,100
Female	300	3.8	2.6 - 5.5	1,100	15.8	13.6 - 18.3	300	4.7	3.4 - 6.4	5,300	75.7	72.7 - 78.5	6,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.4	1.6 - 3.6	1,300	13.2	11.4 - 15.3	400	4.0	3.0 - 5.3	7,800	80.4	77.9 - 82.6	9,700
No	300	7.1	5.3 - 9.6	1,000	23.2	20.1 - 26.6	200	3.9	2.7 - 5.6	2,800	65.8	62.0 - 69.5	4,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.1	3.8 - 7.0	1,300	22.2	19.4 - 25.2	200	3.5	2.5 - 4.8	4,000	69.2	65.9 - 72.4	5,800
131-185%	100	3.6	1.8 - 7.0	300	13.3	10.0 - 17.5	<50	1.9	1.0 - 3.4	1,600	81.2	76.4 - 85.3	2,000
186+%	100	2.2	1.2 - 3.8	600	10.6	8.3 - 13.4	300	5.3	3.6 - 7.5	4,400	82.0	78.6 - 85.0	5,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	200	7.8	5.5 - 10.9	800	24.5	20.6 - 28.8	100	3.4	2.1 - 5.4	2,000	64.3	59.5 - 68.8	3,100
\$15,001-\$25,000	100	2.8	1.4 - 5.6	400	20.2	15.8 - 25.5	100	3.1	1.8 - 5.3	1,600	73.8	68.2 - 78.7	2,100
\$25,001-\$40,000	100	3.0	1.5 - 5.8	400	15.6	12.4 - 19.5	100	3.3	2.1 - 5.0	2,200	78.2	73.8 - 82.0	2,800
>=\$40,001	100	2.0	1.2 - 3.4	500	9.8	7.7 - 12.4	300	4.9	3.3 - 7.1	4,400	83.3	80.0 - 86.2	5,300

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	5.5	2.1 - 13.4	200	20.7	14.4 - 28.7	<50	4.0	1.9 - 8.4	600	69.8	60.6 - 77.6	900
Normal birth weight	500	3.8	2.9 - 4.8	2,100	15.8	14.1 - 17.6	500	3.9	3.0 - 4.9	10,000	76.6	74.5 - 78.6	13,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	10.7	6.3 - 17.7	300	29.5	22.3 - 37.9	<50	3.2	1.6 - 6.5	500	56.6	47.9 - 64.8	900
No	400	3.1	2.3 - 4.1	1,900	15.2	13.5 - 17.0	500	3.9	3.1 - 5.0	9,900	77.9	75.7 - 79.8	12,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	6.4	4.3 - 9.5	700	27.1	22.7 - 32.0	100	4.5	2.7 - 7.6	1,600	62.0	56.8 - 67.0	2,600
No	300	3.0	2.1 - 4.1	1,500	13.4	11.7 - 15.4	400	3.7	2.9 - 4.8	8,800	79.8	77.6 - 81.9	11,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	5.6	4.1 - 7.6	1,200	20.8	18.3 - 23.5	200	3.2	2.3 - 4.4	4,100	70.5	67.3 - 73.4	5,800
No	200	2.4	1.5 - 3.6	1,000	13.1	11.0 - 15.6	400	4.4	3.2 - 6.0	6,400	80.1	77.3 - 82.7	8,000