

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration (grouped), among initiators	Breastfeeding (any) duration (grouped), among initiators

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/28/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	500	3.8	3.0 - 4.9	2,300	16.3	14.7 - 18.1	600	4.0	3.2 - 5.0	10,600	75.9	73.8 - 77.8	14,000
2001	600	4.6	3.7 - 5.6	2,400	17.0	15.3 - 18.9	400	3.0	2.2 - 3.9	10,600	75.4	73.3 - 77.5	14,000
2002	600	4.3	3.2 - 5.7	2,200	15.4	13.4 - 17.7	400	2.6	1.8 - 3.6	11,200	77.7	75.1 - 80.1	14,400
2003	700	4.7	3.6 - 6.0	2,400	16.3	14.3 - 18.6	600	4.4	3.4 - 5.8	11,000	74.6	72.0 - 77.1	14,800
2004	700	4.2	3.4 - 5.3	2,500	15.6	14.0 - 17.4	700	4.2	3.4 - 5.2	12,000	76.0	74.0 - 77.9	15,700
2005	600	4.2	3.2 - 5.3	2,500	16.0	14.3 - 18.0	500	3.4	2.6 - 4.4	11,700	76.4	74.2 - 78.5	15,300
2006	500	2.9	2.1 - 3.9	2,800	17.4	15.5 - 19.5	600	3.5	2.7 - 4.6	12,400	76.2	73.9 - 78.4	16,300
2007	600	3.8	2.9 - 4.9	2,700	16.2	14.4 - 18.1	500	3.3	2.5 - 4.3	12,700	76.7	74.5 - 78.8	16,600
2008	500	2.9	2.1 - 3.8	2,600	15.3	13.6 - 17.2	800	4.6	3.6 - 5.7	12,900	77.3	75.1 - 79.3	16,700
2009	800	4.6	3.4 - 6.3	2,600	15.6	13.4 - 18.2	900	5.3	4.0 - 7.0	12,500	74.4	71.4 - 77.2	16,700
2010	600	3.3	2.3 - 4.7	2,400	14.2	12.0 - 16.7	700	4.3	3.1 - 5.9	13,400	78.3	75.4 - 80.9	17,200
2011	300	1.7	1.1 - 2.8	2,200	12.7	10.7 - 15.1	700	4.0	2.9 - 5.4	14,200	81.5	78.8 - 83.9	17,500
2012	500	2.8	1.8 - 4.2	2,200	13.0	10.9 - 15.4	600	3.2	2.3 - 4.7	14,000	81.0	78.2 - 83.5	17,300
2013	400	2.1	1.2 - 3.4	2,100	12.0	9.8 - 14.5	500	3.0	2.0 - 4.6	14,400	82.9	80.1 - 85.5	17,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	400	2.1	1.2 - 3.4	2,100	12.0	9.8 - 14.5	500	3.0	2.0 - 4.6	14,400	82.9	80.1 - 85.5	17,400

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	2.5	1.2 - 5.3	200	10.2	7.2 - 14.2	100	2.4	1.1 - 5.0	1,900	85.0	80.3 - 88.7	2,200
Honolulu	200	2.0	1.0 - 3.9	1,500	12.0	9.2 - 15.4	400	3.3	1.9 - 5.5	10,400	82.8	78.9 - 86.1	12,500
Kauai	<50	0.9	0.3 - 2.9	100	13.0	9.8 - 17.2	<50	2.9	1.6 - 5.4	700	83.1	78.6 - 86.8	800
Maui	100	2.7	1.4 - 5.3	200	13.3	9.8 - 17.6	<50	2.3	1.1 - 4.8	1,500	81.7	76.8 - 85.7	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.2	0.7 - 6.5	500	9.6	6.0 - 14.8	100	2.7	1.2 - 6.2	4,100	85.5	79.5 - 89.9	4,800
Native Hawaiian	100	2.8	1.4 - 5.6	900	17.2	12.8 - 22.7	200	4.0	2.1 - 7.4	3,800	76.0	70.1 - 81.1	5,000
Chinese	<50	0.9	0.1 - 5.7	100	17.0	7.8 - 33.0	0	0.0	-	700	82.1	66.2 - 91.5	900
Filipino	100	3.4	1.3 - 8.8	300	12.3	8.1 - 18.1	100	4.2	1.6 - 10.6	2,200	80.1	72.5 - 86.0	2,800
Japanese	0	0.0	-	100	3.0	0.9 - 9.3	0	0.0	-	1,700	97.0	90.7 - 99.1	1,800
Other	<50	0.5	0.1 - 3.3	200	10.0	5.2 - 18.3	100	3.6	1.1 - 11.1	1,800	85.9	76.8 - 91.8	2,100

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	1.8	0.5 - 6.6	200	21.8	12.0 - 36.4	<50	5.9	1.3 - 22.6	600	70.5	55.0 - 82.4	800
20-24	100	1.9	0.7 - 5.2	400	9.8	6.4 - 14.8	200	5.2	2.5 - 10.7	3,300	83.0	76.6 - 88.0	3,900
25-34	200	2.3	1.1 - 4.5	1,200	13.1	10.1 - 16.9	200	1.7	0.9 - 3.1	7,800	82.9	78.8 - 86.4	9,400
35+	100	1.7	0.5 - 5.3	300	8.7	5.2 - 14.2	100	3.7	1.5 - 8.4	2,800	86.0	79.5 - 90.6	3,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	200	2.2	1.1 - 4.4	900	10.6	7.9 - 14.0	200	2.5	1.3 - 4.8	7,200	84.7	80.7 - 88.0	8,500
Female	200	1.9	0.9 - 4.0	1,200	13.3	10.2 - 17.1	300	3.5	2.0 - 6.1	7,200	81.3	76.9 - 84.9	8,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.1	1.1 - 4.1	1,300	11.2	8.6 - 14.5	200	2.1	1.2 - 3.8	9,500	84.6	80.9 - 87.6	11,200
No	100	2.0	1.0 - 3.9	800	13.3	10.1 - 17.4	300	4.7	2.6 - 8.3	4,900	80.0	75.1 - 84.1	6,100

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	100	2.4	1.2 - 4.8	900	15.1	11.4 - 19.8	300	4.6	2.6 - 8.1	4,900	77.9	72.6 - 82.4	6,300
131-185%	100	2.4	0.8 - 7.5	300	12.7	7.4 - 21.1	<50	1.8	0.4 - 8.9	1,900	83.0	74.1 - 89.3	2,300
186+%	200	2.1	0.9 - 5.0	500	7.6	5.1 - 11.1	100	1.9	0.9 - 4.1	6,300	88.4	84.2 - 91.6	7,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Income (2012+)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	100	2.1	0.8 - 5.3	600	16.6	11.8 - 22.9	200	4.6	2.1 - 9.7	3,000	76.6	69.7 - 82.4	3,900
\$17,001-\$26,000	<50	0.6	0.1 - 2.7	300	15.0	9.2 - 23.5	100	2.3	0.6 - 8.3	1,900	82.0	73.2 - 88.4	2,300
\$26,001-\$51,000	200	5.0	2.4 - 10.0	300	9.2	5.6 - 14.8	100	3.7	1.6 - 8.1	3,100	82.2	75.3 - 87.5	3,700
>=\$51,001	100	1.2	0.3 - 4.1	500	7.7	5.0 - 11.6	100	1.7	0.7 - 3.9	5,400	89.4	85.1 - 92.6	6,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	1.1	0.4 - 3.1	200	15.1	12.0 - 19.0	100	4.3	2.7 - 6.9	1,000	79.4	75.1 - 83.1	1,200
Normal birth weight	300	2.1	1.3 - 3.6	1,900	11.7	9.5 - 14.4	500	2.9	1.8 - 4.6	13,400	83.2	80.1 - 85.9	16,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.1	2.1 - 21.2	200	21.3	11.7 - 35.7	<50	1.6	0.5 - 4.9	600	69.9	54.9 - 81.6	900
No	300	1.8	1.1 - 3.2	1,900	11.5	9.3 - 14.1	500	3.1	2.0 - 4.8	13,500	83.6	80.6 - 86.2	16,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.6	2.8 - 11.0	500	16.8	11.3 - 24.2	100	3.1	1.1 - 8.5	2,300	74.4	66.3 - 81.2	3,100
No	200	1.3	0.6 - 2.8	1,500	11.0	8.7 - 13.7	400	3.0	1.9 - 4.8	11,800	84.7	81.6 - 87.4	13,900

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.4	1.2 - 4.7	1,200	16.6	12.9 - 21.1	300	4.3	2.5 - 7.1	5,700	76.7	71.8 - 81.0	7,400
No	200	1.8	0.9 - 3.8	800	8.2	6.0 - 11.1	200	2.1	1.1 - 4.2	8,700	87.8	84.4 - 90.6	9,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	2.8	1.8 - 4.2	2,200	13.0	10.9 - 15.4	600	3.2	2.3 - 4.7	14,000	81.0	78.2 - 83.5	17,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	<50	1.2	0.4 - 3.5	300	15.1	11.5 - 19.4	100	5.1	3.1 - 8.2	1,700	78.6	73.8 - 82.8	2,200
Honolulu	400	3.1	1.9 - 5.2	1,600	12.6	9.9 - 15.9	400	2.9	1.7 - 4.9	10,100	81.4	77.6 - 84.7	12,400
Kauai	<50	3.5	1.8 - 7.0	100	11.7	8.0 - 16.6	<50	2.2	0.9 - 5.1	700	82.6	77.0 - 87.1	800
Maui	<50	2.0	0.9 - 4.1	300	13.9	10.7 - 18.0	100	3.8	2.3 - 6.2	1,500	80.3	75.8 - 84.1	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.1	1.3 - 7.2	300	8.8	5.5 - 13.8	100	1.4	0.5 - 4.4	3,400	86.7	81.0 - 90.9	3,900
Native Hawaiian	100	2.1	0.9 - 4.7	900	18.8	14.5 - 24.1	300	5.6	3.5 - 8.9	3,600	73.5	67.8 - 78.5	4,900
Chinese	100	5.2	1.3 - 18.0	100	5.7	1.6 - 17.9	<50	3.2	0.5 - 18.9	900	86.0	71.5 - 93.7	1,000
Filipino	100	4.3	2.0 - 9.1	600	20.1	14.4 - 27.4	100	4.3	1.9 - 9.6	2,200	71.2	63.4 - 78.0	3,100
Japanese	<50	0.3	0.1 - 1.7	100	5.9	2.6 - 13.0	<50	1.0	0.4 - 3.0	1,400	92.7	85.9 - 96.4	1,500
Other	100	2.4	0.7 - 7.8	200	7.3	4.0 - 13.1	<50	1.7	0.4 - 6.5	2,500	88.5	81.8 - 93.0	2,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.2	3.5 - 22.4	200	25.3	15.2 - 39.0	<50	2.7	0.9 - 7.7	600	62.8	48.7 - 75.0	1,000
20-24	100	3.6	1.6 - 8.1	600	17.5	12.5 - 24.1	100	3.7	1.7 - 8.0	2,700	75.1	68.0 - 81.1	3,600
25-34	200	2.0	1.1 - 3.8	1,200	12.0	9.3 - 15.3	400	3.7	2.3 - 6.0	8,000	82.3	78.5 - 85.5	9,700
35+	100	2.2	0.7 - 6.3	200	7.0	4.2 - 11.4	<50	1.3	0.6 - 2.6	2,700	89.5	84.5 - 93.1	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.6	2.1 - 6.0	1,100	13.3	10.4 - 16.8	200	2.7	1.5 - 4.9	7,000	80.4	76.3 - 83.9	8,700
Female	200	2.0	1.0 - 3.9	1,100	12.7	9.9 - 16.3	300	3.8	2.4 - 6.0	7,000	81.5	77.6 - 84.9	8,600

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.4	1.3 - 4.3	1,200	10.8	8.4 - 13.8	300	2.7	1.6 - 4.4	9,100	84.2	80.7 - 87.1	10,700
No	200	3.5	2.0 - 6.1	1,100	16.7	13.0 - 21.1	300	4.2	2.5 - 7.0	4,900	75.6	70.7 - 79.9	6,500

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.5	2.6 - 7.6	1,000	17.3	13.5 - 22.0	200	2.6	1.3 - 5.1	4,500	75.6	70.4 - 80.1	6,000
131-185%	<50	1.4	0.3 - 6.4	400	14.0	9.1 - 21.1	200	5.6	2.8 - 10.9	2,300	79.0	71.2 - 85.1	2,900
186+%	100	1.8	0.7 - 4.2	600	8.7	6.1 - 12.3	200	2.8	1.5 - 5.2	5,900	86.7	82.5 - 89.9	6,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Income (2012+)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	200	4.7	2.5 - 8.9	700	17.2	12.5 - 23.1	100	2.3	0.9 - 5.7	2,900	75.8	69.2 - 81.4	3,800
\$17,001-\$26,000	100	2.4	0.6 - 8.8	300	14.5	9.3 - 21.8	100	6.5	3.2 - 12.5	1,700	76.7	68.2 - 83.5	2,200
\$26,001-\$51,000	100	2.7	1.1 - 6.3	700	16.1	11.4 - 22.2	100	2.8	1.2 - 6.2	3,200	78.4	71.9 - 83.8	4,000
>=\$51,001	100	1.6	0.6 - 4.3	500	7.9	5.3 - 11.6	200	2.9	1.5 - 5.7	5,100	87.6	83.2 - 91.0	5,800

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	2.7	1.4 - 5.2	200	15.0	11.1 - 20.0	<50	3.3	1.8 - 6.1	900	78.9	73.6 - 83.4	1,100
Normal birth weight	500	2.8	1.8 - 4.3	2,100	12.9	10.7 - 15.5	500	3.2	2.2 - 4.8	13,100	81.1	78.1 - 83.7	16,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	3.6	1.4 - 9.1	200	27.3	15.3 - 44.0	100	13.1	4.9 - 30.5	400	56.0	39.6 - 71.1	700
No	500	2.8	1.8 - 4.4	2,000	12.6	10.5 - 15.2	500	2.9	2.0 - 4.3	13,100	81.6	78.7 - 84.2	16,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.6	2.0 - 10.2	500	17.6	12.2 - 24.8	200	7.7	4.1 - 14.0	1,900	70.1	61.8 - 77.2	2,700
No	400	2.5	1.5 - 4.1	1,700	12.4	10.1 - 15.1	400	2.5	1.6 - 3.9	11,600	82.6	79.6 - 85.3	14,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	4.9	3.0 - 7.7	1,400	18.3	14.7 - 22.5	200	3.2	1.9 - 5.3	5,600	73.7	69.0 - 77.9	7,600
No	100	1.1	0.4 - 2.7	800	8.9	6.6 - 11.8	300	3.0	1.8 - 5.0	8,300	87.1	83.7 - 89.8	9,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	300	1.7	1.1 - 2.8	2,200	12.7	10.7 - 15.1	700	4.0	2.9 - 5.4	14,200	81.5	78.8 - 83.9	17,500

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	<50	1.9	0.9 - 4.0	400	16.1	12.5 - 20.5	100	4.2	2.5 - 6.9	1,800	77.7	73.0 - 81.9	2,300
Honolulu	200	1.6	0.8 - 3.1	1,500	12.2	9.6 - 15.5	500	3.8	2.5 - 5.9	10,300	82.3	78.7 - 85.5	12,500
Kauai	<50	1.0	0.3 - 3.5	100	9.7	6.5 - 14.2	<50	4.2	2.3 - 7.6	700	85.0	79.9 - 89.0	800
Maui	100	2.8	1.5 - 4.9	200	13.3	10.4 - 17.0	100	4.8	3.1 - 7.2	1,500	79.1	75.0 - 82.7	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	1.3	0.4 - 4.2	500	11.9	7.9 - 17.4	100	3.3	1.7 - 6.4	3,600	83.5	77.7 - 88.1	4,400
Native Hawaiian	100	2.0	0.9 - 4.4	800	15.8	12.0 - 20.5	300	6.4	3.9 - 10.1	3,700	75.9	70.4 - 80.6	4,900
Chinese	<50	4.5	0.7 - 25.2	<50	2.3	0.8 - 6.3	<50	1.8	0.6 - 5.0	600	91.4	77.4 - 97.1	600
Filipino	<50	0.9	0.4 - 1.9	500	16.1	11.4 - 22.3	100	3.5	1.9 - 6.5	2,500	79.5	73.1 - 84.7	3,100
Japanese	<50	0.4	0.1 - 2.2	200	10.3	5.3 - 19.3	<50	1.7	0.4 - 7.8	1,600	87.6	78.5 - 93.2	1,800
Other	100	3.3	1.3 - 8.3	200	8.6	4.6 - 15.3	100	3.4	1.3 - 8.9	2,200	84.7	76.9 - 90.2	2,600

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	2.0	0.7 - 5.8	400	40.2	28.3 - 53.3	<50	3.4	1.3 - 8.3	500	54.4	41.7 - 66.6	1,000
20-24	100	2.3	1.0 - 5.3	700	16.8	12.2 - 22.8	100	1.3	0.7 - 2.3	3,200	79.5	73.4 - 84.6	4,000
25-34	200	1.9	1.0 - 3.7	1,000	10.6	8.1 - 13.8	500	5.1	3.4 - 7.5	7,700	82.4	78.6 - 85.7	9,400
35+	<50	0.4	0.1 - 1.3	200	5.3	3.0 - 9.0	100	4.5	2.3 - 8.6	2,800	89.9	85.0 - 93.3	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	200	2.6	1.4 - 4.5	1,100	12.4	9.6 - 15.9	400	4.0	2.6 - 6.3	7,200	81.0	77.0 - 84.4	8,900
Female	100	0.9	0.4 - 2.0	1,100	13.1	10.3 - 16.5	300	4.0	2.6 - 6.1	7,000	82.1	78.3 - 85.3	8,600

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.2	1.2 - 3.9	1,000	9.5	7.3 - 12.3	500	4.2	2.9 - 6.2	9,200	84.1	80.8 - 87.0	10,900
No	100	1.0	0.6 - 1.8	1,200	18.2	14.4 - 22.7	200	3.6	2.1 - 6.2	5,000	77.2	72.4 - 81.3	6,500

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	100	2.5	1.3 - 4.7	1,000	16.8	13.2 - 21.2	300	4.2	2.5 - 6.9	4,700	76.6	71.8 - 80.7	6,100
131-185%	<50	1.6	0.4 - 6.3	400	13.7	8.7 - 20.9	100	3.5	1.5 - 8.0	2,100	81.2	73.5 - 87.1	2,600
186+%	100	1.3	0.5 - 3.1	700	8.5	6.0 - 12.0	300	4.2	2.6 - 6.7	6,700	86.0	82.1 - 89.2	7,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	2.9	1.2 - 6.7	600	18.2	13.3 - 24.5	100	2.5	0.9 - 6.5	2,400	76.4	69.5 - 82.1	3,100
\$10,000-\$24,999	100	1.7	0.6 - 5.0	500	16.4	11.4 - 22.9	100	4.3	2.4 - 7.6	2,500	77.6	70.8 - 83.2	3,200
\$25,000-\$49,999	100	1.6	0.5 - 5.0	600	14.0	9.9 - 19.5	200	5.6	3.0 - 10.0	3,300	78.8	72.5 - 83.9	4,200
>=\$50,000	100	1.2	0.5 - 3.1	400	6.5	4.2 - 10.1	200	3.6	2.1 - 6.2	5,300	88.6	84.5 - 91.7	6,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	2.5	1.2 - 5.1	200	18.4	14.3 - 23.4	100	6.0	3.8 - 9.4	900	73.0	67.6 - 77.9	1,200
Normal birth weight	300	1.7	1.0 - 2.8	2,000	12.3	10.2 - 14.9	600	3.9	2.7 - 5.4	13,400	82.1	79.3 - 84.7	16,300

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	0.7	0.1 - 4.1	300	33.4	21.9 - 47.2	100	9.3	3.7 - 21.3	500	56.6	43.0 - 69.4	800
No	300	1.8	1.1 - 2.9	1,900	11.4	9.4 - 13.7	600	3.8	2.7 - 5.2	13,700	83.1	80.4 - 85.5	16,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	0.6	0.2 - 1.6	800	23.1	17.6 - 29.6	200	4.8	2.3 - 9.6	2,300	71.6	64.6 - 77.7	3,300
No	300	2.0	1.2 - 3.3	1,400	10.0	7.9 - 12.5	500	3.8	2.7 - 5.4	11,800	84.2	81.3 - 86.7	14,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	1.8	0.9 - 3.3	1,300	16.9	13.6 - 20.9	300	3.9	2.5 - 6.0	6,000	77.4	73.2 - 81.2	7,800
No	200	1.7	0.8 - 3.4	900	9.2	6.8 - 12.2	400	4.2	2.7 - 6.3	8,100	85.0	81.4 - 88.0	9,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.3	2.3 - 4.7	2,400	14.2	12.0 - 16.7	700	4.3	3.1 - 5.9	13,400	78.3	75.4 - 80.9	17,200

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.9	2.3 - 6.5	300	15.9	12.4 - 20.2	100	2.9	1.5 - 5.4	1,700	77.4	72.6 - 81.5	2,200
Honolulu	400	3.4	2.1 - 5.4	1,700	13.9	11.1 - 17.3	600	4.9	3.3 - 7.2	9,600	77.8	73.9 - 81.3	12,300
Kauai	0	0.0	-	100	11.1	7.8 - 15.5	<50	3.0	1.5 - 5.9	700	85.9	81.2 - 89.6	800
Maui	100	3.3	2.0 - 5.6	300	15.5	12.3 - 19.4	<50	2.1	1.1 - 4.1	1,500	79.0	74.8 - 82.8	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.5	1.6 - 7.6	400	9.5	6.0 - 14.8	300	7.9	4.8 - 12.8	3,300	79.1	72.6 - 84.4	4,200
Native Hawaiian	200	3.9	2.0 - 7.1	900	17.4	13.5 - 22.1	200	4.0	2.2 - 7.2	3,800	74.8	69.4 - 79.5	5,100
Chinese	<50	1.5	0.4 - 5.4	100	7.6	2.2 - 22.9	<50	4.4	0.8 - 20.7	600	86.5	70.8 - 94.4	700
Filipino	100	2.7	1.3 - 5.5	700	22.3	16.4 - 29.5	100	3.1	1.3 - 7.0	2,200	71.9	64.5 - 78.4	3,000
Japanese	<50	0.5	0.1 - 2.9	100	4.1	1.8 - 9.2	<50	1.8	0.3 - 9.0	1,600	93.7	87.3 - 96.9	1,700
Other	100	4.9	2.0 - 11.2	400	14.4	8.9 - 22.4	<50	1.7	0.5 - 5.8	1,900	79.0	70.3 - 85.6	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	1.8	0.6 - 5.1	300	28.5	18.3 - 41.4	100	9.4	3.6 - 22.1	600	60.4	47.4 - 72.0	1,100
20-24	100	2.7	1.2 - 6.4	800	19.9	15.0 - 26.0	100	3.4	1.5 - 7.3	3,000	74.0	67.4 - 79.6	4,100
25-34	300	3.2	1.9 - 5.4	1,100	11.9	9.2 - 15.3	400	4.2	2.7 - 6.6	7,200	80.7	76.7 - 84.1	8,900
35+	100	4.7	2.3 - 9.5	300	8.5	5.2 - 13.6	100	3.7	1.7 - 8.0	2,600	83.1	76.6 - 88.1	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.3	2.0 - 5.6	1,300	15.1	12.0 - 18.7	400	4.4	2.9 - 6.9	6,700	77.1	73.0 - 80.9	8,700
Female	300	3.2	1.9 - 5.4	1,100	13.3	10.4 - 16.9	300	4.1	2.5 - 6.6	6,700	79.4	75.3 - 83.0	8,500

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	1.8 - 4.8	1,200	11.2	8.7 - 14.3	500	4.1	2.7 - 6.3	9,000	81.8	78.2 - 84.9	10,900
No	200	3.9	2.2 - 6.8	1,200	19.5	15.6 - 24.0	300	4.5	2.7 - 7.4	4,500	72.1	67.1 - 76.6	6,200

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	200	3.8	2.1 - 6.6	1,100	18.6	14.8 - 23.2	200	4.2	2.3 - 7.3	4,300	73.4	68.3 - 78.0	5,900
131-185%	200	5.5	2.5 - 11.8	400	14.5	9.4 - 21.8	100	4.8	2.3 - 9.8	2,100	75.2	67.0 - 81.9	2,800
186+%	200	2.3	1.2 - 4.3	700	9.1	6.4 - 12.7	300	4.1	2.4 - 6.7	6,300	84.6	80.4 - 88.0	7,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	4.4	2.2 - 8.7	600	20.3	15.0 - 26.9	100	3.9	1.8 - 8.3	2,200	71.4	64.2 - 77.7	3,100
\$10,000-\$24,999	100	3.1	1.3 - 7.2	400	14.5	10.1 - 20.4	200	5.2	2.5 - 10.4	2,300	77.2	70.2 - 82.9	2,900
\$25,000-\$49,999	200	4.2	2.1 - 8.4	700	16.1	11.5 - 22.0	200	4.8	2.6 - 8.9	3,200	74.8	68.3 - 80.4	4,300
>=\$50,000	100	2.4	1.2 - 5.0	500	8.5	5.7 - 12.5	200	3.6	2.0 - 6.5	5,100	85.5	80.8 - 89.2	6,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	3.3	1.7 - 6.4	200	14.6	10.9 - 19.1	100	6.8	4.3 - 10.7	900	75.3	69.9 - 80.0	1,200
Normal birth weight	500	3.3	2.2 - 4.9	2,300	14.2	11.9 - 16.8	700	4.1	2.8 - 5.8	12,600	78.5	75.4 - 81.3	16,000

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	10.1	4.3 - 22.2	200	19.6	11.5 - 31.3	<50	2.2	0.8 - 6.2	700	68.1	55.1 - 78.8	1,100
No	500	2.9	1.9 - 4.3	2,200	13.9	11.7 - 16.5	700	4.4	3.2 - 6.2	12,500	78.8	75.8 - 81.5	15,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	7.8	4.5 - 13.2	700	21.8	16.3 - 28.6	200	6.1	3.4 - 10.6	2,200	64.3	56.9 - 71.0	3,400
No	300	2.2	1.3 - 3.6	1,700	12.4	10.1 - 15.1	500	3.9	2.6 - 5.7	11,100	81.6	78.5 - 84.3	13,600

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.2	2.6 - 6.8	1,300	17.5	14.1 - 21.5	400	4.9	3.1 - 7.7	5,400	73.4	68.8 - 77.6	7,400
No	300	2.6	1.5 - 4.6	1,100	11.8	9.1 - 15.2	400	3.8	2.4 - 6.0	7,900	81.7	77.9 - 85.1	9,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	800	4.6	3.4 - 6.3	2,600	15.6	13.4 - 18.2	900	5.3	4.0 - 7.0	12,500	74.4	71.4 - 77.2	16,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.2	4.0 - 9.3	400	16.6	12.9 - 21.1	100	4.1	2.4 - 6.8	1,600	73.1	68.1 - 77.7	2,200
Honolulu	600	4.7	3.1 - 7.1	1,900	15.8	12.8 - 19.3	600	5.4	3.7 - 7.8	8,700	74.1	70.0 - 77.9	11,800
Kauai	<50	2.9	1.3 - 6.1	100	13.5	9.8 - 18.4	<50	4.3	2.4 - 7.7	600	79.3	73.8 - 83.9	800
Maui	100	3.6	2.1 - 5.9	300	14.5	11.4 - 18.3	100	6.6	4.6 - 9.4	1,500	75.3	70.9 - 79.2	2,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	3.7	1.7 - 7.9	400	9.1	6.0 - 13.6	200	5.9	3.4 - 10.2	3,300	81.3	75.4 - 86.0	4,100
Native Hawaiian	300	5.9	3.6 - 9.6	1,100	21.9	17.4 - 27.1	200	4.9	2.9 - 7.9	3,400	67.4	61.6 - 72.6	5,100
Chinese	0	0.0	-	100	9.1	2.9 - 25.0	<50	1.4	0.4 - 5.1	600	89.5	74.4 - 96.1	700
Filipino	100	4.5	2.6 - 7.6	600	19.2	13.8 - 25.9	200	6.2	3.3 - 11.2	2,100	70.2	62.9 - 76.6	3,000
Japanese	100	4.0	1.3 - 11.6	100	7.6	3.4 - 16.4	100	3.9	1.4 - 10.6	1,300	84.4	74.6 - 90.9	1,600
Other	100	5.6	2.3 - 13.0	400	16.2	10.3 - 24.7	100	6.3	2.9 - 13.1	1,600	71.8	62.3 - 79.7	2,300

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.5	2.8 - 14.2	500	29.8	21.0 - 40.4	100	6.2	2.6 - 13.8	900	57.6	46.6 - 67.8	1,600
20-24	200	6.0	3.3 - 10.8	700	18.3	13.7 - 24.0	300	7.0	4.2 - 11.5	2,600	68.6	61.9 - 74.6	3,800
25-34	300	3.6	2.2 - 5.9	1,100	12.7	9.9 - 16.1	400	5.0	3.3 - 7.5	6,600	78.7	74.6 - 82.3	8,400
35+	100	4.8	2.3 - 9.7	400	12.5	7.9 - 19.4	100	3.6	1.7 - 7.5	2,300	79.1	71.6 - 85.0	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.9	2.5 - 6.2	1,300	14.8	11.8 - 18.3	600	6.2	4.3 - 8.9	6,600	75.0	70.8 - 78.8	8,800
Female	400	5.4	3.6 - 8.2	1,300	16.6	13.3 - 20.4	300	4.3	2.7 - 6.6	5,800	73.7	69.3 - 77.7	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.4	2.1 - 5.4	1,200	11.3	8.8 - 14.4	500	5.1	3.6 - 7.3	8,200	80.2	76.5 - 83.4	10,200
No	400	6.6	4.3 - 9.9	1,500	22.4	18.4 - 27.0	400	5.6	3.6 - 8.7	4,300	65.4	60.3 - 70.3	6,500

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	7.0	4.5 - 10.5	1,200	21.5	17.2 - 26.4	200	4.0	2.3 - 6.9	3,800	67.6	62.1 - 72.7	5,600
131-185%	100	5.6	2.6 - 11.7	400	16.8	11.5 - 23.9	200	6.5	3.5 - 11.7	1,900	71.2	63.1 - 78.1	2,600
186+%	200	3.2	1.7 - 5.7	700	9.6	6.9 - 13.1	400	5.9	3.9 - 9.0	5,700	81.3	76.9 - 85.0	7,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	5.9	3.0 - 11.4	700	25.1	18.5 - 33.2	100	5.3	2.5 - 10.9	1,700	63.7	55.2 - 71.4	2,700
\$10,000-\$24,999	300	8.1	4.9 - 13.2	600	17.0	12.5 - 22.6	200	4.6	2.4 - 8.5	2,400	70.3	63.6 - 76.2	3,500
\$25,000-\$49,999	200	6.0	3.0 - 11.7	400	13.1	9.0 - 18.7	200	5.9	3.3 - 10.3	2,600	74.9	68.0 - 80.8	3,400
>=\$50,000	100	2.0	0.9 - 4.0	600	10.6	7.5 - 14.8	300	5.5	3.4 - 8.8	4,800	81.9	77.1 - 85.9	5,900

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.3	3.1 - 8.9	200	18.8	14.8 - 23.6	100	5.1	3.1 - 8.1	800	70.8	65.3 - 75.7	1,100
Normal birth weight	700	4.6	3.3 - 6.4	2,400	15.4	13.0 - 18.1	800	5.3	4.0 - 7.2	11,600	74.7	71.5 - 77.6	15,600

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	12.0	6.1 - 22.3	400	26.5	17.8 - 37.5	100	6.2	2.2 - 16.0	700	55.4	43.8 - 66.4	1,300
No	600	4.1	2.9 - 5.8	2,200	14.4	12.1 - 17.0	700	5.0	3.7 - 6.7	11,500	76.5	73.4 - 79.4	15,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	7.5	4.3 - 12.9	800	23.3	17.8 - 30.0	200	6.6	3.8 - 11.4	2,100	62.5	55.2 - 69.2	3,400
No	500	4.0	2.7 - 5.8	1,700	13.4	11.0 - 16.1	600	4.7	3.3 - 6.6	10,100	77.9	74.6 - 80.9	13,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	8.0	5.5 - 11.6	1,400	22.0	18.1 - 26.4	300	5.0	3.2 - 7.7	4,300	65.0	60.0 - 69.8	6,500
No	200	2.5	1.4 - 4.5	1,100	11.5	8.9 - 14.8	500	5.2	3.5 - 7.6	7,800	80.7	76.9 - 84.1	9,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	2.9	2.1 - 3.8	2,600	15.3	13.6 - 17.2	800	4.6	3.6 - 5.7	12,900	77.3	75.1 - 79.3	16,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.1	3.4 - 10.7	300	15.4	10.8 - 21.5	<50	1.8	0.6 - 5.3	1,600	76.7	69.9 - 82.3	2,100
Honolulu	300	2.2	1.5 - 3.3	1,900	16.1	14.0 - 18.4	600	4.8	3.6 - 6.2	9,100	76.9	74.3 - 79.3	11,900
Kauai	<50	1.7	0.4 - 7.0	100	13.3	7.3 - 23.3	100	10.2	5.0 - 19.5	600	74.8	63.8 - 83.3	800
Maui	100	3.7	1.7 - 7.8	200	11.0	7.1 - 16.8	100	4.0	1.9 - 8.1	1,600	81.3	74.7 - 86.5	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.0	1.0 - 3.9	500	13.1	10.1 - 16.8	300	6.3	4.3 - 9.3	3,300	78.6	74.3 - 82.4	4,100
Native Hawaiian	200	4.0	2.4 - 6.7	800	17.8	14.2 - 22.2	300	6.4	4.3 - 9.4	3,300	71.8	66.8 - 76.2	4,500
Chinese	<50	1.2	0.4 - 3.8	100	9.2	5.9 - 14.0	<50	3.9	2.0 - 7.6	500	85.8	80.3 - 89.9	600
Filipino	100	3.9	2.2 - 6.7	500	18.0	14.1 - 22.6	100	3.2	1.7 - 5.8	2,300	75.0	69.9 - 79.5	3,000
Japanese	100	2.6	1.0 - 6.6	200	9.5	5.8 - 14.9	100	2.5	1.1 - 5.8	1,700	85.4	79.3 - 89.9	2,000
Other	<50	1.6	0.7 - 4.1	400	17.2	12.6 - 22.9	<50	1.8	0.6 - 4.9	2,000	79.4	73.4 - 84.3	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.4	2.3 - 11.9	300	20.4	13.7 - 29.1	<50	3.5	1.3 - 9.1	900	70.8	61.4 - 78.6	1,300
20-24	100	2.9	1.6 - 5.5	900	24.0	19.5 - 29.1	300	7.6	5.1 - 11.1	2,300	65.5	60.0 - 70.6	3,600
25-34	200	2.1	1.3 - 3.4	1,100	12.8	10.6 - 15.3	400	4.0	2.8 - 5.7	7,000	81.1	78.2 - 83.7	8,700
35+	100	3.8	2.2 - 6.5	300	10.4	7.4 - 14.5	100	3.1	1.7 - 5.7	2,700	82.7	78.1 - 86.6	3,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.1	2.0 - 4.5	1,400	15.9	13.5 - 18.7	500	5.1	3.8 - 6.9	6,700	75.9	72.8 - 78.7	8,800
Female	200	2.7	1.7 - 4.1	1,200	14.6	12.2 - 17.4	300	4.0	2.7 - 5.7	6,300	78.8	75.7 - 81.6	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.2	1.5 - 3.3	1,600	14.6	12.5 - 16.9	500	4.3	3.2 - 5.8	8,500	78.9	76.3 - 81.3	10,800
No	200	4.1	2.6 - 6.2	1,000	16.6	13.6 - 20.1	300	5.0	3.4 - 7.3	4,400	74.4	70.3 - 78.0	6,000

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	200	3.4	2.1 - 5.5	1,100	19.3	16.0 - 23.0	300	4.5	3.0 - 6.6	4,200	72.8	68.7 - 76.6	5,800
131-185%	<50	1.5	0.5 - 4.5	300	12.1	8.3 - 17.3	100	3.0	1.4 - 6.4	2,000	83.3	77.7 - 87.8	2,400
186+%	200	2.3	1.4 - 3.6	900	12.3	10.0 - 14.9	400	5.2	3.8 - 7.2	6,000	80.2	77.1 - 83.0	7,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	4.4	2.3 - 8.2	500	21.0	16.1 - 27.0	100	3.7	1.9 - 7.0	1,900	70.9	64.6 - 76.6	2,600
\$10,000-\$24,999	100	3.2	1.7 - 5.9	600	17.6	13.7 - 22.4	200	5.7	3.6 - 9.0	2,600	73.4	68.1 - 78.1	3,500
\$25,000-\$49,999	100	2.5	1.4 - 4.7	600	16.6	13.1 - 20.9	200	4.1	2.5 - 6.8	3,000	76.7	72.0 - 80.8	3,800
>=\$50,000	100	1.6	0.9 - 3.1	600	9.8	7.6 - 12.6	300	4.6	3.1 - 6.7	4,900	84.0	80.7 - 86.8	5,800

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	4.3	1.7 - 10.6	200	14.9	9.1 - 23.4	<50	3.6	1.3 - 9.4	900	77.2	68.0 - 84.4	1,100
Normal birth weight	400	2.8	2.0 - 3.8	2,400	15.3	13.5 - 17.3	700	4.6	3.7 - 5.9	12,100	77.3	75.0 - 79.4	15,600

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	1.8	0.5 - 6.6	300	27.6	19.9 - 36.9	100	10.3	5.7 - 17.8	800	60.3	50.7 - 69.2	1,300
No	400	2.9	2.1 - 3.9	2,200	14.3	12.5 - 16.2	600	4.2	3.3 - 5.4	11,900	78.7	76.4 - 80.7	15,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	2.2	0.9 - 5.0	600	22.1	17.4 - 27.7	300	9.7	6.6 - 14.1	1,900	66.0	59.9 - 71.6	2,900
No	400	2.9	2.1 - 4.0	1,900	13.8	12.0 - 15.9	500	3.6	2.7 - 4.8	10,800	79.7	77.4 - 81.8	13,500

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	3.7	2.4 - 5.6	1,400	20.0	17.0 - 23.5	300	4.4	3.0 - 6.4	4,800	71.9	68.1 - 75.5	6,700
No	200	2.4	1.6 - 3.6	1,200	12.0	10.0 - 14.3	500	4.8	3.6 - 6.4	7,900	80.8	78.1 - 83.2	9,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.8	2.9 - 4.9	2,700	16.2	14.4 - 18.1	500	3.3	2.5 - 4.3	12,700	76.7	74.5 - 78.8	16,600

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.3	2.1 - 8.8	200	9.9	6.3 - 15.2	100	2.4	0.9 - 6.0	1,800	83.4	77.1 - 88.2	2,100
Honolulu	400	3.6	2.6 - 5.0	2,000	17.2	15.0 - 19.6	400	3.6	2.6 - 4.8	8,800	75.6	73.0 - 78.1	11,600
Kauai	<50	1.4	0.2 - 8.5	100	14.6	8.4 - 24.1	<50	4.7	1.8 - 11.4	700	79.3	69.3 - 86.7	900
Maui	100	5.4	2.8 - 10.2	300	17.6	12.4 - 24.3	<50	2.1	0.7 - 6.2	1,500	75.0	67.7 - 81.0	2,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.5	2.0 - 6.1	400	11.5	8.6 - 15.4	100	3.8	2.2 - 6.4	3,000	81.2	76.7 - 85.0	3,700
Native Hawaiian	300	6.3	4.2 - 9.5	1,100	24.5	20.3 - 29.3	100	2.4	1.2 - 4.5	3,000	66.8	61.7 - 71.5	4,600
Chinese	<50	3.7	1.7 - 7.7	<50	6.9	4.0 - 11.5	<50	1.7	0.6 - 4.4	600	87.8	82.3 - 91.7	700
Filipino	100	1.9	0.9 - 4.0	600	19.5	15.5 - 24.2	200	5.8	3.7 - 9.0	2,200	72.7	67.6 - 77.3	3,000
Japanese	<50	1.3	0.3 - 5.2	200	7.7	4.6 - 12.7	<50	2.2	0.8 - 5.5	1,800	88.7	83.2 - 92.6	2,000
Other	100	4.0	2.1 - 7.5	300	13.0	9.2 - 18.2	100	2.7	1.3 - 5.7	2,100	80.3	74.5 - 85.0	2,600

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.8	5.3 - 17.6	300	23.7	16.5 - 33.0	100	4.9	2.1 - 11.1	900	61.5	51.7 - 70.4	1,500
20-24	100	3.1	1.7 - 5.5	1,000	23.8	19.7 - 28.6	100	3.2	1.8 - 5.4	2,800	69.9	64.9 - 74.4	4,100
25-34	300	3.5	2.4 - 5.1	1,100	13.9	11.6 - 16.5	200	2.9	1.9 - 4.3	6,500	79.7	76.8 - 82.4	8,100
35+	100	2.6	1.3 - 5.1	200	8.0	5.4 - 11.7	100	3.9	2.2 - 6.8	2,500	85.5	81.0 - 89.1	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	200	2.6	1.7 - 4.1	1,500	17.5	14.9 - 20.3	300	3.4	2.3 - 4.9	6,600	76.5	73.4 - 79.4	8,600
Female	400	5.1	3.7 - 7.0	1,200	14.7	12.3 - 17.5	300	3.2	2.2 - 4.7	6,100	76.9	73.7 - 79.9	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	2.0 - 4.2	1,200	11.8	9.9 - 13.9	300	2.5	1.7 - 3.6	8,700	82.8	80.4 - 85.0	10,500
No	300	5.3	3.6 - 7.8	1,400	23.8	20.3 - 27.8	300	4.8	3.3 - 6.9	4,000	66.1	61.8 - 70.1	6,000

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.4	3.7 - 7.8	1,400	24.0	20.4 - 27.9	200	3.1	1.9 - 5.1	4,000	67.5	63.3 - 71.5	6,000
131-185%	100	3.2	1.6 - 6.6	400	14.4	10.4 - 19.6	100	3.8	2.0 - 7.3	1,900	78.6	72.7 - 83.4	2,400
186+%	200	3.3	2.2 - 5.0	700	9.9	7.8 - 12.4	200	3.0	1.9 - 4.5	5,800	83.9	80.9 - 86.4	7,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	7.1	4.2 - 11.5	700	25.8	20.4 - 32.0	100	4.2	2.2 - 7.8	1,700	63.0	56.4 - 69.2	2,700
\$10,000-\$24,999	100	4.4	2.5 - 7.5	700	20.0	15.8 - 24.9	100	3.2	1.8 - 5.9	2,400	72.4	67.1 - 77.2	3,300
\$25,000-\$49,999	100	2.6	1.4 - 5.0	700	17.8	14.1 - 22.1	100	2.7	1.5 - 5.0	2,900	76.9	72.2 - 81.0	3,800
>=\$50,000	200	3.4	2.2 - 5.3	500	8.1	6.1 - 10.7	200	2.9	1.8 - 4.6	4,800	85.6	82.4 - 88.3	5,600

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	3.9	1.6 - 9.4	300	21.8	15.2 - 30.1	<50	3.1	1.2 - 7.6	1,000	71.3	62.6 - 78.7	1,300
Normal birth weight	600	3.8	2.9 - 5.0	2,400	15.7	13.8 - 17.7	500	3.3	2.5 - 4.4	11,800	77.2	74.9 - 79.3	15,200

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.3	4.2 - 15.5	400	32.4	24.0 - 42.1	100	6.9	3.4 - 13.4	700	52.5	42.9 - 61.9	1,300
No	500	3.5	2.6 - 4.6	2,300	14.9	13.1 - 16.9	500	3.1	2.3 - 4.1	11,900	78.5	76.3 - 80.6	15,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.1	3.0 - 8.6	1,000	31.0	25.7 - 36.9	100	4.6	2.7 - 7.8	1,800	59.3	53.3 - 65.0	3,100
No	500	3.5	2.6 - 4.8	1,700	12.9	11.1 - 14.9	400	3.1	2.3 - 4.2	10,700	80.5	78.2 - 82.6	13,300

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.2	2.8 - 6.4	1,400	21.7	18.5 - 25.3	200	3.3	2.2 - 5.1	4,600	70.7	66.8 - 74.3	6,600
No	300	3.5	2.5 - 4.9	1,200	12.1	10.1 - 14.4	300	3.4	2.4 - 4.8	7,800	81.0	78.4 - 83.4	9,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	2.9	2.1 - 3.9	2,800	17.4	15.5 - 19.5	600	3.5	2.7 - 4.6	12,400	76.2	73.9 - 78.4	16,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.6	1.5 - 8.0	400	19.9	14.4 - 26.7	<50	2.1	0.7 - 6.1	1,500	74.5	67.2 - 80.6	2,000
Honolulu	400	3.2	2.3 - 4.4	2,000	16.9	14.7 - 19.4	400	3.6	2.6 - 4.8	9,100	76.4	73.7 - 78.8	11,900
Kauai	0	0.0	-	100	16.0	8.8 - 27.5	<50	3.4	0.9 - 12.1	600	80.5	68.7 - 88.6	800
Maui	<50	1.2	0.3 - 5.5	300	18.5	12.9 - 25.8	100	4.8	2.2 - 10.1	1,300	75.5	67.6 - 81.9	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.5	1.3 - 4.8	500	13.9	10.6 - 18.1	200	5.2	3.3 - 8.0	3,100	78.4	73.7 - 82.5	3,900
Native Hawaiian	200	3.5	2.0 - 6.2	1,100	24.4	20.0 - 29.4	100	2.5	1.3 - 4.8	3,000	69.6	64.4 - 74.4	4,400
Chinese	<50	2.1	0.8 - 5.6	100	11.3	7.5 - 16.8	<50	2.8	1.2 - 6.5	500	83.7	77.7 - 88.3	600
Filipino	100	2.8	1.4 - 5.6	600	19.5	15.3 - 24.4	100	4.3	2.5 - 7.4	2,200	73.5	68.1 - 78.2	3,000
Japanese	<50	1.9	0.6 - 5.7	200	8.0	4.8 - 13.1	<50	0.6	0.1 - 3.5	1,700	89.5	84.0 - 93.3	1,900
Other	100	3.3	1.6 - 6.9	400	16.8	12.2 - 22.7	100	4.2	2.2 - 7.6	1,900	75.7	69.3 - 81.1	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	4.1	1.6 - 10.2	400	32.3	23.9 - 42.1	<50	1.5	0.3 - 6.7	800	62.1	52.3 - 71.0	1,300
20-24	100	3.3	1.8 - 5.9	1,000	26.1	21.5 - 31.2	200	4.8	2.9 - 7.7	2,600	65.9	60.5 - 70.9	4,000
25-34	200	3.0	2.0 - 4.6	1,100	13.7	11.4 - 16.4	300	3.5	2.4 - 5.0	6,500	79.8	76.7 - 82.6	8,100
35+	<50	1.4	0.6 - 3.3	300	8.9	6.1 - 12.8	100	2.8	1.4 - 5.4	2,500	87.0	82.6 - 90.4	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.2	2.1 - 4.7	1,600	19.4	16.6 - 22.5	400	5.0	3.6 - 6.8	6,100	72.4	69.0 - 75.6	8,400
Female	200	2.6	1.6 - 4.1	1,200	15.2	12.7 - 18.1	200	2.0	1.2 - 3.3	6,400	80.2	77.1 - 83.0	8,000

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.5	1.7 - 3.7	1,300	12.1	10.2 - 14.3	300	3.3	2.4 - 4.6	8,600	82.1	79.6 - 84.4	10,500
No	200	3.5	2.2 - 5.7	1,600	26.8	22.9 - 31.1	200	3.9	2.5 - 6.1	3,900	65.7	61.2 - 70.0	5,900

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.9	3.3 - 7.4	1,400	25.7	21.8 - 30.1	100	2.5	1.4 - 4.4	3,700	66.8	62.3 - 71.1	5,500
131-185%	100	2.5	1.0 - 6.3	300	13.9	9.5 - 20.0	100	5.8	3.2 - 10.5	1,600	77.7	70.9 - 83.3	2,100
186+%	100	1.6	0.9 - 2.9	900	12.0	9.8 - 14.7	300	3.9	2.7 - 5.5	6,300	82.5	79.5 - 85.1	7,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	4.8	2.6 - 8.7	800	30.3	24.3 - 36.9	100	2.5	1.1 - 5.6	1,700	62.5	55.7 - 68.9	2,700
\$10,000-\$24,999	200	5.3	3.2 - 8.7	700	20.7	16.3 - 25.9	100	2.3	1.1 - 4.9	2,400	71.7	66.1 - 76.7	3,400
\$25,000-\$49,999	100	2.1	1.1 - 4.3	700	19.0	15.1 - 23.6	200	5.4	3.4 - 8.3	2,900	73.5	68.5 - 78.0	3,900
>=\$50,000	100	1.1	0.5 - 2.5	400	7.5	5.6 - 10.1	200	3.8	2.5 - 5.8	4,700	87.5	84.4 - 90.1	5,400

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	4.9	2.1 - 10.9	200	15.1	9.4 - 23.5	<50	3.0	1.1 - 7.4	1,000	77.0	68.1 - 84.0	1,300
Normal birth weight	400	2.7	1.9 - 3.8	2,600	17.6	15.6 - 19.8	500	3.6	2.7 - 4.7	11,500	76.1	73.7 - 78.4	15,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.6	3.7 - 14.7	300	23.2	15.7 - 33.0	100	5.0	2.2 - 11.1	800	64.2	54.1 - 73.2	1,200
No	400	2.5	1.7 - 3.5	2,500	16.7	14.7 - 18.9	500	3.4	2.6 - 4.6	11,500	77.4	75.0 - 79.6	14,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.6	2.6 - 8.0	700	22.3	17.5 - 27.9	200	5.3	3.2 - 8.8	2,200	67.8	61.8 - 73.3	3,200
No	300	2.4	1.7 - 3.5	2,100	15.9	13.9 - 18.2	400	3.1	2.3 - 4.3	10,100	78.5	76.0 - 80.8	12,900

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.9	3.4 - 7.1	1,600	23.2	19.8 - 27.0	200	3.6	2.3 - 5.5	4,600	68.3	64.2 - 72.1	6,700
No	100	1.1	0.6 - 1.9	1,300	13.4	11.3 - 15.9	300	3.3	2.3 - 4.7	7,700	82.2	79.5 - 84.6	9,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.2	3.2 - 5.3	2,500	16.0	14.3 - 18.0	500	3.4	2.6 - 4.4	11,700	76.4	74.2 - 78.5	15,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	<50	2.6	1.0 - 6.5	300	15.5	10.8 - 21.7	<50	1.9	0.7 - 5.6	1,500	80.0	73.3 - 85.3	1,900
Honolulu	500	4.4	3.3 - 5.8	1,700	15.2	13.2 - 17.4	400	4.0	3.0 - 5.3	8,600	76.5	73.9 - 78.9	11,200
Kauai	0	0.0	-	100	13.9	7.5 - 24.4	<50	1.9	0.3 - 11.4	600	84.2	73.3 - 91.2	700
Maui	100	6.6	3.5 - 12.0	400	23.7	17.6 - 31.2	<50	1.6	0.5 - 5.0	1,000	68.1	60.3 - 75.0	1,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.4	1.9 - 6.1	300	9.6	6.8 - 13.2	100	4.1	2.5 - 6.8	2,900	82.9	78.5 - 86.6	3,500
Native Hawaiian	300	6.6	4.4 - 9.8	800	19.5	15.6 - 24.1	200	5.1	3.2 - 7.9	2,900	68.8	63.7 - 73.5	4,100
Chinese	<50	1.3	0.4 - 4.2	100	10.4	6.9 - 15.3	<50	3.2	1.5 - 6.7	500	85.1	79.6 - 89.3	600
Filipino	100	4.9	3.1 - 7.8	700	23.2	19.0 - 28.0	100	3.7	2.1 - 6.3	2,000	68.2	63.0 - 73.0	2,900
Japanese	<50	0.5	0.1 - 3.0	200	10.8	7.4 - 15.6	<50	0.9	0.3 - 3.5	1,900	87.8	82.8 - 91.5	2,200
Other	100	4.2	2.1 - 8.1	400	17.2	12.7 - 23.0	<50	1.0	0.3 - 3.6	1,600	77.6	71.4 - 82.7	2,100

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.6	2.6 - 11.6	400	27.5	20.1 - 36.4	<50	3.4	1.3 - 8.4	800	63.6	54.4 - 71.8	1,300
20-24	200	6.2	4.1 - 9.3	900	25.0	20.8 - 29.8	100	3.9	2.3 - 6.5	2,400	64.9	59.8 - 69.7	3,700
25-34	300	3.9	2.7 - 5.6	900	12.0	9.9 - 14.5	300	3.7	2.6 - 5.3	6,100	80.4	77.5 - 83.1	7,600
35+	<50	1.5	0.6 - 3.9	300	9.6	6.7 - 13.4	100	1.9	0.9 - 3.9	2,400	87.1	82.8 - 90.4	2,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.4	3.2 - 6.2	1,200	15.3	12.9 - 18.0	300	4.3	3.1 - 6.0	5,900	76.0	72.9 - 78.9	7,800
Female	300	3.9	2.7 - 5.6	1,300	16.8	14.3 - 19.7	200	2.4	1.6 - 3.8	5,800	76.8	73.6 - 79.7	7,500

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.5	2.5 - 4.8	1,200	12.0	10.1 - 14.1	300	3.3	2.3 - 4.5	8,100	81.3	78.8 - 83.6	10,000
No	300	5.5	3.7 - 7.9	1,300	23.6	20.1 - 27.5	200	3.6	2.3 - 5.6	3,600	67.3	63.1 - 71.3	5,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.0	4.3 - 8.5	1,200	22.3	18.8 - 26.1	200	3.8	2.4 - 5.8	3,600	68.0	63.8 - 71.9	5,300
131-185%	100	4.6	2.2 - 9.7	300	17.5	12.4 - 24.1	<50	2.2	0.7 - 6.3	1,200	75.7	68.3 - 81.8	1,600
186+%	200	2.3	1.4 - 3.8	700	10.3	8.3 - 12.7	300	3.5	2.4 - 5.1	6,000	83.9	81.0 - 86.4	7,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	8.0	5.2 - 12.1	700	25.9	20.7 - 31.8	100	4.6	2.6 - 8.0	1,600	61.6	55.3 - 67.5	2,600
\$10,000-\$24,999	200	6.0	3.8 - 9.3	600	20.6	16.4 - 25.6	100	3.2	1.7 - 5.9	2,100	70.2	64.8 - 75.1	3,000
\$25,000-\$49,999	100	3.2	1.7 - 5.8	500	14.2	10.9 - 18.2	100	2.4	1.2 - 4.6	2,900	80.2	75.8 - 84.1	3,600
>=\$50,000	100	1.6	0.8 - 3.3	400	8.6	6.5 - 11.3	200	4.0	2.6 - 6.1	4,300	85.8	82.5 - 88.6	5,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.5	2.3 - 12.2	200	16.8	10.9 - 25.0	100	4.5	1.9 - 10.4	800	73.2	64.2 - 80.7	1,100
Normal birth weight	600	4.1	3.1 - 5.3	2,300	16.0	14.2 - 18.0	500	3.3	2.5 - 4.4	10,900	76.6	74.4 - 78.8	14,200

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	13.1	7.8 - 21.3	300	26.9	19.2 - 36.4	100	5.6	2.4 - 12.4	600	54.3	44.6 - 63.8	1,100
No	500	3.3	2.5 - 4.4	2,100	15.1	13.4 - 17.1	500	3.3	2.5 - 4.3	11,000	78.3	76.0 - 80.4	14,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	7.4	4.8 - 11.1	700	24.5	19.8 - 29.9	200	6.9	4.5 - 10.5	1,800	61.2	55.4 - 66.7	3,000
No	400	3.2	2.3 - 4.4	1,700	13.8	12.0 - 15.8	300	2.6	1.8 - 3.6	9,700	80.4	78.1 - 82.6	12,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.6	4.0 - 7.7	1,300	20.0	17.0 - 23.4	200	3.4	2.2 - 5.1	4,500	71.1	67.4 - 74.6	6,300
No	200	2.8	1.9 - 4.2	1,200	13.2	11.1 - 15.6	300	3.2	2.3 - 4.6	7,000	80.7	78.0 - 83.2	8,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	700	4.2	3.4 - 5.3	2,500	15.6	14.0 - 17.4	700	4.2	3.4 - 5.2	12,000	76.0	74.0 - 77.9	15,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.6	2.4 - 8.5	300	13.8	9.7 - 19.1	<50	1.6	0.6 - 4.2	1,500	80.1	74.1 - 85.0	1,900
Honolulu	400	3.8	2.9 - 5.0	1,800	15.5	13.6 - 17.6	500	4.6	3.6 - 5.8	8,700	76.2	73.8 - 78.4	11,400
Kauai	<50	5.3	2.1 - 12.8	100	14.5	8.5 - 23.5	<50	1.3	0.2 - 8.1	600	78.9	69.0 - 86.3	700
Maui	100	6.1	3.6 - 10.2	300	19.0	14.0 - 25.1	100	5.7	3.3 - 9.8	1,200	69.2	62.5 - 75.2	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	<50	1.3	0.6 - 2.9	500	13.5	10.5 - 17.0	100	3.9	2.4 - 6.2	3,000	81.4	77.4 - 84.8	3,600
Native Hawaiian	200	5.6	3.7 - 8.4	900	22.4	18.5 - 26.8	200	4.7	3.0 - 7.1	2,800	67.4	62.6 - 71.8	4,100
Chinese	<50	2.9	1.6 - 5.0	<50	6.7	4.7 - 9.5	<50	2.8	1.6 - 4.9	500	87.6	84.1 - 90.4	600
Filipino	200	7.9	5.7 - 10.9	500	17.6	14.3 - 21.6	200	5.3	3.5 - 7.8	2,100	69.2	64.7 - 73.4	3,000
Japanese	100	3.0	1.4 - 6.4	100	5.9	3.4 - 9.8	100	3.9	2.0 - 7.3	1,800	87.3	82.2 - 91.0	2,100
Other	100	3.1	1.4 - 6.4	300	15.6	11.4 - 20.9	100	3.0	1.5 - 5.7	1,800	78.4	72.6 - 83.2	2,200

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.1	4.4 - 14.6	300	27.9	20.5 - 36.8	<50	4.0	1.6 - 9.7	600	60.0	50.8 - 68.5	1,100
20-24	200	5.2	3.4 - 7.8	1,000	25.0	21.0 - 29.4	200	3.9	2.5 - 6.1	2,600	65.9	61.2 - 70.3	3,900
25-34	300	3.6	2.6 - 5.0	900	11.0	9.1 - 13.2	400	4.7	3.5 - 6.2	6,300	80.8	78.1 - 83.2	7,700
35+	100	3.2	1.8 - 5.6	300	10.8	7.9 - 14.5	100	3.3	1.9 - 5.7	2,500	82.8	78.4 - 86.4	3,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	4.3	3.2 - 5.8	1,300	15.8	13.6 - 18.2	400	4.6	3.5 - 6.1	6,400	75.3	72.5 - 77.9	8,500
Female	300	4.1	2.9 - 5.8	1,100	15.4	13.0 - 18.1	300	3.7	2.6 - 5.2	5,500	76.8	73.8 - 79.6	7,200

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.0	2.2 - 4.0	1,500	13.0	11.3 - 15.0	500	4.4	3.4 - 5.6	8,900	79.6	77.4 - 81.7	11,200
No	300	7.3	5.3 - 10.0	1,000	21.9	18.5 - 25.8	200	3.7	2.4 - 5.7	3,100	67.1	62.8 - 71.1	4,600

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.0	4.4 - 8.2	1,300	22.8	19.6 - 26.4	100	1.9	1.1 - 3.3	3,900	69.2	65.4 - 72.8	5,600
131-185%	100	3.1	1.5 - 6.4	200	13.4	9.3 - 18.9	100	4.9	2.7 - 8.8	1,500	78.6	72.5 - 83.7	1,900
186+%	200	2.3	1.4 - 3.5	800	10.8	8.9 - 13.1	400	6.1	4.7 - 7.8	5,800	80.9	78.1 - 83.3	7,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	8.8	5.8 - 13.1	500	23.1	18.3 - 28.7	100	3.2	1.6 - 6.1	1,500	64.9	58.8 - 70.6	2,400
\$10,000-\$24,999	200	5.7	3.9 - 8.3	800	21.2	17.5 - 25.5	100	2.2	1.2 - 4.2	2,600	70.8	66.2 - 75.0	3,700
\$25,000-\$49,999	100	2.1	1.1 - 3.9	600	13.9	11.0 - 17.3	300	6.8	4.9 - 9.3	3,200	77.2	73.3 - 80.8	4,100
>=\$50,000	100	1.7	0.9 - 3.1	400	8.5	6.5 - 11.1	200	4.3	2.9 - 6.4	3,900	85.5	82.3 - 88.2	4,500

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.1	3.6 - 13.7	200	15.6	10.0 - 23.5	<50	3.4	1.3 - 8.5	700	73.9	65.0 - 81.1	1,000
Normal birth weight	600	4.0	3.2 - 5.1	2,300	15.6	13.9 - 17.4	600	4.2	3.4 - 5.3	11,200	76.1	74.0 - 78.1	14,800

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.5	3.8 - 14.5	200	20.0	13.6 - 28.4	<50	3.9	1.5 - 9.5	700	68.6	59.4 - 76.5	1,100
No	600	3.9	3.1 - 5.0	2,200	15.1	13.5 - 17.0	600	4.2	3.3 - 5.2	11,100	76.8	74.7 - 78.7	14,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	3.8	2.2 - 6.5	600	19.3	15.3 - 24.0	200	5.8	3.8 - 8.9	2,200	71.1	65.9 - 75.8	3,000
No	500	4.3	3.3 - 5.4	1,800	14.5	12.8 - 16.5	500	3.7	2.9 - 4.8	9,700	77.5	75.2 - 79.5	12,500

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.7	3.4 - 6.5	1,400	20.3	17.6 - 23.4	200	3.0	2.0 - 4.4	4,900	71.9	68.6 - 75.0	6,900
No	300	3.8	2.8 - 5.2	1,000	11.9	10.0 - 14.1	400	5.2	4.0 - 6.7	6,800	79.1	76.5 - 81.4	8,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	700	4.7	3.6 - 6.0	2,400	16.3	14.3 - 18.6	600	4.4	3.4 - 5.8	11,000	74.6	72.0 - 77.1	14,800

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.6	2.8 - 7.6	300	18.2	14.7 - 22.4	<50	3.2	1.8 - 5.4	1,100	74.0	69.3 - 78.2	1,500
Honolulu	500	4.7	3.4 - 6.6	1,800	16.5	13.9 - 19.5	500	4.2	3.0 - 6.1	8,100	74.5	71.1 - 77.7	10,900
Kauai	<50	5.4	2.7 - 10.4	100	7.2	4.3 - 11.7	100	7.2	4.1 - 12.5	600	80.2	73.4 - 85.6	800
Maui	100	3.9	2.3 - 6.5	300	17.8	14.1 - 22.2	100	5.3	3.4 - 8.2	1,200	73.0	68.1 - 77.4	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.2	0.9 - 5.5	400	12.7	8.6 - 18.3	100	4.3	2.2 - 8.1	2,800	80.9	74.7 - 85.8	3,400
Native Hawaiian	200	6.5	5.0 - 8.5	800	20.6	18.0 - 23.4	200	4.8	3.6 - 6.4	2,600	68.1	64.9 - 71.2	3,800
Chinese	0	0.0	-	<50	6.4	1.7 - 21.6	<50	1.9	0.5 - 6.9	600	91.7	78.0 - 97.2	700
Filipino	200	5.5	3.2 - 9.4	700	21.1	15.9 - 27.3	200	5.9	3.4 - 10.1	2,100	67.5	60.7 - 73.7	3,100
Japanese	100	6.0	2.6 - 13.3	100	6.8	3.3 - 13.7	<50	2.5	0.8 - 8.0	1,500	84.7	76.1 - 90.6	1,800
Other	100	4.4	1.8 - 10.2	400	19.0	12.6 - 27.7	100	4.0	1.6 - 9.9	1,400	72.6	63.3 - 80.2	1,900

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	13.8	7.9 - 22.7	400	34.3	26.0 - 43.6	100	4.1	2.3 - 7.2	600	47.9	39.0 - 57.0	1,300
20-24	100	3.9	2.7 - 5.8	800	25.7	20.7 - 31.4	100	4.0	2.2 - 7.4	2,200	66.4	60.5 - 71.8	3,300
25-34	300	4.6	3.1 - 6.6	900	11.5	9.1 - 14.5	400	4.9	3.4 - 7.0	5,900	79.0	75.4 - 82.2	7,500
35+	<50	1.4	0.4 - 4.6	300	9.7	6.1 - 15.0	100	3.7	1.7 - 7.8	2,300	85.2	79.2 - 89.7	2,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.2	2.9 - 6.1	1,300	17.0	14.2 - 20.3	400	4.6	3.1 - 6.7	5,800	74.2	70.4 - 77.6	7,800
Female	400	5.2	3.6 - 7.4	1,100	15.6	12.7 - 18.9	300	4.2	2.9 - 6.0	5,200	75.1	71.3 - 78.5	6,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.4	1.6 - 3.7	1,300	13.0	10.6 - 15.7	500	4.6	3.3 - 6.5	8,100	80.0	76.8 - 82.8	10,200
No	400	9.6	7.0 - 13.0	1,100	23.8	19.9 - 28.2	200	3.9	2.7 - 5.7	2,900	62.7	58.0 - 67.2	4,600

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.1	4.3 - 8.5	1,500	23.6	20.0 - 27.6	300	4.2	2.9 - 6.2	4,100	66.1	61.8 - 70.2	6,200
131-185%	100	4.3	2.0 - 8.9	300	14.2	9.8 - 20.1	100	4.4	2.0 - 9.3	1,700	77.1	70.1 - 82.9	2,200
186+%	100	2.2	1.0 - 4.6	400	7.8	5.4 - 11.2	200	4.0	2.4 - 6.6	4,400	86.1	81.9 - 89.4	5,100

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	300	8.6	5.7 - 12.6	700	23.9	19.1 - 29.4	100	4.6	2.6 - 8.1	1,900	62.9	56.9 - 68.6	3,100
\$15,001-\$25,000	100	2.9	1.1 - 7.1	400	22.2	16.1 - 30.0	<50	2.2	1.1 - 4.3	1,300	72.7	64.9 - 79.4	1,900
\$25,001-\$40,000	200	6.0	3.5 - 10.2	500	17.5	12.9 - 23.4	200	5.3	3.0 - 9.4	2,100	71.1	64.5 - 76.9	2,900
>=\$40,001	100	1.7	0.8 - 3.5	600	9.3	6.9 - 12.5	300	4.6	3.0 - 7.0	5,100	84.4	80.6 - 87.6	6,100

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	4.4	1.5 - 12.1	200	21.9	13.5 - 33.5	100	5.6	2.0 - 14.8	700	68.1	56.3 - 78.1	1,000
Normal birth weight	600	4.7	3.6 - 6.1	2,200	15.9	13.8 - 18.3	600	4.3	3.3 - 5.7	10,300	75.1	72.4 - 77.6	13,700

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.4	4.4 - 15.7	200	22.1	14.6 - 32.1	<50	3.1	1.4 - 7.0	600	66.3	55.9 - 75.3	1,000
No	600	4.4	3.3 - 5.8	2,200	16.0	13.9 - 18.4	600	4.4	3.3 - 5.9	10,200	75.1	72.4 - 77.7	13,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	9.2	6.0 - 13.9	600	22.1	17.1 - 28.1	200	6.6	4.0 - 10.7	1,600	62.1	55.5 - 68.2	2,600
No	400	3.7	2.6 - 5.1	1,700	14.8	12.6 - 17.3	500	3.9	2.8 - 5.5	9,100	77.6	74.7 - 80.2	11,800

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	6.5	4.6 - 9.1	1,400	24.8	21.1 - 28.9	200	3.9	2.6 - 5.8	3,800	64.8	60.4 - 68.9	5,800
No	300	3.5	2.4 - 5.2	900	10.8	8.5 - 13.6	400	4.6	3.2 - 6.7	7,100	81.1	77.7 - 84.0	8,800

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.3	3.2 - 5.7	2,200	15.4	13.4 - 17.7	400	2.6	1.8 - 3.6	11,200	77.7	75.1 - 80.1	14,400

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.7	3.0 - 7.3	300	19.5	16.0 - 23.5	100	3.6	2.1 - 6.0	1,200	72.3	67.7 - 76.4	1,700
Honolulu	400	4.2	2.9 - 6.1	1,500	14.0	11.4 - 17.0	300	2.6	1.6 - 4.0	8,400	79.3	75.9 - 82.3	10,500
Kauai	<50	1.2	0.2 - 7.2	100	21.6	15.4 - 29.4	<50	2.9	1.0 - 8.0	400	74.3	66.1 - 81.1	600
Maui	100	5.5	3.5 - 8.5	300	18.8	14.9 - 23.3	<50	1.5	0.7 - 3.4	1,200	74.2	69.3 - 78.6	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.0	1.4 - 6.3	300	10.5	6.7 - 16.1	200	5.9	3.3 - 10.1	2,500	80.6	74.3 - 85.6	3,100
Native Hawaiian	200	4.9	3.6 - 6.7	700	19.4	17.0 - 22.2	100	3.4	2.3 - 4.9	2,700	72.2	69.1 - 75.1	3,700
Chinese	<50	6.5	1.9 - 20.3	100	11.4	3.9 - 29.4	<50	3.8	0.6 - 21.8	400	78.2	60.1 - 89.5	500
Filipino	200	5.8	3.1 - 10.7	600	20.6	15.5 - 26.9	<50	1.3	0.4 - 3.9	2,100	72.3	65.4 - 78.2	3,000
Japanese	<50	0.2	0.0 - 1.4	200	7.8	4.4 - 13.6	<50	0.2	0.0 - 1.4	2,100	91.7	86.0 - 95.2	2,300
Other	100	7.3	3.3 - 15.2	300	18.2	11.4 - 27.7	0	0.0	-	1,300	74.5	64.4 - 82.6	1,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	7.4	4.1 - 13.1	400	30.9	22.7 - 40.4	<50	1.1	0.4 - 3.2	700	60.6	51.0 - 69.4	1,200
20-24	300	7.1	4.4 - 11.3	700	19.9	15.5 - 25.1	100	2.8	1.7 - 4.4	2,500	70.3	64.6 - 75.4	3,600
25-34	200	3.0	1.9 - 4.8	900	12.5	10.0 - 15.6	200	2.9	1.8 - 4.7	5,900	81.6	78.1 - 84.7	7,200
35+	100	2.3	0.9 - 6.3	200	9.8	6.4 - 14.7	<50	2.1	0.7 - 6.2	2,000	85.8	79.9 - 90.1	2,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.9	3.2 - 7.4	1,100	15.7	12.9 - 19.0	200	2.4	1.4 - 3.9	5,400	77.0	73.3 - 80.4	7,000
Female	300	3.7	2.5 - 5.4	1,100	15.2	12.4 - 18.5	200	2.8	1.7 - 4.4	5,700	78.3	74.7 - 81.5	7,300

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.2	2.1 - 5.0	1,200	12.2	10.0 - 14.9	300	2.6	1.6 - 4.0	8,200	82.0	78.9 - 84.7	10,000
No	300	6.7	4.6 - 9.7	1,000	22.9	19.0 - 27.3	100	2.6	1.7 - 4.0	2,900	67.8	63.1 - 72.1	4,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.4	4.3 - 9.5	1,000	19.6	16.2 - 23.6	100	2.3	1.5 - 3.5	3,800	71.6	67.2 - 75.7	5,300
131-185%	100	4.5	2.5 - 8.2	500	15.9	11.6 - 21.4	100	2.7	1.2 - 6.1	2,300	76.9	70.7 - 82.0	3,000
186+%	100	1.6	0.8 - 3.4	500	9.1	6.6 - 12.6	200	3.1	1.7 - 5.7	4,300	86.2	82.2 - 89.4	5,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	200	7.6	4.7 - 12.0	500	21.1	16.1 - 27.1	100	3.2	1.9 - 5.3	1,700	68.2	61.8 - 73.9	2,500
\$15,001-\$25,000	100	6.2	3.2 - 11.6	300	15.3	10.8 - 21.2	100	3.0	1.4 - 6.2	1,600	75.6	68.6 - 81.4	2,100
\$25,001-\$40,000	200	5.5	2.9 - 9.9	500	16.4	12.2 - 21.8	100	2.5	1.0 - 5.9	2,300	75.7	69.4 - 81.0	3,000
>=\$40,001	100	1.8	0.9 - 3.5	700	10.9	8.3 - 14.3	100	2.4	1.3 - 4.4	5,100	84.9	81.2 - 88.0	6,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	6.0	1.9 - 17.0	200	22.2	13.4 - 34.5	<50	2.7	0.6 - 11.9	600	69.0	56.1 - 79.6	900
Normal birth weight	600	4.2	3.1 - 5.6	2,000	15.0	12.9 - 17.3	300	2.6	1.8 - 3.6	10,500	78.3	75.7 - 80.7	13,400

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	5.5	2.2 - 13.1	300	30.5	21.3 - 41.6	<50	5.2	2.6 - 10.3	500	58.7	47.8 - 68.9	900
No	600	4.2	3.1 - 5.7	1,900	14.4	12.4 - 16.7	300	2.4	1.7 - 3.6	10,500	78.9	76.3 - 81.4	13,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	7.4	4.3 - 12.2	600	22.8	17.7 - 28.9	100	4.5	2.6 - 7.9	1,700	65.3	58.6 - 71.5	2,600
No	300	3.0	2.1 - 4.4	1,600	14.0	11.9 - 16.5	300	2.2	1.4 - 3.4	9,200	80.7	77.9 - 83.2	11,400

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	6.4	4.3 - 9.4	1,200	22.1	18.4 - 26.2	100	2.5	1.5 - 4.1	3,900	69.0	64.5 - 73.2	5,600
No	300	3.0	2.0 - 4.4	1,000	11.3	9.1 - 13.9	200	2.6	1.6 - 4.1	7,200	83.2	80.1 - 85.8	8,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.6	3.7 - 5.6	2,400	17.0	15.3 - 18.9	400	3.0	2.2 - 3.9	10,600	75.4	73.3 - 77.5	14,000

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.1	3.3 - 7.8	200	15.0	11.9 - 18.9	100	3.5	2.0 - 6.0	1,200	76.4	71.9 - 80.4	1,500
Honolulu	400	4.3	3.3 - 5.6	1,800	17.4	15.1 - 19.8	300	2.8	1.9 - 4.0	7,800	75.6	72.8 - 78.1	10,400
Kauai	<50	4.2	2.0 - 8.4	100	15.0	9.9 - 22.2	<50	4.2	1.9 - 9.2	500	76.6	68.8 - 82.9	700
Maui	100	5.9	3.8 - 9.0	300	17.3	13.5 - 21.8	100	3.4	2.0 - 5.8	1,100	73.4	68.4 - 77.9	1,500

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.2	1.1 - 4.3	400	12.7	9.5 - 16.7	100	1.9	0.9 - 4.2	2,600	83.2	78.8 - 86.8	3,100
Native Hawaiian	300	7.3	6.0 - 8.9	700	20.9	18.7 - 23.2	100	3.2	2.4 - 4.3	2,400	68.6	66.0 - 71.0	3,600
Chinese	<50	3.1	1.1 - 8.5	100	15.8	7.5 - 30.3	<50	5.0	1.3 - 17.3	500	76.1	61.6 - 86.4	600
Filipino	200	5.4	3.4 - 8.5	600	20.4	15.9 - 25.7	100	3.1	1.6 - 6.1	2,100	71.1	65.3 - 76.2	3,000
Japanese	<50	0.9	0.3 - 2.6	200	12.4	8.3 - 18.3	100	4.2	2.0 - 9.0	1,600	82.4	75.9 - 87.4	1,900
Other	100	6.2	3.3 - 11.3	300	16.5	11.4 - 23.4	<50	2.0	0.7 - 5.2	1,400	75.3	67.8 - 81.5	1,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.9	6.6 - 14.7	400	34.0	27.8 - 40.8	<50	0.8	0.3 - 2.1	700	55.3	48.4 - 62.0	1,300
20-24	200	6.6	4.6 - 9.6	800	22.6	18.9 - 26.8	100	3.1	1.8 - 5.2	2,400	67.6	63.1 - 71.9	3,600
25-34	200	3.2	2.3 - 4.2	900	13.4	11.1 - 16.2	200	3.0	2.0 - 4.5	5,500	80.4	77.4 - 83.1	6,800
35+	100	2.5	1.3 - 4.8	200	9.1	5.9 - 13.6	100	4.0	2.1 - 7.5	1,900	84.4	79.2 - 88.5	2,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	5.0	3.7 - 6.5	1,200	17.0	14.6 - 19.8	200	2.8	1.8 - 4.3	5,500	75.2	72.1 - 78.1	7,300
Female	300	4.1	3.1 - 5.5	1,100	17.0	14.6 - 19.7	200	3.2	2.1 - 4.6	5,100	75.7	72.7 - 78.5	6,700

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.2	2.4 - 4.2	1,300	13.8	11.8 - 16.0	300	3.3	2.3 - 4.7	7,700	79.8	77.2 - 82.1	9,600
No	300	7.6	5.8 - 10.0	1,100	24.2	20.9 - 27.8	100	2.2	1.4 - 3.5	2,900	66.0	62.1 - 69.7	4,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.4	4.8 - 8.4	1,000	20.5	17.6 - 23.8	100	2.5	1.6 - 4.0	3,500	70.6	66.9 - 73.9	5,000
131-185%	100	3.0	2.0 - 4.5	500	16.5	13.0 - 20.6	100	2.8	1.7 - 4.6	2,400	77.8	73.4 - 81.6	3,100
186+%	100	2.7	1.7 - 4.1	500	10.0	7.7 - 12.9	200	3.8	2.4 - 6.0	4,200	83.5	80.1 - 86.5	5,000

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	200	8.0	5.7 - 11.3	500	20.1	16.1 - 24.7	100	2.1	1.1 - 4.0	1,800	69.8	64.7 - 74.5	2,600
\$15,001-\$25,000	100	4.9	2.8 - 8.4	400	23.6	18.5 - 29.6	100	4.3	2.2 - 8.4	1,200	67.2	60.8 - 73.0	1,800
\$25,001-\$40,000	100	2.8	2.0 - 4.1	600	18.2	14.6 - 22.4	100	2.5	1.4 - 4.3	2,400	76.5	72.1 - 80.4	3,100
>=\$40,001	200	2.8	2.0 - 4.1	600	10.7	8.5 - 13.5	200	3.6	2.3 - 5.5	4,800	82.8	79.6 - 85.6	5,800

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	4.4	1.8 - 10.3	200	22.7	15.4 - 32.2	<50	2.1	0.8 - 5.4	600	70.8	61.1 - 78.9	900
Normal birth weight	600	4.6	3.7 - 5.6	2,200	16.7	14.9 - 18.6	400	3.0	2.3 - 4.1	10,000	75.7	73.5 - 77.8	13,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	6.3	4.0 - 9.9	300	27.7	20.4 - 36.3	<50	2.8	1.3 - 5.7	700	63.2	54.8 - 70.9	1,100
No	500	4.3	3.4 - 5.3	2,000	16.0	14.3 - 18.0	400	3.0	2.2 - 4.1	9,800	76.6	74.4 - 78.7	12,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	5.1	3.6 - 7.0	700	26.0	21.5 - 31.1	100	3.0	1.8 - 5.1	1,700	65.9	60.7 - 70.7	2,600
No	500	4.2	3.3 - 5.4	1,600	14.8	12.9 - 16.9	300	2.9	2.1 - 4.0	8,500	78.1	75.7 - 80.3	10,900

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	6.8	5.3 - 8.7	1,200	21.7	18.8 - 24.8	200	3.4	2.3 - 5.1	3,900	68.2	64.7 - 71.4	5,700
No	200	3.0	2.2 - 4.1	1,100	13.8	11.6 - 16.3	200	2.7	1.8 - 4.0	6,600	80.5	77.8 - 83.0	8,200

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	3.8	3.0 - 4.9	2,300	16.3	14.7 - 18.1	600	4.0	3.2 - 5.0	10,600	75.9	73.8 - 77.8	14,000

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.0	3.1 - 8.1	400	23.2	19.3 - 27.7	100	3.8	2.3 - 6.4	1,100	67.9	63.0 - 72.5	1,600
Honolulu	400	3.7	2.7 - 5.1	1,500	14.9	13.0 - 17.1	400	4.1	3.1 - 5.4	8,000	77.3	74.7 - 79.7	10,400
Kauai	<50	3.5	1.4 - 8.5	100	15.6	10.2 - 23.0	<50	1.9	0.5 - 6.4	500	79.0	71.1 - 85.2	600
Maui	100	3.5	2.0 - 6.2	300	18.3	14.5 - 22.9	100	4.2	2.4 - 7.1	1,100	73.9	68.9 - 78.4	1,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.6	2.1 - 6.3	400	13.0	9.9 - 17.0	200	4.4	2.9 - 6.8	2,700	78.9	74.4 - 82.8	3,400
Native Hawaiian	200	4.8	3.7 - 6.1	900	24.4	22.0 - 26.9	200	4.4	3.4 - 5.7	2,400	66.4	63.7 - 69.0	3,600
Chinese	<50	2.5	0.4 - 15.1	100	18.0	9.3 - 32.0	<50	2.5	0.4 - 15.1	400	76.9	62.4 - 87.0	600
Filipino	100	4.3	2.4 - 7.6	500	17.4	13.6 - 22.0	100	3.4	2.0 - 5.9	2,100	74.9	69.7 - 79.5	2,900
Japanese	<50	1.0	0.4 - 2.9	100	4.5	2.1 - 9.4	100	4.6	2.0 - 10.1	1,600	89.8	83.7 - 93.8	1,800
Other	100	5.0	2.3 - 10.4	300	15.7	10.5 - 22.7	100	2.9	1.1 - 7.0	1,400	76.5	68.8 - 82.8	1,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.4	5.7 - 12.3	400	28.7	23.3 - 34.8	100	3.7	1.8 - 7.4	900	59.2	52.7 - 65.3	1,600
20-24	200	5.7	3.7 - 8.7	700	21.4	18.0 - 25.3	100	3.4	2.3 - 5.1	2,300	69.5	65.1 - 73.5	3,300
25-34	200	2.8	1.8 - 4.3	900	13.3	11.1 - 15.9	300	4.0	2.9 - 5.5	5,400	79.9	76.9 - 82.6	6,800
35+	<50	1.4	0.5 - 4.1	200	9.6	6.6 - 13.9	100	4.8	2.6 - 8.7	2,000	84.2	79.0 - 88.3	2,400

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.9	2.8 - 5.4	1,200	16.8	14.5 - 19.4	200	3.3	2.3 - 4.5	5,400	76.0	73.1 - 78.7	7,100
Female	300	3.8	2.6 - 5.5	1,100	15.8	13.6 - 18.3	300	4.7	3.4 - 6.4	5,300	75.7	72.7 - 78.5	6,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	2.4	1.6 - 3.6	1,300	13.2	11.4 - 15.3	400	4.0	3.0 - 5.3	7,800	80.4	77.9 - 82.6	9,700
No	300	7.1	5.3 - 9.6	1,000	23.2	20.1 - 26.6	200	3.9	2.7 - 5.6	2,800	65.8	62.0 - 69.5	4,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.1	3.8 - 7.0	1,300	22.2	19.4 - 25.2	200	3.5	2.5 - 4.8	4,000	69.2	65.9 - 72.4	5,800
131-185%	100	3.6	1.8 - 7.0	300	13.3	10.0 - 17.5	<50	1.9	1.0 - 3.4	1,600	81.2	76.4 - 85.3	2,000
186+%	100	2.2	1.2 - 3.8	600	10.6	8.3 - 13.4	300	5.3	3.6 - 7.5	4,400	82.0	78.6 - 85.0	5,300

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	200	7.8	5.5 - 10.9	800	24.5	20.6 - 28.8	100	3.4	2.1 - 5.4	2,000	64.3	59.5 - 68.8	3,100
\$15,001-\$25,000	100	2.8	1.4 - 5.6	400	20.2	15.8 - 25.5	100	3.1	1.8 - 5.3	1,600	73.8	68.2 - 78.7	2,100
\$25,001-\$40,000	100	3.0	1.5 - 5.8	400	15.6	12.4 - 19.5	100	3.3	2.1 - 5.0	2,200	78.2	73.8 - 82.0	2,800
>=\$40,001	100	2.0	1.2 - 3.4	500	9.8	7.7 - 12.4	300	4.9	3.3 - 7.1	4,400	83.3	80.0 - 86.2	5,300

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	5.5	2.1 - 13.4	200	20.7	14.4 - 28.7	<50	4.0	1.9 - 8.4	600	69.8	60.6 - 77.6	900
Normal birth weight	500	3.8	2.9 - 4.8	2,100	15.8	14.1 - 17.6	500	3.9	3.0 - 4.9	10,000	76.6	74.5 - 78.6	13,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	10.7	6.3 - 17.7	300	29.5	22.3 - 37.9	<50	3.2	1.6 - 6.5	500	56.6	47.9 - 64.8	900
No	400	3.1	2.3 - 4.1	1,900	15.2	13.5 - 17.0	500	3.9	3.1 - 5.0	9,900	77.9	75.7 - 79.8	12,700

PRAMS Health Indicator Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	6.4	4.3 - 9.5	700	27.1	22.7 - 32.0	100	4.5	2.7 - 7.6	1,600	62.0	56.8 - 67.0	2,600
No	300	3.0	2.1 - 4.1	1,500	13.4	11.7 - 15.4	400	3.7	2.9 - 4.8	8,800	79.8	77.6 - 81.9	11,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	5.6	4.1 - 7.6	1,200	20.8	18.3 - 23.5	200	3.2	2.3 - 4.4	4,100	70.5	67.3 - 73.4	5,800
No	200	2.4	1.5 - 3.6	1,000	13.1	11.0 - 15.6	400	4.4	3.2 - 6.0	6,400	80.1	77.3 - 82.7	8,000