

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration (grouped)	Breastfeeding (any) duration (grouped)

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/22/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013


Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	1,700	10.4	9.7 - 11.0	600	3.9	3.5 - 4.3	2,400	14.5	13.7 - 15.3	500	3.3	2.9 - 3.7	11,100	68.1	67.0 - 69.1	16,300
2001-2005	1,700	10.2	9.5 - 10.9	600	3.9	3.5 - 4.4	2,400	14.4	13.7 - 15.2	500	3.2	2.8 - 3.6	11,300	68.3	67.2 - 69.3	16,500
2002-2006	1,600	9.7	9.0 - 10.4	600	3.6	3.2 - 4.1	2,500	14.6	13.8 - 15.4	600	3.3	2.9 - 3.7	11,700	68.8	67.7 - 69.8	16,900
2003-2007	1,600	9.1	8.4 - 9.7	600	3.6	3.2 - 4.0	2,600	14.8	14.1 - 15.6	600	3.4	3.0 - 3.8	12,000	69.1	68.1 - 70.1	17,300
2004-2008	1,500	8.5	7.9 - 9.1	600	3.3	2.9 - 3.7	2,600	14.7	14.0 - 15.5	600	3.5	3.1 - 3.9	12,400	70.1	69.1 - 71.0	17,600
2005-2009	1,400	8.0	7.4 - 8.7	600	3.4	3.0 - 3.8	2,600	14.8	14.0 - 15.7	700	3.7	3.3 - 4.2	12,500	70.1	69.0 - 71.2	17,800
2006-2010	1,300	7.2	6.6 - 7.9	600	3.2	2.8 - 3.7	2,600	14.6	13.7 - 15.5	700	3.9	3.4 - 4.4	12,800	71.1	69.9 - 72.2	18,000
2007-2011	1,200	6.5	5.9 - 7.2	600	3.0	2.6 - 3.5	2,500	13.8	12.9 - 14.7	700	4.0	3.5 - 4.6	13,200	72.6	71.4 - 73.8	18,100
2008-2012	1,100	6.0	5.4 - 6.7	500	2.9	2.4 - 3.4	2,400	13.3	12.4 - 14.3	700	4.0	3.5 - 4.6	13,400	73.8	72.6 - 75.0	18,200
2009-2013	900	5.2	4.6 - 5.9	500	2.7	2.3 - 3.3	2,300	12.8	11.8 - 13.8	700	3.8	3.2 - 4.4	13,700	75.5	74.2 - 76.8	18,100

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	900	5.2	4.6 - 5.9	500	2.7	2.3 - 3.3	2,300	12.8	11.8 - 13.8	700	3.8	3.2 - 4.4	13,700	75.5	74.2 - 76.8	18,100

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.4	4.4 - 6.6	100	3.0	2.2 - 3.9	300	14.0	12.4 - 15.7	100	3.5	2.7 - 4.5	1,700	74.2	72.0 - 76.2	2,300
Honolulu	700	5.6	4.8 - 6.6	400	2.8	2.2 - 3.5	1,600	12.5	11.3 - 13.9	500	3.8	3.1 - 4.6	9,800	75.3	73.6 - 76.9	13,000
Kauai	<50	3.7	2.8 - 4.9	<50	1.6	1.0 - 2.5	100	11.3	9.7 - 13.2	<50	3.2	2.4 - 4.3	700	80.1	77.9 - 82.2	800
Maui	100	2.9	2.3 - 3.8	100	2.8	2.1 - 3.6	300	13.7	12.2 - 15.3	100	3.8	3.1 - 4.8	1,500	76.7	74.8 - 78.5	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	3.6	2.6 - 5.0	100	2.6	1.7 - 3.9	400	9.4	7.7 - 11.5	200	4.1	3.0 - 5.5	3,600	80.3	77.7 - 82.6	4,400
Native Hawaiian	400	7.0	5.7 - 8.6	200	3.1	2.3 - 4.2	900	17.0	15.1 - 19.0	200	4.6	3.6 - 5.8	3,700	68.3	65.8 - 70.7	5,400
Chinese	100	6.7	3.7 - 11.7	<50	2.4	1.0 - 5.8	100	8.0	4.9 - 12.9	<50	2.0	0.8 - 5.3	700	80.9	74.5 - 86.0	800
Filipino	100	4.4	3.2 - 6.0	100	3.0	2.1 - 4.3	500	17.3	14.9 - 20.0	100	4.1	2.9 - 5.7	2,300	71.3	68.1 - 74.3	3,200
Japanese	<50	1.3	0.6 - 3.0	<50	1.0	0.4 - 2.4	100	6.1	4.2 - 8.9	<50	1.6	0.8 - 3.2	1,500	89.9	86.7 - 92.4	1,700
Other	200	7.2	5.3 - 9.8	100	3.1	2.0 - 4.9	300	10.3	8.1 - 13.1	100	3.0	1.9 - 4.8	2,000	76.3	72.6 - 79.6	2,600

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.0	2.8 - 8.5	100	4.4	2.6 - 7.3	300	28.0	23.2 - 33.3	100	5.3	3.3 - 8.6	700	57.4	51.7 - 62.9	1,100
20-24	200	5.8	4.5 - 7.5	100	3.1	2.2 - 4.4	600	15.5	13.4 - 17.9	200	3.9	2.8 - 5.3	3,000	71.7	68.8 - 74.4	4,100
25-34	500	5.0	4.2 - 6.1	200	2.4	1.9 - 3.2	1,100	11.4	10.2 - 12.8	400	3.7	3.0 - 4.6	7,500	77.4	75.6 - 79.0	9,700
35+	200	5.0	3.7 - 6.7	100	2.6	1.7 - 3.9	300	7.9	6.3 - 9.9	100	3.2	2.2 - 4.5	2,600	81.3	78.6 - 83.8	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	500	5.3	4.4 - 6.3	300	3.0	2.3 - 3.8	1,200	12.5	11.2 - 14.0	300	3.8	3.1 - 4.7	6,900	75.4	73.6 - 77.2	9,200
Female	500	5.1	4.3 - 6.2	200	2.5	1.9 - 3.3	1,200	13.0	11.7 - 14.5	300	3.7	3.0 - 4.6	6,800	75.6	73.8 - 77.4	9,000

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	3.9	3.2 - 4.7	300	2.5	1.9 - 3.2	1,200	10.4	9.3 - 11.6	400	3.5	2.9 - 4.2	9,000	79.8	78.2 - 81.3	11,200
No	500	7.4	6.2 - 8.8	200	3.2	2.4 - 4.1	1,200	16.7	15.1 - 18.5	300	4.2	3.3 - 5.3	4,700	68.5	66.3 - 70.7	6,900

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.9	5.8 - 8.3	200	3.7	2.9 - 4.7	1,100	16.6	14.9 - 18.4	200	3.7	2.8 - 4.7	4,400	69.2	66.9 - 71.4	6,400
131-185%	200	5.7	4.1 - 7.9	100	3.1	2.0 - 4.9	400	13.6	11.1 - 16.4	100	4.3	3.0 - 6.0	2,100	73.4	69.8 - 76.6	2,800
186+%	200	2.9	2.2 - 3.8	200	2.0	1.5 - 2.9	600	8.5	7.2 - 9.9	300	3.7	2.9 - 4.7	6,200	82.9	81.1 - 84.6	7,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.5	4.5 - 6.8	<50	2.8	2.1 - 3.8	200	15.5	13.8 - 17.3	100	4.8	3.9 - 6.0	900	71.4	69.1 - 73.5	1,200
Normal birth weight	900	5.2	4.5 - 6.0	500	2.7	2.3 - 3.3	2,100	12.6	11.6 - 13.7	600	3.7	3.1 - 4.3	12,800	75.8	74.4 - 77.1	16,900

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	14.0	10.4 - 18.6	100	6.5	4.1 - 10.1	200	21.7	17.5 - 26.7	100	5.1	3.1 - 8.4	600	52.6	46.9 - 58.3	1,100
No	800	4.6	4.0 - 5.4	400	2.5	2.1 - 3.1	2,000	12.1	11.2 - 13.2	600	3.6	3.1 - 4.3	12,800	77.1	75.7 - 78.3	16,700

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	8.3	6.7 - 10.4	200	4.8	3.6 - 6.6	700	19.0	16.5 - 21.7	200	5.2	3.8 - 6.9	2,200	62.7	59.4 - 65.8	3,400
No	600	4.5	3.8 - 5.2	300	2.3	1.8 - 2.8	1,600	11.3	10.2 - 12.3	500	3.4	2.9 - 4.1	11,300	78.6	77.2 - 79.9	14,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	6.8	5.8 - 8.1	300	3.9	3.1 - 4.8	1,300	16.9	15.3 - 18.6	300	3.9	3.2 - 4.9	5,400	68.5	66.4 - 70.5	7,900
No	400	3.9	3.1 - 4.8	200	1.9	1.4 - 2.5	1,000	9.5	8.4 - 10.8	400	3.5	2.8 - 4.3	8,200	81.2	79.6 - 82.7	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,100	6.0	5.4 - 6.7	500	2.9	2.4 - 3.4	2,400	13.3	12.4 - 14.3	700	4.0	3.5 - 4.6	13,400	73.8	72.6 - 75.0	18,200

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.4	5.2 - 7.8	100	3.6	2.7 - 4.7	300	14.8	13.1 - 16.7	100	3.4	2.6 - 4.4	1,700	71.8	69.5 - 74.0	2,300
Honolulu	800	6.4	5.5 - 7.4	400	2.8	2.2 - 3.5	1,700	13.2	12.0 - 14.5	500	4.1	3.4 - 4.8	9,600	73.6	71.9 - 75.1	13,000
Kauai	<50	3.2	2.2 - 4.6	<50	1.8	1.1 - 2.8	100	11.5	9.4 - 13.9	<50	4.7	3.3 - 6.7	700	78.9	75.9 - 81.6	800
Maui	100	4.2	3.3 - 5.5	100	2.9	2.2 - 3.9	300	13.1	11.6 - 14.8	100	4.1	3.3 - 5.1	1,500	75.6	73.5 - 77.7	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	4.2	3.2 - 5.6	100	2.6	1.7 - 3.8	400	10.1	8.4 - 12.0	200	4.8	3.7 - 6.2	3,400	78.4	75.9 - 80.7	4,300
Native Hawaiian	400	7.6	6.3 - 9.2	200	3.3	2.5 - 4.4	900	17.0	15.2 - 18.9	300	5.0	4.0 - 6.2	3,600	67.1	64.7 - 69.4	5,300
Chinese	100	6.7	3.7 - 11.7	<50	2.5	1.0 - 6.2	<50	6.2	3.8 - 10.2	<50	2.8	1.3 - 6.0	600	81.8	75.7 - 86.6	800
Filipino	200	5.6	4.3 - 7.2	100	3.1	2.2 - 4.2	600	18.1	15.7 - 20.7	100	3.8	2.8 - 5.3	2,300	69.5	66.5 - 72.3	3,200
Japanese	<50	1.6	0.8 - 3.1	<50	1.5	0.8 - 2.9	100	7.5	5.5 - 10.2	<50	2.2	1.2 - 3.7	1,500	87.2	84.0 - 89.8	1,700
Other	200	8.8	6.8 - 11.2	100	3.2	2.1 - 4.9	300	11.4	9.3 - 14.0	100	2.7	1.7 - 4.2	2,000	73.9	70.5 - 77.1	2,800

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.6	4.3 - 9.9	100	4.8	3.0 - 7.5	300	26.6	22.3 - 31.4	100	4.8	3.0 - 7.5	700	57.3	52.1 - 62.3	1,300
20-24	300	7.3	5.9 - 9.0	100	3.3	2.3 - 4.5	700	17.8	15.7 - 20.2	200	4.2	3.2 - 5.5	2,800	67.4	64.6 - 70.1	4,100
25-34	600	5.8	4.9 - 6.8	200	2.4	1.9 - 3.1	1,100	11.3	10.1 - 12.5	400	4.1	3.4 - 5.0	7,300	76.4	74.7 - 78.0	9,600
35+	200	4.8	3.6 - 6.4	100	3.0	2.1 - 4.3	300	8.3	6.7 - 10.2	100	3.1	2.2 - 4.3	2,600	80.8	78.2 - 83.2	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	500	5.9	5.0 - 6.9	300	3.1	2.5 - 3.9	1,300	13.5	12.2 - 14.8	400	4.3	3.5 - 5.1	6,800	73.3	71.5 - 75.0	9,300
Female	500	6.1	5.2 - 7.2	200	2.6	2.0 - 3.3	1,200	13.1	11.9 - 14.5	300	3.8	3.1 - 4.6	6,600	74.3	72.6 - 76.0	8,900

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.5	3.8 - 5.3	300	2.5	2.0 - 3.1	1,200	11.0	9.9 - 12.1	400	3.9	3.3 - 4.6	8,800	78.2	76.7 - 79.6	11,200
No	600	8.5	7.3 - 9.9	200	3.5	2.8 - 4.4	1,200	17.1	15.5 - 18.8	300	4.2	3.4 - 5.2	4,600	66.7	64.6 - 68.8	6,900

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	500	8.5	7.2 - 9.9	200	3.8	3.0 - 4.8	1,100	17.1	15.4 - 18.9	200	3.6	2.8 - 4.5	4,300	67.1	64.9 - 69.2	6,400
131-185%	200	6.3	4.7 - 8.3	100	3.0	1.9 - 4.7	400	13.4	11.1 - 16.0	100	4.4	3.2 - 6.1	2,100	73.0	69.6 - 76.1	2,800
186+%	200	3.1	2.4 - 4.0	200	2.1	1.5 - 2.8	700	9.3	8.2 - 10.7	300	4.3	3.5 - 5.3	6,100	81.1	79.4 - 82.8	7,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	6.4	5.0 - 8.0	<50	3.4	2.4 - 4.7	200	15.3	13.3 - 17.5	100	4.7	3.6 - 6.0	900	70.3	67.5 - 72.9	1,200
Normal birth weight	1,000	6.0	5.3 - 6.8	500	2.8	2.4 - 3.4	2,200	13.2	12.2 - 14.2	700	4.0	3.4 - 4.6	12,500	74.1	72.7 - 75.3	16,900

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	15.3	11.8 - 19.5	100	5.3	3.3 - 8.4	300	22.4	18.5 - 26.9	100	6.5	4.4 - 9.7	600	50.4	45.3 - 55.6	1,200
No	900	5.3	4.7 - 6.1	400	2.7	2.3 - 3.2	2,100	12.6	11.6 - 13.6	600	3.8	3.3 - 4.4	12,500	75.6	74.3 - 76.8	16,600

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	9.9	8.1 - 12.0	100	4.2	3.0 - 5.8	700	19.6	17.3 - 22.2	200	6.2	4.8 - 7.9	2,100	60.1	57.0 - 63.2	3,500
No	700	5.1	4.4 - 5.9	400	2.6	2.1 - 3.1	1,700	11.7	10.8 - 12.8	500	3.5	3.0 - 4.1	11,100	77.1	75.8 - 78.4	14,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	8.1	7.0 - 9.4	300	4.1	3.3 - 5.0	1,400	17.3	15.8 - 18.9	300	3.9	3.2 - 4.7	5,200	66.6	64.6 - 68.6	7,900
No	400	4.3	3.6 - 5.2	200	2.0	1.5 - 2.6	1,000	10.2	9.1 - 11.4	400	4.0	3.3 - 4.8	8,000	79.4	77.9 - 80.9	10,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,200	6.5	5.9 - 7.2	600	3.0	2.6 - 3.5	2,500	13.8	12.9 - 14.7	700	4.0	3.5 - 4.6	13,200	72.6	71.4 - 73.8	18,100

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	7.8	6.4 - 9.4	100	4.1	3.1 - 5.3	300	13.7	12.0 - 15.6	100	2.8	2.1 - 3.8	1,700	71.6	69.1 - 74.0	2,300
Honolulu	800	6.6	5.8 - 7.5	400	2.9	2.3 - 3.5	1,800	14.0	12.9 - 15.2	500	4.2	3.6 - 4.9	9,300	72.3	70.8 - 73.8	12,900
Kauai	<50	4.8	3.2 - 7.1	<50	1.3	0.7 - 2.5	100	11.9	9.6 - 14.7	<50	5.0	3.5 - 7.2	700	76.9	73.4 - 80.1	900
Maui	100	5.4	4.2 - 6.9	100	3.6	2.7 - 4.7	300	13.6	11.9 - 15.6	100	3.7	2.9 - 4.8	1,500	73.7	71.3 - 76.0	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.1	4.0 - 6.4	100	2.6	1.8 - 3.7	500	10.5	8.9 - 12.3	200	5.2	4.1 - 6.6	3,300	76.7	74.3 - 78.9	4,300
Native Hawaiian	400	8.1	6.8 - 9.6	200	4.0	3.2 - 5.1	900	17.9	16.1 - 19.8	200	4.4	3.5 - 5.5	3,500	65.6	63.2 - 67.9	5,300
Chinese	<50	5.4	2.8 - 10.2	<50	2.0	0.9 - 4.7	<50	6.7	4.3 - 10.3	<50	2.5	1.3 - 5.0	600	83.4	78.0 - 87.7	700
Filipino	200	6.4	5.2 - 8.0	100	2.6	1.9 - 3.5	600	17.8	15.6 - 20.2	100	4.1	3.1 - 5.4	2,200	69.1	66.3 - 71.8	3,300
Japanese	<50	1.9	1.1 - 3.2	<50	1.7	0.9 - 3.1	100	7.8	5.8 - 10.3	<50	2.3	1.4 - 3.9	1,600	86.3	83.3 - 88.8	1,800
Other	200	9.1	7.2 - 11.5	100	3.5	2.4 - 5.2	300	12.5	10.4 - 15.1	100	2.9	1.9 - 4.3	2,000	72.0	68.6 - 75.1	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.0	5.7 - 11.1	100	5.1	3.5 - 7.5	400	25.7	21.7 - 30.1	100	5.0	3.3 - 7.7	800	56.2	51.5 - 60.9	1,400
20-24	400	8.2	6.8 - 9.9	100	3.1	2.3 - 4.3	800	18.8	16.8 - 21.0	200	4.0	3.1 - 5.2	2,800	65.8	63.1 - 68.3	4,300
25-34	600	6.1	5.3 - 7.1	200	2.7	2.1 - 3.3	1,100	11.6	10.4 - 12.8	400	4.0	3.3 - 4.8	7,000	75.6	74.0 - 77.2	9,300
35+	100	4.6	3.5 - 6.1	100	3.1	2.2 - 4.3	300	8.5	6.9 - 10.4	100	3.6	2.6 - 4.9	2,600	80.2	77.7 - 82.5	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	600	6.4	5.6 - 7.4	300	2.9	2.3 - 3.6	1,300	14.1	12.9 - 15.5	400	4.4	3.7 - 5.2	6,800	72.2	70.5 - 73.8	9,400
Female	600	6.6	5.7 - 7.6	300	3.2	2.6 - 3.9	1,200	13.5	12.2 - 14.8	300	3.6	3.0 - 4.4	6,400	73.1	71.4 - 74.7	8,700

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	4.9	4.2 - 5.7	300	2.6	2.1 - 3.2	1,200	11.1	10.1 - 12.2	400	3.9	3.3 - 4.6	8,700	77.6	76.1 - 78.9	11,200
No	600	9.1	7.9 - 10.4	300	3.8	3.0 - 4.7	1,300	18.3	16.7 - 20.0	300	4.3	3.5 - 5.2	4,400	64.6	62.5 - 66.6	6,900

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	600	9.2	8.0 - 10.6	300	4.0	3.2 - 4.9	1,200	18.2	16.6 - 19.9	200	3.6	2.9 - 4.5	4,200	65.0	62.9 - 67.1	6,500
131-185%	200	6.5	5.0 - 8.5	100	3.3	2.2 - 5.0	400	13.4	11.2 - 16.0	100	4.1	2.9 - 5.6	2,000	72.7	69.5 - 75.6	2,700
186+%	300	3.7	3.0 - 4.6	200	2.3	1.8 - 3.0	700	9.5	8.4 - 10.8	300	4.3	3.5 - 5.2	6,100	80.1	78.4 - 81.7	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	9.0	7.4 - 10.9	100	4.4	3.3 - 5.9	600	20.0	17.6 - 22.6	100	3.5	2.5 - 4.9	2,000	63.1	60.0 - 66.1	3,100
\$10,000-\$24,999	300	8.8	7.2 - 10.7	100	3.8	2.8 - 5.1	600	15.7	13.7 - 17.8	200	4.2	3.2 - 5.5	2,400	67.5	64.7 - 70.2	3,600
\$25,000-\$49,999	300	6.3	5.1 - 7.8	100	3.1	2.2 - 4.4	600	14.6	12.7 - 16.6	200	4.3	3.3 - 5.7	3,000	71.6	69.0 - 74.1	4,200
>=\$50,000	200	3.5	2.7 - 4.4	100	2.1	1.5 - 2.8	500	8.4	7.2 - 9.8	200	3.9	3.1 - 4.9	5,000	82.2	80.4 - 83.9	6,100

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.4	5.8 - 9.3	<50	3.6	2.5 - 5.0	200	16.5	14.2 - 19.1	100	4.5	3.4 - 5.9	900	68.0	64.9 - 71.0	1,300
Normal birth weight	1,100	6.4	5.8 - 7.2	500	3.0	2.6 - 3.5	2,300	13.6	12.7 - 14.6	700	4.0	3.5 - 4.6	12,300	73.0	71.7 - 74.2	16,800

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	16.6	13.3 - 20.6	100	5.9	3.9 - 8.7	300	23.1	19.4 - 27.1	100	5.8	4.0 - 8.3	700	48.7	44.0 - 53.4	1,400
No	900	5.7	5.1 - 6.4	500	2.8	2.4 - 3.3	2,100	13.0	12.1 - 13.9	600	3.9	3.4 - 4.4	12,300	74.7	73.5 - 75.9	16,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	11.0	9.3 - 13.1	200	4.2	3.1 - 5.7	800	21.5	19.2 - 24.0	200	5.6	4.4 - 7.1	2,100	57.6	54.7 - 60.5	3,600
No	800	5.4	4.7 - 6.1	400	2.8	2.3 - 3.3	1,700	11.8	10.9 - 12.8	500	3.6	3.1 - 4.2	10,900	76.5	75.2 - 77.7	14,200

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	8.9	7.8 - 10.1	300	3.9	3.2 - 4.8	1,400	17.8	16.3 - 19.3	300	3.9	3.2 - 4.8	5,000	65.6	63.6 - 67.4	7,700
No	500	4.7	4.0 - 5.6	200	2.4	1.9 - 3.0	1,100	10.8	9.7 - 12.0	400	4.1	3.4 - 4.8	7,900	78.0	76.5 - 79.4	10,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,300	7.2	6.6 - 7.9	600	3.2	2.8 - 3.7	2,600	14.6	13.7 - 15.5	700	3.9	3.4 - 4.4	12,800	71.1	69.9 - 72.2	18,000

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	8.3	6.8 - 10.1	100	4.4	3.4 - 5.8	300	14.2	12.3 - 16.3	100	2.4	1.7 - 3.5	1,600	70.6	68.0 - 73.2	2,300
Honolulu	900	7.3	6.5 - 8.2	400	3.2	2.6 - 3.8	1,900	14.8	13.7 - 15.9	500	4.1	3.5 - 4.8	9,100	70.6	69.2 - 72.1	12,800
Kauai	<50	5.7	3.9 - 8.3	<50	1.1	0.6 - 2.4	100	12.9	10.3 - 16.2	<50	4.9	3.3 - 7.2	700	75.3	71.3 - 78.9	900
Maui	100	6.3	4.9 - 8.0	100	3.3	2.4 - 4.5	300	14.4	12.4 - 16.6	100	3.7	2.7 - 4.9	1,500	72.4	69.7 - 75.0	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.0	4.9 - 7.4	100	2.8	2.0 - 3.9	500	10.7	9.2 - 12.4	200	5.5	4.4 - 6.9	3,200	74.9	72.6 - 77.1	4,300
Native Hawaiian	500	9.2	7.9 - 10.7	200	4.3	3.4 - 5.4	1,000	19.2	17.4 - 21.1	200	3.7	2.9 - 4.7	3,300	63.7	61.3 - 65.9	5,200
Chinese	<50	4.0	2.0 - 8.1	<50	1.6	1.0 - 2.7	100	8.4	5.8 - 12.0	<50	2.7	1.4 - 5.2	600	83.2	78.4 - 87.2	700
Filipino	200	6.9	5.6 - 8.4	100	2.9	2.2 - 3.9	600	18.3	16.2 - 20.6	100	4.2	3.2 - 5.5	2,200	67.7	65.0 - 70.2	3,200
Japanese	<50	2.5	1.6 - 3.9	<50	2.0	1.1 - 3.4	100	7.3	5.6 - 9.5	<50	2.1	1.2 - 3.5	1,600	86.1	83.3 - 88.5	1,900
Other	300	9.8	7.9 - 12.0	100	3.5	2.4 - 5.1	400	14.0	11.8 - 16.5	100	3.0	2.0 - 4.3	1,900	69.8	66.6 - 72.8	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.4	6.2 - 11.2	100	5.3	3.6 - 7.7	400	24.7	21.0 - 28.8	100	4.5	2.9 - 7.1	800	57.0	52.5 - 61.4	1,500
20-24	400	8.7	7.3 - 10.3	100	3.3	2.4 - 4.4	900	20.5	18.5 - 22.6	200	4.7	3.7 - 5.9	2,700	62.9	60.4 - 65.4	4,300
25-34	600	6.8	5.9 - 7.7	300	2.9	2.3 - 3.6	1,100	12.1	11.0 - 13.3	300	3.7	3.1 - 4.4	6,800	74.6	73.0 - 76.1	9,100
35+	200	6.1	4.8 - 7.7	100	3.2	2.3 - 4.5	300	9.1	7.5 - 10.9	100	3.2	2.4 - 4.4	2,500	78.4	75.9 - 80.7	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	600	7.0	6.1 - 8.0	300	3.0	2.4 - 3.7	1,400	15.4	14.1 - 16.6	400	4.5	3.8 - 5.3	6,500	70.2	68.5 - 71.7	9,300
Female	700	7.5	6.6 - 8.5	300	3.5	2.9 - 4.3	1,200	13.7	12.6 - 15.0	300	3.2	2.7 - 4.0	6,300	72.0	70.4 - 73.6	8,700

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	5.7	5.0 - 6.4	300	2.6	2.2 - 3.2	1,300	11.5	10.5 - 12.5	400	3.6	3.1 - 4.3	8,600	76.6	75.2 - 77.9	11,200
No	700	9.8	8.6 - 11.2	300	4.3	3.5 - 5.2	1,300	19.7	18.1 - 21.3	300	4.3	3.5 - 5.2	4,200	62.0	60.0 - 64.0	6,800

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	600	10.1	8.9 - 11.4	300	4.4	3.6 - 5.4	1,300	19.6	18.0 - 21.3	200	3.3	2.6 - 4.1	4,000	62.7	60.6 - 64.7	6,400
131-185%	200	7.6	6.0 - 9.5	100	3.5	2.3 - 5.2	400	13.3	11.2 - 15.8	100	4.4	3.3 - 6.0	1,900	71.2	68.0 - 74.1	2,700
186+%	300	4.1	3.4 - 4.9	200	2.4	1.9 - 3.1	800	10.2	9.1 - 11.4	300	4.2	3.5 - 5.1	6,000	79.1	77.5 - 80.6	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	10.1	8.4 - 12.1	100	4.7	3.6 - 6.3	700	21.9	19.5 - 24.6	100	3.5	2.5 - 4.9	1,800	59.7	56.7 - 62.7	3,100
\$10,000-\$24,999	300	9.3	7.7 - 11.1	200	4.4	3.4 - 5.8	600	16.4	14.5 - 18.4	100	3.8	2.9 - 5.0	2,400	66.1	63.5 - 68.7	3,700
\$25,000-\$49,999	300	6.8	5.6 - 8.2	100	3.2	2.3 - 4.5	600	15.5	13.7 - 17.5	200	4.3	3.3 - 5.5	2,900	70.3	67.8 - 72.6	4,100
>=\$50,000	200	4.1	3.3 - 5.0	100	2.0	1.5 - 2.7	500	8.6	7.4 - 9.9	200	3.9	3.2 - 4.9	4,900	81.4	79.6 - 83.0	6,000

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.0	7.1 - 11.3	100	3.9	2.8 - 5.6	200	15.6	13.2 - 18.4	100	3.9	2.8 - 5.3	900	67.6	64.2 - 70.8	1,300
Normal birth weight	1,200	7.1	6.4 - 7.8	500	3.2	2.7 - 3.7	2,400	14.5	13.6 - 15.5	700	3.9	3.4 - 4.5	11,900	71.3	70.1 - 72.5	16,700

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	18.8	15.5 - 22.5	100	6.5	4.5 - 9.2	300	21.2	17.9 - 24.9	100	5.0	3.5 - 7.2	700	48.6	44.2 - 53.0	1,500
No	1,000	6.2	5.6 - 6.9	500	3.0	2.5 - 3.4	2,300	13.9	13.0 - 14.8	600	3.8	3.3 - 4.3	11,900	73.2	72.0 - 74.3	16,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	12.4	10.6 - 14.4	200	4.9	3.7 - 6.4	800	21.1	18.9 - 23.5	200	5.6	4.5 - 7.1	2,000	56.0	53.2 - 58.8	3,600
No	800	5.9	5.3 - 6.7	400	2.8	2.4 - 3.3	1,800	12.9	11.9 - 13.8	500	3.4	3.0 - 4.0	10,600	74.9	73.7 - 76.1	14,100

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	9.8	8.7 - 11.0	300	4.5	3.7 - 5.4	1,400	18.8	17.3 - 20.3	300	3.8	3.1 - 4.7	4,800	63.1	61.2 - 65.0	7,500
No	500	5.4	4.7 - 6.2	200	2.3	1.8 - 2.9	1,200	11.5	10.5 - 12.6	400	3.9	3.3 - 4.6	7,800	76.9	75.5 - 78.3	10,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,400	8.0	7.4 - 8.7	600	3.4	3.0 - 3.8	2,600	14.8	14.0 - 15.7	700	3.7	3.3 - 4.2	12,500	70.1	69.0 - 71.2	17,800

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	9.9	8.2 - 11.9	100	4.1	3.1 - 5.5	300	13.9	11.9 - 16.1	100	2.2	1.5 - 3.3	1,600	69.8	67.0 - 72.5	2,300
Honolulu	1,000	8.0	7.2 - 8.8	400	3.3	2.8 - 3.9	1,900	14.9	13.9 - 16.0	500	3.9	3.4 - 4.5	8,900	69.8	68.5 - 71.2	12,700
Kauai	100	6.2	4.2 - 9.1	<50	1.2	0.6 - 2.4	100	13.4	10.5 - 17.0	<50	4.7	3.1 - 7.2	600	74.5	70.1 - 78.4	800
Maui	100	6.8	5.3 - 8.7	100	3.8	2.7 - 5.2	300	15.6	13.4 - 18.0	100	3.6	2.7 - 4.9	1,400	70.2	67.2 - 73.0	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	6.0	4.9 - 7.2	100	2.8	2.1 - 3.8	400	10.8	9.4 - 12.4	200	4.8	3.9 - 6.0	3,100	75.6	73.5 - 77.6	4,100
Native Hawaiian	500	10.6	9.2 - 12.1	200	4.7	3.8 - 5.8	1,000	19.4	17.6 - 21.2	200	3.8	3.0 - 4.8	3,100	61.6	59.3 - 63.8	5,100
Chinese	<50	6.0	3.6 - 9.8	<50	1.5	0.9 - 2.5	100	8.8	6.5 - 11.7	<50	2.4	1.6 - 3.5	500	81.3	77.2 - 84.8	700
Filipino	200	7.5	6.3 - 8.9	100	3.3	2.6 - 4.3	600	18.3	16.4 - 20.4	100	4.3	3.3 - 5.5	2,100	66.6	64.1 - 69.0	3,200
Japanese	100	3.4	2.3 - 4.8	<50	1.9	1.1 - 3.2	200	8.5	6.8 - 10.7	<50	1.9	1.1 - 3.1	1,700	84.4	81.7 - 86.7	2,000
Other	300	10.9	9.1 - 13.0	100	3.3	2.3 - 4.7	400	14.3	12.2 - 16.6	100	2.9	2.0 - 4.2	1,800	68.6	65.7 - 71.4	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.5	7.3 - 12.3	100	5.7	4.0 - 8.1	400	24.3	20.8 - 28.1	100	3.6	2.3 - 5.6	900	56.8	52.6 - 60.9	1,600
20-24	400	9.4	8.1 - 10.9	200	3.9	3.0 - 5.0	900	21.3	19.4 - 23.3	200	4.7	3.8 - 5.9	2,600	60.7	58.3 - 63.1	4,200
25-34	700	7.7	6.8 - 8.6	300	3.0	2.4 - 3.6	1,100	12.0	11.0 - 13.1	300	3.5	3.0 - 4.2	6,500	73.8	72.3 - 75.2	8,900
35+	200	6.3	5.1 - 7.7	100	2.7	1.9 - 3.7	300	9.3	7.7 - 11.0	100	2.9	2.1 - 3.9	2,500	78.9	76.6 - 81.1	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	7.8	6.9 - 8.7	300	3.2	2.6 - 3.8	1,400	15.3	14.1 - 16.5	400	4.4	3.8 - 5.2	6,400	69.3	67.8 - 70.8	9,200
Female	700	8.3	7.4 - 9.2	300	3.6	3.0 - 4.3	1,200	14.3	13.2 - 15.5	300	2.9	2.4 - 3.5	6,100	70.9	69.4 - 72.4	8,600

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	6.4	5.7 - 7.1	300	2.7	2.3 - 3.2	1,300	11.6	10.7 - 12.5	400	3.5	3.0 - 4.0	8,400	75.9	74.6 - 77.1	11,100
No	700	10.7	9.5 - 12.0	300	4.5	3.7 - 5.4	1,300	20.2	18.6 - 21.8	300	4.1	3.4 - 5.0	4,000	60.5	58.6 - 62.4	6,700

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	800	11.7	10.5 - 13.1	300	4.7	3.9 - 5.6	1,300	19.9	18.3 - 21.5	200	3.2	2.5 - 3.9	3,900	60.5	58.6 - 62.5	6,400
131-185%	200	8.8	7.1 - 10.8	100	3.2	2.1 - 4.7	300	13.5	11.5 - 15.9	100	4.0	2.9 - 5.5	1,700	70.5	67.4 - 73.3	2,400
186+%	300	4.2	3.6 - 5.0	200	2.4	1.9 - 3.0	800	10.4	9.4 - 11.5	300	4.1	3.5 - 4.9	6,000	78.8	77.4 - 80.2	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	400	12.0	10.2 - 14.1	200	5.3	4.1 - 6.8	700	22.6	20.2 - 25.1	100	3.5	2.6 - 4.8	1,700	56.6	53.6 - 59.5	3,000
\$10,000-\$24,999	400	10.3	8.8 - 12.0	200	4.8	3.8 - 6.1	600	17.2	15.3 - 19.1	100	3.4	2.6 - 4.5	2,400	64.3	61.8 - 66.7	3,700
\$25,000-\$49,999	300	7.3	6.1 - 8.7	100	3.0	2.2 - 4.1	600	15.1	13.4 - 16.9	200	3.8	3.0 - 4.9	2,800	70.8	68.6 - 73.0	4,000
>=\$50,000	300	4.4	3.7 - 5.3	100	1.9	1.4 - 2.5	500	8.6	7.5 - 9.8	200	4.0	3.2 - 4.9	4,700	81.1	79.5 - 82.7	5,800

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.0	7.1 - 11.5	100	4.3	3.0 - 6.2	200	16.0	13.4 - 19.0	<50	3.5	2.4 - 5.0	900	67.2	63.6 - 70.6	1,300
Normal birth weight	1,300	7.9	7.3 - 8.6	500	3.3	2.9 - 3.8	2,400	14.7	13.9 - 15.6	600	3.7	3.3 - 4.2	11,600	70.3	69.2 - 71.4	16,400

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	19.5	16.4 - 23.0	100	6.8	5.0 - 9.3	300	22.0	18.8 - 25.6	100	5.5	3.9 - 7.7	700	46.2	42.1 - 50.3	1,500
No	1,100	7.0	6.4 - 7.6	500	3.0	2.6 - 3.5	2,200	14.0	13.2 - 14.9	600	3.5	3.1 - 4.0	11,500	72.4	71.3 - 73.5	15,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	13.7	12.0 - 15.7	200	4.7	3.6 - 6.0	800	21.2	19.2 - 23.5	200	5.7	4.6 - 7.0	2,000	54.6	52.0 - 57.3	3,600
No	900	6.6	6.0 - 7.3	400	3.0	2.6 - 3.5	1,800	13.0	12.2 - 13.9	400	3.2	2.8 - 3.7	10,300	74.2	73.0 - 75.3	13,900

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	10.9	9.8 - 12.1	300	4.7	3.9 - 5.6	1,400	19.1	17.7 - 20.5	300	3.5	2.9 - 4.2	4,600	61.8	60.0 - 63.6	7,400
No	600	6.0	5.3 - 6.8	200	2.3	1.9 - 2.8	1,200	11.7	10.7 - 12.7	400	3.8	3.2 - 4.4	7,600	76.3	74.9 - 77.6	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,500	8.5	7.9 - 9.1	600	3.3	2.9 - 3.7	2,600	14.7	14.0 - 15.5	600	3.5	3.1 - 3.9	12,400	70.1	69.1 - 71.0	17,600

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	10.9	9.1 - 13.0	100	3.8	2.7 - 5.2	300	13.2	11.3 - 15.5	<50	1.8	1.1 - 2.8	1,600	70.3	67.4 - 73.1	2,200
Honolulu	1,100	8.4	7.7 - 9.1	400	3.1	2.7 - 3.6	1,900	14.8	13.9 - 15.7	500	3.7	3.3 - 4.2	8,800	69.9	68.7 - 71.0	12,700
Kauai	<50	5.8	3.8 - 8.8	<50	1.6	0.8 - 3.3	100	13.6	10.5 - 17.5	<50	4.2	2.6 - 6.8	600	74.7	70.1 - 78.9	800
Maui	100	7.1	5.5 - 9.1	100	4.3	3.1 - 5.8	300	16.4	14.1 - 19.0	100	3.4	2.4 - 4.8	1,300	68.8	65.7 - 71.8	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.6	5.5 - 7.8	100	2.4	1.8 - 3.1	500	11.6	10.2 - 13.1	200	4.4	3.6 - 5.4	3,000	75.1	73.1 - 77.0	4,100
Native Hawaiian	500	10.9	9.6 - 12.4	200	4.6	3.8 - 5.7	900	19.4	17.7 - 21.2	200	3.7	3.0 - 4.6	3,000	61.4	59.2 - 63.5	4,900
Chinese	<50	4.5	3.5 - 5.8	<50	2.2	1.5 - 3.2	100	8.4	6.9 - 10.2	<50	2.7	2.0 - 3.8	500	82.2	79.9 - 84.3	600
Filipino	300	8.1	6.9 - 9.5	100	3.9	3.1 - 4.9	600	17.9	16.2 - 19.8	100	4.1	3.3 - 5.1	2,100	65.9	63.7 - 68.1	3,200
Japanese	100	4.1	3.0 - 5.6	<50	1.8	1.1 - 2.8	200	8.0	6.5 - 9.9	<50	1.9	1.3 - 2.9	1,800	84.1	81.7 - 86.3	2,100
Other	300	11.7	10.0 - 13.7	100	2.8	2.0 - 3.9	400	14.0	12.2 - 16.1	100	2.3	1.6 - 3.2	1,900	69.1	66.4 - 71.7	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	11.0	8.7 - 13.8	100	5.9	4.2 - 8.1	300	23.4	20.2 - 27.0	<50	3.1	2.0 - 4.8	800	56.6	52.6 - 60.6	1,500
20-24	400	9.9	8.6 - 11.4	200	3.7	3.0 - 4.7	1,000	22.3	20.5 - 24.3	200	4.1	3.3 - 5.1	2,600	59.9	57.7 - 62.0	4,300
25-34	700	7.9	7.1 - 8.8	300	2.9	2.5 - 3.5	1,000	11.7	10.7 - 12.7	300	3.5	3.0 - 4.0	6,500	74.0	72.6 - 75.3	8,700
35+	200	6.8	5.6 - 8.1	100	2.4	1.7 - 3.2	300	8.9	7.6 - 10.4	100	2.8	2.1 - 3.7	2,500	79.2	77.1 - 81.1	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	800	8.3	7.5 - 9.1	300	3.2	2.7 - 3.8	1,400	15.4	14.3 - 16.5	400	4.1	3.6 - 4.7	6,300	69.0	67.6 - 70.4	9,200
Female	700	8.7	7.8 - 9.6	300	3.3	2.8 - 4.0	1,200	14.0	13.0 - 15.1	200	2.8	2.3 - 3.3	6,000	71.2	69.7 - 72.6	8,500

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	7.1	6.4 - 7.8	300	2.6	2.2 - 3.1	1,300	11.8	11.0 - 12.7	400	3.3	2.9 - 3.8	8,600	75.2	74.0 - 76.3	11,400
No	700	11.0	9.8 - 12.2	300	4.5	3.8 - 5.3	1,300	20.1	18.6 - 21.6	200	3.8	3.1 - 4.5	3,800	60.7	58.9 - 62.5	6,200

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	800	12.1	10.9 - 13.4	300	4.5	3.8 - 5.3	1,300	20.0	18.6 - 21.6	200	2.8	2.2 - 3.4	3,900	60.6	58.8 - 62.4	6,400
131-185%	200	9.8	8.2 - 11.8	100	2.6	1.8 - 3.8	300	12.7	10.8 - 14.8	100	3.6	2.6 - 4.9	1,600	71.3	68.5 - 73.9	2,300
186+%	400	4.7	4.1 - 5.4	200	2.2	1.8 - 2.8	800	10.6	9.6 - 11.6	300	4.1	3.6 - 4.8	6,000	78.4	77.1 - 79.7	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	11.9	10.2 - 13.8	200	5.8	4.6 - 7.2	700	22.3	20.0 - 24.7	100	3.2	2.4 - 4.3	1,700	56.9	54.1 - 59.6	2,900
\$10,000-\$24,999	400	11.2	9.8 - 12.8	200	4.4	3.5 - 5.4	700	17.8	16.1 - 19.7	100	3.0	2.3 - 3.9	2,400	63.7	61.4 - 65.9	3,800
\$25,000-\$49,999	300	7.3	6.2 - 8.6	100	2.3	1.7 - 3.1	600	15.1	13.6 - 16.8	200	4.0	3.3 - 5.0	3,000	71.2	69.2 - 73.2	4,200
>=\$50,000	300	5.1	4.3 - 6.0	100	1.8	1.4 - 2.4	500	8.1	7.1 - 9.2	200	3.7	3.1 - 4.5	4,500	81.3	79.8 - 82.7	5,600

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.1	7.0 - 11.7	100	4.6	3.2 - 6.6	200	15.5	12.7 - 18.6	<50	3.2	2.1 - 4.8	900	67.7	63.8 - 71.3	1,300
Normal birth weight	1,400	8.4	7.8 - 9.1	500	3.2	2.8 - 3.6	2,400	14.7	13.9 - 15.5	600	3.5	3.1 - 3.9	11,500	70.2	69.2 - 71.2	16,400

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	19.6	16.7 - 23.0	100	6.0	4.4 - 8.1	300	21.1	18.0 - 24.4	100	5.2	3.7 - 7.2	700	48.1	44.2 - 52.0	1,500
No	1,200	7.5	6.9 - 8.1	500	3.0	2.6 - 3.4	2,200	14.1	13.3 - 14.9	500	3.4	3.0 - 3.8	11,500	72.1	71.1 - 73.1	15,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	13.4	11.8 - 15.2	100	4.0	3.1 - 5.1	700	20.6	18.7 - 22.7	200	5.6	4.5 - 6.8	2,000	56.4	53.9 - 58.8	3,500
No	1,000	7.2	6.6 - 7.9	400	3.0	2.6 - 3.5	1,800	13.2	12.4 - 14.0	400	3.0	2.6 - 3.4	10,200	73.6	72.5 - 74.6	13,900

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	11.2	10.2 - 12.4	300	4.1	3.5 - 4.8	1,400	18.7	17.4 - 20.1	200	3.1	2.6 - 3.8	4,700	62.9	61.2 - 64.5	7,500
No	600	6.3	5.7 - 7.1	200	2.5	2.1 - 3.0	1,200	11.7	10.8 - 12.7	400	3.7	3.2 - 4.3	7,400	75.7	74.5 - 76.9	9,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,600	9.1	8.4 - 9.7	600	3.6	3.2 - 4.0	2,600	14.8	14.1 - 15.6	600	3.4	3.0 - 3.8	12,000	69.1	68.1 - 70.1	17,300

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	12.0	10.2 - 14.1	100	3.4	2.5 - 4.8	300	13.4	11.5 - 15.6	<50	1.9	1.3 - 2.9	1,500	69.2	66.4 - 71.9	2,100
Honolulu	1,100	9.0	8.3 - 9.8	400	3.6	3.1 - 4.1	1,900	14.8	13.9 - 15.8	500	3.6	3.2 - 4.1	8,600	69.0	67.8 - 70.2	12,500
Kauai	100	7.2	5.1 - 10.2	<50	2.2	1.3 - 3.9	100	12.3	9.5 - 15.8	<50	3.5	2.2 - 5.6	600	74.7	70.4 - 78.6	800
Maui	100	6.8	5.4 - 8.6	100	4.3	3.2 - 5.8	300	17.9	15.6 - 20.5	100	3.6	2.6 - 5.0	1,200	67.3	64.3 - 70.2	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.9	5.8 - 8.2	100	2.4	1.8 - 3.3	400	11.4	10.0 - 13.1	200	4.0	3.1 - 5.0	2,900	75.3	73.2 - 77.3	3,900
Native Hawaiian	600	11.9	10.6 - 13.3	200	5.0	4.2 - 6.0	900	19.7	18.1 - 21.4	200	3.4	2.7 - 4.2	2,900	60.1	58.0 - 62.1	4,800
Chinese	<50	4.7	3.3 - 6.7	<50	1.9	1.2 - 2.9	100	7.8	5.9 - 10.4	<50	2.3	1.6 - 3.5	500	83.2	80.1 - 85.9	600
Filipino	300	9.2	7.9 - 10.8	100	4.2	3.3 - 5.3	600	18.3	16.5 - 20.3	200	4.5	3.6 - 5.7	2,100	63.7	61.3 - 66.1	3,300
Japanese	100	5.0	3.7 - 6.7	<50	2.3	1.4 - 3.7	200	7.5	6.0 - 9.4	<50	1.9	1.2 - 3.0	1,800	83.3	80.7 - 85.6	2,100
Other	300	11.4	9.5 - 13.5	100	3.3	2.4 - 4.6	400	14.3	12.3 - 16.6	100	2.7	1.9 - 3.8	1,700	68.3	65.4 - 71.1	2,600

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	11.8	9.5 - 14.5	100	7.3	5.4 - 9.8	400	25.7	22.3 - 29.3	<50	3.1	2.1 - 4.7	800	52.1	48.1 - 56.0	1,500
20-24	500	10.8	9.4 - 12.2	200	3.9	3.1 - 4.8	1,000	22.4	20.6 - 24.3	100	3.5	2.8 - 4.4	2,500	59.5	57.2 - 61.7	4,300
25-34	700	8.5	7.6 - 9.4	300	3.4	2.9 - 4.0	1,000	11.4	10.4 - 12.4	300	3.6	3.0 - 4.2	6,200	73.2	71.8 - 74.6	8,500
35+	200	7.1	5.8 - 8.5	100	1.9	1.3 - 2.7	300	8.7	7.4 - 10.3	100	2.9	2.2 - 3.9	2,400	79.4	77.3 - 81.4	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	800	9.0	8.1 - 9.9	300	3.4	2.9 - 4.0	1,400	15.5	14.4 - 16.6	400	4.0	3.4 - 4.6	6,200	68.2	66.8 - 69.6	9,000
Female	800	9.2	8.3 - 10.1	300	3.8	3.2 - 4.4	1,200	14.1	13.0 - 15.3	200	2.8	2.3 - 3.3	5,800	70.2	68.7 - 71.6	8,300

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	7.4	6.7 - 8.1	300	2.6	2.3 - 3.1	1,300	11.5	10.6 - 12.4	400	3.3	2.9 - 3.9	8,500	75.2	74.0 - 76.3	11,300
No	700	12.2	11.0 - 13.5	300	5.3	4.5 - 6.2	1,300	21.2	19.6 - 22.8	200	3.5	2.9 - 4.2	3,500	57.8	55.9 - 59.7	6,000

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	900	13.0	11.8 - 14.3	300	5.0	4.2 - 5.8	1,400	20.6	19.2 - 22.1	200	2.7	2.2 - 3.4	3,900	58.7	56.9 - 60.5	6,600
131-185%	200	9.7	8.0 - 11.7	100	3.2	2.2 - 4.5	300	13.1	11.2 - 15.4	100	3.9	2.8 - 5.3	1,600	70.1	67.2 - 72.8	2,300
186+%	300	4.6	3.9 - 5.3	200	2.2	1.8 - 2.8	700	9.9	8.9 - 10.9	300	3.9	3.3 - 4.6	5,700	79.5	78.1 - 80.8	7,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.5	7.4 - 12.1	100	4.6	3.1 - 6.7	200	16.6	13.6 - 20.0	<50	3.5	2.2 - 5.3	800	65.9	61.9 - 69.7	1,300
Normal birth weight	1,400	9.0	8.4 - 9.7	600	3.5	3.1 - 3.9	2,400	14.7	13.9 - 15.5	500	3.4	3.0 - 3.8	11,100	69.4	68.3 - 70.4	16,000

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	23.7	20.5 - 27.2	100	6.8	5.1 - 9.0	300	19.2	16.3 - 22.5	100	3.8	2.6 - 5.5	700	46.5	42.6 - 50.4	1,500
No	1,200	7.7	7.1 - 8.3	500	3.2	2.8 - 3.7	2,200	14.4	13.6 - 15.2	500	3.4	3.0 - 3.8	11,100	71.3	70.2 - 72.3	15,600

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	15.1	13.3 - 17.0	200	5.0	4.0 - 6.2	700	20.3	18.3 - 22.4	200	4.9	4.0 - 6.1	1,900	54.7	52.2 - 57.1	3,500
No	1,000	7.4	6.8 - 8.1	400	3.2	2.7 - 3.6	1,800	13.3	12.5 - 14.2	400	3.0	2.7 - 3.5	9,900	73.1	71.9 - 74.1	13,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	12.1	11.0 - 13.3	300	4.5	3.9 - 5.3	1,400	19.3	18.0 - 20.7	200	3.0	2.5 - 3.6	4,500	61.1	59.4 - 62.7	7,400
No	600	6.7	6.0 - 7.5	300	2.7	2.3 - 3.3	1,100	11.5	10.6 - 12.4	400	3.7	3.2 - 4.3	7,300	75.4	74.2 - 76.7	9,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,600	9.7	9.0 - 10.4	600	3.6	3.2 - 4.1	2,500	14.6	13.8 - 15.4	600	3.3	2.9 - 3.7	11,700	68.8	67.7 - 69.8	16,900

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	12.0	10.3 - 13.9	100	3.5	2.6 - 4.7	300	15.2	13.3 - 17.4	<50	2.1	1.5 - 3.1	1,400	67.2	64.5 - 69.7	2,000
Honolulu	1,200	9.9	9.0 - 10.7	500	3.6	3.1 - 4.2	1,700	14.1	13.1 - 15.1	400	3.4	3.0 - 3.9	8,600	69.0	67.7 - 70.3	12,400
Kauai	100	7.3	5.3 - 10.0	<50	2.3	1.3 - 4.0	100	13.2	10.4 - 16.6	<50	3.2	1.9 - 5.1	600	74.0	69.8 - 77.7	800
Maui	100	6.9	5.5 - 8.5	100	4.3	3.3 - 5.7	300	18.2	15.9 - 20.6	100	3.6	2.6 - 4.9	1,200	67.1	64.2 - 69.8	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.8	5.6 - 8.2	100	2.3	1.6 - 3.2	400	11.3	9.8 - 13.1	200	4.3	3.4 - 5.5	2,800	75.3	73.0 - 77.4	3,800
Native Hawaiian	600	13.4	12.2 - 14.8	200	4.7	3.9 - 5.6	900	18.5	17.0 - 20.0	200	3.5	2.9 - 4.3	2,800	59.9	58.0 - 61.8	4,700
Chinese	<50	7.2	4.6 - 11.0	<50	2.2	1.2 - 4.2	100	8.4	6.0 - 11.8	<50	2.7	1.5 - 4.6	500	79.5	74.8 - 83.4	600
Filipino	300	9.2	7.8 - 10.9	200	4.9	3.9 - 6.1	600	18.5	16.6 - 20.6	100	3.7	2.9 - 4.8	2,100	63.6	61.1 - 66.1	3,300
Japanese	100	4.4	3.3 - 6.0	<50	2.1	1.2 - 3.4	200	7.6	6.0 - 9.6	<50	1.5	0.9 - 2.5	1,800	84.4	81.8 - 86.7	2,200
Other	300	13.1	10.9 - 15.6	100	3.7	2.6 - 5.3	400	15.0	12.8 - 17.6	100	2.2	1.5 - 3.3	1,600	66.0	62.7 - 69.1	2,400

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	13.3	10.8 - 16.3	100	6.7	5.0 - 9.1	400	26.6	23.2 - 30.3	<50	2.4	1.6 - 3.7	700	50.9	47.0 - 54.9	1,400
20-24	500	10.9	9.5 - 12.4	200	4.6	3.7 - 5.6	900	21.7	19.9 - 23.7	100	3.5	2.7 - 4.4	2,500	59.4	57.1 - 61.6	4,200
25-34	800	9.1	8.2 - 10.1	300	3.3	2.7 - 3.9	900	11.0	10.1 - 12.1	300	3.6	3.0 - 4.2	6,100	73.0	71.5 - 74.4	8,400
35+	200	7.9	6.5 - 9.5	100	1.8	1.2 - 2.7	300	9.0	7.6 - 10.6	100	2.6	1.8 - 3.6	2,300	78.8	76.5 - 80.9	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	10.1	9.1 - 11.1	300	3.8	3.2 - 4.4	1,300	15.0	13.9 - 16.2	300	3.8	3.2 - 4.4	5,900	67.4	65.9 - 68.8	8,800
Female	800	9.3	8.4 - 10.3	300	3.5	2.9 - 4.1	1,200	14.2	13.1 - 15.4	200	2.7	2.2 - 3.3	5,700	70.3	68.8 - 71.8	8,200

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	7.8	7.0 - 8.6	300	2.7	2.3 - 3.2	1,300	11.5	10.6 - 12.4	400	3.4	2.9 - 3.9	8,400	74.7	73.5 - 75.9	11,200
No	800	13.5	12.2 - 14.9	300	5.5	4.7 - 6.4	1,200	20.7	19.2 - 22.4	200	3.1	2.5 - 3.8	3,300	57.2	55.3 - 59.1	5,700

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	900	14.1	12.8 - 15.4	300	5.1	4.3 - 6.0	1,300	19.6	18.2 - 21.1	200	2.6	2.1 - 3.1	3,800	58.7	56.8 - 60.5	6,500
131-185%	200	10.0	8.3 - 12.1	100	3.5	2.5 - 4.8	300	13.5	11.5 - 15.7	100	3.6	2.6 - 5.0	1,700	69.4	66.5 - 72.3	2,400
186+%	300	4.2	3.6 - 5.0	100	1.9	1.5 - 2.5	700	9.8	8.8 - 10.9	300	4.0	3.4 - 4.7	5,400	80.1	78.6 - 81.4	6,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.1	7.1 - 11.7	100	5.0	3.4 - 7.4	200	16.5	13.3 - 20.1	<50	3.5	2.2 - 5.5	800	65.9	61.6 - 70.0	1,200
Normal birth weight	1,500	9.7	9.0 - 10.5	600	3.5	3.1 - 4.0	2,300	14.5	13.7 - 15.3	500	3.3	2.9 - 3.7	10,900	69.0	67.9 - 70.1	15,800

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	26.2	22.9 - 29.9	100	6.3	4.6 - 8.4	300	18.0	15.1 - 21.2	<50	3.4	2.3 - 4.9	700	46.1	42.3 - 50.1	1,400
No	1,300	8.2	7.6 - 8.9	500	3.3	2.9 - 3.8	2,200	14.2	13.4 - 15.1	500	3.3	2.9 - 3.7	10,800	71.0	69.9 - 72.0	15,300

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	16.0	14.2 - 18.0	200	5.3	4.3 - 6.6	600	18.6	16.7 - 20.7	200	4.9	4.0 - 6.1	1,900	55.1	52.6 - 57.6	3,400
No	1,100	8.0	7.3 - 8.7	400	3.0	2.6 - 3.5	1,800	13.5	12.6 - 14.4	400	2.9	2.5 - 3.3	9,600	72.6	71.4 - 73.7	13,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	13.2	12.0 - 14.5	300	4.8	4.1 - 5.7	1,400	19.1	17.8 - 20.5	200	2.8	2.4 - 3.4	4,300	60.0	58.3 - 61.7	7,200
No	700	7.0	6.3 - 7.8	200	2.6	2.2 - 3.1	1,100	11.3	10.4 - 12.3	300	3.5	3.0 - 4.1	7,200	75.6	74.2 - 76.9	9,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,700	10.2	9.5 - 10.9	600	3.9	3.5 - 4.4	2,400	14.4	13.7 - 15.2	500	3.2	2.8 - 3.6	11,300	68.3	67.2 - 69.3	16,500

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	13.4	11.7 - 15.3	100	3.7	2.8 - 4.8	300	14.1	12.5 - 16.0	<50	2.3	1.7 - 3.2	1,300	66.5	64.0 - 68.8	2,000
Honolulu	1,200	10.3	9.5 - 11.2	500	3.8	3.3 - 4.4	1,700	14.1	13.1 - 15.1	400	3.3	2.8 - 3.8	8,300	68.5	67.2 - 69.8	12,100
Kauai	100	6.9	5.1 - 9.3	<50	3.2	2.0 - 4.9	100	13.1	10.5 - 16.1	<50	3.3	2.2 - 5.1	500	73.6	69.8 - 77.0	700
Maui	100	7.0	5.7 - 8.4	100	5.2	4.1 - 6.6	300	18.0	15.9 - 20.2	100	3.3	2.5 - 4.4	1,100	66.6	63.9 - 69.1	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	6.8	5.6 - 8.3	100	2.2	1.6 - 3.1	400	11.0	9.5 - 12.8	100	3.7	2.9 - 4.9	2,700	76.2	73.9 - 78.3	3,600
Native Hawaiian	700	14.5	13.3 - 15.8	200	5.3	4.6 - 6.1	800	17.6	16.3 - 18.9	200	3.6	3.0 - 4.4	2,700	59.0	57.3 - 60.7	4,500
Chinese	<50	7.7	5.0 - 11.7	<50	2.4	1.3 - 4.4	100	9.2	6.3 - 13.3	<50	3.1	1.7 - 5.6	500	77.6	72.4 - 82.1	600
Filipino	300	9.4	8.0 - 11.1	200	5.4	4.3 - 6.6	600	18.6	16.7 - 20.8	100	3.5	2.7 - 4.6	2,100	63.1	60.5 - 65.6	3,300
Japanese	100	4.3	3.2 - 5.9	<50	1.9	1.1 - 3.2	200	8.4	6.7 - 10.4	<50	2.2	1.4 - 3.3	1,800	83.3	80.6 - 85.6	2,200
Other	300	14.2	11.9 - 16.8	100	4.2	3.0 - 5.9	300	14.8	12.5 - 17.4	<50	1.8	1.1 - 2.8	1,500	65.0	61.7 - 68.3	2,300

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	14.8	12.4 - 17.6	100	7.7	5.9 - 9.9	400	26.4	23.3 - 29.8	<50	2.3	1.5 - 3.4	700	48.8	45.2 - 52.5	1,400
20-24	500	11.5	10.2 - 13.0	200	5.2	4.3 - 6.3	900	20.9	19.2 - 22.8	100	3.1	2.5 - 4.0	2,400	59.3	57.1 - 61.4	4,100
25-34	800	9.6	8.7 - 10.6	300	3.3	2.8 - 3.9	900	10.9	9.9 - 11.9	300	3.5	2.9 - 4.1	5,900	72.7	71.3 - 74.2	8,200
35+	200	7.6	6.2 - 9.2	100	2.0	1.4 - 2.9	300	9.1	7.6 - 10.8	100	2.8	2.0 - 3.8	2,200	78.6	76.2 - 80.7	2,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	10.5	9.6 - 11.5	300	4.1	3.5 - 4.7	1,200	14.5	13.4 - 15.6	300	3.4	2.9 - 4.0	5,800	67.6	66.1 - 69.0	8,600
Female	800	9.8	8.9 - 10.8	300	3.8	3.2 - 4.4	1,100	14.4	13.3 - 15.6	200	2.9	2.4 - 3.5	5,500	69.1	67.6 - 70.5	7,900

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	7.7	7.0 - 8.5	300	2.8	2.4 - 3.3	1,300	11.8	10.9 - 12.7	400	3.4	2.9 - 3.9	8,200	74.3	73.1 - 75.5	11,000
No	800	15.1	13.8 - 16.5	300	6.2	5.3 - 7.2	1,100	19.8	18.3 - 21.3	200	2.8	2.3 - 3.3	3,100	56.2	54.3 - 58.0	5,500

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,000	14.8	13.6 - 16.2	300	5.3	4.5 - 6.1	1,200	18.6	17.2 - 20.0	200	2.5	2.1 - 3.1	3,800	58.7	57.0 - 60.5	6,400
131-185%	300	9.9	8.3 - 11.8	100	3.5	2.6 - 4.7	400	14.0	12.2 - 16.1	100	3.0	2.1 - 4.1	1,800	69.6	66.9 - 72.2	2,600
186+%	300	4.1	3.5 - 4.9	100	2.1	1.7 - 2.7	600	9.3	8.3 - 10.5	200	4.0	3.4 - 4.8	4,900	80.4	78.9 - 81.8	6,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.5	7.5 - 12.0	100	5.0	3.3 - 7.4	200	17.8	14.6 - 21.6	<50	3.4	2.1 - 5.5	700	64.3	60.0 - 68.4	1,100
Normal birth weight	1,600	10.2	9.5 - 10.9	600	3.9	3.4 - 4.3	2,200	14.2	13.4 - 15.0	500	3.1	2.8 - 3.6	10,600	68.6	67.5 - 69.6	15,400

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	26.6	23.3 - 30.1	100	6.1	4.6 - 8.1	300	18.6	15.8 - 21.7	<50	3.0	2.1 - 4.4	600	45.8	42.0 - 49.5	1,400
No	1,300	8.7	8.0 - 9.3	500	3.7	3.2 - 4.1	2,100	14.0	13.2 - 14.9	500	3.2	2.8 - 3.6	10,500	70.5	69.4 - 71.5	14,900

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	16.5	14.7 - 18.4	200	5.4	4.4 - 6.7	600	19.1	17.2 - 21.1	200	4.5	3.6 - 5.6	1,800	54.4	52.0 - 56.9	3,300
No	1,100	8.5	7.8 - 9.2	400	3.4	2.9 - 3.8	1,700	13.2	12.3 - 14.1	400	2.8	2.4 - 3.3	9,300	72.2	71.0 - 73.3	12,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	13.9	12.8 - 15.1	400	5.1	4.4 - 5.9	1,300	18.7	17.4 - 20.0	200	2.8	2.3 - 3.4	4,200	59.5	57.8 - 61.2	7,100
No	700	7.2	6.5 - 8.1	300	3.0	2.5 - 3.5	1,000	11.3	10.4 - 12.3	300	3.4	2.9 - 4.0	6,900	75.1	73.7 - 76.3	9,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,700	10.4	9.7 - 11.0	600	3.9	3.5 - 4.3	2,400	14.5	13.7 - 15.3	500	3.3	2.9 - 3.7	11,100	68.1	67.0 - 69.1	16,300

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	12.9	11.4 - 14.6	100	4.2	3.3 - 5.2	300	15.5	14.0 - 17.2	100	2.7	2.1 - 3.5	1,200	64.7	62.4 - 66.9	1,900
Honolulu	1,300	10.6	9.7 - 11.4	400	3.7	3.2 - 4.3	1,700	14.0	13.0 - 15.0	400	3.3	2.8 - 3.8	8,200	68.5	67.1 - 69.7	12,000
Kauai	<50	6.8	5.1 - 8.9	<50	3.8	2.5 - 5.7	100	13.4	11.0 - 16.2	<50	3.4	2.2 - 5.0	500	72.7	69.2 - 76.0	700
Maui	100	7.5	6.4 - 8.9	100	4.6	3.7 - 5.8	300	16.9	15.1 - 18.9	100	3.8	2.9 - 4.9	1,100	67.2	64.8 - 69.5	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	7.8	6.5 - 9.3	100	2.2	1.6 - 3.1	400	11.6	10.0 - 13.3	100	3.8	2.9 - 4.9	2,700	74.7	72.4 - 76.8	3,600
Native Hawaiian	600	14.6	13.5 - 15.7	200	5.0	4.4 - 5.7	800	18.4	17.3 - 19.6	200	3.5	3.0 - 4.1	2,600	58.6	57.1 - 60.0	4,400
Chinese	<50	7.4	4.6 - 11.7	<50	2.6	1.4 - 5.0	100	10.6	7.3 - 15.3	<50	2.9	1.5 - 5.7	500	76.4	70.6 - 81.3	600
Filipino	300	9.2	7.7 - 10.9	200	5.3	4.2 - 6.6	600	17.7	15.7 - 19.8	100	3.5	2.7 - 4.5	2,100	64.4	61.8 - 66.9	3,300
Japanese	100	4.8	3.5 - 6.4	<50	2.0	1.2 - 3.3	100	7.2	5.6 - 9.2	100	2.8	1.9 - 4.2	1,700	83.2	80.4 - 85.7	2,100
Other	300	14.1	11.7 - 16.8	100	4.4	3.1 - 6.1	300	14.6	12.2 - 17.3	<50	2.1	1.3 - 3.2	1,400	64.9	61.4 - 68.2	2,200

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	14.6	12.3 - 17.2	100	8.2	6.5 - 10.3	400	26.6	23.7 - 29.8	<50	2.3	1.6 - 3.4	700	48.3	44.9 - 51.7	1,500
20-24	500	11.8	10.5 - 13.2	200	5.1	4.2 - 6.1	800	20.2	18.5 - 22.1	100	3.0	2.4 - 3.8	2,400	59.9	57.8 - 62.0	4,000
25-34	800	9.6	8.7 - 10.6	200	3.1	2.6 - 3.7	900	11.1	10.2 - 12.2	300	3.5	3.0 - 4.2	5,800	72.6	71.2 - 74.1	8,000
35+	200	8.2	6.7 - 9.9	100	2.0	1.4 - 2.9	300	9.0	7.6 - 10.7	100	3.3	2.4 - 4.5	2,100	77.5	75.1 - 79.7	2,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	11.0	10.0 - 12.0	300	4.0	3.4 - 4.6	1,200	14.7	13.6 - 15.8	300	3.2	2.7 - 3.8	5,700	67.2	65.8 - 68.7	8,500
Female	800	9.7	8.8 - 10.7	300	3.8	3.2 - 4.4	1,100	14.2	13.2 - 15.4	300	3.3	2.8 - 3.9	5,400	69.0	67.5 - 70.4	7,800

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	8.2	7.5 - 9.0	300	2.6	2.2 - 3.1	1,300	12.0	11.1 - 12.9	400	3.5	3.0 - 4.0	8,100	73.8	72.5 - 75.0	11,000
No	800	14.9	13.6 - 16.3	300	6.5	5.7 - 7.5	1,000	19.7	18.3 - 21.2	100	2.8	2.3 - 3.3	2,900	56.0	54.2 - 57.8	5,200

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	900	14.5	13.3 - 15.7	300	5.1	4.4 - 6.0	1,200	18.7	17.4 - 20.0	200	2.5	2.1 - 3.0	3,900	59.2	57.5 - 60.9	6,500
131-185%	300	9.8	8.2 - 11.6	100	3.4	2.5 - 4.5	400	13.5	11.7 - 15.4	100	2.9	2.1 - 4.0	1,900	70.5	67.9 - 72.9	2,700
186+%	200	4.3	3.6 - 5.2	100	2.1	1.6 - 2.7	500	9.3	8.3 - 10.5	300	4.4	3.7 - 5.2	4,600	79.9	78.3 - 81.4	5,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	10.8	8.7 - 13.4	100	4.9	3.2 - 7.4	200	18.3	15.1 - 22.1	<50	3.2	2.0 - 5.2	700	62.7	58.3 - 66.9	1,000
Normal birth weight	1,600	10.3	9.6 - 11.0	600	3.8	3.4 - 4.3	2,200	14.2	13.4 - 15.0	500	3.2	2.9 - 3.7	10,400	68.5	67.4 - 69.5	15,200

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	27.7	24.5 - 31.1	100	5.6	4.2 - 7.4	300	18.6	15.9 - 21.7	<50	2.6	1.8 - 3.7	600	45.5	41.9 - 49.2	1,300
No	1,300	8.7	8.1 - 9.4	500	3.6	3.2 - 4.1	2,000	14.0	13.2 - 14.8	500	3.3	2.9 - 3.7	10,300	70.4	69.3 - 71.4	14,600

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	16.9	15.2 - 18.8	200	5.2	4.2 - 6.4	600	19.4	17.6 - 21.4	100	4.1	3.3 - 5.1	1,800	54.4	52.0 - 56.8	3,200
No	1,100	8.5	7.8 - 9.3	400	3.3	2.9 - 3.8	1,700	13.1	12.3 - 14.0	400	3.0	2.6 - 3.5	9,100	72.0	70.8 - 73.1	12,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped)- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	14.2	13.1 - 15.5	400	5.1	4.4 - 5.9	1,300	18.8	17.5 - 20.1	200	2.7	2.3 - 3.3	4,100	59.2	57.5 - 60.8	7,000
No	700	7.3	6.5 - 8.1	300	2.9	2.5 - 3.4	1,000	11.3	10.3 - 12.3	300	3.6	3.1 - 4.2	6,800	74.9	73.6 - 76.2	9,100