

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- for the State of Hawaii, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 3-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, income data does not exist for some 3-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Binge drinking 3 mos before pregnancy (2000-2008)	Binge drinking 3 mos before pregnancy (2000-2008)

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 9/30/15

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**


profiles@hhdw.org

(855) 946-5899 x15

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- for the State of Hawaii, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	
2000-2002	1,900	11.9	11.1 - 12.9	13,700	88.1	87.1 - 88.9	15,500
2001-2003	1,900	11.8	10.9 - 12.8	13,800	88.2	87.2 - 89.1	15,700
2002-2004	2,200	13.3	12.4 - 14.4	14,300	86.7	85.6 - 87.6	16,500
2003-2005	2,700	15.8	14.8 - 16.9	14,200	84.2	83.1 - 85.2	16,900
2004-2006	3,200	18.1	17.1 - 19.2	14,400	81.9	80.8 - 82.9	17,600
2005-2007	3,400	19.2	18.1 - 20.3	14,400	80.8	79.7 - 81.9	17,900
2006-2008	3,500	19.3	18.2 - 20.5	14,800	80.7	79.5 - 81.8	18,300



PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,500	19.3	18.2 - 20.5	14,800	80.7	79.5 - 81.8	18,300

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	18.7	15.8 - 22.0	1,900	81.3	78.0 - 84.2	2,300
Honolulu	2,500	19.0	17.7 - 20.3	10,500	81.0	79.7 - 82.3	13,000
Kauai	200	23.5	18.6 - 29.3	700	76.5	70.7 - 81.4	900
Maui	400	20.5	17.3 - 24.1	1,600	79.5	75.9 - 82.7	2,100

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,000	24.4	22.0 - 26.9	3,200	75.6	73.1 - 78.0	4,300
Native Hawaiian	1,200	23.5	21.1 - 26.0	3,800	76.5	74.0 - 78.9	5,000
Chinese	100	8.8	6.8 - 11.3	600	91.2	88.7 - 93.2	600
Filipino	400	13.0	11.1 - 15.1	3,000	87.0	84.9 - 88.9	3,400
Japanese	400	17.7	14.8 - 21.1	1,700	82.3	78.9 - 85.2	2,100
Other	500	15.7	13.3 - 18.4	2,500	84.3	81.6 - 86.7	2,900

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	17.4	13.8 - 21.7	1,300	82.6	78.3 - 86.2	1,500
20-24	1,200	26.8	24.3 - 29.5	3,200	73.2	70.5 - 75.7	4,300
25-34	1,700	18.7	17.3 - 20.3	7,400	81.3	79.7 - 82.7	9,100
35+	400	11.9	10.0 - 14.1	2,900	88.1	85.9 - 90.0	3,300

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,800	19.5	18.0 - 21.1	7,600	80.5	78.9 - 82.0	9,500
Female	1,700	19.1	17.6 - 20.8	7,100	80.9	79.2 - 82.4	8,800

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	16.7	15.5 - 18.0	9,600	83.3	82.0 - 84.5	11,500
No	1,600	23.7	21.8 - 25.8	5,200	76.3	74.2 - 78.2	6,800

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,200	18.1	16.4 - 20.1	5,500	81.9	79.9 - 83.6	6,700
131-185%	500	20.0	17.2 - 23.2	2,100	80.0	76.8 - 82.8	2,600
186+%	1,600	20.6	19.0 - 22.3	6,200	79.4	77.7 - 81.0	7,800

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	500	17.0	14.4 - 19.9	2,600	83.0	80.1 - 85.6	3,100
\$10,000-\$24,999	900	22.4	19.9 - 25.1	3,000	77.6	74.9 - 80.1	3,900
\$25,000-\$49,999	1,000	22.7	20.4 - 25.3	3,300	77.3	74.7 - 79.6	4,200
>=\$50,000	1,000	17.0	15.4 - 18.9	5,000	83.0	81.1 - 84.6	6,000

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	14.5	11.3 - 18.4	1,200	85.5	81.6 - 88.7	1,400
Normal birth weight	3,300	19.8	18.6 - 21.0	13,500	80.2	79.0 - 81.4	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	35.0	30.5 - 39.9	1,000	65.0	60.1 - 69.5	1,600
No	3,000	17.7	16.7 - 18.9	13,700	82.3	81.1 - 83.3	16,600

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	41.5	38.3 - 44.7	2,100	58.5	55.3 - 61.7	3,700
No	2,000	13.7	12.6 - 14.8	12,600	86.3	85.2 - 87.4	14,600

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	19.6	17.9 - 21.5	6,200	80.4	78.5 - 82.1	7,700
No	2,000	19.1	17.7 - 20.6	8,300	80.9	79.4 - 82.3	10,300

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,400	19.2	18.1 - 20.3	14,400	80.8	79.7 - 81.9	17,900

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	19.2	16.2 - 22.5	1,800	80.8	77.5 - 83.8	2,300
Honolulu	2,400	18.6	17.4 - 20.0	10,400	81.4	80.0 - 82.6	12,800
Kauai	200	20.8	16.0 - 26.5	700	79.2	73.5 - 84.0	900
Maui	400	22.1	18.7 - 26.0	1,500	77.9	74.0 - 81.3	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,000	24.2	21.7 - 26.7	3,100	75.8	73.3 - 78.3	4,100
Native Hawaiian	1,200	24.7	22.3 - 27.2	3,700	75.3	72.8 - 77.7	4,900
Chinese	<50	6.6	4.9 - 8.8	600	93.4	91.2 - 95.1	600
Filipino	400	11.9	10.1 - 14.0	2,900	88.1	86.0 - 89.9	3,300
Japanese	300	15.8	13.1 - 18.9	1,800	84.2	81.1 - 86.9	2,200
Other	500	16.4	13.9 - 19.2	2,300	83.6	80.8 - 86.1	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	18.4	14.8 - 22.8	1,300	81.6	77.2 - 85.2	1,600
20-24	1,200	26.7	24.2 - 29.3	3,200	73.3	70.7 - 75.8	4,400
25-34	1,600	18.1	16.7 - 19.7	7,200	81.9	80.3 - 83.3	8,800
35+	400	12.0	10.0 - 14.2	2,700	88.0	85.8 - 90.0	3,100

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,800	19.5	17.9 - 21.1	7,400	80.5	78.9 - 82.1	9,200
Female	1,600	18.9	17.4 - 20.5	7,100	81.1	79.5 - 82.6	8,700

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	15.9	14.7 - 17.3	9,500	84.1	82.7 - 85.3	11,300
No	1,600	24.7	22.7 - 26.9	5,000	75.3	73.1 - 77.3	6,600

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,200	18.5	16.7 - 20.5	5,200	81.5	79.5 - 83.3	6,400
131-185%	500	22.7	19.6 - 26.2	1,800	77.3	73.8 - 80.4	2,300
186+%	1,500	19.8	18.2 - 21.5	6,200	80.2	78.5 - 81.8	7,800

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	500	18.3	15.7 - 21.3	2,400	81.7	78.7 - 84.3	3,000
\$10,000-\$24,999	800	21.9	19.4 - 24.6	2,900	78.1	75.4 - 80.6	3,700
\$25,000-\$49,999	900	22.4	20.1 - 24.9	3,200	77.6	75.1 - 79.9	4,200
>=\$50,000	1,000	17.2	15.4 - 19.0	4,700	82.8	81.0 - 84.6	5,700

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	17.5	14.0 - 21.7	1,200	82.5	78.3 - 86.0	1,400
Normal birth weight	3,200	19.3	18.2 - 20.5	13,200	80.7	79.5 - 81.8	16,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	36.6	32.0 - 41.5	1,000	63.4	58.5 - 68.0	1,500
No	2,800	17.4	16.3 - 18.6	13,400	82.6	81.4 - 83.7	16,200

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	41.2	38.2 - 44.4	2,200	58.8	55.6 - 61.8	3,700
No	1,900	13.3	12.3 - 14.4	12,200	86.7	85.6 - 87.7	14,100

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	20.0	18.2 - 21.8	6,000	80.0	78.2 - 81.8	7,500
No	1,900	18.6	17.2 - 20.0	8,200	81.4	80.0 - 82.8	10,000

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,200	18.1	17.1 - 19.2	14,400	81.9	80.8 - 82.9	17,600

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	20.4	17.4 - 23.8	1,700	79.6	76.2 - 82.6	2,200
Honolulu	2,200	17.2	16.0 - 18.4	10,600	82.8	81.6 - 84.0	12,800
Kauai	100	18.1	13.5 - 23.9	600	81.9	76.1 - 86.5	800
Maui	400	21.7	18.3 - 25.5	1,400	78.3	74.5 - 81.7	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	21.9	19.6 - 24.3	3,100	78.1	75.7 - 80.4	4,000
Native Hawaiian	1,200	24.6	22.3 - 27.1	3,700	75.4	72.9 - 77.7	4,900
Chinese	<50	5.8	4.3 - 7.6	600	94.2	92.4 - 95.7	600
Filipino	400	11.0	9.3 - 13.0	2,900	89.0	87.0 - 90.7	3,200
Japanese	300	13.7	11.3 - 16.5	1,900	86.3	83.5 - 88.7	2,200
Other	400	15.7	13.3 - 18.5	2,200	84.3	81.5 - 86.7	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	18.5	14.8 - 22.8	1,200	81.5	77.2 - 85.2	1,400
20-24	1,100	25.2	22.8 - 27.7	3,200	74.8	72.3 - 77.2	4,300
25-34	1,500	17.2	15.8 - 18.7	7,200	82.8	81.3 - 84.2	8,600
35+	300	10.6	8.8 - 12.7	2,800	89.4	87.3 - 91.2	3,200

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,700	18.3	16.9 - 19.8	7,500	81.7	80.2 - 83.1	9,200
Female	1,500	17.9	16.4 - 19.4	6,900	82.1	80.6 - 83.6	8,400

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	14.8	13.7 - 16.0	9,900	85.2	84.0 - 86.3	11,600
No	1,500	24.4	22.4 - 26.6	4,500	75.6	73.4 - 77.6	6,000

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,100	18.1	16.3 - 20.0	5,200	81.9	80.0 - 83.7	6,300
131-185%	500	23.5	20.2 - 27.1	1,600	76.5	72.9 - 79.8	2,100
186+%	1,400	18.1	16.6 - 19.7	6,400	81.9	80.3 - 83.4	7,800

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	500	18.9	16.2 - 21.8	2,300	81.1	78.2 - 83.8	2,900
\$10,000-\$24,999	800	20.7	18.4 - 23.2	3,100	79.3	76.8 - 81.6	3,900
\$25,000-\$49,999	900	20.8	18.6 - 23.1	3,400	79.2	76.9 - 81.4	4,200
>=\$50,000	800	15.8	14.1 - 17.6	4,500	84.2	82.4 - 85.9	5,400

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	17.2	13.6 - 21.5	1,100	82.8	78.5 - 86.4	1,300
Normal birth weight	3,000	18.2	17.1 - 19.3	13,300	81.8	80.7 - 82.9	16,300

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	35.5	30.9 - 40.3	1,000	64.5	59.7 - 69.1	1,500
No	2,600	16.4	15.4 - 17.5	13,400	83.6	82.5 - 84.6	16,000

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	38.3	35.3 - 41.3	2,300	61.7	58.7 - 64.7	3,700
No	1,700	12.7	11.7 - 13.7	12,100	87.3	86.3 - 88.3	13,800

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	19.1	17.4 - 20.8	6,100	80.9	79.2 - 82.6	7,500
No	1,700	17.4	16.1 - 18.8	8,000	82.6	81.2 - 83.9	9,700

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	2,700	15.8	14.8 - 16.9	14,200	84.2	83.1 - 85.2	16,900

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	18.5	15.8 - 21.6	1,700	81.5	78.4 - 84.2	2,100
Honolulu	1,900	15.1	13.9 - 16.3	10,500	84.9	83.7 - 86.1	12,300
Kauai	100	13.8	10.1 - 18.6	700	86.2	81.4 - 89.9	800
Maui	300	19.0	16.1 - 22.3	1,400	81.0	77.7 - 83.9	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	20.3	17.9 - 22.9	2,900	79.7	77.1 - 82.1	3,700
Native Hawaiian	1,000	22.5	20.4 - 24.7	3,600	77.5	75.3 - 79.6	4,600
Chinese	<50	3.7	2.6 - 5.2	600	96.3	94.8 - 97.4	600
Filipino	300	10.2	8.4 - 12.3	3,000	89.8	87.7 - 91.6	3,400
Japanese	200	10.9	8.6 - 13.5	1,900	89.1	86.5 - 91.4	2,200
Other	300	12.0	9.7 - 14.7	2,200	88.0	85.3 - 90.3	2,400

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	15.4	12.2 - 19.2	1,200	84.6	80.8 - 87.8	1,400
20-24	900	22.5	20.2 - 25.0	3,200	77.5	75.0 - 79.8	4,100
25-34	1,200	14.8	13.4 - 16.3	7,200	85.2	83.7 - 86.6	8,400
35+	300	9.9	8.1 - 12.1	2,700	90.1	87.9 - 91.9	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,400	15.8	14.5 - 17.3	7,500	84.2	82.7 - 85.5	8,900
Female	1,300	15.8	14.4 - 17.4	6,700	84.2	82.6 - 85.6	8,000

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	12.4	11.3 - 13.6	9,900	87.6	86.4 - 88.7	11,300
No	1,300	22.7	20.8 - 24.8	4,300	77.3	75.2 - 79.2	5,600

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,100	16.4	14.8 - 18.2	5,500	83.6	81.8 - 85.2	6,500
131-185%	400	17.5	14.7 - 20.7	1,700	82.5	79.3 - 85.3	2,100
186+%	1,100	16.2	14.6 - 17.9	5,700	83.8	82.1 - 85.4	6,800

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	14.8	11.3 - 19.2	1,000	85.2	80.8 - 88.7	1,200
Normal birth weight	2,500	15.9	14.9 - 17.0	13,200	84.1	83.0 - 85.1	15,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	30.1	25.7 - 34.9	1,000	69.9	65.1 - 74.3	1,400
No	2,200	14.5	13.5 - 15.5	13,200	85.5	84.5 - 86.5	15,400

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	33.6	30.7 - 36.7	2,300	66.4	63.3 - 69.3	3,400
No	1,500	11.2	10.2 - 12.2	11,800	88.8	87.8 - 89.8	13,300

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	16.5	15.0 - 18.1	6,000	83.5	81.9 - 85.0	7,200
No	1,500	15.4	14.1 - 16.8	8,000	84.6	83.2 - 85.9	9,400

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	2,200	13.3	12.4 - 14.4	14,300	86.7	85.6 - 87.6	16,500

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	15.8	13.5 - 18.4	1,600	84.2	81.6 - 86.5	1,900
Honolulu	1,500	12.7	11.5 - 14.0	10,500	87.3	86.0 - 88.5	12,000
Kauai	100	13.5	10.2 - 17.6	600	86.5	82.4 - 89.8	700
Maui	300	14.8	12.6 - 17.4	1,500	85.2	82.6 - 87.4	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	17.1	14.7 - 19.9	2,900	82.9	80.1 - 85.3	3,600
Native Hawaiian	800	19.1	17.4 - 20.9	3,500	80.9	79.1 - 82.6	4,400
Chinese	<50	3.6	1.9 - 6.7	600	96.4	93.3 - 98.1	600
Filipino	300	7.7	6.0 - 9.7	3,100	92.3	90.3 - 94.0	3,400
Japanese	200	10.9	8.4 - 14.0	1,900	89.1	86.0 - 91.6	2,200
Other	200	10.1	7.6 - 13.2	2,100	89.9	86.8 - 92.4	2,400

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	10.8	8.3 - 13.8	1,200	89.2	86.2 - 91.7	1,400
20-24	700	18.9	16.6 - 21.3	3,200	81.1	78.7 - 83.4	3,900
25-34	1,100	13.4	12.0 - 14.9	7,100	86.6	85.1 - 88.0	8,200
35+	200	7.1	5.5 - 9.1	2,700	92.9	90.9 - 94.5	2,900

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,100	13.2	11.9 - 14.6	7,500	86.8	85.4 - 88.1	8,600
Female	1,100	13.5	12.1 - 15.0	6,800	86.5	85.0 - 87.9	7,900

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	10.9	9.8 - 12.1	10,100	89.1	87.9 - 90.2	11,300
No	1,000	18.6	16.8 - 20.6	4,200	81.4	79.4 - 83.2	5,200

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	900	13.8	12.3 - 15.5	5,600	86.2	84.5 - 87.7	6,500
131-185%	300	13.5	11.2 - 16.2	2,200	86.5	83.8 - 88.8	2,600
186+%	800	13.7	12.1 - 15.5	5,100	86.3	84.5 - 87.9	5,900

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	10.0	7.1 - 13.8	1,000	90.0	86.2 - 92.9	1,100
Normal birth weight	2,100	13.5	12.5 - 14.6	13,200	86.5	85.4 - 87.5	15,300

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	25.5	21.3 - 30.3	1,000	74.5	69.7 - 78.7	1,300
No	1,800	12.2	11.2 - 13.2	13,200	87.8	86.8 - 88.8	15,000

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	30.1	27.1 - 33.3	2,200	69.9	66.7 - 72.9	3,200
No	1,200	9.1	8.2 - 10.1	11,900	90.9	89.9 - 91.8	13,100

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	14.1	12.6 - 15.7	6,000	85.9	84.3 - 87.4	7,000
No	1,200	12.9	11.6 - 14.2	8,100	87.1	85.8 - 88.4	9,300

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,900	11.8	10.9 - 12.8	13,800	88.2	87.2 - 89.1	15,700

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	13.6	11.9 - 15.6	1,500	86.4	84.4 - 88.1	1,700
Honolulu	1,300	11.5	10.3 - 12.8	10,200	88.5	87.2 - 89.7	11,600
Kauai	100	11.2	8.7 - 14.4	600	88.8	85.6 - 91.3	700
Maui	200	12.5	10.7 - 14.6	1,500	87.5	85.4 - 89.3	1,700

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	500	15.5	13.1 - 18.4	2,800	84.5	81.6 - 86.9	3,300
Native Hawaiian	700	16.4	15.2 - 17.8	3,400	83.6	82.2 - 84.8	4,100
Chinese	<50	4.0	1.7 - 8.9	600	96.0	91.1 - 98.3	600
Filipino	200	5.9	4.4 - 7.9	3,200	94.1	92.1 - 95.6	3,400
Japanese	200	9.9	7.4 - 13.1	1,800	90.1	86.9 - 92.6	2,100
Other	200	10.7	8.0 - 14.1	2,000	89.3	85.9 - 92.0	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.6	6.8 - 10.8	1,300	91.4	89.2 - 93.2	1,400
20-24	600	17.1	14.9 - 19.4	3,100	82.9	80.6 - 85.1	3,700
25-34	900	11.7	10.4 - 13.2	6,900	88.3	86.8 - 89.6	7,900
35+	200	6.4	4.8 - 8.6	2,500	93.6	91.4 - 95.2	2,700

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	11.5	10.3 - 13.0	7,100	88.5	87.0 - 89.7	8,000
Female	900	12.1	10.8 - 13.5	6,800	87.9	86.5 - 89.2	7,700

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	10.0	8.9 - 11.2	9,600	90.0	88.8 - 91.1	10,700
No	800	15.7	14.1 - 17.5	4,300	84.3	82.5 - 85.9	5,000

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	800	13.0	11.5 - 14.6	5,400	87.0	85.4 - 88.5	6,300
131-185%	300	10.5	8.6 - 12.6	2,700	89.5	87.4 - 91.4	3,000
186+%	600	12.1	10.4 - 14.0	4,500	87.9	86.0 - 89.6	5,100

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Income (2000-2003)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	400	13.6	11.6 - 15.8	2,800	86.4	84.2 - 88.4	3,300
\$15,001-\$25,000	300	13.8	11.3 - 16.8	1,800	86.2	83.2 - 88.7	2,100
\$25,001-\$40,000	400	11.7	9.7 - 14.1	2,900	88.3	85.9 - 90.3	3,200
>=\$40,001	700	10.9	9.4 - 12.6	5,400	89.1	87.4 - 90.6	6,100

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.5	5.0 - 11.1	1,000	92.5	88.9 - 95.0	1,100
Normal birth weight	1,800	12.1	11.1 - 13.1	12,800	87.9	86.9 - 88.9	14,600

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	21.2	17.4 - 25.5	1,000	78.8	74.5 - 82.6	1,300
No	1,500	10.8	9.8 - 11.8	12,700	89.2	88.2 - 90.2	14,200

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	27.1	24.2 - 30.2	2,100	72.9	69.8 - 75.8	2,900
No	1,000	8.0	7.1 - 8.9	11,500	92.0	91.1 - 92.9	12,500

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	13.0	11.6 - 14.5	5,700	87.0	85.5 - 88.4	6,500
No	1,000	11.0	9.8 - 12.4	8,100	89.0	87.6 - 90.2	9,100

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,900	11.9	11.1 - 12.9	13,700	88.1	87.1 - 88.9	15,500

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	14.5	12.6 - 16.5	1,500	85.5	83.5 - 87.4	1,700
Honolulu	1,300	11.5	10.4 - 12.6	10,200	88.5	87.4 - 89.6	11,500
Kauai	100	13.6	10.8 - 17.1	600	86.4	82.9 - 89.2	600
Maui	200	12.0	10.2 - 14.1	1,400	88.0	85.9 - 89.8	1,600

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	500	14.9	12.8 - 17.4	2,900	85.1	82.6 - 87.2	3,400
Native Hawaiian	700	17.0	15.8 - 18.3	3,300	83.0	81.7 - 84.2	4,000
Chinese	<50	4.3	2.0 - 9.2	600	95.7	90.8 - 98.0	600
Filipino	200	5.1	3.8 - 6.8	3,100	94.9	93.2 - 96.2	3,200
Japanese	200	11.4	8.7 - 14.7	1,800	88.6	85.3 - 91.3	2,000
Other	200	10.7	8.1 - 14.0	1,900	89.3	86.0 - 91.9	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	10.8	8.7 - 13.3	1,300	89.2	86.7 - 91.3	1,500
20-24	600	17.2	15.2 - 19.4	3,100	82.8	80.6 - 84.8	3,700
25-34	900	11.7	10.4 - 13.1	6,700	88.3	86.9 - 89.6	7,600
35+	200	5.9	4.5 - 7.7	2,500	94.1	92.3 - 95.5	2,700

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,000	12.3	11.1 - 13.6	6,900	87.7	86.4 - 88.9	7,800
Female	900	11.6	10.3 - 12.9	6,800	88.4	87.1 - 89.7	7,700

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	10.4	9.4 - 11.5	9,600	89.6	88.5 - 90.6	10,800
No	700	15.5	13.9 - 17.2	4,000	84.5	82.8 - 86.1	4,800

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	800	12.5	11.2 - 13.9	5,300	87.5	86.1 - 88.8	6,100
131-185%	300	11.8	9.9 - 14.0	2,500	88.2	86.0 - 90.1	2,900
186+%	700	12.4	10.8 - 14.1	4,700	87.6	85.9 - 89.2	5,300

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Income (2000-2003)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	500	14.2	12.3 - 16.3	2,700	85.8	83.7 - 87.7	3,200
\$15,001-\$25,000	300	12.3	10.2 - 14.8	2,000	87.7	85.2 - 89.8	2,300
\$25,001-\$40,000	400	12.5	10.6 - 14.7	2,800	87.5	85.3 - 89.4	3,200
>=\$40,001	700	11.0	9.6 - 12.5	5,300	89.0	87.5 - 90.4	6,000

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	8.5	6.0 - 11.9	900	91.5	88.1 - 94.0	1,000
Normal birth weight	1,700	12.1	11.2 - 13.1	12,700	87.9	86.9 - 88.8	14,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	23.4	19.8 - 27.6	1,000	76.6	72.4 - 80.2	1,300
No	1,500	10.8	9.9 - 11.8	12,600	89.2	88.2 - 90.1	14,100

PRAMS Health Indicator 3-Year Aggregate Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	28.1	25.3 - 31.1	2,000	71.9	68.9 - 74.7	2,800
No	1,000	8.1	7.3 - 8.9	11,400	91.9	91.1 - 92.7	12,400

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	13.1	11.7 - 14.5	5,700	86.9	85.5 - 88.3	6,500
No	1,000	11.1	10.0 - 12.4	8,000	88.9	87.6 - 90.0	8,900