

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Drinking alcohol 3 months before pregnancy	Drinking alcohol 3 months before pregnancy

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/21/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15

**PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- for the State of Hawaii,
for the Year(s)**

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013


Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	7,000	41.5	40.4 - 42.5	9,800	58.5	57.5 - 59.6	16,800
2001-2005	7,200	42.3	41.2 - 43.3	9,800	57.7	56.7 - 58.8	16,900
2002-2006	7,500	43.4	42.3 - 44.5	9,800	56.6	55.5 - 57.7	17,300
2003-2007	7,900	44.9	43.8 - 46.0	9,700	55.1	54.0 - 56.2	17,700
2004-2008	8,300	46.5	45.5 - 47.5	9,600	53.5	52.5 - 54.5	17,900
2005-2009	8,700	48.0	46.9 - 49.1	9,400	52.0	50.9 - 53.1	18,100
2006-2010	9,000	49.2	48.0 - 50.4	9,300	50.8	49.6 - 52.0	18,300
2007-2011	9,300	50.3	49.1 - 51.6	9,100	49.7	48.4 - 50.9	18,400
2008-2012	9,400	51.5	50.1 - 52.9	8,900	48.5	47.1 - 49.9	18,300
2009-2013	9,500	52.3	50.8 - 53.7	8,700	47.7	46.3 - 49.2	18,200

**PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- for the State of Hawaii,
for the Year(s)**

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,500	52.3	50.8 - 53.7	8,700	47.7	46.3 - 49.2	18,200

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,300	53.5	51.2 - 55.9	1,100	46.5	44.1 - 48.8	2,300
Honolulu	6,700	51.3	49.3 - 53.2	6,400	48.7	46.8 - 50.7	13,100
Kauai	500	58.1	55.4 - 60.7	300	41.9	39.3 - 44.6	800
Maui	1,100	55.1	52.8 - 57.2	900	44.9	42.8 - 47.2	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	66.4	63.5 - 69.1	1,500	33.6	30.9 - 36.5	4,400
Native Hawaiian	3,000	54.9	52.3 - 57.5	2,400	45.1	42.5 - 47.7	5,400
Chinese	400	43.1	36.2 - 50.3	500	56.9	49.7 - 63.8	800
Filipino	1,300	40.3	36.9 - 43.7	1,900	59.7	56.3 - 63.1	3,200
Japanese	900	53.8	48.9 - 58.6	800	46.2	41.4 - 51.1	1,700
Other	1,100	39.6	35.8 - 43.6	1,600	60.4	56.4 - 64.2	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	27.8	23.2 - 33.0	900	72.2	67.0 - 76.8	1,200
20-24	2,300	54.9	51.8 - 58.0	1,900	45.1	42.0 - 48.2	4,100
25-34	5,300	55.5	53.4 - 57.5	4,300	44.5	42.5 - 46.6	9,600
35+	1,600	48.4	45.0 - 51.8	1,700	51.6	48.2 - 55.0	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,800	51.6	49.6 - 53.7	4,500	48.4	46.3 - 50.4	9,200
Female	4,800	52.9	50.8 - 55.0	4,200	47.1	45.0 - 49.2	9,000

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,700	50.9	49.0 - 52.8	5,500	49.1	47.2 - 51.0	11,200
No	3,800	54.5	52.2 - 56.8	3,200	45.5	43.2 - 47.8	7,000

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,900	45.0	42.6 - 47.4	3,600	55.0	52.6 - 57.4	6,500
131-185%	1,600	55.4	51.6 - 59.2	1,200	44.6	40.8 - 48.4	2,800
186+%	4,500	60.6	58.3 - 62.8	2,900	39.4	37.2 - 41.7	7,400

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	48.6	46.3 - 51.0	700	51.4	49.0 - 53.7	1,300
Normal birth weight	8,900	52.6	51.0 - 54.2	8,000	47.4	45.8 - 49.0	16,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	56.7	51.1 - 62.2	500	43.3	37.8 - 48.9	1,200
No	8,800	52.0	50.5 - 53.6	8,100	48.0	46.4 - 49.5	16,900

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	73.2	70.2 - 75.9	900	26.8	24.1 - 29.8	3,500
No	6,900	47.3	45.7 - 49.0	7,600	52.7	51.0 - 54.3	14,500

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,700	47.0	44.8 - 49.1	4,200	53.0	50.9 - 55.2	8,000
No	5,700	56.7	54.7 - 58.6	4,300	43.3	41.4 - 45.3	10,000

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,400	51.5	50.1 - 52.9	8,900	48.5	47.1 - 49.9	18,300

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	52.5	50.0 - 55.0	1,100	47.5	45.0 - 50.0	2,400
Honolulu	6,600	50.3	48.5 - 52.1	6,500	49.7	47.9 - 51.5	13,100
Kauai	500	58.7	55.4 - 61.9	300	41.3	38.1 - 44.6	800
Maui	1,100	55.0	52.5 - 57.3	900	45.0	42.7 - 47.5	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	67.2	64.5 - 69.7	1,400	32.8	30.3 - 35.5	4,400
Native Hawaiian	2,900	54.1	51.6 - 56.6	2,500	45.9	43.4 - 48.4	5,400
Chinese	300	43.0	36.5 - 49.7	400	57.0	50.3 - 63.5	800
Filipino	1,300	39.4	36.3 - 42.5	2,000	60.6	57.5 - 63.7	3,300
Japanese	900	52.1	47.6 - 56.5	800	47.9	43.5 - 52.4	1,700
Other	1,100	38.2	34.6 - 41.8	1,700	61.8	58.2 - 65.4	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	30.9	26.4 - 35.8	900	69.1	64.2 - 73.6	1,300
20-24	2,400	57.0	54.0 - 59.9	1,800	43.0	40.1 - 46.0	4,100
25-34	5,100	53.3	51.4 - 55.2	4,500	46.7	44.8 - 48.6	9,600
35+	1,500	47.4	44.3 - 50.6	1,700	52.6	49.4 - 55.7	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,800	51.3	49.4 - 53.2	4,600	48.7	46.8 - 50.6	9,400
Female	4,600	51.7	49.7 - 53.7	4,300	48.3	46.3 - 50.3	8,900

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,600	49.9	48.2 - 51.7	5,600	50.1	48.3 - 51.8	11,300
No	3,800	54.0	51.8 - 56.2	3,200	46.0	43.8 - 48.2	7,000

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,900	44.2	42.0 - 46.5	3,700	55.8	53.5 - 58.0	6,600
131-185%	1,500	53.2	49.6 - 56.7	1,300	46.8	43.3 - 50.4	2,800
186+%	4,500	60.0	57.9 - 62.0	3,000	40.0	38.0 - 42.1	7,500

**PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	47.1	44.3 - 49.9	700	52.9	50.1 - 55.7	1,300
Normal birth weight	8,800	51.9	50.4 - 53.3	8,200	48.1	46.7 - 49.6	17,000

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	57.6	52.5 - 62.5	500	42.4	37.5 - 47.5	1,300
No	8,600	51.1	49.7 - 52.5	8,200	48.9	47.5 - 50.3	16,800

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	73.7	70.9 - 76.3	900	26.3	23.7 - 29.1	3,500
No	6,700	46.2	44.7 - 47.7	7,800	53.8	52.3 - 55.3	14,600

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,700	46.1	44.0 - 48.2	4,300	53.9	51.8 - 56.0	7,900
No	5,600	55.8	53.9 - 57.6	4,500	44.2	42.4 - 46.1	10,100

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,300	50.3	49.1 - 51.6	9,100	49.7	48.4 - 50.9	18,400

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	51.4	48.8 - 54.0	1,200	48.6	46.0 - 51.2	2,400
Honolulu	6,400	49.0	47.3 - 50.7	6,700	51.0	49.3 - 52.7	13,000
Kauai	500	58.4	54.6 - 62.1	400	41.6	37.9 - 45.4	900
Maui	1,100	54.2	51.6 - 56.8	900	45.8	43.2 - 48.4	2,100

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	67.2	64.7 - 69.7	1,400	32.8	30.3 - 35.3	4,400
Native Hawaiian	2,800	52.3	49.9 - 54.8	2,500	47.7	45.2 - 50.1	5,300
Chinese	300	40.1	34.2 - 46.4	400	59.9	53.6 - 65.8	700
Filipino	1,300	38.9	36.0 - 41.8	2,000	61.1	58.2 - 64.0	3,300
Japanese	900	50.3	46.3 - 54.3	900	49.7	45.7 - 53.7	1,900
Other	1,000	36.4	33.2 - 39.8	1,800	63.6	60.2 - 66.8	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	32.0	27.8 - 36.4	1,000	68.0	63.6 - 72.2	1,400
20-24	2,400	55.4	52.6 - 58.1	1,900	44.6	41.9 - 47.4	4,300
25-34	4,900	52.2	50.4 - 54.0	4,500	47.8	46.0 - 49.6	9,400
35+	1,500	46.2	43.2 - 49.2	1,700	53.8	50.8 - 56.8	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,800	50.1	48.3 - 51.9	4,800	49.9	48.1 - 51.7	9,500
Female	4,500	50.6	48.8 - 52.5	4,400	49.4	47.5 - 51.2	8,900

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,600	49.6	48.0 - 51.2	5,700	50.4	48.8 - 52.0	11,400
No	3,600	51.6	49.4 - 53.7	3,400	48.4	46.3 - 50.6	7,000

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,800	42.1	40.0 - 44.2	3,800	57.9	55.8 - 60.0	6,600
131-185%	1,500	53.1	49.8 - 56.5	1,300	46.9	43.5 - 50.2	2,800
186+%	4,600	59.7	57.8 - 61.7	3,100	40.3	38.3 - 42.2	7,700

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	46.6	43.5 - 49.7	700	53.4	50.3 - 56.5	1,400
Normal birth weight	8,600	50.7	49.3 - 52.0	8,400	49.3	48.0 - 50.7	17,000

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	54.8	50.2 - 59.4	600	45.2	40.6 - 49.8	1,400
No	8,400	50.0	48.7 - 51.3	8,400	50.0	48.7 - 51.3	16,800

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	71.2	68.5 - 73.7	1,100	28.8	26.3 - 31.5	3,700
No	6,500	45.1	43.6 - 46.5	8,000	54.9	53.5 - 56.4	14,500

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,500	45.2	43.2 - 47.1	4,300	54.8	52.9 - 56.8	7,800
No	5,600	54.3	52.6 - 56.1	4,700	45.7	43.9 - 47.4	10,200

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
>=\$50,000	3,500	57.4	55.2 - 59.6	2,600	42.6	40.4 - 44.8	6,100
\$25,000-\$49,999	2,400	55.7	53.0 - 58.3	1,900	44.3	41.7 - 47.0	4,200
\$10,000-\$24,999	1,800	49.4	46.6 - 52.3	1,800	50.6	47.7 - 53.4	3,700
<\$10,000	1,200	38.5	35.5 - 41.5	2,000	61.5	58.5 - 64.5	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,000	49.2	48.0 - 50.4	9,300	50.8	49.6 - 52.0	18,300

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	51.5	48.7 - 54.3	1,100	48.5	45.7 - 51.3	2,400
Honolulu	6,200	47.7	46.2 - 49.3	6,800	52.3	50.7 - 53.8	13,000
Kauai	500	57.3	53.0 - 61.4	400	42.7	38.6 - 47.0	900
Maui	1,100	52.6	49.8 - 55.5	1,000	47.4	44.5 - 50.2	2,100

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	66.2	63.8 - 68.6	1,500	33.8	31.4 - 36.2	4,300
Native Hawaiian	2,700	51.7	49.3 - 54.0	2,600	48.3	46.0 - 50.7	5,300
Chinese	300	40.4	35.0 - 46.1	400	59.6	53.9 - 65.0	700
Filipino	1,200	36.2	33.6 - 38.9	2,100	63.8	61.1 - 66.4	3,300
Japanese	900	48.6	44.9 - 52.3	1,000	51.4	47.7 - 55.1	1,900
Other	1,000	36.2	33.1 - 39.3	1,800	63.8	60.7 - 66.9	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	33.5	29.4 - 37.8	1,000	66.5	62.2 - 70.6	1,500
20-24	2,400	54.6	52.0 - 57.1	2,000	45.4	42.9 - 48.0	4,400
25-34	4,700	50.8	49.1 - 52.5	4,500	49.2	47.5 - 50.9	9,200
35+	1,500	45.0	42.2 - 47.8	1,800	55.0	52.2 - 57.8	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,700	48.9	47.2 - 50.6	4,900	51.1	49.4 - 52.8	9,500
Female	4,400	49.5	47.8 - 51.3	4,500	50.5	48.7 - 52.2	8,800

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,600	49.0	47.5 - 50.5	5,800	51.0	49.5 - 52.5	11,400
No	3,500	49.6	47.5 - 51.6	3,500	50.4	48.4 - 52.5	7,000

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,700	41.2	39.2 - 43.3	3,900	58.8	56.7 - 60.8	6,600
131-185%	1,400	52.4	49.2 - 55.6	1,300	47.6	44.4 - 50.8	2,700
186+%	4,500	58.1	56.2 - 59.9	3,200	41.9	40.1 - 43.8	7,700

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	43.7	40.4 - 47.0	800	56.3	53.0 - 59.6	1,400
Normal birth weight	8,400	49.7	48.4 - 50.9	8,500	50.3	49.1 - 51.6	16,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	54.8	50.5 - 59.1	700	45.2	40.9 - 49.5	1,600
No	8,100	48.7	47.4 - 50.0	8,500	51.3	50.0 - 52.6	16,600

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	69.9	67.3 - 72.4	1,100	30.1	27.6 - 32.7	3,800
No	6,300	43.8	42.5 - 45.2	8,100	56.2	54.8 - 57.5	14,400

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,400	44.3	42.5 - 46.2	4,300	55.7	53.8 - 57.5	7,700
No	5,400	52.9	51.3 - 54.5	4,800	47.1	45.5 - 48.7	10,300

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
>=\$50,000	3,400	56.2	54.1 - 58.2	2,700	43.8	41.8 - 45.9	6,100
\$25,000-\$49,999	2,300	54.6	52.1 - 57.1	1,900	45.4	42.9 - 47.9	4,200
\$10,000-\$24,999	1,800	48.8	46.2 - 51.5	1,900	51.2	48.5 - 53.8	3,700
<\$10,000	1,200	36.6	33.8 - 39.6	2,000	63.4	60.4 - 66.2	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,700	48.0	46.9 - 49.1	9,400	52.0	50.9 - 53.1	18,100

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	49.8	46.9 - 52.7	1,200	50.2	47.3 - 53.1	2,300
Honolulu	6,000	46.6	45.2 - 48.0	6,900	53.4	52.0 - 54.8	12,900
Kauai	500	54.7	50.1 - 59.2	400	45.3	40.8 - 49.9	900
Maui	1,000	52.0	48.9 - 55.0	1,000	48.0	45.0 - 51.1	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,800	66.4	64.1 - 68.6	1,400	33.6	31.4 - 35.9	4,200
Native Hawaiian	2,600	49.7	47.4 - 52.0	2,600	50.3	48.0 - 52.6	5,200
Chinese	200	36.9	32.5 - 41.6	400	63.1	58.4 - 67.5	700
Filipino	1,100	34.4	32.1 - 36.9	2,200	65.6	63.1 - 67.9	3,300
Japanese	1,000	47.2	43.9 - 50.6	1,100	52.8	49.4 - 56.1	2,000
Other	1,000	36.5	33.7 - 39.4	1,700	63.5	60.6 - 66.3	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	35.6	31.7 - 39.8	1,000	64.4	60.2 - 68.3	1,600
20-24	2,300	53.5	51.1 - 55.9	2,000	46.5	44.1 - 48.9	4,300
25-34	4,400	49.1	47.5 - 50.7	4,600	50.9	49.3 - 52.5	9,000
35+	1,400	43.7	41.2 - 46.3	1,800	56.3	53.7 - 58.8	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,500	47.6	46.0 - 49.2	4,900	52.4	50.8 - 54.0	9,400
Female	4,200	48.4	46.8 - 50.1	4,500	51.6	49.9 - 53.2	8,700

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,400	47.6	46.2 - 49.0	5,900	52.4	51.0 - 53.8	11,300
No	3,300	48.7	46.8 - 50.7	3,500	51.3	49.3 - 53.2	6,800

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,600	39.7	37.8 - 41.6	3,900	60.3	58.4 - 62.2	6,500
131-185%	1,200	50.2	47.1 - 53.4	1,200	49.8	46.6 - 52.9	2,500
186+%	4,400	57.4	55.7 - 59.0	3,300	42.6	41.0 - 44.3	7,700

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	43.4	39.8 - 47.0	800	56.6	53.0 - 60.2	1,400
Normal birth weight	8,100	48.4	47.2 - 49.6	8,600	51.6	50.4 - 52.8	16,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	55.0	51.0 - 59.0	700	45.0	41.0 - 49.0	1,600
No	7,700	47.3	46.2 - 48.5	8,600	52.7	51.5 - 53.8	16,300

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	68.9	66.4 - 71.2	1,200	31.1	28.8 - 33.6	3,700
No	6,000	42.6	41.3 - 43.8	8,200	57.4	56.2 - 58.7	14,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,200	42.3	40.5 - 44.1	4,400	57.7	55.9 - 59.5	7,600
No	5,300	52.4	50.9 - 53.8	4,800	47.6	46.2 - 49.1	10,200

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
>=\$50,000	3,300	55.3	53.3 - 57.2	2,600	44.7	42.8 - 46.7	5,900
\$25,000-\$49,999	2,200	53.7	51.3 - 56.0	1,900	46.3	44.0 - 48.7	4,100
\$10,000-\$24,999	1,700	45.6	43.1 - 48.1	2,100	54.4	51.9 - 56.9	3,800
<\$10,000	1,100	36.7	33.9 - 39.6	1,900	63.3	60.4 - 66.1	3,100

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,300	46.5	45.5 - 47.5	9,600	53.5	52.5 - 54.5	17,900

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	47.5	44.5 - 50.5	1,200	52.5	49.5 - 55.5	2,300
Honolulu	5,800	45.2	44.0 - 46.4	7,000	54.8	53.6 - 56.0	12,800
Kauai	400	52.3	47.5 - 57.2	400	47.7	42.8 - 52.5	900
Maui	1,000	51.1	47.9 - 54.4	900	48.9	45.6 - 52.1	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	65.8	63.7 - 67.8	1,400	34.2	32.2 - 36.3	4,100
Native Hawaiian	2,400	48.3	46.2 - 50.5	2,600	51.7	49.5 - 53.8	4,900
Chinese	200	32.9	30.3 - 35.6	400	67.1	64.4 - 69.7	600
Filipino	1,100	32.7	30.6 - 34.8	2,200	67.3	65.2 - 69.4	3,300
Japanese	1,000	47.1	44.1 - 50.1	1,100	52.9	49.9 - 55.9	2,200
Other	900	34.1	31.6 - 36.7	1,800	65.9	63.3 - 68.4	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	34.0	30.3 - 37.9	1,000	66.0	62.1 - 69.7	1,500
20-24	2,300	52.0	49.8 - 54.2	2,100	48.0	45.8 - 50.2	4,300
25-34	4,200	47.1	45.7 - 48.6	4,700	52.9	51.4 - 54.3	8,900
35+	1,400	43.1	40.8 - 45.4	1,800	56.9	54.6 - 59.2	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,300	46.3	44.8 - 47.7	5,000	53.7	52.3 - 55.2	9,300
Female	4,000	46.8	45.3 - 48.3	4,600	53.2	51.7 - 54.7	8,600

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,300	46.1	44.9 - 47.4	6,200	53.9	52.6 - 55.1	11,500
No	3,000	47.1	45.3 - 49.0	3,400	52.9	51.0 - 54.7	6,400

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,400	37.3	35.6 - 39.1	4,100	62.7	60.9 - 64.4	6,500
131-185%	1,100	48.6	45.6 - 51.5	1,200	51.4	48.5 - 54.4	2,300
186+%	4,400	56.4	54.9 - 57.9	3,400	43.6	42.1 - 45.1	7,700

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	41.7	37.9 - 45.6	800	58.3	54.4 - 62.1	1,300
Normal birth weight	7,800	46.9	45.8 - 48.0	8,800	53.1	52.0 - 54.2	16,600

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	55.3	51.5 - 59.1	700	44.7	40.9 - 48.5	1,500
No	7,400	45.7	44.6 - 46.7	8,900	54.3	53.3 - 55.4	16,300

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	68.1	65.7 - 70.3	1,200	31.9	29.7 - 34.3	3,600
No	5,800	41.0	39.9 - 42.1	8,400	59.0	57.9 - 60.1	14,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,000	39.9	38.3 - 41.6	4,600	60.1	58.4 - 61.7	7,600
No	5,200	51.6	50.3 - 53.0	4,800	48.4	47.0 - 49.7	10,000

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
≥\$50,000	3,100	54.9	53.1 - 56.6	2,500	45.1	43.4 - 46.9	5,600
\$25,000-\$49,999	2,200	51.8	49.6 - 53.9	2,000	48.2	46.1 - 50.4	4,200
\$10,000-\$24,999	1,700	42.6	40.4 - 44.9	2,200	57.4	55.1 - 59.6	3,900
<\$10,000	1,000	35.2	32.7 - 37.9	1,900	64.8	62.1 - 67.3	3,000

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	7,900	44.9	43.8 - 46.0	9,700	55.1	54.0 - 56.2	17,700

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	46.4	43.5 - 49.4	1,200	53.6	50.6 - 56.5	2,200
Honolulu	5,600	43.8	42.5 - 45.1	7,200	56.2	54.9 - 57.5	12,700
Kauai	400	48.6	44.0 - 53.3	400	51.4	46.7 - 56.0	900
Maui	900	48.8	45.8 - 51.9	1,000	51.2	48.1 - 54.2	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,600	64.6	62.3 - 66.8	1,400	35.4	33.2 - 37.7	4,000
Native Hawaiian	2,300	47.7	45.7 - 49.8	2,500	52.3	50.2 - 54.3	4,800
Chinese	200	31.0	27.2 - 34.9	400	69.0	65.1 - 72.8	600
Filipino	1,000	30.3	28.1 - 32.6	2,400	69.7	67.4 - 71.9	3,400
Japanese	900	43.4	40.3 - 46.5	1,200	56.6	53.5 - 59.7	2,200
Other	900	33.8	31.0 - 36.6	1,800	66.2	63.4 - 69.0	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	32.1	28.6 - 35.8	1,000	67.9	64.2 - 71.4	1,500
20-24	2,100	49.0	46.8 - 51.2	2,200	51.0	48.8 - 53.2	4,300
25-34	4,000	46.3	44.8 - 47.8	4,700	53.7	52.2 - 55.2	8,700
35+	1,300	41.5	39.0 - 43.9	1,800	58.5	56.1 - 61.0	3,100

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,100	44.6	43.2 - 46.1	5,100	55.4	53.9 - 56.8	9,200
Female	3,800	45.2	43.6 - 46.7	4,600	54.8	53.3 - 56.4	8,400

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,100	44.6	43.3 - 45.9	6,400	55.4	54.1 - 56.7	11,500
No	2,800	45.5	43.6 - 47.3	3,400	54.5	52.7 - 56.4	6,200

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,400	36.4	34.6 - 38.1	4,200	63.6	61.9 - 65.4	6,700
131-185%	1,100	47.3	44.2 - 50.3	1,200	52.7	49.7 - 55.8	2,300
186+%	4,000	55.0	53.4 - 56.7	3,300	45.0	43.3 - 46.6	7,300

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	39.4	35.6 - 43.4	800	60.6	56.6 - 64.4	1,300
Normal birth weight	7,400	45.3	44.2 - 46.4	8,900	54.7	53.6 - 55.8	16,300

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	53.4	49.6 - 57.2	700	46.6	42.8 - 50.4	1,500
No	7,100	44.1	43.0 - 45.2	9,000	55.9	54.8 - 57.0	16,000

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	65.5	63.2 - 67.8	1,300	34.5	32.2 - 36.8	3,600
No	5,500	39.4	38.3 - 40.6	8,400	60.6	59.4 - 61.7	13,900

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	39.3	37.6 - 40.9	4,600	60.7	59.1 - 62.4	7,500
No	4,900	49.3	47.9 - 50.7	5,000	50.7	49.3 - 52.1	9,900

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	7,500	43.4	42.3 - 44.5	9,800	56.6	55.5 - 57.7	17,300

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	900	45.2	42.4 - 47.9	1,100	54.8	52.1 - 57.6	2,100
Honolulu	5,400	42.5	41.1 - 43.8	7,300	57.5	56.2 - 58.9	12,600
Kauai	400	45.2	40.8 - 49.7	400	54.8	50.3 - 59.2	800
Maui	800	46.8	43.9 - 49.7	1,000	53.2	50.3 - 56.1	1,800

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,400	61.9	59.4 - 64.2	1,500	38.1	35.8 - 40.6	3,800
Native Hawaiian	2,200	47.1	45.2 - 49.0	2,500	52.9	51.0 - 54.8	4,700
Chinese	200	30.2	25.7 - 35.0	400	69.8	65.0 - 74.3	600
Filipino	1,000	28.5	26.3 - 30.9	2,400	71.5	69.1 - 73.7	3,400
Japanese	900	41.1	37.9 - 44.3	1,300	58.9	55.7 - 62.1	2,200
Other	800	33.3	30.3 - 36.3	1,700	66.7	63.7 - 69.7	2,500

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	31.8	28.4 - 35.4	1,000	68.2	64.6 - 71.6	1,500
20-24	2,000	48.1	45.8 - 50.3	2,200	51.9	49.7 - 54.2	4,200
25-34	3,800	44.6	43.1 - 46.2	4,700	55.4	53.8 - 56.9	8,600
35+	1,200	38.7	36.2 - 41.3	1,900	61.3	58.7 - 63.8	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,900	43.6	42.1 - 45.2	5,100	56.4	54.8 - 57.9	9,000
Female	3,600	43.1	41.5 - 44.7	4,700	56.9	55.3 - 58.5	8,300

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,900	42.7	41.3 - 44.0	6,600	57.3	56.0 - 58.7	11,500
No	2,600	44.7	42.8 - 46.6	3,200	55.3	53.4 - 57.2	5,800

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,400	36.7	35.0 - 38.5	4,200	63.3	61.5 - 65.0	6,600
131-185%	1,100	44.6	41.5 - 47.7	1,300	55.4	52.3 - 58.5	2,400
186+%	3,600	52.2	50.5 - 54.0	3,300	47.8	46.0 - 49.5	6,800

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	38.2	34.1 - 42.4	800	61.8	57.6 - 65.9	1,200
Normal birth weight	7,000	43.7	42.6 - 44.9	9,000	56.3	55.1 - 57.4	16,100

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	52.5	48.6 - 56.3	700	47.5	43.7 - 51.4	1,500
No	6,700	42.5	41.3 - 43.6	9,000	57.5	56.4 - 58.7	15,700

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	64.4	62.0 - 66.8	1,300	35.6	33.2 - 38.0	3,600
No	5,100	37.7	36.5 - 38.9	8,500	62.3	61.1 - 63.5	13,600

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	38.6	36.9 - 40.3	4,500	61.4	59.7 - 63.1	7,400
No	4,600	47.1	45.6 - 48.6	5,100	52.9	51.4 - 54.4	9,700

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	7,200	42.3	41.2 - 43.3	9,800	57.7	56.7 - 58.8	16,900

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	900	44.3	41.8 - 46.9	1,100	55.7	53.1 - 58.2	2,000
Honolulu	5,200	41.5	40.2 - 42.9	7,300	58.5	57.1 - 59.8	12,400
Kauai	300	42.6	38.6 - 46.7	400	57.4	53.3 - 61.4	800
Maui	800	44.8	42.1 - 47.4	1,000	55.2	52.6 - 57.9	1,700

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,200	60.8	58.3 - 63.2	1,400	39.2	36.8 - 41.7	3,700
Native Hawaiian	2,100	45.3	43.6 - 47.0	2,500	54.7	53.0 - 56.4	4,600
Chinese	200	28.4	23.7 - 33.7	500	71.6	66.3 - 76.3	700
Filipino	900	27.9	25.6 - 30.3	2,400	72.1	69.7 - 74.4	3,400
Japanese	900	41.4	38.1 - 44.7	1,300	58.6	55.3 - 61.9	2,200
Other	800	33.1	30.1 - 36.2	1,600	66.9	63.8 - 69.9	2,400

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	33.2	30.0 - 36.6	1,000	66.8	63.4 - 70.0	1,500
20-24	1,900	46.2	44.0 - 48.4	2,300	53.8	51.6 - 56.0	4,200
25-34	3,600	43.5	41.9 - 45.0	4,700	56.5	55.0 - 58.1	8,400
35+	1,100	37.8	35.2 - 40.4	1,800	62.2	59.6 - 64.8	2,900

**PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,700	42.5	41.0 - 44.0	5,000	57.5	56.0 - 59.0	8,800
Female	3,400	42.0	40.5 - 43.6	4,700	58.0	56.4 - 59.5	8,200

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,700	41.3	40.0 - 42.7	6,700	58.7	57.3 - 60.0	11,300
No	2,500	44.1	42.3 - 46.0	3,100	55.9	54.0 - 57.7	5,600

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,400	36.1	34.4 - 37.7	4,200	63.9	62.3 - 65.6	6,600
131-185%	1,100	41.8	39.0 - 44.7	1,500	58.2	55.3 - 61.0	2,600
186+%	3,300	52.1	50.2 - 53.9	3,000	47.9	46.1 - 49.8	6,300

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	36.3	32.2 - 40.6	700	63.7	59.4 - 67.8	1,100
Normal birth weight	6,700	42.7	41.5 - 43.8	9,000	57.3	56.2 - 58.5	15,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	53.0	49.3 - 56.6	700	47.0	43.4 - 50.7	1,400
No	6,300	41.2	40.1 - 42.3	9,000	58.8	57.7 - 59.9	15,300

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	63.3	60.9 - 65.6	1,300	36.7	34.4 - 39.1	3,400
No	4,800	36.6	35.4 - 37.8	8,400	63.4	62.2 - 64.6	13,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	37.5	35.9 - 39.2	4,500	62.5	60.8 - 64.1	7,200
No	4,400	45.9	44.5 - 47.4	5,200	54.1	52.6 - 55.5	9,500

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	7,000	41.5	40.4 - 42.5	9,800	58.5	57.5 - 59.6	16,800

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	900	44.6	42.3 - 46.9	1,100	55.4	53.1 - 57.7	1,900
Honolulu	5,000	40.6	39.3 - 42.0	7,300	59.4	58.0 - 60.7	12,300
Kauai	300	43.5	39.8 - 47.3	400	56.5	52.7 - 60.2	700
Maui	800	42.9	40.5 - 45.3	1,000	57.1	54.7 - 59.5	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,200	59.4	57.0 - 61.8	1,500	40.6	38.2 - 43.0	3,700
Native Hawaiian	2,000	44.6	43.1 - 46.0	2,500	55.4	54.0 - 56.9	4,500
Chinese	200	28.2	23.1 - 33.9	500	71.8	66.1 - 76.9	700
Filipino	900	26.8	24.6 - 29.2	2,500	73.2	70.8 - 75.4	3,400
Japanese	900	41.7	38.3 - 45.2	1,200	58.3	54.8 - 61.7	2,100
Other	700	32.0	29.0 - 35.3	1,600	68.0	64.7 - 71.0	2,300

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	33.4	30.4 - 36.6	1,000	66.6	63.4 - 69.6	1,500
20-24	1,900	45.9	43.8 - 48.1	2,200	54.1	51.9 - 56.2	4,200
25-34	3,500	42.1	40.5 - 43.7	4,800	57.9	56.3 - 59.5	8,200
35+	1,100	37.5	34.9 - 40.1	1,800	62.5	59.9 - 65.1	2,900

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,700	42.1	40.6 - 43.6	5,100	57.9	56.4 - 59.4	8,700
Female	3,300	40.7	39.2 - 42.3	4,800	59.3	57.7 - 60.8	8,100

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,600	40.3	39.0 - 41.7	6,800	59.7	58.3 - 61.0	11,400
No	2,300	43.9	42.1 - 45.7	3,000	56.1	54.3 - 57.9	5,300

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,400	35.4	33.8 - 37.0	4,300	64.6	63.0 - 66.2	6,700
131-185%	1,200	43.0	40.3 - 45.8	1,500	57.0	54.2 - 59.7	2,700
186+%	3,000	50.6	48.7 - 52.5	2,900	49.4	47.5 - 51.3	5,900

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight							
Low birth weight	400	34.6	30.6 - 38.8	700	65.4	61.2 - 69.4	1,100
Normal birth weight	6,600	41.9	40.8 - 43.0	9,100	58.1	57.0 - 59.2	15,700

	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester							
Yes	800	52.5	48.9 - 56.1	700	47.5	43.9 - 51.1	1,400
No	6,100	40.3	39.2 - 41.5	9,100	59.7	58.5 - 60.8	15,200

	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant							
Yes	2,100	62.6	60.3 - 64.9	1,300	37.4	35.1 - 39.7	3,400
No	4,700	35.7	34.6 - 36.9	8,500	64.3	63.1 - 65.4	13,200

PRAMS Health Indicator 5-Year Aggregate Report - Drinking alcohol 3 months before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	37.4	35.9 - 39.0	4,500	62.6	61.0 - 64.1	7,200
No	4,200	44.6	43.2 - 46.1	5,200	55.4	53.9 - 56.8	9,500