

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Binge drinking 3 mos before pregnancy (2000-2008)	Binge drinking 3 mos before pregnancy (2000-2008)

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 9/4/15

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	1,900	12.4	11.1 - 14.0	13,600	87.6	86.0 - 88.9	15,600
2001	1,800	11.7	10.4 - 13.2	13,400	88.3	86.8 - 89.6	15,200
2002	1,800	11.6	9.9 - 13.5	14,000	88.4	86.5 - 90.1	15,800
2003	1,900	12.1	10.4 - 14.0	14,200	87.9	86.0 - 89.6	16,100
2004	2,800	16.1	14.5 - 17.8	14,600	83.9	82.2 - 85.5	17,500
2005	3,300	19.1	17.3 - 21.1	13,900	80.9	78.9 - 82.7	17,200
2006	3,400	19.1	17.2 - 21.1	14,600	80.9	78.9 - 82.8	18,100
2007	3,600	19.4	17.6 - 21.3	14,800	80.6	78.7 - 82.4	18,400
2008	3,600	19.5	17.7 - 21.5	14,800	80.5	78.5 - 82.3	18,500

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,600	19.5	17.7 - 21.5	14,800	80.5	78.5 - 82.3	18,500

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	19.7	14.8 - 25.7	1,900	80.3	74.3 - 85.2	2,400
Honolulu	2,500	19.2	17.1 - 21.5	10,500	80.8	78.5 - 82.9	13,000
Kauai	200	24.4	16.0 - 35.3	600	75.6	64.7 - 84.0	900
Maui	400	19.3	14.3 - 25.6	1,700	80.7	74.4 - 85.7	2,100

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,100	24.3	20.4 - 28.7	3,300	75.7	71.3 - 79.6	4,400
Native Hawaiian	1,100	22.3	18.5 - 26.7	4,000	77.7	73.3 - 81.5	5,100
Chinese	100	12.5	8.6 - 17.8	500	87.5	82.2 - 91.4	600
Filipino	500	15.3	11.9 - 19.4	2,900	84.7	80.6 - 88.1	3,500
Japanese	400	17.8	12.9 - 24.1	1,600	82.2	75.9 - 87.1	2,000
Other	400	15.1	11.4 - 19.8	2,500	84.9	80.2 - 88.6	2,900

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	18.6	12.6 - 26.6	1,200	81.4	73.4 - 87.4	1,400
20-24	1,200	28.9	24.4 - 33.9	2,900	71.1	66.1 - 75.6	4,000
25-34	1,700	17.9	15.6 - 20.6	7,900	82.1	79.4 - 84.4	9,600
35+	400	13.2	9.9 - 17.4	2,900	86.8	82.6 - 90.1	3,400

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,900	19.6	17.1 - 22.4	7,800	80.4	77.6 - 82.9	9,700
Female	1,700	19.4	16.8 - 22.3	7,100	80.6	77.7 - 83.2	8,800

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	16.5	14.4 - 18.8	9,700	83.5	81.2 - 85.6	11,600
No	1,700	24.7	21.4 - 28.3	5,200	75.3	71.7 - 78.6	6,900

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,300	19.2	16.2 - 22.6	5,600	80.8	77.4 - 83.8	6,900
131-185%	400	16.0	11.8 - 21.2	2,200	84.0	78.8 - 88.2	2,600
186+%	1,600	21.0	18.3 - 24.1	6,100	79.0	75.9 - 81.7	7,800

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	600	19.0	14.7 - 24.2	2,500	81.0	75.8 - 85.3	3,100
\$10,000-\$24,999	900	21.6	17.5 - 26.2	3,100	78.4	73.8 - 82.5	4,000
\$25,000-\$49,999	1,000	22.9	19.0 - 27.4	3,200	77.1	72.6 - 81.0	4,200
>=\$50,000	1,000	16.8	14.0 - 20.0	5,100	83.2	80.0 - 86.0	6,200

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	12.1	7.3 - 19.5	1,100	87.9	80.5 - 92.7	1,200
Normal birth weight	3,500	20.1	18.2 - 22.1	13,700	79.9	77.9 - 81.8	17,200

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	31.2	23.9 - 39.7	1,100	68.8	60.3 - 76.1	1,600
No	3,100	18.4	16.6 - 20.4	13,700	81.6	79.6 - 83.4	16,900

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	39.6	34.3 - 45.2	2,100	60.4	54.8 - 65.7	3,500
No	2,200	14.8	13.0 - 16.8	12,700	85.2	83.2 - 87.0	14,900

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	19.3	16.5 - 22.5	6,400	80.7	77.5 - 83.5	7,900
No	2,100	19.9	17.5 - 22.5	8,300	80.1	77.5 - 82.5	10,300

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,600	19.4	17.6 - 21.3	14,800	80.6	78.7 - 82.4	18,400

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	17.9	13.3 - 23.6	2,000	82.1	76.4 - 86.7	2,400
Honolulu	2,400	19.1	17.0 - 21.4	10,300	80.9	78.6 - 83.0	12,700
Kauai	300	24.5	16.9 - 34.0	800	75.5	66.0 - 83.1	1,100
Maui	400	20.0	14.8 - 26.4	1,700	80.0	73.6 - 85.2	2,200

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,100	25.9	21.7 - 30.5	3,100	74.1	69.5 - 78.3	4,100
Native Hawaiian	1,200	24.2	20.2 - 28.6	3,800	75.8	71.4 - 79.8	5,000
Chinese	100	7.7	4.9 - 12.1	600	92.3	87.9 - 95.1	700
Filipino	400	12.0	9.1 - 15.8	3,000	88.0	84.2 - 90.9	3,400
Japanese	400	20.2	15.2 - 26.3	1,700	79.8	73.7 - 84.8	2,200
Other	400	12.8	9.4 - 17.3	2,600	87.2	82.7 - 90.6	3,000

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	16.2	10.7 - 23.8	1,400	83.8	76.2 - 89.3	1,700
20-24	1,200	25.6	21.5 - 30.1	3,400	74.4	69.9 - 78.5	4,600
25-34	1,700	19.0	16.6 - 21.8	7,300	81.0	78.2 - 83.4	9,000
35+	400	12.9	9.7 - 17.1	2,700	87.1	82.9 - 90.3	3,100

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,800	19.4	16.9 - 22.1	7,700	80.6	77.9 - 83.1	9,500
Female	1,700	19.4	16.8 - 22.2	7,100	80.6	77.8 - 83.2	8,900

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	16.5	14.4 - 18.8	9,500	83.5	81.2 - 85.6	11,400
No	1,700	24.1	20.9 - 27.7	5,300	75.9	72.3 - 79.1	7,000

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,200	17.1	14.3 - 20.4	5,700	82.9	79.6 - 85.7	6,900
131-185%	500	18.6	14.3 - 23.8	2,200	81.4	76.2 - 85.7	2,700
186+%	1,700	22.2	19.3 - 25.4	5,800	77.8	74.6 - 80.7	7,500

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	500	16.2	12.1 - 21.3	2,600	83.8	78.7 - 87.9	3,100
\$10,000-\$24,999	800	21.1	17.1 - 25.7	3,000	78.9	74.3 - 82.9	3,800
\$25,000-\$49,999	900	21.1	17.4 - 25.5	3,300	78.9	74.5 - 82.6	4,200
>=\$50,000	1,200	19.3	16.3 - 22.7	4,800	80.7	77.3 - 83.7	6,000

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	16.7	11.3 - 24.0	1,300	83.3	76.0 - 88.7	1,600
Normal birth weight	3,300	19.6	17.7 - 21.7	13,500	80.4	78.3 - 82.3	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	36.8	29.0 - 45.5	1,000	63.2	54.5 - 71.0	1,500
No	3,000	17.7	15.9 - 19.7	13,800	82.3	80.3 - 84.1	16,700

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	44.1	38.8 - 49.7	2,000	55.9	50.3 - 61.2	3,700
No	1,900	13.1	11.4 - 15.0	12,700	86.9	85.0 - 88.6	14,600

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	19.1	16.3 - 22.3	6,100	80.9	77.7 - 83.7	7,600
No	2,000	19.4	17.1 - 22.0	8,400	80.6	78.0 - 82.9	10,400

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,400	19.1	17.2 - 21.1	14,600	80.9	78.9 - 82.8	18,100

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	18.6	13.6 - 24.9	1,800	81.4	75.1 - 86.4	2,200
Honolulu	2,400	18.5	16.4 - 20.9	10,700	81.5	79.1 - 83.6	13,100
Kauai	200	21.4	13.4 - 32.5	700	78.6	67.5 - 86.6	900
Maui	400	22.3	16.5 - 29.5	1,500	77.7	70.5 - 83.5	1,900

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,000	22.9	19.0 - 27.4	3,300	77.1	72.6 - 81.0	4,300
Native Hawaiian	1,200	23.9	19.8 - 28.5	3,800	76.1	71.5 - 80.2	4,900
Chinese	<50	6.2	3.5 - 10.6	600	93.8	89.4 - 96.5	600
Filipino	400	11.5	8.4 - 15.5	2,900	88.5	84.5 - 91.6	3,300
Japanese	300	15.1	10.6 - 21.1	1,800	84.9	78.9 - 89.4	2,100
Other	600	19.3	14.8 - 24.8	2,300	80.7	75.2 - 85.2	2,900

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	17.6	11.5 - 25.8	1,300	82.4	74.2 - 88.5	1,500
20-24	1,200	26.2	21.8 - 31.0	3,300	73.8	69.0 - 78.2	4,400
25-34	1,700	19.3	16.7 - 22.2	7,200	80.7	77.8 - 83.3	8,900
35+	300	9.6	6.8 - 13.4	3,000	90.4	86.6 - 93.2	3,300

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,800	19.5	16.9 - 22.4	7,400	80.5	77.6 - 83.1	9,200
Female	1,700	18.7	16.0 - 21.6	7,200	81.3	78.4 - 84.0	8,800

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	17.2	15.1 - 19.6	9,600	82.8	80.4 - 84.9	11,500
No	1,500	22.4	19.0 - 26.2	5,100	77.6	73.8 - 81.0	6,500

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,100	18.1	14.9 - 21.7	5,200	81.9	78.3 - 85.1	6,300
131-185%	600	26.0	20.4 - 32.5	1,800	74.0	67.5 - 79.6	2,400
186+%	1,500	18.7	16.0 - 21.6	6,600	81.3	78.4 - 84.0	8,100

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	500	15.7	11.5 - 21.0	2,500	84.3	79.0 - 88.5	3,000
\$10,000-\$24,999	900	24.5	20.0 - 29.7	2,800	75.5	70.3 - 80.0	3,800
\$25,000-\$49,999	1,000	24.1	20.0 - 28.8	3,300	75.9	71.2 - 80.0	4,300
>=\$50,000	900	15.0	12.3 - 18.2	4,900	85.0	81.8 - 87.7	5,800

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	14.0	9.0 - 21.1	1,300	86.0	78.9 - 91.0	1,500
Normal birth weight	3,200	19.6	17.6 - 21.7	13,300	80.4	78.3 - 82.4	16,500

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	36.9	29.1 - 45.5	1,100	63.1	54.5 - 70.9	1,700
No	2,800	17.1	15.2 - 19.2	13,500	82.9	80.8 - 84.8	16,300

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	40.6	35.3 - 46.2	2,300	59.4	53.8 - 64.7	3,800
No	1,900	13.1	11.3 - 15.1	12,300	86.9	84.9 - 88.7	14,200

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	20.4	17.4 - 23.8	6,100	79.6	76.2 - 82.6	7,600
No	1,800	18.0	15.6 - 20.6	8,300	82.0	79.4 - 84.4	10,200

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,300	19.1	17.3 - 21.1	13,900	80.9	78.9 - 82.7	17,200

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	21.2	16.2 - 27.3	1,700	78.8	72.7 - 83.8	2,200
Honolulu	2,300	18.3	16.2 - 20.5	10,300	81.7	79.5 - 83.8	12,600
Kauai	100	14.4	7.7 - 25.4	600	85.6	74.6 - 92.3	700
Maui	400	24.6	18.4 - 32.1	1,300	75.4	67.9 - 81.6	1,700

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	23.6	19.5 - 28.3	2,900	76.4	71.7 - 80.5	3,700
Native Hawaiian	1,300	26.0	21.9 - 30.5	3,600	74.0	69.5 - 78.1	4,900
Chinese	<50	5.8	3.5 - 9.6	600	94.2	90.4 - 96.5	600
Filipino	400	12.2	9.2 - 16.0	2,800	87.8	84.0 - 90.8	3,200
Japanese	300	12.3	8.7 - 17.1	2,000	87.7	82.9 - 91.3	2,300
Other	400	17.3	13.1 - 22.4	2,100	82.7	77.6 - 86.9	2,500

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	21.9	15.5 - 29.9	1,100	78.1	70.1 - 84.5	1,500
20-24	1,200	28.5	24.2 - 33.2	3,000	71.5	66.8 - 75.8	4,200
25-34	1,400	16.0	13.6 - 18.6	7,200	84.0	81.4 - 86.4	8,600
35+	400	13.5	10.1 - 17.9	2,600	86.5	82.1 - 89.9	3,000

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,700	19.5	17.0 - 22.3	7,000	80.5	77.7 - 83.0	8,700
Female	1,600	18.7	16.1 - 21.5	6,900	81.3	78.5 - 83.9	8,400

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	14.1	12.1 - 16.3	9,400	85.9	83.7 - 87.9	10,900
No	1,700	27.9	24.4 - 31.7	4,500	72.1	68.3 - 75.6	6,200

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,300	20.6	17.5 - 24.1	4,900	79.4	75.9 - 82.5	6,200
131-185%	400	24.5	18.6 - 31.5	1,400	75.5	68.5 - 81.4	1,800
186+%	1,400	18.7	16.1 - 21.7	6,300	81.3	78.3 - 83.9	7,700

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	700	23.4	18.7 - 28.9	2,200	76.6	71.1 - 81.3	2,900
\$10,000-\$24,999	700	20.0	16.1 - 24.5	2,900	80.0	75.5 - 83.9	3,600
\$25,000-\$49,999	900	21.9	18.0 - 26.4	3,100	78.1	73.6 - 82.0	4,000
>=\$50,000	900	17.1	14.1 - 20.5	4,500	82.9	79.5 - 85.9	5,400

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	22.9	16.0 - 31.7	1,000	77.1	68.3 - 84.0	1,200
Normal birth weight	3,000	18.8	16.9 - 20.8	12,900	81.2	79.2 - 83.1	15,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	36.0	28.2 - 44.6	900	64.0	55.4 - 71.8	1,400
No	2,700	17.5	15.6 - 19.5	12,900	82.5	80.5 - 84.4	15,600

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	38.9	33.9 - 44.3	2,200	61.1	55.7 - 66.1	3,600
No	1,800	13.7	11.9 - 15.7	11,600	86.3	84.3 - 88.1	13,400

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	20.4	17.5 - 23.6	5,800	79.6	76.4 - 82.5	7,300
No	1,700	18.2	15.9 - 20.8	7,800	81.8	79.2 - 84.1	9,600

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	2,800	16.1	14.5 - 17.8	14,600	83.9	82.2 - 85.5	17,500

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	21.4	16.6 - 27.3	1,700	78.6	72.7 - 83.4	2,200
Honolulu	1,900	14.7	13.0 - 16.7	10,800	85.3	83.3 - 87.0	12,700
Kauai	100	17.9	11.1 - 27.5	600	82.1	72.5 - 88.9	800
Maui	300	18.3	13.5 - 24.2	1,500	81.7	75.8 - 86.5	1,800

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	19.0	15.6 - 22.8	3,100	81.0	77.2 - 84.4	3,900
Native Hawaiian	1,100	24.0	20.4 - 28.2	3,600	76.0	71.8 - 79.6	4,800
Chinese	<50	5.4	3.6 - 7.9	600	94.6	92.1 - 96.4	700
Filipino	300	9.4	7.1 - 12.5	3,000	90.6	87.5 - 92.9	3,300
Japanese	300	13.8	10.0 - 18.8	2,000	86.2	81.2 - 90.0	2,300
Other	300	10.2	7.3 - 14.2	2,300	89.8	85.8 - 92.7	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	15.7	10.5 - 22.9	1,100	84.3	77.1 - 89.5	1,300
20-24	900	21.1	17.6 - 25.0	3,500	78.9	75.0 - 82.4	4,400
25-34	1,400	16.3	14.1 - 18.8	7,100	83.7	81.2 - 85.9	8,500
35+	300	8.9	6.5 - 12.1	3,000	91.1	87.9 - 93.5	3,300

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,500	16.1	14.0 - 18.4	8,000	83.9	81.6 - 86.0	9,500
Female	1,300	16.1	13.8 - 18.6	6,700	83.9	81.4 - 86.2	8,000

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	13.2	11.5 - 15.1	10,600	86.8	84.9 - 88.5	12,300
No	1,200	22.9	19.6 - 26.6	4,000	77.1	73.4 - 80.4	5,200

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,000	15.7	13.1 - 18.6	5,400	84.3	81.4 - 86.9	6,400
131-185%	400	19.5	14.9 - 25.1	1,600	80.5	74.9 - 85.1	2,000
186+%	1,300	16.8	14.5 - 19.4	6,300	83.2	80.6 - 85.5	7,600

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	500	17.5	13.4 - 22.6	2,200	82.5	77.4 - 86.6	2,700
\$10,000-\$24,999	800	17.8	14.6 - 21.5	3,500	82.2	78.5 - 85.4	4,200
\$25,000-\$49,999	700	16.4	13.4 - 19.9	3,700	83.6	80.1 - 86.6	4,400
>=\$50,000	700	15.2	12.5 - 18.4	4,100	84.8	81.6 - 87.5	4,900

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	15.0	9.6 - 22.6	900	85.0	77.4 - 90.4	1,100
Normal birth weight	2,600	16.2	14.6 - 17.9	13,700	83.8	82.1 - 85.4	16,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	33.2	26.0 - 41.3	900	66.8	58.7 - 74.0	1,400
No	2,300	14.6	13.1 - 16.3	13,600	85.4	83.7 - 86.9	16,000

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	35.1	30.5 - 40.1	2,300	64.9	59.9 - 69.5	3,600
No	1,500	11.2	9.7 - 12.8	12,300	88.8	87.2 - 90.3	13,800

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	16.5	14.1 - 19.2	6,400	83.5	80.8 - 85.9	7,700
No	1,500	15.9	13.9 - 18.2	7,900	84.1	81.8 - 86.1	9,400

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,900	12.1	10.4 - 14.0	14,200	87.9	86.0 - 89.6	16,100

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	11.8	9.2 - 15.1	1,600	88.2	84.9 - 90.8	1,800
Honolulu	1,400	12.0	9.8 - 14.5	10,300	88.0	85.5 - 90.2	11,700
Kauai	100	9.2	5.6 - 14.8	700	90.8	85.2 - 94.4	800
Maui	300	14.5	11.3 - 18.3	1,500	85.5	81.7 - 88.7	1,800

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	18.2	13.6 - 23.9	2,800	81.8	76.1 - 86.4	3,500
Native Hawaiian	700	16.6	14.3 - 19.1	3,500	83.4	80.9 - 85.7	4,200
Chinese	0	0.0	-	700	100.0	-	700
Filipino	300	9.1	5.9 - 13.7	3,300	90.9	86.3 - 94.1	3,700
Japanese	100	5.7	2.8 - 11.2	1,800	94.3	88.8 - 97.2	1,900
Other	200	8.2	4.5 - 14.4	2,000	91.8	85.6 - 95.5	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.3	5.0 - 13.4	1,300	91.7	86.6 - 95.0	1,400
20-24	600	17.4	13.7 - 21.9	3,000	82.6	78.1 - 86.3	3,700
25-34	1,000	12.0	9.7 - 14.8	7,200	88.0	85.2 - 90.3	8,200
35+	200	7.4	4.5 - 11.9	2,600	92.6	88.1 - 95.5	2,900

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,000	11.7	9.5 - 14.4	7,500	88.3	85.6 - 90.5	8,500
Female	900	12.5	10.1 - 15.3	6,700	87.5	84.7 - 89.9	7,600

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	9.8	7.9 - 12.2	9,700	90.2	87.8 - 92.1	10,700
No	900	16.6	13.7 - 19.9	4,500	83.4	80.1 - 86.3	5,400

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,000	13.5	11.0 - 16.5	6,100	86.5	83.5 - 89.0	7,100
131-185%	300	10.5	7.2 - 14.9	2,100	89.5	85.1 - 92.8	2,400
186+%	600	11.5	8.5 - 15.3	4,500	88.5	84.7 - 91.5	5,100

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	500	12.2	9.1 - 16.1	3,400	87.8	83.9 - 90.9	3,800
\$15,001-\$25,000	300	15.4	11.0 - 21.3	1,700	84.6	78.7 - 89.0	2,000
\$25,001-\$40,000	400	14.2	10.2 - 19.5	2,600	85.8	80.5 - 89.8	3,000
>=\$40,001	600	10.4	7.9 - 13.5	5,600	89.6	86.5 - 92.1	6,200

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	6.1	2.7 - 13.4	1,100	93.9	86.6 - 97.3	1,200
Normal birth weight	1,900	12.6	10.8 - 14.5	13,100	87.4	85.5 - 89.2	14,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	20.8	14.2 - 29.5	1,100	79.2	70.5 - 85.8	1,400
No	1,600	11.1	9.4 - 13.0	13,000	88.9	87.0 - 90.6	14,600

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	25.8	20.9 - 31.4	2,300	74.2	68.6 - 79.1	3,100
No	1,100	8.5	6.9 - 10.4	11,700	91.5	89.6 - 93.1	12,800

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	12.2	9.9 - 15.0	5,900	87.8	85.0 - 90.1	6,700
No	1,100	12.1	9.8 - 14.7	8,200	87.9	85.3 - 90.2	9,400

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,800	11.6	9.9 - 13.5	14,000	88.4	86.5 - 90.1	15,800

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	13.1	10.3 - 16.5	1,500	86.9	83.5 - 89.7	1,800
Honolulu	1,300	11.3	9.2 - 13.8	10,400	88.7	86.2 - 90.8	11,700
Kauai	100	13.2	8.6 - 19.7	500	86.8	80.3 - 91.4	600
Maui	200	11.5	8.6 - 15.1	1,500	88.5	84.9 - 91.4	1,700

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	500	13.9	9.8 - 19.3	2,900	86.1	80.7 - 90.2	3,300
Native Hawaiian	600	15.8	13.6 - 18.3	3,400	84.2	81.7 - 86.4	4,100
Chinese	<50	5.6	1.6 - 18.3	500	94.4	81.7 - 98.4	600
Filipino	100	4.4	2.4 - 7.7	3,100	95.6	92.3 - 97.6	3,300
Japanese	300	12.4	7.9 - 18.9	2,000	87.6	81.1 - 92.1	2,300
Other	300	11.8	6.9 - 19.4	2,000	88.2	80.6 - 93.1	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.7	5.8 - 12.9	1,300	91.3	87.1 - 94.2	1,400
20-24	700	17.7	13.8 - 22.5	3,100	82.3	77.5 - 86.2	3,800
25-34	900	11.6	9.2 - 14.4	7,000	88.4	85.6 - 90.8	8,000
35+	100	4.5	2.4 - 8.1	2,500	95.5	91.9 - 97.6	2,600

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	11.3	9.0 - 14.0	6,900	88.7	86.0 - 91.0	7,800
Female	1,000	11.9	9.6 - 14.7	7,000	88.1	85.3 - 90.4	8,000

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	9.5	7.6 - 11.8	9,900	90.5	88.2 - 92.4	10,900
No	800	16.3	13.2 - 20.0	4,100	83.7	80.0 - 86.8	4,900

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	700	12.3	9.7 - 15.3	5,300	87.7	84.7 - 90.3	6,000
131-185%	400	11.9	8.4 - 16.7	2,900	88.1	83.3 - 91.6	3,200
186+%	600	11.5	8.6 - 15.2	4,500	88.5	84.8 - 91.4	5,100

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	500	15.7	11.9 - 20.5	2,500	84.3	79.5 - 88.1	3,000
\$15,001-\$25,000	300	11.1	7.2 - 16.5	2,100	88.9	83.5 - 92.8	2,300
\$25,001-\$40,000	300	10.0	6.8 - 14.3	3,100	90.0	85.7 - 93.2	3,500
>=\$40,001	700	11.4	8.8 - 14.7	5,400	88.6	85.3 - 91.2	6,000

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.2	4.8 - 16.9	1,000	90.8	83.1 - 95.2	1,100
Normal birth weight	1,700	11.6	9.9 - 13.6	13,000	88.4	86.4 - 90.1	14,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	22.1	15.1 - 31.2	900	77.9	68.8 - 84.9	1,200
No	1,500	10.6	9.0 - 12.6	12,900	89.4	87.4 - 91.0	14,400

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	28.5	23.0 - 34.8	2,000	71.5	65.2 - 77.0	2,800
No	1,000	7.5	6.0 - 9.3	11,800	92.5	90.7 - 94.0	12,700

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	13.2	10.6 - 16.3	5,700	86.8	83.7 - 89.4	6,500
No	1,000	10.6	8.5 - 13.1	8,300	89.4	86.9 - 91.5	9,200

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,800	11.7	10.4 - 13.2	13,400	88.3	86.8 - 89.6	15,200

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	16.3	13.0 - 20.2	1,400	83.7	79.8 - 87.0	1,600
Honolulu	1,300	11.1	9.5 - 13.0	10,000	88.9	87.0 - 90.5	11,300
Kauai	100	11.7	7.7 - 17.4	600	88.3	82.6 - 92.3	700
Maui	200	11.4	8.5 - 15.1	1,400	88.6	84.9 - 91.5	1,600

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	500	14.4	11.2 - 18.4	2,800	85.6	81.6 - 88.8	3,200
Native Hawaiian	700	16.9	15.0 - 18.9	3,300	83.1	81.1 - 85.0	4,000
Chinese	<50	6.3	2.1 - 17.3	700	93.7	82.7 - 97.9	700
Filipino	100	3.9	2.3 - 6.6	3,100	96.1	93.4 - 97.7	3,200
Japanese	200	11.1	7.1 - 17.1	1,700	88.9	82.9 - 92.9	1,900
Other	300	12.3	8.3 - 17.7	1,900	87.7	82.3 - 91.7	2,100

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.7	6.3 - 11.9	1,300	91.3	88.1 - 93.7	1,400
20-24	600	16.1	13.2 - 19.5	3,200	83.9	80.5 - 86.8	3,800
25-34	900	11.6	9.7 - 13.8	6,500	88.4	86.2 - 90.3	7,400
35+	200	7.4	4.7 - 11.3	2,400	92.6	88.7 - 95.3	2,600

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	900	11.6	9.7 - 13.7	6,700	88.4	86.3 - 90.3	7,600
Female	900	11.9	10.0 - 14.0	6,700	88.1	86.0 - 90.0	7,600

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	10.6	9.0 - 12.5	9,200	89.4	87.5 - 91.0	10,300
No	700	14.2	11.9 - 16.7	4,200	85.8	83.3 - 88.1	4,900

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	700	13.1	11.0 - 15.5	5,000	86.9	84.5 - 89.0	5,700
131-185%	300	9.0	7.0 - 11.6	3,000	91.0	88.4 - 93.0	3,300
186+%	700	13.3	10.7 - 16.4	4,400	86.7	83.6 - 89.3	5,100

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	400	13.3	10.5 - 16.6	2,600	86.7	83.4 - 89.5	3,000
\$15,001-\$25,000	300	15.4	11.6 - 20.2	1,700	84.6	79.8 - 88.4	2,100
\$25,001-\$40,000	400	11.4	8.8 - 14.6	2,900	88.6	85.4 - 91.2	3,300
>=\$40,001	700	11.0	8.9 - 13.6	5,300	89.0	86.4 - 91.1	6,000

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.3	3.8 - 13.3	900	92.7	86.7 - 96.2	1,000
Normal birth weight	1,700	12.1	10.7 - 13.6	12,500	87.9	86.4 - 89.3	14,200

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	20.7	16.0 - 26.4	1,100	79.3	73.6 - 84.0	1,400
No	1,500	10.7	9.3 - 12.2	12,200	89.3	87.8 - 90.7	13,700

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	27.0	22.9 - 31.7	2,100	73.0	68.3 - 77.1	2,900
No	1,000	8.0	6.7 - 9.4	11,000	92.0	90.6 - 93.3	12,000

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	13.6	11.5 - 16.0	5,500	86.4	84.0 - 88.5	6,400
No	900	10.4	8.7 - 12.3	7,900	89.6	87.7 - 91.3	8,800

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,900	12.4	11.1 - 14.0	13,600	87.6	86.0 - 88.9	15,600

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	14.2	11.2 - 17.9	1,500	85.8	82.1 - 88.8	1,700
Honolulu	1,400	11.9	10.3 - 13.8	10,200	88.1	86.2 - 89.7	11,600
Kauai	100	16.5	11.1 - 23.8	500	83.5	76.2 - 88.9	600
Maui	200	13.1	9.9 - 17.0	1,400	86.9	83.0 - 90.1	1,600

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	16.3	13.1 - 20.2	3,000	83.7	79.8 - 86.9	3,600
Native Hawaiian	700	18.4	16.4 - 20.6	3,300	81.6	79.4 - 83.6	4,000
Chinese	<50	0.9	0.1 - 5.3	600	99.1	94.7 - 99.9	600
Filipino	200	7.0	4.6 - 10.5	3,000	93.0	89.5 - 95.4	3,200
Japanese	200	10.5	6.6 - 16.2	1,700	89.5	83.8 - 93.4	1,900
Other	200	8.1	4.8 - 13.4	2,000	91.9	86.6 - 95.2	2,100

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	14.5	10.2 - 20.3	1,300	85.5	79.7 - 89.8	1,600
20-24	600	17.8	14.8 - 21.3	3,000	82.2	78.7 - 85.2	3,600
25-34	900	11.9	9.9 - 14.1	6,600	88.1	85.9 - 90.1	7,500
35+	200	5.9	4.0 - 8.7	2,700	94.1	91.3 - 96.0	2,800

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,100	13.9	12.0 - 16.2	7,000	86.1	83.8 - 88.0	8,100
Female	800	10.8	9.0 - 13.0	6,700	89.2	87.0 - 91.0	7,500

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	11.0	9.5 - 12.8	9,800	89.0	87.2 - 90.5	11,000
No	700	15.9	13.4 - 18.8	3,800	84.1	81.2 - 86.6	4,500

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	800	12.1	10.3 - 14.3	5,700	87.9	85.7 - 89.7	6,500
131-185%	300	16.1	12.2 - 20.9	1,700	83.9	79.1 - 87.8	2,100
186+%	700	12.3	10.0 - 15.1	5,100	87.7	84.9 - 90.0	5,800

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	500	13.7	11.0 - 17.1	3,000	86.3	82.9 - 89.0	3,500
\$15,001-\$25,000	300	11.0	8.3 - 14.4	2,200	89.0	85.6 - 91.7	2,500
\$25,001-\$40,000	500	16.8	13.2 - 21.1	2,400	83.2	78.9 - 86.8	2,900
>=\$40,001	600	10.5	8.5 - 13.0	5,300	89.5	87.0 - 91.5	5,900

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.0	5.4 - 14.6	900	91.0	85.4 - 94.6	1,000
Normal birth weight	1,800	12.7	11.2 - 14.2	12,600	87.3	85.8 - 88.8	14,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	27.9	21.5 - 35.3	900	72.1	64.7 - 78.5	1,200
No	1,600	11.1	9.8 - 12.7	12,600	88.9	87.3 - 90.2	14,200

PRAMS Health Indicator Report - Binge drinking 3 mos before pregnancy (2000-2008)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	28.7	24.3 - 33.6	2,000	71.3	66.4 - 75.7	2,800
No	1,100	8.7	7.4 - 10.2	11,400	91.3	89.8 - 92.6	12,500

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	12.5	10.6 - 14.8	5,800	87.5	85.2 - 89.4	6,700
No	1,100	12.5	10.6 - 14.6	7,700	87.5	85.4 - 89.4	8,800