

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- for the State of Hawaii,
for the Year(s)**

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 3-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, income data does not exist for some 3-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Discuss breastfeeding with HCW during PNC	Discuss breastfeeding with HCW during PNC

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 9/16/15

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- for the State of Hawaii, for the Year(s)

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012


Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2002	14,200	85.6	84.6 - 86.6	2,400	14.4	13.4 - 15.4	16,600
2001-2003	14,400	86.1	85.1 - 87.2	2,300	13.9	12.8 - 14.9	16,700
2002-2004	14,600	85.7	84.6 - 86.7	2,400	14.3	13.3 - 15.4	17,000
2003-2005	14,700	85.8	84.8 - 86.7	2,400	14.2	13.3 - 15.2	17,100
2004-2006	14,700	84.8	83.9 - 85.8	2,600	15.2	14.2 - 16.1	17,300
2005-2007	14,900	84.7	83.7 - 85.7	2,700	15.3	14.3 - 16.3	17,600
2006-2008	15,100	83.9	82.9 - 84.9	2,900	16.1	15.1 - 17.1	18,000
2007-2009	15,100	83.5	82.4 - 84.6	3,000	16.5	15.4 - 17.6	18,100
2008-2010	15,100	83.4	82.1 - 84.6	3,000	16.6	15.4 - 17.9	18,200
2009-2011	15,200	84.0	82.6 - 85.3	2,900	16.0	14.7 - 17.4	18,100
2010-2012	15,400	84.5	83.1 - 85.8	2,800	15.5	14.2 - 16.9	18,200

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- for the State of Hawaii, for the Year(s)

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	15,400	84.5	83.1 - 85.8	2,800	15.5	14.2 - 16.9	18,200

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	2,000	86.3	84.1 - 88.2	300	13.7	11.8 - 15.9	2,300
Honolulu	11,000	84.0	82.1 - 85.7	2,100	16.0	14.3 - 17.9	13,100
Kauai	700	84.7	81.9 - 87.0	100	15.3	13.0 - 18.1	800
Maui	1,700	86.2	84.1 - 88.0	300	13.8	12.0 - 15.9	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,700	84.3	81.5 - 86.8	700	15.7	13.2 - 18.5	4,400
Native Hawaiian	4,600	87.8	85.3 - 89.9	600	12.2	10.1 - 14.7	5,300
Chinese	600	75.8	67.3 - 82.7	200	24.2	17.3 - 32.7	800
Filipino	2,800	85.1	81.6 - 88.0	500	14.9	12.0 - 18.4	3,300
Japanese	1,200	69.9	63.8 - 75.4	500	30.1	24.6 - 36.2	1,700
Other	2,500	89.5	86.1 - 92.2	300	10.5	7.8 - 13.9	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	92.9	87.4 - 96.1	100	7.1	3.9 - 12.6	1,000
20-24	3,800	91.0	88.4 - 93.1	400	9.0	6.9 - 11.6	4,200
25-34	8,200	84.0	82.1 - 85.8	1,600	16.0	14.2 - 17.9	9,800
35+	2,400	75.2	71.2 - 78.8	800	24.8	21.2 - 28.8	3,200

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,800	85.1	83.1 - 86.8	1,400	14.9	13.2 - 16.9	9,100
Female	7,600	84.0	82.0 - 85.9	1,400	16.0	14.1 - 18.0	9,100

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,300	81.6	79.7 - 83.4	2,100	18.4	16.6 - 20.3	11,400
No	6,100	89.4	87.5 - 91.1	700	10.6	8.9 - 12.5	6,800

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,700	89.6	87.6 - 91.3	700	10.4	8.7 - 12.4	6,400
131-185%	2,500	85.6	81.8 - 88.8	400	14.4	11.2 - 18.2	2,900
186+%	6,000	79.3	76.9 - 81.6	1,600	20.7	18.4 - 23.1	7,600

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	81.4	78.8 - 83.8	200	18.6	16.2 - 21.2	1,300
Normal birth weight	14,300	84.8	83.3 - 86.2	2,600	15.2	13.8 - 16.7	16,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	86.4	79.9 - 91.1	100	13.6	8.9 - 20.1	1,000
No	14,200	84.5	83.0 - 85.8	2,600	15.5	14.2 - 17.0	16,800

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	84.9	81.5 - 87.7	500	15.1	12.3 - 18.5	3,400
No	12,300	84.5	83.0 - 86.0	2,200	15.5	14.0 - 17.0	14,500

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,300	90.7	89.0 - 92.1	800	9.3	7.9 - 11.0	8,100
No	7,900	79.6	77.5 - 81.6	2,000	20.4	18.4 - 22.5	10,000

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	15,200	84.0	82.6 - 85.3	2,900	16.0	14.7 - 17.4	18,100

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	2,000	84.8	82.6 - 86.7	400	15.2	13.3 - 17.4	2,300
Honolulu	10,800	83.6	81.6 - 85.3	2,100	16.4	14.7 - 18.4	13,000
Kauai	700	84.1	81.4 - 86.5	100	15.9	13.5 - 18.6	800
Maui	1,700	85.8	83.8 - 87.6	300	14.2	12.4 - 16.2	2,000

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,700	84.5	81.8 - 86.8	700	15.5	13.2 - 18.2	4,400
Native Hawaiian	4,800	86.6	84.1 - 88.7	700	13.4	11.3 - 15.9	5,500
Chinese	500	69.8	60.4 - 77.8	200	30.2	22.2 - 39.6	700
Filipino	2,800	86.3	83.0 - 89.1	400	13.7	10.9 - 17.0	3,200
Japanese	1,200	72.0	66.2 - 77.2	500	28.0	22.8 - 33.8	1,700
Other	2,200	86.7	82.7 - 89.8	300	13.3	10.2 - 17.3	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	91.6	86.6 - 94.9	100	8.4	5.1 - 13.4	1,300
20-24	3,900	89.9	87.2 - 92.0	400	10.1	8.0 - 12.8	4,300
25-34	7,900	83.5	81.5 - 85.3	1,600	16.5	14.7 - 18.5	9,400
35+	2,300	74.3	70.2 - 77.9	800	25.7	22.1 - 29.8	3,100

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,900	84.1	82.2 - 85.9	1,500	15.9	14.1 - 17.8	9,300
Female	7,400	83.8	81.8 - 85.7	1,400	16.2	14.3 - 18.2	8,800

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,000	80.7	78.8 - 82.5	2,200	19.3	17.5 - 21.2	11,200
No	6,200	89.3	87.3 - 91.0	800	10.7	9.0 - 12.7	7,000

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,600	89.2	87.1 - 91.0	700	10.8	9.0 - 12.9	6,300
131-185%	2,400	86.3	82.5 - 89.3	400	13.7	10.7 - 17.5	2,800
186+%	6,000	78.4	76.0 - 80.6	1,700	21.6	19.4 - 24.0	7,700

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,900	90.7	87.8 - 92.9	300	9.3	7.1 - 12.2	3,200
\$10,000-\$24,999	3,100	88.8	85.9 - 91.2	400	11.2	8.8 - 14.1	3,400
\$25,000-\$49,999	3,600	83.5	80.4 - 86.2	700	16.5	13.8 - 19.6	4,300
>=\$50,000	4,700	77.3	74.6 - 79.9	1,400	22.7	20.1 - 25.4	6,100

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,100	81.9	79.4 - 84.1	200	18.1	15.9 - 20.6	1,300
Normal birth weight	14,100	84.2	82.7 - 85.6	2,700	15.8	14.4 - 17.3	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	85.9	80.4 - 90.0	200	14.1	10.0 - 19.6	1,300
No	13,900	83.9	82.4 - 85.3	2,700	16.1	14.7 - 17.6	16,500

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,200	84.8	81.7 - 87.5	600	15.2	12.5 - 18.3	3,700
No	11,900	83.9	82.3 - 85.3	2,300	16.1	14.7 - 17.7	14,200

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,000	89.9	88.1 - 91.4	800	10.1	8.6 - 11.9	7,800
No	8,000	79.6	77.5 - 81.6	2,100	20.4	18.4 - 22.5	10,100

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	15,100	83.4	82.1 - 84.6	3,000	16.6	15.4 - 17.9	18,200

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	2,000	84.4	81.8 - 86.6	400	15.6	13.4 - 18.2	2,400
Honolulu	10,800	83.3	81.6 - 84.9	2,200	16.7	15.1 - 18.4	12,900
Kauai	700	82.1	78.2 - 85.4	200	17.9	14.6 - 21.8	800
Maui	1,700	83.1	80.5 - 85.5	300	16.9	14.5 - 19.5	2,100

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,600	83.4	80.8 - 85.7	700	16.6	14.3 - 19.2	4,300
Native Hawaiian	4,700	86.2	83.8 - 88.2	800	13.8	11.8 - 16.2	5,500
Chinese	500	68.9	60.5 - 76.2	200	31.1	23.8 - 39.5	700
Filipino	2,700	84.1	80.8 - 86.9	500	15.9	13.1 - 19.2	3,300
Japanese	1,300	73.5	68.5 - 77.9	500	26.5	22.1 - 31.5	1,800
Other	2,300	87.3	84.0 - 90.0	300	12.7	10.0 - 16.0	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	90.1	85.5 - 93.4	100	9.9	6.6 - 14.5	1,400
20-24	3,800	89.5	86.9 - 91.6	400	10.5	8.4 - 13.1	4,200
25-34	7,700	82.4	80.6 - 84.1	1,600	17.6	15.9 - 19.4	9,300
35+	2,400	75.3	71.7 - 78.5	800	24.7	21.5 - 28.3	3,200

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,800	82.7	80.8 - 84.3	1,600	17.3	15.7 - 19.2	9,500
Female	7,300	84.2	82.3 - 85.9	1,400	15.8	14.1 - 17.7	8,700

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,000	80.5	78.7 - 82.1	2,200	19.5	17.9 - 21.3	11,200
No	6,100	88.0	86.1 - 89.7	800	12.0	10.3 - 13.9	7,000

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,600	87.8	85.8 - 89.6	800	12.2	10.4 - 14.2	6,400
131-185%	2,400	86.4	83.0 - 89.2	400	13.6	10.8 - 17.0	2,800
186+%	5,900	78.0	75.8 - 80.0	1,700	22.0	20.0 - 24.2	7,600

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,700	89.3	86.3 - 91.6	300	10.7	8.4 - 13.7	3,100
\$10,000-\$24,999	3,200	87.5	84.8 - 89.8	500	12.5	10.2 - 15.2	3,600
\$25,000-\$49,999	3,400	82.9	80.0 - 85.5	700	17.1	14.5 - 20.0	4,100
>=\$50,000	4,700	77.3	74.9 - 79.6	1,400	22.7	20.4 - 25.1	6,100

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	80.9	77.6 - 83.7	200	19.1	16.3 - 22.4	1,300
Normal birth weight	14,100	83.5	82.2 - 84.8	2,800	16.5	15.2 - 17.8	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	86.3	81.5 - 90.0	200	13.7	10.0 - 18.5	1,500
No	13,600	83.2	81.8 - 84.5	2,700	16.8	15.5 - 18.2	16,300

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,100	84.7	81.7 - 87.3	600	15.3	12.7 - 18.3	3,700
No	11,800	83.1	81.7 - 84.5	2,400	16.9	15.5 - 18.3	14,100

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,800	88.5	86.7 - 90.1	900	11.5	9.9 - 13.3	7,700
No	8,100	79.7	77.8 - 81.4	2,100	20.3	18.6 - 22.2	10,200

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	15,100	83.5	82.4 - 84.6	3,000	16.5	15.4 - 17.6	18,100

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,900	81.4	78.4 - 84.0	400	18.6	16.0 - 21.6	2,400
Honolulu	10,600	84.0	82.6 - 85.4	2,000	16.0	14.6 - 17.4	12,600
Kauai	800	84.0	79.6 - 87.6	100	16.0	12.4 - 20.4	900
Maui	1,700	82.8	79.8 - 85.4	400	17.2	14.6 - 20.2	2,100

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,600	84.3	82.1 - 86.4	700	15.7	13.6 - 17.9	4,200
Native Hawaiian	4,600	86.5	84.3 - 88.4	700	13.5	11.6 - 15.7	5,300
Chinese	500	71.1	64.2 - 77.1	200	28.9	22.9 - 35.8	700
Filipino	2,800	84.9	82.2 - 87.3	500	15.1	12.7 - 17.8	3,300
Japanese	1,400	72.2	67.9 - 76.1	500	27.8	23.9 - 32.1	1,900
Other	2,300	85.9	82.9 - 88.4	400	14.1	11.6 - 17.1	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,500	90.3	86.3 - 93.2	200	9.7	6.8 - 13.7	1,600
20-24	3,800	89.9	87.7 - 91.6	400	10.1	8.4 - 12.3	4,200
25-34	7,500	82.6	81.0 - 84.2	1,600	17.4	15.8 - 19.0	9,100
35+	2,300	74.1	70.9 - 77.2	800	25.9	22.8 - 29.1	3,100

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,900	82.8	81.2 - 84.3	1,600	17.2	15.7 - 18.8	9,500
Female	7,200	84.3	82.7 - 85.9	1,300	15.7	14.1 - 17.3	8,600

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,000	80.9	79.4 - 82.3	2,100	19.1	17.7 - 20.6	11,100
No	6,100	87.7	85.9 - 89.3	900	12.3	10.7 - 14.1	6,900

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,800	88.2	86.4 - 89.8	800	11.8	10.2 - 13.6	6,500
131-185%	2,300	85.4	82.4 - 87.9	400	14.6	12.1 - 17.6	2,700
186+%	5,800	78.2	76.3 - 80.1	1,600	21.8	19.9 - 23.7	7,500

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,700	89.6	86.8 - 91.8	300	10.4	8.2 - 13.2	3,000
\$10,000-\$24,999	3,400	87.8	85.4 - 89.8	500	12.2	10.2 - 14.6	3,900
\$25,000-\$49,999	3,300	83.3	80.8 - 85.5	700	16.7	14.5 - 19.2	4,000
>=\$50,000	4,600	77.1	74.9 - 79.2	1,400	22.9	20.8 - 25.1	6,000

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,100	80.4	76.7 - 83.7	300	19.6	16.3 - 23.3	1,400
Normal birth weight	14,000	83.8	82.5 - 84.9	2,700	16.2	15.1 - 17.5	16,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	88.8	85.1 - 91.6	200	11.2	8.4 - 14.9	1,500
No	13,400	83.0	81.8 - 84.2	2,700	17.0	15.8 - 18.2	16,200

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,100	87.2	84.7 - 89.3	500	12.8	10.7 - 15.3	3,600
No	11,700	82.6	81.3 - 83.9	2,500	17.4	16.1 - 18.7	14,100

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,700	88.6	87.0 - 90.1	900	11.4	9.9 - 13.0	7,600
No	8,200	79.8	78.2 - 81.4	2,100	20.2	18.6 - 21.8	10,300

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	15,100	83.9	82.9 - 84.9	2,900	16.1	15.1 - 17.1	18,000

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,900	82.3	79.0 - 85.1	400	17.7	14.9 - 21.0	2,300
Honolulu	10,800	84.4	83.2 - 85.5	2,000	15.6	14.5 - 16.8	12,700
Kauai	800	85.8	80.9 - 89.5	100	14.2	10.5 - 19.1	900
Maui	1,700	82.4	78.9 - 85.3	400	17.6	14.7 - 21.1	2,100

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,500	83.3	81.0 - 85.3	700	16.7	14.7 - 19.0	4,200
Native Hawaiian	4,300	86.7	84.7 - 88.6	700	13.3	11.4 - 15.3	5,000
Chinese	500	74.9	71.3 - 78.2	200	25.1	21.8 - 28.7	600
Filipino	2,900	86.5	84.3 - 88.4	500	13.5	11.6 - 15.7	3,300
Japanese	1,500	72.6	68.9 - 76.0	600	27.4	24.0 - 31.1	2,100
Other	2,400	87.3	84.8 - 89.5	400	12.7	10.5 - 15.2	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,400	90.0	86.3 - 92.7	200	10.0	7.3 - 13.7	1,500
20-24	3,800	88.7	86.7 - 90.4	500	11.3	9.6 - 13.3	4,300
25-34	7,500	83.4	81.9 - 84.8	1,500	16.6	15.2 - 18.1	9,000
35+	2,500	76.3	73.6 - 78.9	800	23.7	21.1 - 26.4	3,200

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,800	84.0	82.6 - 85.4	1,500	16.0	14.6 - 17.4	9,300
Female	7,300	83.9	82.3 - 85.3	1,400	16.1	14.7 - 17.7	8,700

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,300	81.9	80.6 - 83.2	2,100	18.1	16.8 - 19.4	11,400
No	5,800	87.4	85.8 - 88.9	800	12.6	11.1 - 14.2	6,600

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,700	88.1	86.5 - 89.6	800	11.9	10.4 - 13.5	6,500
131-185%	2,200	86.6	83.9 - 88.9	300	13.4	11.1 - 16.1	2,500
186+%	6,100	79.0	77.3 - 80.6	1,600	21.0	19.4 - 22.7	7,700

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,600	89.1	86.6 - 91.2	300	10.9	8.8 - 13.4	3,000
\$10,000-\$24,999	3,300	87.6	85.4 - 89.5	500	12.4	10.5 - 14.6	3,800
\$25,000-\$49,999	3,500	84.4	82.2 - 86.4	600	15.6	13.6 - 17.8	4,100
>=\$50,000	4,700	77.9	75.9 - 79.7	1,300	22.1	20.3 - 24.1	6,000

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,100	80.4	76.2 - 84.0	300	19.6	16.0 - 23.8	1,400
Normal birth weight	14,000	84.2	83.2 - 85.3	2,600	15.8	14.7 - 16.8	16,600

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	87.2	83.6 - 90.2	200	12.8	9.8 - 16.4	1,500
No	13,500	83.6	82.5 - 84.6	2,700	16.4	15.4 - 17.5	16,200

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,100	87.0	84.7 - 89.1	500	13.0	10.9 - 15.3	3,500
No	11,800	83.1	81.9 - 84.2	2,400	16.9	15.8 - 18.1	14,200

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,800	88.7	87.2 - 90.0	900	11.3	10.0 - 12.8	7,700
No	8,300	80.3	78.9 - 81.7	2,000	19.7	18.3 - 21.1	10,300

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	14,900	84.7	83.7 - 85.7	2,700	15.3	14.3 - 16.3	17,600

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,800	82.4	79.2 - 85.3	400	17.6	14.7 - 20.8	2,200
Honolulu	10,700	85.0	83.8 - 86.1	1,900	15.0	13.9 - 16.2	12,600
Kauai	800	88.4	83.7 - 91.8	100	11.6	8.2 - 16.3	900
Maui	1,600	83.8	80.4 - 86.8	300	16.2	13.2 - 19.6	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,400	84.0	81.8 - 86.0	600	16.0	14.0 - 18.2	4,000
Native Hawaiian	4,300	88.3	86.3 - 90.0	600	11.7	10.0 - 13.7	4,900
Chinese	500	77.4	73.9 - 80.5	100	22.6	19.5 - 26.1	600
Filipino	2,800	86.7	84.5 - 88.6	400	13.3	11.4 - 15.5	3,300
Japanese	1,600	73.7	70.1 - 77.0	600	26.3	23.0 - 29.9	2,100
Other	2,300	87.2	84.7 - 89.5	300	12.8	10.5 - 15.3	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,400	91.2	87.8 - 93.7	100	8.8	6.3 - 12.2	1,600
20-24	3,800	88.5	86.5 - 90.2	500	11.5	9.8 - 13.5	4,300
25-34	7,300	84.5	83.0 - 85.8	1,300	15.5	14.2 - 17.0	8,600
35+	2,300	76.7	73.9 - 79.2	700	23.3	20.8 - 26.1	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,700	85.6	84.2 - 86.9	1,300	14.4	13.1 - 15.8	9,000
Female	7,200	83.8	82.2 - 85.2	1,400	16.2	14.8 - 17.8	8,600

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,200	82.9	81.6 - 84.2	1,900	17.1	15.8 - 18.4	11,100
No	5,600	87.7	86.0 - 89.2	800	12.3	10.8 - 14.0	6,400

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,600	88.2	86.6 - 89.7	700	11.8	10.3 - 13.4	6,300
131-185%	2,000	86.4	83.6 - 88.8	300	13.6	11.2 - 16.4	2,300
186+%	6,200	80.5	78.9 - 82.1	1,500	19.5	17.9 - 21.1	7,700

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,600	89.6	87.2 - 91.6	300	10.4	8.4 - 12.8	2,900
\$10,000-\$24,999	3,200	87.9	85.7 - 89.8	400	12.1	10.2 - 14.3	3,700
\$25,000-\$49,999	3,400	84.5	82.3 - 86.5	600	15.5	13.5 - 17.7	4,100
>=\$50,000	4,500	79.3	77.3 - 81.1	1,200	20.7	18.9 - 22.7	5,700

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,100	81.2	77.0 - 84.7	300	18.8	15.3 - 23.0	1,400
Normal birth weight	13,700	85.0	84.0 - 86.0	2,400	15.0	14.0 - 16.0	16,200

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	88.3	84.6 - 91.1	200	11.7	8.9 - 15.4	1,500
No	13,300	84.3	83.2 - 85.3	2,500	15.7	14.7 - 16.8	15,800

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,200	88.6	86.4 - 90.5	400	11.4	9.5 - 13.6	3,600
No	11,500	83.6	82.5 - 84.7	2,200	16.4	15.3 - 17.5	13,700

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,700	88.9	87.4 - 90.2	800	11.1	9.8 - 12.6	7,500
No	8,100	81.4	80.0 - 82.8	1,800	18.6	17.2 - 20.0	9,900

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	14,700	84.8	83.9 - 85.8	2,600	15.2	14.2 - 16.1	17,300

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,800	82.9	79.7 - 85.7	400	17.1	14.3 - 20.3	2,100
Honolulu	10,700	85.1	84.0 - 86.2	1,900	14.9	13.8 - 16.0	12,600
Kauai	700	89.1	84.4 - 92.4	100	10.9	7.6 - 15.6	800
Maui	1,500	83.5	80.1 - 86.4	300	16.5	13.6 - 19.9	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,300	83.4	81.2 - 85.4	700	16.6	14.6 - 18.8	3,900
Native Hawaiian	4,200	87.5	85.6 - 89.3	600	12.5	10.7 - 14.4	4,800
Chinese	500	75.9	72.8 - 78.8	100	24.1	21.2 - 27.2	600
Filipino	2,800	88.0	85.9 - 89.8	400	12.0	10.2 - 14.1	3,200
Japanese	1,600	73.9	70.4 - 77.0	600	26.1	23.0 - 29.6	2,200
Other	2,300	89.6	87.3 - 91.5	300	10.4	8.5 - 12.7	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	90.6	87.2 - 93.2	100	9.4	6.8 - 12.8	1,400
20-24	3,800	88.9	87.0 - 90.6	500	11.1	9.4 - 13.0	4,300
25-34	7,200	84.5	83.1 - 85.9	1,300	15.5	14.1 - 16.9	8,500
35+	2,400	77.4	74.8 - 79.9	700	22.6	20.1 - 25.2	3,100

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,800	85.9	84.6 - 87.2	1,300	14.1	12.8 - 15.4	9,100
Female	6,900	83.7	82.2 - 85.1	1,400	16.3	14.9 - 17.8	8,300

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,500	83.3	82.1 - 84.5	1,900	16.7	15.5 - 17.9	11,400
No	5,200	87.8	86.1 - 89.3	700	12.2	10.7 - 13.9	5,900

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,400	87.8	86.2 - 89.2	800	12.2	10.8 - 13.8	6,200
131-185%	1,800	89.3	86.6 - 91.5	200	10.7	8.5 - 13.4	2,100
186+%	6,300	80.9	79.3 - 82.4	1,500	19.1	17.6 - 20.7	7,700

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,500	88.1	85.6 - 90.2	300	11.9	9.8 - 14.4	2,800
\$10,000-\$24,999	3,400	88.0	86.0 - 89.8	500	12.0	10.2 - 14.0	3,800
\$25,000-\$49,999	3,600	86.0	84.0 - 87.8	600	14.0	12.2 - 16.0	4,200
>=\$50,000	4,200	79.4	77.4 - 81.2	1,100	20.6	18.8 - 22.6	5,300

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	82.8	78.6 - 86.3	200	17.2	13.7 - 21.4	1,200
Normal birth weight	13,700	85.0	84.0 - 86.0	2,400	15.0	14.0 - 16.0	16,100

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	88.8	85.4 - 91.6	200	11.2	8.4 - 14.6	1,400
No	13,200	84.4	83.4 - 85.4	2,400	15.6	14.6 - 16.6	15,600

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,100	87.7	85.5 - 89.6	400	12.3	10.4 - 14.5	3,600
No	11,300	84.0	82.9 - 85.1	2,200	16.0	14.9 - 17.1	13,500

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,700	88.7	87.3 - 90.0	900	11.3	10.0 - 12.7	7,600
No	7,900	81.7	80.3 - 83.0	1,800	18.3	17.0 - 19.7	9,600

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	14,700	85.8	84.8 - 86.7	2,400	14.2	13.3 - 15.2	17,100

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,800	85.0	82.2 - 87.4	300	15.0	12.6 - 17.8	2,100
Honolulu	10,700	86.0	84.8 - 87.1	1,700	14.0	12.9 - 15.2	12,500
Kauai	700	89.0	85.0 - 92.1	100	11.0	7.9 - 15.0	800
Maui	1,500	83.7	80.8 - 86.3	300	16.3	13.7 - 19.2	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,200	84.7	82.4 - 86.7	600	15.3	13.3 - 17.6	3,800
Native Hawaiian	4,200	88.1	86.4 - 89.6	600	11.9	10.4 - 13.6	4,700
Chinese	500	79.6	74.7 - 83.7	100	20.4	16.3 - 25.3	600
Filipino	3,000	87.5	85.2 - 89.5	400	12.5	10.5 - 14.8	3,400
Japanese	1,600	76.1	72.4 - 79.4	500	23.9	20.6 - 27.6	2,200
Other	2,200	90.7	88.2 - 92.6	200	9.3	7.4 - 11.8	2,400

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	91.9	88.9 - 94.2	100	8.1	5.8 - 11.1	1,400
20-24	3,800	91.3	89.6 - 92.7	400	8.7	7.3 - 10.4	4,200
25-34	7,100	84.5	83.0 - 85.9	1,300	15.5	14.1 - 17.0	8,400
35+	2,400	78.5	75.7 - 81.0	600	21.5	19.0 - 24.3	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,800	86.3	84.9 - 87.5	1,200	13.7	12.5 - 15.1	9,100
Female	6,800	85.1	83.6 - 86.5	1,200	14.9	13.5 - 16.4	8,000

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,600	84.3	83.0 - 85.5	1,800	15.7	14.5 - 17.0	11,400
No	5,100	88.7	87.2 - 90.1	600	11.3	9.9 - 12.8	5,700

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,900	88.9	87.4 - 90.3	700	11.1	9.7 - 12.6	6,700
131-185%	1,800	86.4	83.3 - 89.0	300	13.6	11.0 - 16.7	2,100
186+%	5,700	82.5	80.8 - 84.0	1,200	17.5	16.0 - 19.2	6,900

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	83.0	78.5 - 86.7	200	17.0	13.3 - 21.5	1,200
Normal birth weight	13,700	86.0	84.9 - 86.9	2,200	14.0	13.1 - 15.1	15,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	88.7	85.1 - 91.5	200	11.3	8.5 - 14.9	1,400
No	13,200	85.5	84.4 - 86.5	2,200	14.5	13.5 - 15.6	15,400

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,000	87.5	85.2 - 89.4	400	12.5	10.6 - 14.8	3,500
No	11,300	85.4	84.2 - 86.4	1,900	14.6	13.6 - 15.8	13,300

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,700	89.8	88.4 - 91.0	800	10.2	9.0 - 11.6	7,500
No	7,800	82.5	81.1 - 83.9	1,600	17.5	16.1 - 18.9	9,400

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	14,600	85.7	84.6 - 86.7	2,400	14.3	13.3 - 15.4	17,000

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,700	84.8	82.3 - 87.0	300	15.2	13.0 - 17.7	2,000
Honolulu	10,700	85.8	84.5 - 87.1	1,800	14.2	12.9 - 15.5	12,400
Kauai	700	90.0	86.6 - 92.6	100	10.0	7.4 - 13.4	800
Maui	1,500	84.1	81.6 - 86.4	300	15.9	13.6 - 18.4	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,100	84.0	81.5 - 86.2	600	16.0	13.8 - 18.5	3,700
Native Hawaiian	4,100	87.4	85.9 - 88.7	600	12.6	11.3 - 14.1	4,700
Chinese	500	79.9	73.5 - 85.1	100	20.1	14.9 - 26.5	600
Filipino	3,100	88.5	86.1 - 90.6	400	11.5	9.4 - 13.9	3,500
Japanese	1,700	76.3	72.2 - 80.0	500	23.7	20.0 - 27.8	2,200
Other	2,100	91.4	88.6 - 93.5	200	8.6	6.5 - 11.4	2,300

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	91.1	87.8 - 93.6	100	8.9	6.4 - 12.2	1,400
20-24	3,800	91.0	89.3 - 92.5	400	9.0	7.5 - 10.7	4,200
25-34	7,200	84.8	83.2 - 86.3	1,300	15.2	13.7 - 16.8	8,400
35+	2,300	78.1	75.0 - 80.9	600	21.9	19.1 - 25.0	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,700	86.0	84.6 - 87.4	1,200	14.0	12.6 - 15.4	8,900
Female	6,900	85.3	83.7 - 86.8	1,200	14.7	13.2 - 16.3	8,100

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,800	84.0	82.6 - 85.3	1,900	16.0	14.7 - 17.4	11,600
No	4,800	89.3	87.7 - 90.7	600	10.7	9.3 - 12.3	5,400

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	6,000	88.7	87.1 - 90.1	800	11.3	9.9 - 12.9	6,800
131-185%	2,300	85.2	82.1 - 87.9	400	14.8	12.1 - 17.9	2,600
186+%	5,100	83.0	81.1 - 84.8	1,000	17.0	15.2 - 18.9	6,100

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	83.8	79.1 - 87.6	200	16.2	12.4 - 20.9	1,100
Normal birth weight	13,600	85.9	84.8 - 86.9	2,200	14.1	13.1 - 15.2	15,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	86.9	83.0 - 90.0	200	13.1	10.0 - 17.0	1,400
No	13,100	85.6	84.5 - 86.7	2,200	14.4	13.3 - 15.5	15,300

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	86.1	83.6 - 88.3	500	13.9	11.7 - 16.4	3,400
No	11,300	85.6	84.4 - 86.7	1,900	14.4	13.3 - 15.6	13,200

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,600	90.0	88.5 - 91.2	700	10.0	8.8 - 11.5	7,400
No	7,800	82.4	80.8 - 83.9	1,700	17.6	16.1 - 19.2	9,500

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	14,400	86.1	85.1 - 87.2	2,300	13.9	12.8 - 14.9	16,700

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,700	86.7	84.7 - 88.4	300	13.3	11.6 - 15.3	1,900
Honolulu	10,600	86.0	84.6 - 87.3	1,700	14.0	12.7 - 15.4	12,300
Kauai	700	89.4	86.4 - 91.9	100	10.6	8.1 - 13.6	800
Maui	1,500	85.3	83.1 - 87.2	300	14.7	12.8 - 16.9	1,700

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,000	83.5	80.8 - 85.8	600	16.5	14.2 - 19.2	3,600
Native Hawaiian	4,000	88.2	87.1 - 89.2	500	11.8	10.8 - 12.9	4,500
Chinese	600	83.5	76.0 - 89.0	100	16.5	11.0 - 24.0	700
Filipino	3,200	89.1	86.6 - 91.2	400	10.9	8.8 - 13.4	3,600
Japanese	1,700	78.3	73.9 - 82.1	500	21.7	17.9 - 26.1	2,100
Other	2,000	89.9	86.7 - 92.4	200	10.1	7.6 - 13.3	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,400	92.0	89.1 - 94.2	100	8.0	5.8 - 10.9	1,500
20-24	3,700	90.8	89.1 - 92.2	400	9.2	7.8 - 10.9	4,100
25-34	7,100	86.1	84.5 - 87.5	1,200	13.9	12.5 - 15.5	8,300
35+	2,100	76.2	72.7 - 79.4	700	23.8	20.6 - 27.3	2,800

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,400	86.2	84.6 - 87.6	1,200	13.8	12.4 - 15.4	8,500
Female	7,000	86.1	84.6 - 87.6	1,100	13.9	12.4 - 15.4	8,200

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,500	84.2	82.7 - 85.5	1,800	15.8	14.5 - 17.3	11,200
No	4,900	90.2	88.7 - 91.5	500	9.8	8.5 - 11.3	5,500

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	6,000	89.4	87.9 - 90.7	700	10.6	9.3 - 12.1	6,700
131-185%	2,600	83.9	81.1 - 86.4	500	16.1	13.6 - 18.9	3,200
186+%	4,500	83.8	81.6 - 85.7	900	16.2	14.3 - 18.4	5,400

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Income (2000-2003)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	3,200	90.0	88.1 - 91.6	400	10.0	8.4 - 11.9	3,500
\$15,001-\$25,000	2,000	88.2	85.4 - 90.5	300	11.8	9.5 - 14.6	2,300
\$25,001-\$40,000	3,000	86.6	84.0 - 88.8	500	13.4	11.2 - 16.0	3,400
>=\$40,001	5,300	82.9	80.9 - 84.7	1,100	17.1	15.3 - 19.1	6,400

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	86.1	81.5 - 89.7	200	13.9	10.3 - 18.5	1,100
Normal birth weight	13,400	86.3	85.1 - 87.3	2,100	13.7	12.7 - 14.9	15,600

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	86.1	82.4 - 89.1	200	13.9	10.9 - 17.6	1,400
No	12,900	86.3	85.1 - 87.4	2,100	13.7	12.6 - 14.9	15,000

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	86.5	84.0 - 88.6	400	13.5	11.4 - 16.0	3,300
No	11,100	86.1	84.8 - 87.2	1,800	13.9	12.8 - 15.2	12,900

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,400	90.5	89.1 - 91.7	700	9.5	8.3 - 10.9	7,100
No	7,900	82.9	81.3 - 84.4	1,600	17.1	15.6 - 18.7	9,500

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	14,200	85.6	84.6 - 86.6	2,400	14.4	13.4 - 15.4	16,600

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,600	84.8	82.7 - 86.6	300	15.2	13.4 - 17.3	1,900
Honolulu	10,500	85.3	84.0 - 86.6	1,800	14.7	13.4 - 16.0	12,300
Kauai	600	90.0	87.0 - 92.4	100	10.0	7.6 - 13.0	700
Maui	1,500	86.7	84.5 - 88.6	200	13.3	11.4 - 15.5	1,700

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,000	82.1	79.6 - 84.4	700	17.9	15.6 - 20.4	3,700
Native Hawaiian	4,000	88.6	87.6 - 89.6	500	11.4	10.4 - 12.4	4,500
Chinese	500	80.1	72.7 - 85.9	100	19.9	14.1 - 27.3	700
Filipino	3,000	89.6	87.2 - 91.6	400	10.4	8.4 - 12.8	3,400
Japanese	1,600	77.1	73.0 - 80.7	500	22.9	19.3 - 27.0	2,100
Other	2,000	89.2	86.1 - 91.6	200	10.8	8.4 - 13.9	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,500	92.5	90.0 - 94.4	100	7.5	5.6 - 10.0	1,600
20-24	3,700	90.0	88.4 - 91.4	400	10.0	8.6 - 11.6	4,100
25-34	7,000	86.0	84.5 - 87.4	1,100	14.0	12.6 - 15.5	8,100
35+	2,000	73.9	70.5 - 77.0	700	26.1	23.0 - 29.5	2,700

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,300	86.2	84.8 - 87.5	1,200	13.8	12.5 - 15.2	8,400
Female	6,900	85.0	83.5 - 86.4	1,200	15.0	13.6 - 16.5	8,100

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,400	83.3	81.9 - 84.6	1,900	16.7	15.4 - 18.1	11,300
No	4,800	90.7	89.3 - 91.8	500	9.3	8.2 - 10.7	5,300

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,800	89.1	87.7 - 90.3	700	10.9	9.7 - 12.3	6,500
131-185%	2,600	85.5	82.9 - 87.7	400	14.5	12.3 - 17.1	3,100
186+%	4,600	81.4	79.4 - 83.3	1,000	18.6	16.7 - 20.6	5,600

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Income (2000-2003)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	3,100	89.3	87.4 - 90.9	400	10.7	9.1 - 12.6	3,400
\$15,001-\$25,000	2,200	88.3	85.8 - 90.3	300	11.7	9.7 - 14.2	2,400
\$25,001-\$40,000	3,000	87.2	84.9 - 89.2	400	12.8	10.8 - 15.1	3,500
>=\$40,001	5,200	81.4	79.5 - 83.2	1,200	18.6	16.8 - 20.5	6,400

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	86.6	82.3 - 89.9	100	13.4	10.1 - 17.7	1,100
Normal birth weight	13,200	85.6	84.5 - 86.6	2,200	14.4	13.4 - 15.5	15,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	86.1	82.7 - 88.9	200	13.9	11.1 - 17.3	1,400
No	12,700	85.7	84.6 - 86.8	2,100	14.3	13.2 - 15.4	14,800

PRAMS Health Indicator 3-Year Aggregate Report - Discuss breastfeeding with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	87.0	84.9 - 88.9	400	13.0	11.1 - 15.1	3,200
No	10,900	85.2	84.0 - 86.3	1,900	14.8	13.7 - 16.0	12,800

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,400	90.2	88.9 - 91.4	700	9.8	8.6 - 11.1	7,000
No	7,700	82.3	80.8 - 83.8	1,700	17.7	16.2 - 19.2	9,400