

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- for the State of Hawaii, for the Year(s)

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 3-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, income data does not exist for some 3-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration (grouped), among initiators	Breastfeeding (any) duration (grouped), among initiators

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 9/21/15

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- for the State of Hawaii, for the Year(s)

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012


Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2002	600	4.2	3.7 - 4.9	2,300	16.2	15.2 - 17.4	400	3.2	2.7 - 3.7	10,800	76.4	75.1 - 77.6	14,100
2001-2003	600	4.5	3.9 - 5.2	2,300	16.3	15.1 - 17.5	500	3.3	2.8 - 4.0	10,900	75.9	74.5 - 77.3	14,400
2002-2004	700	4.4	3.8 - 5.1	2,400	15.8	14.7 - 17.0	600	3.7	3.2 - 4.4	11,400	76.1	74.7 - 77.4	15,000
2003-2005	700	4.3	3.8 - 5.0	2,400	16.0	14.9 - 17.1	600	4.0	3.4 - 4.6	11,600	75.7	74.4 - 76.9	15,300
2004-2006	600	3.7	3.2 - 4.3	2,600	16.4	15.3 - 17.5	600	3.7	3.2 - 4.3	12,000	76.2	75.0 - 77.4	15,800
2005-2007	600	3.6	3.1 - 4.2	2,700	16.5	15.5 - 17.7	500	3.4	2.9 - 4.0	12,300	76.5	75.2 - 77.7	16,100
2006-2008	500	3.2	2.7 - 3.8	2,700	16.3	15.2 - 17.4	600	3.8	3.3 - 4.4	12,700	76.7	75.5 - 78.0	16,500
2007-2009	600	3.8	3.2 - 4.5	2,600	15.7	14.5 - 16.9	700	4.4	3.8 - 5.1	12,700	76.1	74.7 - 77.5	16,700
2008-2010	600	3.6	3.0 - 4.4	2,500	15.0	13.8 - 16.3	800	4.7	4.0 - 5.5	12,900	76.7	75.1 - 78.1	16,900
2009-2011	500	3.2	2.6 - 4.0	2,400	14.2	12.9 - 15.6	800	4.5	3.8 - 5.4	13,400	78.1	76.5 - 79.7	17,100
2010-2012	400	2.6	2.0 - 3.3	2,300	13.3	12.1 - 14.7	700	3.8	3.2 - 4.6	13,900	80.3	78.7 - 81.7	17,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- for the State of Hawaii, for the Year(s)

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	400	2.6	2.0 - 3.3	2,300	13.3	12.1 - 14.7	700	3.8	3.2 - 4.6	13,900	80.3	78.7 - 81.7	17,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	2.3	1.6 - 3.5	300	15.7	13.6 - 18.1	100	4.1	3.0 - 5.5	1,700	77.9	75.2 - 80.4	2,200
Honolulu	300	2.7	2.0 - 3.7	1,600	12.9	11.3 - 14.8	500	3.9	3.0 - 5.0	10,000	80.5	78.4 - 82.5	12,400
Kauai	<50	1.5	0.8 - 2.8	100	10.8	8.7 - 13.3	<50	3.2	2.1 - 4.7	700	84.5	81.7 - 87.0	800
Maui	100	2.7	1.9 - 3.8	300	14.3	12.4 - 16.4	100	3.6	2.7 - 4.7	1,500	79.5	77.1 - 81.7	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.6	1.5 - 4.4	400	10.1	7.8 - 12.9	200	4.3	2.9 - 6.2	3,500	83.0	79.7 - 85.9	4,200
Native Hawaiian	100	2.6	1.7 - 4.0	900	17.3	14.9 - 20.1	300	5.3	3.9 - 7.1	3,700	74.7	71.6 - 77.6	5,000
Chinese	<50	3.9	1.4 - 9.9	<50	5.4	2.5 - 11.3	<50	3.2	1.1 - 9.2	700	87.6	79.9 - 92.6	800
Filipino	100	2.6	1.6 - 4.3	600	19.5	16.1 - 23.3	100	3.7	2.3 - 5.7	2,300	74.2	70.1 - 77.9	3,100
Japanese	<50	0.4	0.1 - 1.1	100	6.9	4.4 - 10.8	<50	1.5	0.6 - 3.8	1,500	91.2	87.1 - 94.0	1,700
Other	100	3.5	2.0 - 6.1	300	10.0	7.2 - 13.6	100	2.3	1.2 - 4.4	2,200	84.3	80.1 - 87.7	2,600

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	4.3	2.1 - 8.7	300	31.2	24.6 - 38.7	100	5.3	2.7 - 9.9	600	59.2	51.6 - 66.5	1,000
20-24	100	2.9	1.8 - 4.6	700	18.1	15.1 - 21.5	100	2.8	1.7 - 4.4	3,000	76.2	72.6 - 79.6	3,900
25-34	200	2.4	1.7 - 3.3	1,100	11.5	9.9 - 13.3	400	4.3	3.4 - 5.6	7,600	81.8	79.7 - 83.8	9,300
35+	100	2.4	1.4 - 4.3	200	6.9	5.1 - 9.3	100	3.2	2.0 - 5.0	2,700	87.5	84.4 - 90.0	3,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.2	2.3 - 4.3	1,200	13.6	11.8 - 15.6	300	3.7	2.8 - 4.9	7,000	79.5	77.2 - 81.6	8,700
Female	200	2.0	1.4 - 2.9	1,100	13.0	11.3 - 15.0	300	3.9	3.0 - 5.1	6,900	81.0	78.8 - 83.0	8,600

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.5	1.8 - 3.4	1,100	10.5	9.1 - 12.1	400	3.7	2.9 - 4.7	9,100	83.4	81.4 - 85.1	10,900
No	200	2.8	2.0 - 4.0	1,200	18.1	15.8 - 20.6	300	4.1	3.1 - 5.5	4,800	75.0	72.3 - 77.6	6,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	200	3.6	2.5 - 5.0	1,100	17.6	15.3 - 20.1	200	3.7	2.6 - 5.1	4,500	75.2	72.4 - 77.8	6,000
131-185%	100	2.9	1.5 - 5.4	400	14.1	10.9 - 17.9	100	4.7	3.0 - 7.1	2,200	78.4	74.0 - 82.2	2,800
186+%	100	1.8	1.1 - 2.8	600	8.8	7.2 - 10.7	300	3.7	2.7 - 5.0	6,300	85.7	83.5 - 87.7	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	2.9	1.9 - 4.2	200	16.0	13.7 - 18.7	100	5.4	4.1 - 7.2	900	75.7	72.7 - 78.5	1,200
Normal birth weight	400	2.6	2.0 - 3.3	2,100	13.1	11.8 - 14.6	600	3.7	3.0 - 4.6	13,000	80.6	78.9 - 82.2	16,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	5.5	2.7 - 10.8	200	25.9	19.5 - 33.6	100	7.3	4.0 - 13.0	500	61.3	53.1 - 68.9	900
No	400	2.5	1.9 - 3.2	2,000	12.6	11.3 - 14.0	600	3.7	3.0 - 4.5	13,100	81.2	79.6 - 82.7	16,100

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.4	2.8 - 6.8	700	21.1	17.7 - 24.8	200	6.1	4.2 - 8.7	2,100	68.5	64.2 - 72.5	3,100
No	300	2.2	1.7 - 3.0	1,600	11.6	10.2 - 13.0	500	3.4	2.7 - 4.3	11,500	82.8	81.1 - 84.4	13,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.6	2.7 - 4.9	1,300	17.6	15.5 - 19.8	300	4.0	3.0 - 5.2	5,700	74.9	72.3 - 77.3	7,600
No	200	1.8	1.2 - 2.7	1,000	10.0	8.5 - 11.7	400	3.7	2.8 - 4.8	8,100	84.6	82.6 - 86.4	9,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	3.2	2.6 - 4.0	2,400	14.2	12.9 - 15.6	800	4.5	3.8 - 5.4	13,400	78.1	76.5 - 79.7	17,100

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.0	2.9 - 5.3	400	16.2	14.1 - 18.7	100	3.7	2.7 - 5.1	1,700	76.1	73.4 - 78.6	2,200
Honolulu	400	3.2	2.4 - 4.2	1,700	13.9	12.2 - 15.8	600	4.7	3.7 - 5.9	9,500	78.2	76.0 - 80.2	12,200
Kauai	<50	1.3	0.7 - 2.5	100	11.4	9.3 - 13.9	<50	3.9	2.7 - 5.5	700	83.4	80.6 - 85.9	800
Maui	100	3.2	2.4 - 4.4	300	14.5	12.6 - 16.6	100	4.5	3.5 - 5.8	1,500	77.8	75.4 - 80.0	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.8	1.7 - 4.6	400	10.2	8.0 - 13.0	200	5.7	4.1 - 7.8	3,400	81.3	78.0 - 84.3	4,200
Native Hawaiian	200	3.9	2.8 - 5.6	900	18.4	15.9 - 21.1	300	5.0	3.7 - 6.8	3,700	72.6	69.5 - 75.6	5,000
Chinese	<50	2.0	0.5 - 7.7	<50	6.5	3.1 - 13.1	<50	2.6	0.9 - 7.5	600	89.0	81.5 - 93.7	700
Filipino	100	2.7	1.8 - 4.0	600	19.2	15.9 - 22.9	100	4.2	2.9 - 6.3	2,300	73.9	69.9 - 77.6	3,100
Japanese	<50	1.5	0.6 - 3.9	100	7.5	4.7 - 11.5	<50	2.4	1.1 - 5.1	1,500	88.6	84.0 - 92.0	1,700
Other	100	4.6	2.7 - 7.6	300	12.9	9.7 - 17.0	100	3.8	2.2 - 6.4	1,900	78.8	74.0 - 82.9	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	<50	4.0	2.1 - 7.4	400	32.2	26.0 - 39.0	100	6.3	3.6 - 10.9	700	57.5	50.6 - 64.2	1,200
20-24	100	3.6	2.4 - 5.6	700	18.4	15.5 - 21.6	200	3.9	2.6 - 5.6	2,900	74.1	70.5 - 77.5	4,000
25-34	300	2.9	2.1 - 3.9	1,000	11.7	10.1 - 13.5	400	4.8	3.7 - 6.1	7,200	80.7	78.5 - 82.7	8,900
35+	100	3.2	2.0 - 5.3	300	8.7	6.5 - 11.5	100	3.9	2.6 - 6.0	2,500	84.1	80.6 - 87.1	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.3	2.4 - 4.4	1,200	14.1	12.3 - 16.1	400	4.9	3.9 - 6.2	6,800	77.7	75.4 - 79.9	8,800
Female	300	3.1	2.3 - 4.2	1,200	14.3	12.5 - 16.3	300	4.1	3.2 - 5.3	6,500	78.5	76.2 - 80.7	8,300

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.8	2.1 - 3.8	1,100	10.7	9.2 - 12.3	500	4.5	3.6 - 5.6	8,800	82.1	80.1 - 83.9	10,700
No	200	3.8	2.8 - 5.2	1,300	20.0	17.7 - 22.6	300	4.6	3.4 - 6.1	4,600	71.6	68.7 - 74.2	6,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.3	3.2 - 5.8	1,100	18.9	16.6 - 21.5	200	4.1	3.0 - 5.6	4,300	72.6	69.7 - 75.4	5,900
131-185%	100	4.3	2.5 - 7.1	400	15.0	11.8 - 18.9	100	4.9	3.2 - 7.4	2,000	75.8	71.3 - 79.8	2,700
186+%	200	2.2	1.5 - 3.3	700	9.1	7.5 - 11.0	300	4.7	3.6 - 6.1	6,200	84.1	81.7 - 86.1	7,400

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	4.3	2.8 - 6.6	600	21.0	17.6 - 24.9	100	3.8	2.4 - 6.1	2,100	70.8	66.6 - 74.7	3,000
\$10,000-\$24,999	100	4.5	3.0 - 6.6	500	16.0	13.2 - 19.4	100	4.7	3.2 - 6.8	2,400	74.8	71.0 - 78.3	3,200
\$25,000-\$49,999	200	3.8	2.4 - 6.0	600	14.5	11.8 - 17.6	200	5.4	3.8 - 7.6	3,000	76.3	72.6 - 79.6	4,000
>=\$50,000	100	1.9	1.2 - 3.0	500	8.5	6.8 - 10.6	300	4.2	3.1 - 5.7	5,100	85.3	82.8 - 87.5	5,900

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	3.7	2.6 - 5.3	200	17.3	14.9 - 19.9	100	6.0	4.6 - 7.8	900	73.1	70.0 - 75.9	1,200
Normal birth weight	500	3.2	2.5 - 4.0	2,200	13.9	12.6 - 15.4	700	4.4	3.6 - 5.3	12,500	78.5	76.8 - 80.1	15,900

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.5	5.1 - 14.1	300	25.9	20.1 - 32.6	100	5.6	3.0 - 10.1	700	60.0	52.6 - 66.9	1,100
No	500	2.9	2.3 - 3.6	2,100	13.2	11.9 - 14.6	700	4.4	3.6 - 5.3	12,600	79.6	77.9 - 81.1	15,800

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.4	3.7 - 7.8	800	22.7	19.4 - 26.5	200	5.8	4.1 - 8.2	2,200	66.1	61.9 - 70.0	3,300
No	400	2.7	2.1 - 3.5	1,600	11.9	10.5 - 13.3	600	4.1	3.3 - 5.1	11,000	81.3	79.6 - 82.9	13,500

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.5	3.4 - 5.9	1,300	18.6	16.5 - 21.0	300	4.5	3.5 - 5.9	5,200	72.3	69.7 - 74.8	7,200
No	200	2.3	1.6 - 3.2	1,000	10.9	9.3 - 12.6	400	4.4	3.4 - 5.6	7,900	82.5	80.4 - 84.4	9,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.6	3.0 - 4.4	2,500	15.0	13.8 - 16.3	800	4.7	4.0 - 5.5	12,900	76.7	75.1 - 78.1	16,900

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.4	4.0 - 7.2	300	16.0	13.6 - 18.8	100	2.9	2.0 - 4.3	1,600	75.7	72.6 - 78.6	2,200
Honolulu	400	3.4	2.6 - 4.5	1,800	15.2	13.6 - 17.0	600	5.0	4.1 - 6.1	9,200	76.3	74.3 - 78.2	12,000
Kauai	<50	1.5	0.7 - 3.2	100	12.7	9.7 - 16.4	<50	5.9	3.8 - 9.2	600	79.9	75.5 - 83.6	800
Maui	100	3.5	2.5 - 5.0	300	13.7	11.6 - 16.1	100	4.3	3.1 - 5.8	1,500	78.5	75.6 - 81.1	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.0	1.9 - 4.8	400	10.6	8.5 - 13.0	300	6.7	5.1 - 8.9	3,300	79.7	76.5 - 82.5	4,100
Native Hawaiian	200	4.6	3.4 - 6.3	900	19.1	16.6 - 21.8	200	5.0	3.8 - 6.6	3,500	71.3	68.2 - 74.2	4,900
Chinese	<50	0.9	0.4 - 2.3	100	8.6	4.8 - 14.8	<50	3.2	1.3 - 7.7	600	87.3	80.6 - 91.9	700
Filipino	100	3.7	2.6 - 5.2	600	19.8	16.7 - 23.3	100	4.1	2.8 - 6.2	2,200	72.4	68.6 - 75.9	3,000
Japanese	<50	2.3	1.1 - 4.7	100	7.2	4.9 - 10.5	<50	2.7	1.4 - 5.1	1,500	87.8	83.8 - 90.9	1,700
Other	100	4.0	2.3 - 6.8	400	16.0	12.6 - 20.0	100	3.2	1.8 - 5.6	1,800	76.8	72.3 - 80.8	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	4.9	2.8 - 8.3	400	26.4	21.1 - 32.4	100	6.1	3.5 - 10.4	800	62.6	56.2 - 68.6	1,300
20-24	100	3.9	2.6 - 5.8	800	20.6	17.8 - 23.8	200	5.9	4.4 - 7.9	2,700	69.5	66.0 - 72.9	3,800
25-34	300	3.0	2.2 - 4.0	1,100	12.4	10.9 - 14.2	400	4.4	3.5 - 5.6	6,900	80.2	78.1 - 82.1	8,700
35+	100	4.4	3.0 - 6.5	300	10.4	8.1 - 13.3	100	3.5	2.3 - 5.3	2,500	81.7	78.3 - 84.7	3,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.4	2.6 - 4.5	1,300	15.3	13.6 - 17.1	500	5.3	4.3 - 6.5	6,700	76.0	73.8 - 78.1	8,800
Female	300	3.8	2.9 - 4.9	1,200	14.8	13.1 - 16.7	300	4.1	3.2 - 5.3	6,300	77.4	75.1 - 79.4	8,100

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.8	2.2 - 3.7	1,300	12.4	11.0 - 13.9	500	4.5	3.7 - 5.5	8,500	80.3	78.4 - 82.0	10,600
No	300	4.9	3.7 - 6.4	1,200	19.6	17.4 - 22.0	300	5.1	3.9 - 6.5	4,400	70.5	67.8 - 73.1	6,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.7	3.5 - 6.2	1,100	19.8	17.5 - 22.3	200	4.2	3.1 - 5.6	4,100	71.3	68.5 - 74.0	5,800
131-185%	100	4.3	2.6 - 7.1	400	14.6	11.5 - 18.2	100	4.8	3.2 - 7.2	2,000	76.3	72.0 - 80.1	2,600
186+%	200	2.6	1.8 - 3.6	800	10.3	8.8 - 12.1	400	5.1	4.0 - 6.4	6,000	82.1	79.9 - 84.1	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	4.9	3.3 - 7.2	600	22.1	18.6 - 25.9	100	4.3	2.8 - 6.5	1,900	68.8	64.6 - 72.8	2,800
\$10,000-\$24,999	200	4.9	3.4 - 7.0	500	16.5	13.9 - 19.5	200	5.2	3.7 - 7.2	2,400	73.4	69.9 - 76.7	3,300
\$25,000-\$49,999	200	4.2	2.8 - 6.3	600	15.4	12.9 - 18.3	200	4.9	3.5 - 6.9	2,900	75.5	72.1 - 78.6	3,800
>=\$50,000	100	2.0	1.3 - 3.0	600	9.6	7.9 - 11.6	300	4.5	3.4 - 6.0	4,900	83.8	81.4 - 86.0	5,900

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	4.3	2.8 - 6.5	200	16.1	13.2 - 19.4	100	5.2	3.7 - 7.3	900	74.4	70.6 - 77.9	1,200
Normal birth weight	600	3.5	2.9 - 4.4	2,400	15.0	13.7 - 16.4	700	4.7	3.9 - 5.6	12,100	76.8	75.2 - 78.4	15,700

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.0	4.8 - 12.9	300	24.8	19.7 - 30.7	100	6.4	3.9 - 10.3	700	60.8	54.2 - 67.0	1,200
No	500	3.3	2.6 - 4.0	2,200	14.2	12.9 - 15.5	700	4.5	3.8 - 5.4	12,000	78.0	76.4 - 79.5	15,400

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	6.0	4.2 - 8.6	700	22.4	19.2 - 26.0	200	7.4	5.5 - 9.7	2,100	64.2	60.2 - 68.0	3,200
No	400	3.0	2.4 - 3.8	1,800	13.2	11.9 - 14.6	500	4.0	3.3 - 4.9	10,700	79.8	78.1 - 81.3	13,400

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.2	4.1 - 6.7	1,400	19.7	17.7 - 22.0	300	4.7	3.7 - 6.1	4,800	70.3	67.7 - 72.7	6,900
No	200	2.5	1.9 - 3.4	1,100	11.8	10.3 - 13.4	400	4.6	3.7 - 5.7	7,900	81.1	79.1 - 82.9	9,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.8	3.2 - 4.5	2,600	15.7	14.5 - 16.9	700	4.4	3.8 - 5.1	12,700	76.1	74.7 - 77.5	16,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.5	4.0 - 7.6	300	14.0	11.6 - 16.9	100	2.8	1.8 - 4.3	1,700	77.7	74.3 - 80.7	2,100
Honolulu	400	3.5	2.8 - 4.4	1,900	16.3	14.9 - 17.9	500	4.6	3.8 - 5.6	8,900	75.6	73.7 - 77.3	11,700
Kauai	<50	1.9	0.9 - 4.0	100	13.9	10.3 - 18.4	100	6.4	4.0 - 9.9	700	77.8	72.7 - 82.3	800
Maui	100	4.2	2.9 - 6.1	300	14.4	11.8 - 17.4	100	4.2	3.0 - 6.0	1,500	77.1	73.7 - 80.3	2,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.0	2.0 - 4.6	400	11.2	9.4 - 13.4	200	5.4	4.0 - 7.2	3,200	80.3	77.6 - 82.8	4,000
Native Hawaiian	300	5.4	4.1 - 7.1	1,000	21.4	19.0 - 24.1	200	4.5	3.4 - 6.0	3,200	68.6	65.6 - 71.5	4,700
Chinese	<50	1.6	0.8 - 3.1	100	8.4	5.2 - 13.2	<50	2.3	1.3 - 3.8	600	87.7	82.9 - 91.3	600
Filipino	100	3.4	2.4 - 4.8	600	18.9	16.2 - 21.9	200	5.1	3.6 - 7.0	2,200	72.6	69.3 - 75.7	3,000
Japanese	<50	2.5	1.3 - 4.8	200	8.3	6.0 - 11.5	100	2.8	1.6 - 4.8	1,600	86.4	82.6 - 89.4	1,900
Other	100	3.7	2.2 - 6.0	400	15.4	12.5 - 18.9	100	3.5	2.1 - 5.8	1,900	77.4	73.4 - 80.9	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	7.3	4.7 - 11.0	400	25.0	20.2 - 30.5	100	5.0	2.9 - 8.3	900	62.8	56.9 - 68.3	1,500
20-24	200	4.0	2.8 - 5.8	800	22.0	19.4 - 24.9	200	5.8	4.4 - 7.7	2,600	68.1	64.8 - 71.2	3,800
25-34	300	3.1	2.3 - 4.0	1,100	13.1	11.7 - 14.7	300	4.0	3.1 - 5.0	6,700	79.9	78.0 - 81.6	8,400
35+	100	3.7	2.5 - 5.5	300	10.3	8.1 - 13.0	100	3.5	2.4 - 5.1	2,500	82.5	79.4 - 85.2	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.2	2.5 - 4.1	1,400	16.0	14.5 - 17.8	400	4.9	4.0 - 6.0	6,600	75.8	73.8 - 77.7	8,800
Female	300	4.4	3.5 - 5.5	1,200	15.3	13.7 - 17.1	300	3.8	3.0 - 4.8	6,100	76.5	74.4 - 78.4	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.8	2.2 - 3.6	1,300	12.6	11.3 - 14.0	400	4.0	3.2 - 4.8	8,500	80.6	79.0 - 82.2	10,500
No	300	5.4	4.2 - 6.8	1,300	21.0	18.8 - 23.3	300	5.1	4.0 - 6.5	4,200	68.5	65.9 - 71.0	6,200

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.2	4.1 - 6.7	1,300	21.6	19.4 - 24.0	200	3.9	2.9 - 5.1	4,000	69.3	66.7 - 71.8	5,800
131-185%	100	3.5	2.1 - 5.8	400	14.5	11.8 - 17.8	100	4.5	3.0 - 6.7	1,900	77.5	73.7 - 80.9	2,500
186+%	200	2.9	2.1 - 3.9	800	10.6	9.2 - 12.2	300	4.7	3.8 - 5.9	5,800	81.8	79.8 - 83.6	7,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	5.8	4.1 - 8.2	600	24.0	20.6 - 27.8	100	4.4	2.9 - 6.5	1,800	65.8	61.7 - 69.7	2,700
\$10,000-\$24,999	200	5.3	3.8 - 7.3	600	18.2	15.6 - 21.0	200	4.5	3.3 - 6.2	2,500	72.0	68.7 - 75.1	3,400
\$25,000-\$49,999	100	3.6	2.4 - 5.5	600	15.9	13.7 - 18.5	200	4.2	3.0 - 5.8	2,800	76.2	73.2 - 79.0	3,700
>=\$50,000	100	2.3	1.7 - 3.2	600	9.5	8.0 - 11.3	300	4.3	3.3 - 5.6	4,800	83.8	81.7 - 85.8	5,800

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	4.5	2.8 - 7.0	200	18.7	15.2 - 22.8	<50	3.9	2.5 - 6.0	900	73.0	68.5 - 77.0	1,200
Normal birth weight	600	3.7	3.1 - 4.5	2,400	15.5	14.3 - 16.7	700	4.4	3.8 - 5.2	11,800	76.4	74.9 - 77.8	15,500

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.5	4.7 - 11.7	400	28.8	23.7 - 34.4	100	7.7	5.0 - 11.7	700	56.0	50.0 - 61.9	1,300
No	500	3.5	2.9 - 4.2	2,200	14.5	13.4 - 15.8	600	4.1	3.5 - 4.8	11,800	77.9	76.5 - 79.3	15,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.1	3.5 - 7.3	800	25.5	22.3 - 28.9	200	6.9	5.2 - 9.1	1,900	62.5	58.8 - 66.1	3,100
No	500	3.5	2.8 - 4.2	1,800	13.4	12.2 - 14.6	500	3.8	3.1 - 4.5	10,500	79.4	77.9 - 80.8	13,300

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.3	4.2 - 6.7	1,400	21.2	19.2 - 23.4	300	4.2	3.3 - 5.4	4,600	69.3	66.8 - 71.6	6,600
No	300	2.8	2.2 - 3.6	1,200	11.9	10.6 - 13.3	400	4.5	3.6 - 5.5	7,800	80.8	79.1 - 82.5	9,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	3.2	2.7 - 3.8	2,700	16.3	15.2 - 17.4	600	3.8	3.3 - 4.4	12,700	76.7	75.5 - 78.0	16,500

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.7	3.1 - 6.9	300	15.0	12.1 - 18.3	<50	2.1	1.2 - 3.8	1,600	78.2	74.5 - 81.6	2,100
Honolulu	400	3.0	2.5 - 3.7	2,000	16.7	15.5 - 18.1	500	4.0	3.4 - 4.7	9,000	76.3	74.8 - 77.7	11,800
Kauai	<50	1.1	0.3 - 3.4	100	14.6	10.5 - 20.0	100	6.1	3.6 - 10.1	700	78.2	72.2 - 83.2	800
Maui	100	3.5	2.2 - 5.7	300	15.6	12.5 - 19.2	100	3.6	2.2 - 5.7	1,400	77.3	73.3 - 80.9	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.6	1.8 - 3.8	500	12.9	11.0 - 15.0	200	5.1	4.0 - 6.6	3,100	79.3	76.8 - 81.7	3,900
Native Hawaiian	200	4.6	3.5 - 6.1	1,000	22.2	19.8 - 24.9	200	3.8	2.8 - 5.1	3,100	69.4	66.5 - 72.1	4,500
Chinese	<50	2.4	1.4 - 4.0	100	9.0	7.0 - 11.7	<50	2.8	1.7 - 4.4	500	85.8	82.8 - 88.4	600
Filipino	100	2.9	2.0 - 4.2	600	19.0	16.6 - 21.6	100	4.4	3.3 - 6.0	2,200	73.7	70.8 - 76.5	3,000
Japanese	<50	1.9	1.0 - 3.7	200	8.4	6.3 - 11.1	<50	1.8	1.0 - 3.2	1,700	87.9	84.8 - 90.4	2,000
Other	100	3.0	1.9 - 4.6	400	15.6	13.0 - 18.7	100	2.9	1.9 - 4.4	2,000	78.5	75.1 - 81.5	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.5	4.2 - 10.0	300	25.5	20.9 - 30.7	<50	3.3	1.8 - 6.0	900	64.6	59.1 - 69.8	1,400
20-24	100	3.1	2.2 - 4.4	1,000	24.6	22.0 - 27.5	200	5.1	3.9 - 6.6	2,600	67.2	64.1 - 70.0	3,900
25-34	200	2.9	2.2 - 3.7	1,100	13.4	12.1 - 14.9	300	3.5	2.8 - 4.3	6,700	80.2	78.5 - 81.8	8,300
35+	100	2.6	1.8 - 3.9	300	9.1	7.4 - 11.2	100	3.3	2.3 - 4.6	2,600	85.0	82.5 - 87.2	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	2.9	2.3 - 3.7	1,500	17.6	16.0 - 19.2	400	4.5	3.7 - 5.4	6,400	75.0	73.2 - 76.7	8,600
Female	300	3.4	2.7 - 4.3	1,200	14.9	13.4 - 16.4	200	3.1	2.4 - 3.9	6,200	78.7	76.9 - 80.3	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.5	2.0 - 3.2	1,400	12.8	11.7 - 14.1	400	3.4	2.8 - 4.1	8,600	81.3	79.8 - 82.6	10,600
No	300	4.3	3.4 - 5.5	1,300	22.4	20.3 - 24.6	300	4.6	3.6 - 5.7	4,100	68.7	66.3 - 71.1	6,000

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.6	3.6 - 5.8	1,300	23.0	20.9 - 25.2	200	3.4	2.6 - 4.4	4,000	69.1	66.6 - 71.4	5,800
131-185%	100	2.4	1.5 - 4.0	300	13.5	11.0 - 16.4	100	4.2	2.8 - 6.1	1,800	79.9	76.5 - 82.9	2,300
186+%	200	2.4	1.8 - 3.1	800	11.4	10.1 - 12.9	300	4.0	3.3 - 5.0	6,000	82.2	80.5 - 83.7	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	5.4	3.9 - 7.5	700	25.7	22.5 - 29.3	100	3.4	2.3 - 5.1	1,700	65.4	61.6 - 69.0	2,700
\$10,000-\$24,999	100	4.3	3.1 - 5.9	700	19.4	16.9 - 22.2	100	3.8	2.7 - 5.2	2,500	72.5	69.5 - 75.4	3,400
\$25,000-\$49,999	100	2.4	1.7 - 3.6	700	17.8	15.6 - 20.3	200	4.1	3.0 - 5.5	2,900	75.7	73.0 - 78.2	3,800
>=\$50,000	100	2.1	1.5 - 2.9	500	8.5	7.2 - 10.0	200	3.8	2.9 - 4.8	4,800	85.7	83.9 - 87.3	5,600

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	4.4	2.6 - 7.2	200	17.4	13.6 - 22.0	<50	3.2	1.8 - 5.5	900	75.0	70.0 - 79.4	1,200
Normal birth weight	500	3.1	2.6 - 3.7	2,500	16.2	15.1 - 17.3	600	3.9	3.3 - 4.5	11,800	76.9	75.6 - 78.2	15,300

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	5.9	3.7 - 9.1	300	27.8	23.0 - 33.1	100	7.4	5.0 - 10.8	700	59.0	53.3 - 64.4	1,200
No	400	2.9	2.5 - 3.5	2,300	15.3	14.2 - 16.4	500	3.6	3.0 - 4.2	11,800	78.2	76.9 - 79.4	15,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.0	2.8 - 5.7	800	25.2	22.2 - 28.4	200	6.5	5.0 - 8.4	2,000	64.4	60.9 - 67.7	3,100
No	400	3.0	2.4 - 3.6	1,900	14.2	13.1 - 15.4	400	3.3	2.7 - 3.9	10,500	79.6	78.2 - 80.9	13,200

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.3	3.4 - 5.4	1,400	21.7	19.7 - 23.7	300	3.8	3.0 - 4.8	4,700	70.3	68.1 - 72.5	6,700
No	200	2.3	1.8 - 3.0	1,200	12.5	11.3 - 13.8	400	3.9	3.2 - 4.6	7,800	81.3	79.8 - 82.8	9,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.6	3.1 - 4.2	2,700	16.5	15.5 - 17.7	500	3.4	2.9 - 4.0	12,300	76.5	75.2 - 77.7	16,100

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.5	2.2 - 5.6	300	15.0	12.1 - 18.3	<50	2.2	1.2 - 3.9	1,600	79.3	75.6 - 82.6	2,000
Honolulu	400	3.7	3.1 - 4.4	1,900	16.4	15.2 - 17.8	400	3.7	3.1 - 4.4	8,800	76.2	74.6 - 77.6	11,600
Kauai	<50	0.5	0.1 - 3.5	100	14.9	10.6 - 20.4	<50	3.5	1.7 - 6.9	600	81.1	75.2 - 85.9	800
Maui	100	4.4	2.8 - 6.8	300	19.7	16.3 - 23.6	<50	2.8	1.6 - 4.9	1,300	73.1	68.8 - 77.0	1,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.1	2.2 - 4.4	400	11.8	9.9 - 13.9	200	4.4	3.3 - 5.8	3,000	80.7	78.2 - 83.0	3,700
Native Hawaiian	200	5.5	4.2 - 7.1	1,000	22.9	20.4 - 25.6	100	3.3	2.4 - 4.5	3,000	68.4	65.5 - 71.2	4,400
Chinese	<50	2.4	1.4 - 4.1	100	9.4	7.3 - 12.1	<50	2.5	1.6 - 4.1	500	85.6	82.5 - 88.2	600
Filipino	100	3.2	2.2 - 4.5	600	20.7	18.2 - 23.4	100	4.6	3.4 - 6.2	2,100	71.5	68.6 - 74.3	3,000
Japanese	<50	1.2	0.6 - 2.6	200	8.9	6.8 - 11.5	<50	1.2	0.6 - 2.5	1,800	88.6	85.8 - 91.0	2,100
Other	100	3.8	2.6 - 5.6	400	15.6	12.9 - 18.6	100	2.7	1.7 - 4.3	1,900	77.9	74.5 - 81.0	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.6	4.3 - 10.0	400	27.7	23.1 - 33.0	<50	3.3	1.8 - 5.9	800	62.3	56.8 - 67.5	1,400
20-24	200	4.2	3.1 - 5.6	1,000	25.0	22.4 - 27.7	200	3.9	2.9 - 5.3	2,600	66.9	64.0 - 69.8	3,900
25-34	300	3.5	2.8 - 4.3	1,000	13.2	11.9 - 14.7	300	3.4	2.7 - 4.2	6,300	80.0	78.3 - 81.6	7,900
35+	100	1.8	1.1 - 2.9	300	8.8	7.1 - 10.9	100	2.9	2.0 - 4.2	2,500	86.5	84.1 - 88.6	2,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.4	2.7 - 4.2	1,400	17.4	15.9 - 19.1	300	4.2	3.5 - 5.1	6,200	75.0	73.2 - 76.7	8,300
Female	300	3.9	3.1 - 4.8	1,200	15.6	14.1 - 17.2	200	2.6	2.0 - 3.3	6,100	78.0	76.2 - 79.7	7,800

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.0	2.4 - 3.7	1,200	11.9	10.8 - 13.1	300	3.0	2.5 - 3.7	8,500	82.1	80.7 - 83.4	10,300
No	300	4.8	3.8 - 6.0	1,400	24.8	22.6 - 27.1	200	4.1	3.2 - 5.2	3,800	66.4	63.9 - 68.7	5,800

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.4	4.4 - 6.8	1,300	24.0	21.9 - 26.3	200	3.1	2.4 - 4.1	3,800	67.4	65.0 - 69.8	5,600
131-185%	100	3.4	2.1 - 5.3	300	15.0	12.3 - 18.3	100	4.1	2.7 - 6.1	1,600	77.5	73.8 - 80.8	2,100
186+%	200	2.4	1.8 - 3.2	800	10.8	9.5 - 12.2	300	3.5	2.8 - 4.3	6,100	83.4	81.7 - 84.9	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	6.6	4.9 - 8.8	700	27.3	24.0 - 30.9	100	3.7	2.5 - 5.4	1,600	62.4	58.6 - 66.0	2,600
\$10,000-\$24,999	200	5.2	3.9 - 6.9	700	20.4	17.9 - 23.2	100	2.9	2.0 - 4.2	2,300	71.5	68.4 - 74.4	3,300
\$25,000-\$49,999	100	2.6	1.8 - 3.8	600	17.0	14.9 - 19.5	100	3.5	2.6 - 4.8	2,900	76.8	74.1 - 79.3	3,800
>=\$50,000	100	2.1	1.5 - 2.9	400	8.1	6.8 - 9.5	200	3.5	2.7 - 4.6	4,600	86.3	84.5 - 87.9	5,400

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	4.7	2.9 - 7.6	200	18.0	14.2 - 22.6	<50	3.5	2.0 - 5.8	900	73.8	68.8 - 78.2	1,200
Normal birth weight	500	3.5	3.0 - 4.1	2,400	16.4	15.3 - 17.6	500	3.4	2.9 - 4.0	11,400	76.7	75.3 - 77.9	14,800

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	9.5	6.6 - 13.4	300	27.6	22.7 - 33.0	100	5.9	3.7 - 9.1	700	57.1	51.4 - 62.6	1,200
No	500	3.1	2.6 - 3.7	2,300	15.6	14.5 - 16.8	500	3.3	2.8 - 3.8	11,400	78.1	76.8 - 79.3	14,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.7	4.3 - 7.5	800	25.9	22.9 - 29.1	200	5.6	4.2 - 7.4	1,900	62.8	59.4 - 66.1	3,100
No	400	3.1	2.5 - 3.7	1,800	14.2	13.1 - 15.4	400	2.9	2.4 - 3.5	10,200	79.8	78.5 - 81.1	12,700

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.9	3.9 - 6.1	1,400	21.7	19.8 - 23.7	200	3.4	2.7 - 4.4	4,600	70.0	67.8 - 72.1	6,500
No	200	2.5	1.9 - 3.1	1,200	12.9	11.7 - 14.2	300	3.3	2.7 - 4.1	7,500	81.3	79.8 - 82.8	9,200

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.7	3.2 - 4.3	2,600	16.4	15.3 - 17.5	600	3.7	3.2 - 4.3	12,000	76.2	75.0 - 77.4	15,800

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.6	2.3 - 5.6	300	16.4	13.5 - 19.8	<50	1.9	1.0 - 3.5	1,500	78.1	74.4 - 81.5	1,900
Honolulu	400	3.8	3.2 - 4.5	1,800	15.9	14.7 - 17.2	500	4.0	3.4 - 4.7	8,800	76.3	74.9 - 77.7	11,500
Kauai	<50	1.8	0.7 - 4.6	100	14.8	10.6 - 20.4	<50	2.2	0.9 - 5.6	600	81.1	75.2 - 85.9	700
Maui	100	4.6	3.1 - 6.8	300	20.3	16.9 - 24.1	100	4.1	2.7 - 6.3	1,200	71.0	66.8 - 74.8	1,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.4	1.6 - 3.5	500	12.4	10.6 - 14.5	200	4.4	3.3 - 5.8	3,000	80.8	78.3 - 83.0	3,700
Native Hawaiian	200	5.2	4.0 - 6.7	900	22.1	19.7 - 24.8	200	4.0	3.0 - 5.4	2,900	68.6	65.7 - 71.4	4,200
Chinese	<50	2.1	1.3 - 3.4	100	9.4	7.5 - 11.8	<50	3.0	1.9 - 4.5	500	85.5	82.7 - 87.9	600
Filipino	200	5.2	4.0 - 6.7	600	20.0	17.7 - 22.6	100	4.4	3.3 - 5.9	2,100	70.3	67.5 - 73.0	3,000
Japanese	<50	1.8	1.0 - 3.2	200	8.3	6.3 - 10.7	<50	1.8	1.0 - 3.2	1,800	88.1	85.3 - 90.5	2,100
Other	100	3.5	2.3 - 5.3	400	16.5	13.8 - 19.7	100	2.8	1.8 - 4.3	1,700	77.2	73.7 - 80.3	2,300

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.8	3.7 - 8.8	400	29.4	24.6 - 34.6	<50	2.9	1.6 - 5.2	800	62.0	56.6 - 67.1	1,200
20-24	200	4.9	3.7 - 6.3	1,000	25.4	22.8 - 28.1	200	4.2	3.1 - 5.6	2,500	65.6	62.7 - 68.4	3,900
25-34	300	3.5	2.8 - 4.3	1,000	12.2	11.0 - 13.6	300	4.0	3.3 - 4.8	6,300	80.3	78.7 - 81.9	7,800
35+	100	2.0	1.3 - 3.1	300	9.7	8.0 - 11.8	100	2.7	1.8 - 3.9	2,500	85.5	83.2 - 87.6	2,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.0	3.3 - 4.8	1,400	16.9	15.4 - 18.4	400	4.6	3.9 - 5.5	6,100	74.6	72.8 - 76.2	8,200
Female	300	3.5	2.8 - 4.4	1,200	15.8	14.3 - 17.4	200	2.7	2.1 - 3.4	5,900	78.0	76.2 - 79.7	7,600

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.0	2.4 - 3.6	1,300	12.4	11.3 - 13.6	400	3.7	3.1 - 4.3	8,500	81.0	79.6 - 82.3	10,500
No	300	5.3	4.2 - 6.6	1,300	24.3	22.1 - 26.6	200	3.8	2.9 - 4.8	3,500	66.7	64.2 - 69.1	5,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.7	4.6 - 6.9	1,300	23.6	21.5 - 25.8	100	2.7	2.0 - 3.6	3,700	68.0	65.6 - 70.3	5,500
131-185%	100	3.3	2.1 - 5.3	300	14.8	12.0 - 18.1	100	4.5	3.0 - 6.6	1,400	77.4	73.6 - 80.8	1,900
186+%	200	2.1	1.5 - 2.7	800	11.1	9.8 - 12.4	300	4.5	3.7 - 5.4	6,000	82.4	80.8 - 83.9	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	7.1	5.4 - 9.3	700	26.6	23.4 - 30.0	100	3.4	2.3 - 5.0	1,600	62.9	59.3 - 66.5	2,500
\$10,000-\$24,999	200	5.7	4.4 - 7.3	700	20.9	18.4 - 23.6	100	2.5	1.7 - 3.7	2,400	70.9	68.0 - 73.7	3,400
\$25,000-\$49,999	100	2.4	1.7 - 3.5	600	15.7	13.7 - 17.9	200	4.9	3.9 - 6.3	3,000	76.9	74.4 - 79.3	3,900
>=\$50,000	100	1.4	1.0 - 2.2	400	8.2	6.9 - 9.6	200	4.0	3.1 - 5.1	4,300	86.3	84.6 - 87.9	5,000

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.7	3.6 - 8.9	200	15.8	12.2 - 20.3	<50	3.6	2.1 - 6.1	800	74.8	69.8 - 79.3	1,100
Normal birth weight	500	3.6	3.1 - 4.2	2,400	16.4	15.3 - 17.6	500	3.7	3.2 - 4.3	11,200	76.3	75.0 - 77.5	14,700

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	9.3	6.5 - 13.1	300	23.4	18.9 - 28.5	100	4.9	3.0 - 7.9	700	62.5	56.9 - 67.8	1,100
No	500	3.2	2.7 - 3.8	2,300	15.7	14.6 - 16.8	500	3.6	3.1 - 4.2	11,200	77.5	76.2 - 78.7	14,400

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.2	3.9 - 7.0	700	22.0	19.3 - 25.0	200	6.0	4.6 - 7.8	2,000	66.8	63.5 - 69.8	3,100
No	400	3.3	2.8 - 3.9	1,800	14.8	13.7 - 16.0	400	3.2	2.7 - 3.8	9,800	78.8	77.4 - 80.0	12,500

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	5.1	4.1 - 6.2	1,400	21.2	19.4 - 23.1	200	3.3	2.6 - 4.2	4,700	70.4	68.3 - 72.5	6,600
No	200	2.5	2.0 - 3.2	1,100	12.9	11.6 - 14.2	300	3.9	3.3 - 4.7	7,200	80.7	79.2 - 82.1	8,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	700	4.3	3.8 - 5.0	2,400	16.0	14.9 - 17.1	600	4.0	3.4 - 4.6	11,600	75.7	74.4 - 76.9	15,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.9	2.6 - 5.7	300	15.7	13.1 - 18.7	<50	2.2	1.3 - 3.5	1,400	78.3	75.0 - 81.3	1,800
Honolulu	500	4.3	3.6 - 5.1	1,800	15.7	14.4 - 17.1	500	4.3	3.6 - 5.1	8,500	75.7	74.1 - 77.3	11,200
Kauai	<50	3.7	2.1 - 6.5	100	11.7	8.4 - 16.1	<50	3.6	2.0 - 6.2	600	81.0	76.0 - 85.1	700
Maui	100	5.5	3.9 - 7.7	300	20.1	17.1 - 23.5	100	4.3	3.0 - 6.0	1,100	70.1	66.4 - 73.6	1,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.3	1.5 - 3.5	400	11.9	9.9 - 14.3	100	4.1	3.0 - 5.6	2,900	81.7	79.0 - 84.1	3,500
Native Hawaiian	300	6.2	5.0 - 7.7	800	20.8	18.7 - 23.1	200	4.8	3.8 - 6.1	2,700	68.1	65.5 - 70.6	4,000
Chinese	<50	1.4	0.8 - 2.3	<50	7.7	5.0 - 11.8	<50	2.6	1.6 - 4.3	500	88.3	84.2 - 91.4	600
Filipino	200	6.1	4.8 - 7.9	600	20.6	18.0 - 23.5	100	5.0	3.7 - 6.7	2,100	68.3	65.1 - 71.4	3,000
Japanese	100	3.0	1.7 - 5.2	200	7.9	5.9 - 10.6	<50	2.4	1.4 - 4.1	1,800	86.7	83.4 - 89.4	2,000
Other	100	3.8	2.5 - 6.0	400	17.2	14.1 - 20.8	100	2.6	1.5 - 4.5	1,600	76.3	72.4 - 79.9	2,100

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.2	6.4 - 13.2	400	30.0	25.3 - 35.2	<50	3.8	2.4 - 6.0	700	57.0	51.6 - 62.2	1,200
20-24	200	5.2	4.0 - 6.6	900	25.2	22.6 - 28.0	100	3.9	2.9 - 5.3	2,400	65.7	62.7 - 68.5	3,700
25-34	300	4.0	3.2 - 4.9	900	11.5	10.2 - 12.9	300	4.4	3.6 - 5.4	6,100	80.1	78.3 - 81.7	7,600
35+	100	2.1	1.3 - 3.3	300	10.0	8.1 - 12.3	100	3.0	2.0 - 4.4	2,400	84.9	82.3 - 87.3	2,800

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.3	3.6 - 5.2	1,300	16.0	14.6 - 17.6	400	4.5	3.7 - 5.4	6,000	75.2	73.3 - 76.9	8,000
Female	300	4.4	3.6 - 5.4	1,200	15.9	14.4 - 17.6	200	3.4	2.7 - 4.2	5,500	76.3	74.4 - 78.1	7,200

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	2.4 - 3.6	1,300	12.7	11.5 - 13.9	400	4.1	3.4 - 4.9	8,400	80.3	78.8 - 81.7	10,400
No	400	7.4	6.1 - 8.9	1,100	23.1	21.0 - 25.4	200	3.7	2.9 - 4.8	3,200	65.8	63.3 - 68.2	4,800

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.1	5.0 - 7.3	1,300	22.9	20.9 - 25.1	200	3.3	2.6 - 4.3	3,800	67.7	65.3 - 70.0	5,700
131-185%	100	4.0	2.6 - 6.2	300	14.9	12.1 - 18.1	100	3.9	2.5 - 6.1	1,500	77.2	73.4 - 80.6	1,900
186+%	100	2.3	1.6 - 3.1	600	9.8	8.6 - 11.3	300	4.6	3.7 - 5.6	5,400	83.3	81.6 - 85.0	6,500

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.6	3.5 - 9.0	200	18.1	13.9 - 23.2	<50	4.5	2.6 - 7.8	700	71.8	66.1 - 76.8	1,000
Normal birth weight	600	4.3	3.7 - 4.9	2,300	15.8	14.7 - 17.0	600	3.9	3.4 - 4.6	10,800	76.0	74.6 - 77.2	14,200

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	9.7	6.8 - 13.6	200	23.0	18.6 - 28.2	<50	4.2	2.5 - 7.0	700	63.0	57.4 - 68.3	1,000
No	500	3.9	3.3 - 4.5	2,200	15.4	14.3 - 16.6	600	4.0	3.4 - 4.6	10,800	76.7	75.4 - 78.0	14,000

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	6.7	5.1 - 8.7	600	21.9	19.2 - 24.9	200	6.4	4.9 - 8.3	1,900	64.9	61.6 - 68.1	2,900
No	400	3.7	3.1 - 4.4	1,700	14.4	13.2 - 15.6	400	3.4	2.9 - 4.1	9,500	78.5	77.1 - 79.8	12,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.5	4.6 - 6.7	1,400	21.6	19.7 - 23.5	200	3.4	2.7 - 4.3	4,400	69.5	67.3 - 71.6	6,300
No	300	3.4	2.7 - 4.2	1,000	12.0	10.7 - 13.3	400	4.4	3.6 - 5.3	7,000	80.3	78.6 - 81.8	8,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	700	4.4	3.8 - 5.1	2,400	15.8	14.7 - 17.0	600	3.7	3.2 - 4.4	11,400	76.1	74.7 - 77.4	15,000

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.6	3.4 - 6.3	300	17.0	14.7 - 19.6	<50	2.7	1.9 - 3.9	1,300	75.7	72.7 - 78.4	1,700
Honolulu	500	4.2	3.5 - 5.1	1,700	15.3	13.9 - 16.9	400	3.8	3.1 - 4.6	8,400	76.6	74.9 - 78.3	10,900
Kauai	<50	4.2	2.4 - 7.2	100	13.7	10.5 - 17.7	<50	3.9	2.4 - 6.4	500	78.1	73.5 - 82.2	700
Maui	100	5.2	3.8 - 6.9	300	18.5	16.0 - 21.4	100	4.3	3.0 - 5.9	1,200	72.0	68.8 - 75.1	1,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.1	1.3 - 3.4	400	12.3	10.1 - 15.0	200	4.6	3.3 - 6.4	2,700	81.0	77.9 - 83.7	3,400
Native Hawaiian	200	5.7	4.7 - 6.9	800	20.8	19.0 - 22.8	200	4.3	3.4 - 5.3	2,700	69.2	67.0 - 71.3	3,900
Chinese	<50	2.9	1.2 - 6.4	<50	8.0	4.4 - 14.1	<50	2.8	1.2 - 6.3	500	86.4	79.9 - 91.1	600
Filipino	200	6.4	4.8 - 8.4	600	19.8	17.0 - 22.9	100	4.2	3.0 - 5.9	2,100	69.6	66.1 - 72.9	3,000
Japanese	100	2.8	1.6 - 5.0	100	6.9	4.8 - 9.7	<50	2.1	1.2 - 3.8	1,800	88.2	84.8 - 90.9	2,100
Other	100	4.8	3.0 - 7.6	300	17.5	13.9 - 21.7	<50	2.4	1.3 - 4.4	1,500	75.3	70.7 - 79.5	2,000

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.9	7.0 - 13.9	400	31.2	26.4 - 36.5	<50	3.0	1.9 - 4.9	700	55.8	50.4 - 61.1	1,200
20-24	200	5.4	4.2 - 7.1	800	23.5	20.9 - 26.4	100	3.6	2.6 - 4.8	2,400	67.5	64.4 - 70.4	3,600
25-34	300	3.7	3.0 - 4.7	900	11.7	10.3 - 13.2	300	4.2	3.4 - 5.1	6,000	80.5	78.6 - 82.2	7,500
35+	100	2.3	1.4 - 3.8	300	10.1	8.1 - 12.6	100	3.1	2.0 - 4.8	2,300	84.5	81.5 - 87.0	2,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.5	3.6 - 5.5	1,300	16.2	14.6 - 17.9	300	3.9	3.2 - 4.8	5,900	75.4	73.5 - 77.3	7,800
Female	300	4.3	3.5 - 5.3	1,100	15.4	13.8 - 17.1	300	3.5	2.8 - 4.4	5,500	76.8	74.8 - 78.6	7,200

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	2.3 - 3.6	1,300	12.7	11.5 - 14.1	400	3.9	3.2 - 4.7	8,400	80.5	78.9 - 82.0	10,400
No	400	7.9	6.5 - 9.6	1,000	22.9	20.7 - 25.3	200	3.4	2.7 - 4.3	3,000	65.8	63.2 - 68.3	4,500

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.2	5.0 - 7.6	1,300	22.1	20.1 - 24.3	200	2.9	2.2 - 3.7	3,900	68.8	66.5 - 71.1	5,700
131-185%	100	4.1	2.7 - 6.1	300	14.7	12.0 - 17.8	100	3.8	2.5 - 5.8	1,800	77.4	73.8 - 80.7	2,300
186+%	100	2.0	1.4 - 2.9	500	9.4	8.0 - 11.0	300	4.6	3.6 - 5.7	4,800	83.9	82.0 - 85.7	5,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.8	3.4 - 9.8	200	19.9	15.1 - 25.8	<50	4.0	2.0 - 7.6	700	70.3	64.0 - 76.0	1,000
Normal birth weight	600	4.3	3.7 - 5.0	2,200	15.5	14.4 - 16.7	500	3.7	3.2 - 4.4	10,700	76.5	75.1 - 77.8	14,000

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.2	4.7 - 10.8	200	23.9	19.2 - 29.3	<50	4.0	2.5 - 6.4	600	64.9	59.1 - 70.2	1,000
No	600	4.2	3.5 - 4.9	2,100	15.2	14.0 - 16.4	500	3.7	3.1 - 4.4	10,600	76.9	75.5 - 78.3	13,800

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	6.7	5.0 - 8.8	600	21.3	18.5 - 24.4	200	5.7	4.3 - 7.5	1,800	66.4	62.9 - 69.7	2,700
No	400	3.7	3.1 - 4.4	1,700	14.5	13.2 - 15.8	400	3.3	2.8 - 4.0	9,400	78.5	77.1 - 80.0	11,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.8	4.7 - 7.1	1,400	22.3	20.3 - 24.4	200	3.1	2.5 - 4.0	4,200	68.8	66.5 - 71.0	6,100
No	300	3.4	2.8 - 4.2	1,000	11.3	10.0 - 12.7	400	4.1	3.4 - 5.1	7,000	81.1	79.4 - 82.7	8,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.5	3.9 - 5.2	2,300	16.3	15.1 - 17.5	500	3.3	2.8 - 4.0	10,900	75.9	74.5 - 77.3	14,400

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.8	3.7 - 6.2	300	17.7	15.6 - 19.9	100	3.4	2.5 - 4.7	1,200	74.1	71.5 - 76.6	1,600
Honolulu	500	4.4	3.7 - 5.3	1,700	15.9	14.5 - 17.5	300	3.2	2.5 - 4.0	8,100	76.4	74.6 - 78.2	10,600
Kauai	<50	3.8	2.3 - 6.1	100	13.9	11.0 - 17.4	<50	5.0	3.3 - 7.6	500	77.3	73.1 - 81.0	700
Maui	100	5.0	3.9 - 6.6	300	18.0	15.7 - 20.5	100	3.4	2.5 - 4.7	1,100	73.6	70.7 - 76.2	1,500

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.4	1.5 - 3.9	400	12.0	9.7 - 14.8	100	4.0	2.7 - 5.8	2,600	81.5	78.4 - 84.3	3,200
Native Hawaiian	200	6.2	5.4 - 7.2	700	20.3	18.9 - 21.8	100	3.8	3.2 - 4.6	2,600	69.6	67.9 - 71.3	3,700
Chinese	<50	2.9	1.2 - 6.9	100	11.0	6.3 - 18.6	<50	3.5	1.4 - 8.6	500	82.5	74.4 - 88.4	600
Filipino	200	5.6	4.0 - 7.6	600	20.7	17.7 - 24.0	100	3.5	2.3 - 5.2	2,100	70.2	66.6 - 73.7	3,000
Japanese	<50	2.2	1.1 - 4.4	200	9.0	6.6 - 12.2	<50	2.2	1.2 - 4.1	1,700	86.7	83.0 - 89.6	2,000
Other	100	5.9	3.8 - 9.1	300	17.9	14.1 - 22.5	<50	2.1	1.0 - 4.2	1,400	74.1	69.1 - 78.6	1,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	10.4	7.6 - 14.1	400	33.1	28.6 - 38.0	<50	2.0	1.3 - 3.2	700	54.5	49.5 - 59.4	1,300
20-24	200	5.9	4.6 - 7.7	800	22.7	20.1 - 25.5	100	3.3	2.4 - 4.5	2,400	68.1	65.1 - 71.0	3,500
25-34	300	3.6	2.9 - 4.5	900	12.5	11.0 - 14.1	300	3.6	2.8 - 4.6	5,800	80.3	78.4 - 82.1	7,200
35+	100	2.0	1.2 - 3.5	200	9.5	7.4 - 12.2	100	3.3	2.1 - 5.2	2,100	85.1	82.0 - 87.8	2,400

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.7	3.8 - 5.7	1,200	16.6	15.0 - 18.4	200	3.3	2.6 - 4.2	5,600	75.4	73.4 - 77.3	7,400
Female	300	4.3	3.5 - 5.3	1,100	15.9	14.3 - 17.6	200	3.4	2.7 - 4.2	5,300	76.4	74.4 - 78.3	7,000

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	2.3 - 3.7	1,300	13.0	11.7 - 14.4	300	3.5	2.8 - 4.4	8,000	80.6	78.9 - 82.2	9,900
No	400	8.0	6.6 - 9.6	1,100	23.6	21.4 - 26.0	100	2.9	2.3 - 3.7	2,900	65.4	62.9 - 67.9	4,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.3	5.2 - 7.7	1,200	21.4	19.4 - 23.5	200	3.1	2.4 - 4.0	3,800	69.2	66.9 - 71.5	5,500
131-185%	100	3.9	2.7 - 5.5	400	15.7	13.2 - 18.5	100	3.2	2.1 - 4.8	2,100	77.3	74.1 - 80.2	2,800
186+%	100	2.1	1.5 - 3.1	500	9.0	7.5 - 10.7	200	3.6	2.7 - 4.9	4,300	85.3	83.1 - 87.2	5,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	200	8.1	6.4 - 10.2	600	21.8	19.0 - 24.8	100	3.4	2.4 - 4.8	1,800	66.7	63.4 - 69.9	2,700
\$15,001-\$25,000	100	4.7	3.1 - 7.0	400	20.2	16.9 - 23.8	100	3.1	2.1 - 4.8	1,400	72.0	68.0 - 75.7	1,900
\$25,001-\$40,000	100	4.8	3.4 - 6.6	500	17.4	14.9 - 20.2	100	3.4	2.3 - 5.0	2,200	74.5	71.2 - 77.5	3,000
>=\$40,001	100	2.1	1.5 - 2.9	600	10.3	8.8 - 12.0	200	3.5	2.7 - 4.7	5,000	84.1	82.1 - 85.9	5,900

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	4.9	2.7 - 8.9	200	22.2	17.1 - 28.4	<50	3.6	1.7 - 7.2	600	69.3	62.7 - 75.1	900
Normal birth weight	600	4.5	3.9 - 5.2	2,100	15.9	14.7 - 17.1	400	3.3	2.8 - 4.0	10,300	76.4	74.9 - 77.7	13,400

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	6.8	4.6 - 9.8	300	26.7	21.8 - 32.2	<50	3.6	2.4 - 5.6	600	62.9	57.3 - 68.2	1,000
No	600	4.3	3.7 - 5.0	2,000	15.5	14.3 - 16.8	400	3.3	2.8 - 4.0	10,100	76.9	75.4 - 78.3	13,200

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	7.2	5.5 - 9.4	600	23.6	20.7 - 26.9	100	4.7	3.4 - 6.4	1,700	64.4	60.9 - 67.8	2,600
No	400	3.6	3.0 - 4.4	1,700	14.5	13.3 - 15.9	300	3.0	2.5 - 3.7	9,000	78.8	77.3 - 80.3	11,400

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	6.6	5.4 - 7.9	1,300	22.9	20.8 - 25.0	200	3.3	2.6 - 4.2	3,900	67.3	64.9 - 69.6	5,700
No	300	3.2	2.5 - 3.9	1,000	11.9	10.6 - 13.4	300	3.3	2.6 - 4.2	7,000	81.6	79.9 - 83.2	8,500

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.2	3.7 - 4.9	2,300	16.2	15.2 - 17.4	400	3.2	2.7 - 3.7	10,800	76.4	75.1 - 77.6	14,100

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.9	3.8 - 6.4	300	19.3	17.2 - 21.7	100	3.6	2.7 - 4.9	1,200	72.1	69.4 - 74.6	1,600
Honolulu	400	4.1	3.4 - 4.9	1,600	15.4	14.1 - 16.9	300	3.1	2.5 - 3.8	8,100	77.4	75.7 - 78.9	10,400
Kauai	<50	3.1	1.8 - 5.2	100	17.2	13.8 - 21.3	<50	3.0	1.7 - 5.3	500	76.7	72.3 - 80.6	600
Maui	100	5.0	3.8 - 6.5	300	18.1	15.8 - 20.7	<50	3.0	2.1 - 4.3	1,100	73.9	71.0 - 76.5	1,500

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.9	2.0 - 4.3	400	12.1	10.0 - 14.6	100	4.1	2.9 - 5.7	2,600	80.8	78.0 - 83.4	3,200
Native Hawaiian	200	5.7	4.9 - 6.5	800	21.6	20.2 - 23.0	100	3.7	3.1 - 4.4	2,500	69.1	67.5 - 70.7	3,600
Chinese	<50	3.9	1.8 - 8.6	100	15.2	9.7 - 23.1	<50	3.8	1.5 - 9.5	400	77.0	68.6 - 83.7	600
Filipino	200	5.2	3.7 - 7.1	600	19.5	16.8 - 22.5	100	2.6	1.8 - 3.9	2,100	72.7	69.3 - 75.8	2,900
Japanese	<50	0.7	0.4 - 1.4	200	8.3	6.1 - 11.2	100	2.8	1.6 - 4.8	1,800	88.2	84.9 - 90.8	2,000
Other	100	6.2	4.0 - 9.3	300	16.8	13.2 - 21.1	<50	1.6	0.8 - 3.2	1,400	75.4	70.7 - 79.6	1,800

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.6	6.6 - 11.1	400	31.0	27.2 - 35.2	<50	2.0	1.2 - 3.5	800	58.4	54.0 - 62.6	1,400
20-24	200	6.5	5.1 - 8.3	700	21.3	19.0 - 23.8	100	3.1	2.4 - 4.0	2,400	69.1	66.3 - 71.8	3,500
25-34	200	3.0	2.3 - 3.8	900	13.1	11.7 - 14.6	200	3.3	2.6 - 4.1	5,600	80.7	78.9 - 82.3	6,900
35+	<50	2.1	1.2 - 3.5	200	9.5	7.5 - 11.9	100	3.6	2.4 - 5.5	2,000	84.8	81.8 - 87.3	2,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.6	3.8 - 5.6	1,200	16.5	15.0 - 18.1	200	2.8	2.2 - 3.6	5,400	76.1	74.2 - 77.8	7,100
Female	300	3.9	3.2 - 4.7	1,100	16.0	14.5 - 17.6	200	3.5	2.8 - 4.4	5,400	76.6	74.8 - 78.4	7,000

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	2.3 - 3.7	1,300	13.0	11.8 - 14.4	300	3.3	2.7 - 4.0	7,900	80.7	79.2 - 82.2	9,800
No	300	7.2	6.0 - 8.6	1,000	23.4	21.4 - 25.6	100	2.9	2.3 - 3.7	2,900	66.5	64.1 - 68.8	4,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.0	4.9 - 7.2	1,100	20.8	19.0 - 22.8	200	2.8	2.2 - 3.5	3,800	70.4	68.3 - 72.5	5,400
131-185%	100	3.7	2.6 - 5.2	400	15.5	13.1 - 18.1	100	2.5	1.7 - 3.8	2,100	78.3	75.3 - 81.0	2,700
186+%	100	2.1	1.5 - 3.0	500	9.9	8.5 - 11.6	200	4.1	3.1 - 5.3	4,300	83.9	81.8 - 85.7	5,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Income (2000-2003)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	200	7.8	6.3 - 9.7	600	22.0	19.5 - 24.8	100	2.9	2.1 - 4.0	1,800	67.2	64.2 - 70.1	2,700
\$15,001-\$25,000	100	4.6	3.1 - 6.7	400	19.5	16.7 - 22.7	100	3.4	2.3 - 5.0	1,500	72.4	68.8 - 75.8	2,000
\$25,001-\$40,000	100	3.8	2.6 - 5.4	500	16.8	14.5 - 19.3	100	2.7	1.9 - 3.9	2,300	76.7	73.9 - 79.4	3,000
>=\$40,001	100	2.2	1.6 - 3.0	600	10.5	9.1 - 12.1	200	3.6	2.8 - 4.7	4,800	83.7	81.8 - 85.4	5,700

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	5.3	2.9 - 9.4	200	21.9	17.2 - 27.5	<50	3.0	1.6 - 5.5	600	69.9	63.8 - 75.3	900
Normal birth weight	600	4.2	3.6 - 4.8	2,100	15.8	14.7 - 17.0	400	3.2	2.7 - 3.7	10,200	76.9	75.6 - 78.1	13,200

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.5	5.3 - 10.5	300	29.1	24.4 - 34.4	<50	3.7	2.4 - 5.5	600	59.7	54.4 - 64.8	900
No	500	3.9	3.3 - 4.5	2,000	15.2	14.1 - 16.4	400	3.1	2.6 - 3.7	10,100	77.8	76.5 - 79.1	12,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	6.3	4.8 - 8.1	700	25.3	22.5 - 28.3	100	4.0	2.9 - 5.5	1,700	64.4	61.1 - 67.5	2,600
No	400	3.4	2.9 - 4.1	1,600	14.1	13.0 - 15.3	300	2.9	2.4 - 3.5	8,800	79.6	78.1 - 80.9	11,100

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	6.2	5.2 - 7.5	1,200	21.5	19.7 - 23.4	200	3.0	2.4 - 3.8	4,000	69.2	67.1 - 71.3	5,700
No	200	2.8	2.2 - 3.5	1,100	12.7	11.4 - 14.1	300	3.2	2.6 - 4.0	6,700	81.3	79.7 - 82.8	8,300