

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 3-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, income data does not exist for some 3-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration 8+ weeks, among initiators	Breastfeeding (any) duration 8+ weeks, among initiators

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 9/21/15

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012


Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2002	3,300	23.1	21.8 - 24.3	10,900	76.9	75.7 - 78.2	14,100
2001-2003	3,400	23.4	22.0 - 24.8	11,000	76.6	75.2 - 78.0	14,400
2002-2004	3,500	23.3	22.0 - 24.6	11,500	76.7	75.4 - 78.0	15,000
2003-2005	3,600	23.5	22.3 - 24.8	11,700	76.5	75.2 - 77.7	15,300
2004-2006	3,700	23.2	22.0 - 24.4	12,100	76.8	75.6 - 78.0	15,800
2005-2007	3,700	23.1	21.8 - 24.3	12,400	76.9	75.7 - 78.2	16,100
2006-2008	3,700	22.6	21.4 - 23.8	12,800	77.4	76.2 - 78.6	16,500
2007-2009	3,800	23.1	21.7 - 24.5	12,800	76.9	75.5 - 78.3	16,700
2008-2010	3,800	22.4	21.0 - 24.0	13,100	77.6	76.0 - 79.0	16,900
2009-2011	3,700	21.3	19.8 - 22.9	13,500	78.7	77.1 - 80.2	17,100
2010-2012	3,300	19.1	17.7 - 20.7	14,000	80.9	79.3 - 82.3	17,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,300	19.1	17.7 - 20.7	14,000	80.9	79.3 - 82.3	17,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	21.1	18.7 - 23.8	1,700	78.9	76.2 - 81.3	2,200
Honolulu	2,300	18.9	17.0 - 21.0	10,100	81.1	79.0 - 83.0	12,400
Kauai	100	15.0	12.5 - 17.8	700	85.0	82.2 - 87.5	800
Maui	400	20.0	17.8 - 22.3	1,500	80.0	77.7 - 82.2	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	16.6	13.7 - 19.9	3,500	83.4	80.1 - 86.3	4,200
Native Hawaiian	1,200	24.5	21.7 - 27.6	3,700	75.5	72.4 - 78.3	5,000
Chinese	100	10.7	6.2 - 18.1	700	89.3	81.9 - 93.8	800
Filipino	800	24.7	21.0 - 28.7	2,300	75.3	71.3 - 79.0	3,100
Japanese	100	8.7	5.9 - 12.8	1,500	91.3	87.2 - 94.1	1,700
Other	400	15.6	12.1 - 19.8	2,200	84.4	80.2 - 87.9	2,600

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	40.5	33.3 - 48.1	600	59.5	51.9 - 66.7	1,000
20-24	900	23.3	20.0 - 27.0	3,000	76.7	73.0 - 80.0	3,900
25-34	1,600	17.4	15.5 - 19.5	7,700	82.6	80.5 - 84.5	9,300
35+	400	12.1	9.6 - 15.1	2,700	87.9	84.9 - 90.4	3,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,700	19.9	17.8 - 22.1	7,000	80.1	77.9 - 82.2	8,700
Female	1,600	18.4	16.4 - 20.6	7,000	81.6	79.4 - 83.6	8,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	16.0	14.2 - 17.9	9,100	84.0	82.1 - 85.8	10,900
No	1,600	24.5	22.0 - 27.2	4,800	75.5	72.8 - 78.0	6,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,400	24.2	21.6 - 27.0	4,500	75.8	73.0 - 78.4	6,000
131-185%	600	21.2	17.3 - 25.6	2,200	78.8	74.4 - 82.7	2,800
186+%	1,000	13.7	11.8 - 16.0	6,300	86.3	84.0 - 88.2	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight							
Low birth weight	300	23.6	20.9 - 26.7	900	76.4	73.3 - 79.1	1,200
Normal birth weight	3,000	18.8	17.3 - 20.5	13,100	81.2	79.5 - 82.7	16,100

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester							
Yes	300	38.2	30.6 - 46.4	500	61.8	53.6 - 69.4	900
No	2,900	18.2	16.7 - 19.8	13,200	81.8	80.2 - 83.3	16,100

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant							
Yes	1,000	30.5	26.6 - 34.8	2,200	69.5	65.2 - 73.4	3,100
No	2,300	16.7	15.1 - 18.3	11,600	83.3	81.7 - 84.9	13,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	24.5	22.1 - 27.0	5,700	75.5	73.0 - 77.9	7,600
No	1,400	14.8	13.0 - 16.8	8,200	85.2	83.2 - 87.0	9,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,700	21.3	19.8 - 22.9	13,500	78.7	77.1 - 80.2	17,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	23.4	20.9 - 26.1	1,700	76.6	73.9 - 79.1	2,200
Honolulu	2,600	21.3	19.3 - 23.5	9,600	78.7	76.5 - 80.7	12,200
Kauai	100	15.9	13.5 - 18.7	700	84.1	81.3 - 86.5	800
Maui	400	21.5	19.3 - 23.9	1,500	78.5	76.1 - 80.7	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	18.1	15.2 - 21.4	3,500	81.9	78.6 - 84.8	4,200
Native Hawaiian	1,300	26.4	23.5 - 29.5	3,700	73.6	70.5 - 76.5	5,000
Chinese	100	10.6	6.0 - 18.2	600	89.4	81.8 - 94.0	700
Filipino	800	25.5	21.9 - 29.5	2,300	74.5	70.5 - 78.1	3,100
Japanese	200	11.4	8.0 - 16.0	1,500	88.6	84.0 - 92.0	1,700
Other	500	21.1	17.0 - 25.9	1,900	78.9	74.1 - 83.0	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	42.5	35.8 - 49.4	700	57.5	50.6 - 64.2	1,200
20-24	1,000	25.6	22.3 - 29.3	3,000	74.4	70.7 - 77.7	4,000
25-34	1,700	18.6	16.6 - 20.8	7,200	81.4	79.2 - 83.4	8,900
35+	500	15.1	12.2 - 18.6	2,600	84.9	81.4 - 87.8	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,900	21.6	19.4 - 23.9	6,900	78.4	76.1 - 80.6	8,800
Female	1,800	21.1	18.9 - 23.4	6,600	78.9	76.6 - 81.1	8,300

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	17.2	15.4 - 19.2	8,900	82.8	80.8 - 84.6	10,700
No	1,800	28.2	25.5 - 31.0	4,600	71.8	69.0 - 74.5	6,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,600	26.9	24.2 - 29.8	4,300	73.1	70.2 - 75.8	5,900
131-185%	600	23.8	19.8 - 28.3	2,000	76.2	71.7 - 80.2	2,700
186+%	1,100	15.1	13.1 - 17.4	6,300	84.9	82.6 - 86.9	7,400

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	29.0	25.1 - 33.3	2,100	71.0	66.7 - 74.9	3,000
\$10,000-\$24,999	800	25.0	21.5 - 28.8	2,400	75.0	71.2 - 78.5	3,200
\$25,000-\$49,999	900	22.8	19.5 - 26.4	3,100	77.2	73.6 - 80.5	4,000
>=\$50,000	800	13.9	11.7 - 16.3	5,100	86.1	83.7 - 88.3	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	26.2	23.4 - 29.2	900	73.8	70.8 - 76.6	1,200
Normal birth weight	3,300	21.0	19.3 - 22.7	12,600	79.0	77.3 - 80.7	15,900

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	39.5	32.6 - 46.8	700	60.5	53.2 - 67.4	1,100
No	3,100	19.9	18.3 - 21.5	12,700	80.1	78.5 - 81.7	15,800

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	33.3	29.4 - 37.5	2,200	66.7	62.5 - 70.6	3,300
No	2,500	18.1	16.5 - 19.9	11,100	81.9	80.1 - 83.5	13,500

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	27.1	24.7 - 29.8	5,300	72.9	70.2 - 75.3	7,200
No	1,600	16.9	15.0 - 19.0	8,000	83.1	81.0 - 85.0	9,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,800	22.4	21.0 - 24.0	13,100	77.6	76.0 - 79.0	16,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	23.7	20.8 - 26.8	1,600	76.3	73.2 - 79.2	2,200
Honolulu	2,700	22.8	20.9 - 24.8	9,300	77.2	75.2 - 79.1	12,000
Kauai	200	19.2	15.6 - 23.5	600	80.8	76.5 - 84.4	800
Maui	400	20.3	17.8 - 23.1	1,500	79.7	76.9 - 82.2	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	18.9	16.2 - 22.0	3,400	81.1	78.0 - 83.8	4,100
Native Hawaiian	1,300	27.2	24.4 - 30.2	3,600	72.8	69.8 - 75.6	4,900
Chinese	100	12.4	7.8 - 19.0	600	87.6	81.0 - 92.2	700
Filipino	800	27.2	23.8 - 31.0	2,200	72.8	69.0 - 76.2	3,000
Japanese	200	12.2	9.1 - 16.2	1,500	87.8	83.8 - 90.9	1,700
Other	600	23.0	19.0 - 27.5	1,800	77.0	72.5 - 81.0	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	36.8	30.8 - 43.2	800	63.2	56.8 - 69.2	1,300
20-24	1,100	29.5	26.2 - 33.1	2,700	70.5	66.9 - 73.8	3,800
25-34	1,600	18.9	17.0 - 20.9	7,000	81.1	79.1 - 83.0	8,700
35+	500	17.5	14.6 - 20.9	2,500	82.5	79.1 - 85.4	3,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	22.8	20.8 - 24.9	6,800	77.2	75.1 - 79.2	8,800
Female	1,800	22.1	20.0 - 24.3	6,300	77.9	75.7 - 80.0	8,100

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	18.7	17.0 - 20.5	8,600	81.3	79.5 - 83.0	10,600
No	1,800	28.9	26.3 - 31.6	4,400	71.1	68.4 - 73.7	6,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,600	27.8	25.2 - 30.6	4,200	72.2	69.4 - 74.8	5,800
131-185%	600	23.1	19.3 - 27.4	2,000	76.9	72.6 - 80.7	2,600
186+%	1,200	16.8	14.8 - 18.9	6,100	83.2	81.1 - 85.2	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	30.7	26.8 - 34.9	1,900	69.3	65.1 - 73.2	2,800
\$10,000-\$24,999	800	25.7	22.5 - 29.3	2,400	74.3	70.7 - 77.5	3,300
\$25,000-\$49,999	900	23.4	20.3 - 26.8	2,900	76.6	73.2 - 79.7	3,800
>=\$50,000	900	15.0	13.0 - 17.4	5,000	85.0	82.6 - 87.0	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	24.6	21.2 - 28.4	900	75.4	71.6 - 78.8	1,200
Normal birth weight	3,500	22.3	20.7 - 23.9	12,200	77.7	76.1 - 79.3	15,700

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	37.4	31.3 - 44.0	800	62.6	56.0 - 68.7	1,200
No	3,200	21.1	19.6 - 22.7	12,100	78.9	77.3 - 80.4	15,400

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	34.8	31.0 - 38.8	2,100	65.2	61.2 - 69.0	3,200
No	2,600	19.4	17.8 - 21.0	10,800	80.6	79.0 - 82.2	13,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	28.9	26.4 - 31.4	4,900	71.1	68.6 - 73.6	6,900
No	1,700	18.0	16.2 - 19.9	7,900	82.0	80.1 - 83.8	9,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,800	23.1	21.7 - 24.5	12,800	76.9	75.5 - 78.3	16,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	21.9	18.9 - 25.3	1,700	78.1	74.7 - 81.1	2,100
Honolulu	2,800	23.6	21.9 - 25.5	9,000	76.4	74.5 - 78.1	11,700
Kauai	200	20.9	16.6 - 26.0	700	79.1	74.0 - 83.4	800
Maui	400	21.8	18.7 - 25.2	1,500	78.2	74.8 - 81.3	2,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	18.5	16.0 - 21.2	3,200	81.5	78.8 - 84.0	4,000
Native Hawaiian	1,400	30.1	27.3 - 33.1	3,300	69.9	66.9 - 72.7	4,700
Chinese	100	12.1	8.5 - 16.9	600	87.9	83.1 - 91.5	600
Filipino	800	26.9	23.8 - 30.2	2,200	73.1	69.8 - 76.2	3,000
Japanese	300	13.5	10.4 - 17.2	1,600	86.5	82.8 - 89.6	1,900
Other	600	22.5	18.9 - 26.5	1,900	77.5	73.5 - 81.1	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	36.6	31.1 - 42.5	900	63.4	57.5 - 68.9	1,500
20-24	1,200	31.0	27.9 - 34.3	2,600	69.0	65.7 - 72.1	3,800
25-34	1,600	19.3	17.6 - 21.2	6,800	80.7	78.8 - 82.4	8,400
35+	500	16.9	14.2 - 20.0	2,500	83.1	80.0 - 85.8	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	23.3	21.4 - 25.2	6,700	76.7	74.8 - 78.6	8,800
Female	1,800	22.9	20.9 - 24.9	6,100	77.1	75.1 - 79.1	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.4	16.9 - 20.1	8,600	81.6	79.9 - 83.1	10,500
No	1,900	30.9	28.4 - 33.5	4,300	69.1	66.5 - 71.6	6,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	30.0	27.5 - 32.6	4,100	70.0	67.4 - 72.5	5,800
131-185%	500	22.0	18.6 - 25.8	1,900	78.0	74.2 - 81.4	2,500
186+%	1,200	17.1	15.4 - 19.1	5,900	82.9	80.9 - 84.6	7,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	33.7	29.8 - 37.9	1,800	66.3	62.1 - 70.2	2,700
\$10,000-\$24,999	900	27.2	24.1 - 30.5	2,500	72.8	69.5 - 75.9	3,400
\$25,000-\$49,999	800	23.0	20.2 - 26.0	2,800	77.0	74.0 - 79.8	3,700
>=\$50,000	900	15.1	13.2 - 17.2	4,900	84.9	82.8 - 86.8	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	26.2	22.2 - 30.7	900	73.8	69.3 - 77.8	1,200
Normal birth weight	3,500	22.8	21.4 - 24.3	11,900	77.2	75.7 - 78.6	15,500

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	42.5	36.7 - 48.5	700	57.5	51.5 - 63.3	1,300
No	3,200	21.3	20.0 - 22.8	11,900	78.7	77.2 - 80.0	15,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	36.6	33.0 - 40.4	2,000	63.4	59.6 - 67.0	3,100
No	2,600	19.8	18.4 - 21.3	10,600	80.2	78.7 - 81.6	13,300

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	30.1	27.7 - 32.5	4,600	69.9	67.5 - 72.3	6,600
No	1,800	18.2	16.6 - 20.0	7,900	81.8	80.0 - 83.4	9,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,700	22.6	21.4 - 23.8	12,800	77.4	76.2 - 78.6	16,500

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	21.3	18.0 - 25.0	1,600	78.7	75.0 - 82.0	2,100
Honolulu	2,700	23.0	21.6 - 24.5	9,100	77.0	75.5 - 78.4	11,800
Kauai	200	20.9	16.0 - 26.8	700	79.1	73.2 - 84.0	800
Maui	400	22.0	18.5 - 26.0	1,400	78.0	74.0 - 81.5	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	19.5	17.3 - 22.0	3,200	80.5	78.0 - 82.7	3,900
Native Hawaiian	1,300	29.7	27.0 - 32.6	3,200	70.3	67.4 - 73.0	4,500
Chinese	100	13.8	11.2 - 16.8	500	86.2	83.2 - 88.8	600
Filipino	800	25.8	23.1 - 28.7	2,200	74.2	71.3 - 76.9	3,000
Japanese	200	11.9	9.4 - 15.0	1,700	88.1	85.0 - 90.6	2,000
Other	500	21.2	18.2 - 24.6	2,000	78.8	75.4 - 81.8	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	34.7	29.6 - 40.2	900	65.3	59.8 - 70.4	1,400
20-24	1,200	31.8	29.0 - 34.8	2,600	68.2	65.2 - 71.0	3,900
25-34	1,600	19.1	17.5 - 20.7	6,700	80.9	79.3 - 82.5	8,300
35+	400	14.7	12.6 - 17.2	2,600	85.3	82.8 - 87.4	3,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,100	24.2	22.5 - 26.0	6,500	75.8	74.0 - 77.5	8,600
Female	1,600	20.8	19.1 - 22.5	6,300	79.2	77.5 - 80.9	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.1	16.8 - 19.5	8,700	81.9	80.5 - 83.2	10,600
No	1,800	30.4	28.1 - 32.8	4,100	69.6	67.2 - 71.9	6,000

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	30.2	27.9 - 32.7	4,000	69.8	67.3 - 72.1	5,800
131-185%	400	19.2	16.3 - 22.5	1,900	80.8	77.5 - 83.7	2,300
186+%	1,300	17.2	15.6 - 18.8	6,100	82.8	81.2 - 84.4	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	34.2	30.6 - 38.0	1,700	65.8	62.0 - 69.4	2,700
\$10,000-\$24,999	900	26.6	23.8 - 29.6	2,500	73.4	70.4 - 76.2	3,400
\$25,000-\$49,999	900	23.6	21.1 - 26.3	2,900	76.4	73.7 - 78.9	3,800
>=\$50,000	800	13.6	12.0 - 15.3	4,900	86.4	84.7 - 88.0	5,600

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	24.1	19.8 - 29.1	900	75.9	70.9 - 80.2	1,200
Normal birth weight	3,400	22.4	21.2 - 23.7	11,900	77.6	76.3 - 78.8	15,300

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	39.7	34.3 - 45.3	700	60.3	54.7 - 65.7	1,200
No	3,200	21.2	19.9 - 22.4	11,900	78.8	77.6 - 80.1	15,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	34.6	31.4 - 38.1	2,000	65.4	61.9 - 68.6	3,100
No	2,600	19.8	18.5 - 21.1	10,600	80.2	78.9 - 81.5	13,200

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	28.9	26.8 - 31.1	4,700	71.1	68.9 - 73.2	6,700
No	1,700	18.1	16.7 - 19.6	7,900	81.9	80.4 - 83.3	9,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,700	23.1	21.8 - 24.3	12,400	76.9	75.7 - 78.2	16,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	20.4	17.1 - 24.1	1,600	79.6	75.9 - 82.9	2,000
Honolulu	2,700	23.3	21.9 - 24.8	8,900	76.7	75.2 - 78.1	11,600
Kauai	100	18.0	13.3 - 23.8	600	82.0	76.2 - 86.7	800
Maui	500	26.9	23.0 - 31.2	1,300	73.1	68.8 - 77.0	1,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	18.7	16.5 - 21.3	3,000	81.3	78.7 - 83.5	3,700
Native Hawaiian	1,400	31.1	28.3 - 34.0	3,000	68.9	66.0 - 71.7	4,400
Chinese	100	14.2	11.6 - 17.2	500	85.8	82.8 - 88.4	600
Filipino	800	27.7	24.9 - 30.6	2,100	72.3	69.4 - 75.1	3,000
Japanese	200	11.2	8.9 - 14.1	1,800	88.8	85.9 - 91.1	2,100
Other	500	21.8	18.8 - 25.3	1,900	78.2	74.7 - 81.2	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	37.7	32.5 - 43.2	800	62.3	56.8 - 67.5	1,400
20-24	1,300	32.7	29.9 - 35.7	2,600	67.3	64.3 - 70.1	3,900
25-34	1,500	19.3	17.8 - 21.0	6,400	80.7	79.0 - 82.2	7,900
35+	400	13.2	11.1 - 15.5	2,500	86.8	84.5 - 88.9	2,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	24.5	22.8 - 26.3	6,200	75.5	73.7 - 77.2	8,300
Female	1,700	21.6	19.9 - 23.4	6,100	78.4	76.6 - 80.1	7,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	17.4	16.1 - 18.8	8,500	82.6	81.2 - 83.9	10,300
No	1,900	33.2	30.8 - 35.7	3,800	66.8	64.3 - 69.2	5,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,800	32.4	30.0 - 34.8	3,800	67.6	65.2 - 70.0	5,600
131-185%	400	21.9	18.6 - 25.5	1,600	78.1	74.5 - 81.4	2,100
186+%	1,200	16.0	14.5 - 17.6	6,100	84.0	82.4 - 85.5	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,000	37.5	33.8 - 41.2	1,700	62.5	58.8 - 66.2	2,600
\$10,000-\$24,999	900	28.3	25.4 - 31.4	2,300	71.7	68.6 - 74.6	3,300
\$25,000-\$49,999	900	22.8	20.3 - 25.4	2,900	77.2	74.6 - 79.7	3,800
>=\$50,000	700	12.9	11.3 - 14.6	4,700	87.1	85.4 - 88.7	5,400

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	25.5	21.1 - 30.4	900	74.5	69.6 - 78.9	1,200
Normal birth weight	3,400	22.9	21.6 - 24.2	11,400	77.1	75.8 - 78.4	14,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	42.6	37.0 - 48.3	700	57.4	51.7 - 63.0	1,200
No	3,100	21.4	20.2 - 22.7	11,500	78.6	77.3 - 79.8	14,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	36.7	33.4 - 40.1	2,000	63.3	59.9 - 66.6	3,100
No	2,500	19.7	18.4 - 21.0	10,200	80.3	79.0 - 81.6	12,700

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	29.6	27.5 - 31.9	4,600	70.4	68.1 - 72.5	6,500
No	1,700	18.1	16.7 - 19.7	7,600	81.9	80.3 - 83.3	9,200

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,700	23.2	22.0 - 24.4	12,100	76.8	75.6 - 78.0	15,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	21.6	18.2 - 25.3	1,500	78.4	74.7 - 81.8	1,900
Honolulu	2,600	23.0	21.6 - 24.4	8,900	77.0	75.6 - 78.4	11,500
Kauai	100	18.3	13.6 - 24.2	600	81.7	75.8 - 86.4	700
Maui	500	28.9	25.0 - 33.1	1,200	71.1	66.9 - 75.0	1,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	18.6	16.4 - 21.0	3,000	81.4	79.0 - 83.6	3,700
Native Hawaiian	1,300	30.7	28.0 - 33.6	2,900	69.3	66.4 - 72.0	4,200
Chinese	100	14.1	11.8 - 16.9	500	85.9	83.1 - 88.2	600
Filipino	900	28.7	26.0 - 31.5	2,100	71.3	68.5 - 74.0	3,000
Japanese	200	11.7	9.4 - 14.5	1,800	88.3	85.5 - 90.6	2,100
Other	500	22.5	19.4 - 26.0	1,800	77.5	74.0 - 80.6	2,300

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	38.0	32.9 - 43.4	800	62.0	56.6 - 67.1	1,200
20-24	1,300	34.1	31.3 - 37.0	2,600	65.9	63.0 - 68.7	3,900
25-34	1,500	18.8	17.2 - 20.4	6,300	81.2	79.6 - 82.8	7,800
35+	400	14.2	12.1 - 16.6	2,500	85.8	83.4 - 87.9	2,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	24.7	23.0 - 26.4	6,200	75.3	73.6 - 77.0	8,200
Female	1,600	21.6	19.9 - 23.4	5,900	78.4	76.6 - 80.1	7,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.4	17.1 - 19.8	8,600	81.6	80.2 - 82.9	10,500
No	1,700	32.8	30.4 - 35.3	3,500	67.2	64.7 - 69.6	5,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	31.7	29.4 - 34.1	3,700	68.3	65.9 - 70.6	5,500
131-185%	400	22.1	18.7 - 25.8	1,400	77.9	74.2 - 81.3	1,900
186+%	1,200	16.6	15.2 - 18.2	6,100	83.4	81.8 - 84.8	7,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	36.8	33.2 - 40.5	1,600	63.2	59.5 - 66.8	2,500
\$10,000-\$24,999	1,000	28.8	26.1 - 31.8	2,400	71.2	68.2 - 73.9	3,400
\$25,000-\$49,999	900	22.2	19.9 - 24.7	3,000	77.8	75.3 - 80.1	3,900
>=\$50,000	600	12.7	11.2 - 14.5	4,300	87.3	85.5 - 88.8	5,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	24.4	20.0 - 29.3	800	75.6	70.7 - 80.0	1,100
Normal birth weight	3,400	23.1	21.9 - 24.4	11,300	76.9	75.6 - 78.1	14,700

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	37.2	31.9 - 42.8	700	62.8	57.2 - 68.1	1,100
No	3,200	21.9	20.7 - 23.2	11,300	78.1	76.8 - 79.3	14,400

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	32.7	29.6 - 35.9	2,100	67.3	64.1 - 70.4	3,100
No	2,600	20.6	19.4 - 21.9	9,900	79.4	78.1 - 80.6	12,500

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	29.1	27.0 - 31.2	4,700	70.9	68.8 - 73.0	6,600
No	1,700	18.7	17.2 - 20.2	7,200	81.3	79.8 - 82.8	8,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,600	23.5	22.3 - 24.8	11,700	76.5	75.2 - 77.7	15,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	21.4	18.4 - 24.7	1,400	78.6	75.3 - 81.6	1,800
Honolulu	2,600	23.4	21.9 - 25.0	8,500	76.6	75.0 - 78.1	11,200
Kauai	100	17.3	13.4 - 22.2	600	82.7	77.8 - 86.6	700
Maui	500	29.4	25.9 - 33.1	1,100	70.6	66.9 - 74.1	1,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	17.6	15.2 - 20.3	2,900	82.4	79.7 - 84.8	3,500
Native Hawaiian	1,300	31.1	28.6 - 33.6	2,800	68.9	66.4 - 71.4	4,000
Chinese	100	11.3	8.2 - 15.4	500	88.7	84.6 - 91.8	600
Filipino	900	30.4	27.4 - 33.5	2,100	69.6	66.5 - 72.6	3,000
Japanese	300	12.7	10.1 - 16.0	1,800	87.3	84.0 - 89.9	2,000
Other	500	23.3	19.7 - 27.2	1,600	76.7	72.8 - 80.3	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	42.9	37.7 - 48.3	700	57.1	51.7 - 62.3	1,200
20-24	1,200	34.0	31.2 - 37.0	2,400	66.0	63.0 - 68.8	3,700
25-34	1,400	18.9	17.3 - 20.6	6,200	81.1	79.4 - 82.7	7,600
35+	400	14.0	11.7 - 16.6	2,400	86.0	83.4 - 88.3	2,800

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,900	24.0	22.3 - 25.8	6,100	76.0	74.2 - 77.7	8,000
Female	1,700	23.0	21.2 - 24.9	5,600	77.0	75.1 - 78.8	7,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	18.8	17.4 - 20.3	8,500	81.2	79.7 - 82.6	10,400
No	1,600	33.8	31.3 - 36.3	3,200	66.2	63.7 - 68.7	4,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,800	31.6	29.3 - 33.9	3,900	68.4	66.1 - 70.7	5,700
131-185%	400	22.7	19.3 - 26.5	1,500	77.3	73.5 - 80.7	1,900
186+%	1,000	15.5	14.0 - 17.2	5,500	84.5	82.8 - 86.0	6,500

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	26.9	22.0 - 32.5	800	73.1	67.5 - 78.0	1,000
Normal birth weight	3,300	23.3	22.0 - 24.6	10,900	76.7	75.4 - 78.0	14,200

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	36.2	31.0 - 41.8	700	63.8	58.2 - 69.0	1,000
No	3,200	22.5	21.2 - 23.8	10,900	77.5	76.2 - 78.8	14,000

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	34.3	31.1 - 37.7	1,900	65.7	62.3 - 68.9	2,900
No	2,500	20.7	19.4 - 22.1	9,600	79.3	77.9 - 80.6	12,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	30.0	28.0 - 32.2	4,400	70.0	67.8 - 72.0	6,300
No	1,600	18.7	17.2 - 20.4	7,100	81.3	79.6 - 82.8	8,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,500	23.3	22.0 - 24.6	11,500	76.7	75.4 - 78.0	15,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	23.8	21.1 - 26.7	1,300	76.2	73.3 - 78.9	1,700
Honolulu	2,500	22.8	21.1 - 24.5	8,400	77.2	75.5 - 78.9	10,900
Kauai	100	20.2	16.3 - 24.8	500	79.8	75.2 - 83.7	700
Maui	400	27.4	24.4 - 30.6	1,200	72.6	69.4 - 75.6	1,600

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	18.3	15.6 - 21.3	2,800	81.7	78.7 - 84.4	3,400
Native Hawaiian	1,200	30.3	28.2 - 32.5	2,700	69.7	67.5 - 71.8	3,900
Chinese	100	13.1	8.5 - 19.7	500	86.9	80.3 - 91.5	600
Filipino	900	29.4	26.1 - 32.8	2,100	70.6	67.2 - 73.9	3,000
Japanese	200	11.3	8.6 - 14.6	1,800	88.7	85.4 - 91.4	2,100
Other	500	24.2	20.1 - 28.8	1,500	75.8	71.2 - 79.9	2,000

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	44.0	38.7 - 49.4	700	56.0	50.6 - 61.3	1,200
20-24	1,200	32.1	29.1 - 35.1	2,400	67.9	64.9 - 70.9	3,600
25-34	1,400	18.8	17.1 - 20.6	6,100	81.2	79.4 - 82.9	7,500
35+	400	14.6	12.2 - 17.5	2,300	85.4	82.5 - 87.8	2,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,900	23.9	22.1 - 25.9	5,900	76.1	74.1 - 77.9	7,800
Female	1,600	22.5	20.7 - 24.5	5,500	77.5	75.5 - 79.3	7,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	18.7	17.2 - 20.3	8,500	81.3	79.7 - 82.8	10,400
No	1,500	33.8	31.3 - 36.4	3,000	66.2	63.6 - 68.7	4,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	30.5	28.2 - 32.8	4,000	69.5	67.2 - 71.8	5,700
131-185%	500	22.4	19.1 - 26.0	1,800	77.6	74.0 - 80.9	2,300
186+%	900	15.2	13.4 - 17.1	4,900	84.8	82.9 - 86.6	5,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight							
Low birth weight	300	28.6	23.0 - 35.0	700	71.4	65.0 - 77.0	1,000
Normal birth weight	3,200	22.9	21.6 - 24.3	10,800	77.1	75.7 - 78.4	14,000

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester							
Yes	300	34.5	29.1 - 40.2	600	65.5	59.8 - 70.9	1,000
No	3,100	22.4	21.1 - 23.8	10,700	77.6	76.2 - 78.9	13,800

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant							
Yes	900	32.8	29.5 - 36.3	1,800	67.2	63.7 - 70.5	2,700
No	2,500	20.8	19.4 - 22.3	9,400	79.2	77.7 - 80.6	11,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	30.7	28.5 - 33.0	4,200	69.3	67.0 - 71.5	6,100
No	1,600	18.2	16.6 - 19.8	7,100	81.8	80.2 - 83.4	8,700

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,400	23.4	22.0 - 24.8	11,000	76.6	75.2 - 78.0	14,400

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	25.2	22.7 - 27.7	1,200	74.8	72.3 - 77.3	1,600
Honolulu	2,400	22.9	21.2 - 24.7	8,200	77.1	75.3 - 78.8	10,600
Kauai	100	21.0	17.4 - 25.0	500	79.0	75.0 - 82.6	700
Maui	400	25.8	23.2 - 28.6	1,100	74.2	71.4 - 76.8	1,500

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	17.6	14.8 - 20.7	2,600	82.4	79.3 - 85.2	3,200
Native Hawaiian	1,100	29.7	28.1 - 31.4	2,600	70.3	68.6 - 71.9	3,700
Chinese	100	17.1	11.3 - 25.2	500	82.9	74.8 - 88.7	600
Filipino	900	28.9	25.5 - 32.5	2,200	71.1	67.5 - 74.5	3,000
Japanese	300	12.7	9.8 - 16.3	1,800	87.3	83.7 - 90.2	2,000
Other	500	25.4	21.0 - 30.4	1,400	74.6	69.6 - 79.0	1,800

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	45.3	40.4 - 50.3	700	54.7	49.7 - 59.6	1,300
20-24	1,100	31.3	28.4 - 34.4	2,400	68.7	65.6 - 71.6	3,500
25-34	1,400	18.9	17.2 - 20.8	5,800	81.1	79.2 - 82.8	7,200
35+	300	13.6	11.1 - 16.6	2,100	86.4	83.4 - 88.9	2,400

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Infant Gender							
Male	1,800	24.0	22.1 - 26.0	5,600	76.0	74.0 - 77.9	7,400
Female	1,600	22.7	20.8 - 24.6	5,400	77.3	75.4 - 79.2	7,000

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Married							
Yes	1,800	18.5	17.0 - 20.2	8,100	81.5	79.8 - 83.0	9,900
No	1,500	34.1	31.7 - 36.7	2,900	65.9	63.3 - 68.3	4,400

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Poverty Level							
0-130%	1,600	29.9	27.7 - 32.3	3,900	70.1	67.7 - 72.3	5,500
131-185%	600	22.2	19.3 - 25.4	2,200	77.8	74.6 - 80.7	2,800
186+%	700	13.9	12.1 - 16.0	4,300	86.1	84.0 - 87.9	5,000

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	900	32.7	29.5 - 36.0	1,800	67.3	64.0 - 70.5	2,700
\$15,001-\$25,000	500	27.1	23.5 - 31.1	1,400	72.9	68.9 - 76.5	1,900
\$25,001-\$40,000	700	24.9	21.9 - 28.1	2,300	75.1	71.9 - 78.1	3,000
>=\$40,001	900	15.1	13.3 - 17.1	5,000	84.9	82.9 - 86.7	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	29.6	23.8 - 36.2	700	70.4	63.8 - 76.2	900
Normal birth weight	3,100	22.9	21.6 - 24.4	10,400	77.1	75.6 - 78.4	13,400

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	36.2	30.9 - 41.8	600	63.8	58.2 - 69.1	1,000
No	3,000	22.4	21.0 - 23.9	10,200	77.6	76.1 - 79.0	13,200

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	34.7	31.4 - 38.3	1,700	65.3	61.7 - 68.6	2,600
No	2,300	20.5	19.1 - 22.0	9,100	79.5	78.0 - 80.9	11,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	32.1	29.8 - 34.4	3,900	67.9	65.6 - 70.2	5,700
No	1,500	17.6	16.0 - 19.3	7,000	82.4	80.7 - 84.0	8,500

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,300	23.1	21.8 - 24.3	10,900	76.9	75.7 - 78.2	14,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	27.3	24.8 - 29.9	1,200	72.7	70.1 - 75.2	1,600
Honolulu	2,300	22.0	20.4 - 23.6	8,100	78.0	76.4 - 79.6	10,400
Kauai	100	22.7	18.9 - 27.1	500	77.3	72.9 - 81.1	600
Maui	400	25.8	23.1 - 28.6	1,100	74.2	71.4 - 76.9	1,500

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	18.3	15.8 - 21.1	2,600	81.7	78.9 - 84.2	3,200
Native Hawaiian	1,100	30.2	28.6 - 31.8	2,500	69.8	68.2 - 71.4	3,600
Chinese	100	22.1	15.5 - 30.5	400	77.9	69.5 - 84.5	600
Filipino	800	27.1	23.9 - 30.4	2,100	72.9	69.6 - 76.1	2,900
Japanese	200	11.3	8.7 - 14.5	1,800	88.7	85.5 - 91.3	2,000
Other	400	24.1	20.0 - 28.9	1,400	75.9	71.1 - 80.0	1,800

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	41.5	37.2 - 45.8	800	58.5	54.2 - 62.8	1,400
20-24	1,100	30.5	27.8 - 33.3	2,400	69.5	66.7 - 72.2	3,500
25-34	1,300	18.6	16.9 - 20.4	5,700	81.4	79.6 - 83.1	6,900
35+	300	14.5	12.0 - 17.4	2,000	85.5	82.6 - 88.0	2,300

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,700	23.5	21.7 - 25.3	5,500	76.5	74.7 - 78.3	7,100
Female	1,600	22.6	20.9 - 24.5	5,400	77.4	75.5 - 79.1	7,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	18.5	17.1 - 20.0	8,000	81.5	80.0 - 82.9	9,800
No	1,400	33.2	30.9 - 35.6	2,900	66.8	64.4 - 69.1	4,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,600	29.1	27.0 - 31.3	3,800	70.9	68.7 - 73.0	5,400
131-185%	600	21.2	18.5 - 24.1	2,100	78.8	75.9 - 81.5	2,700
186+%	800	15.3	13.5 - 17.3	4,300	84.7	82.7 - 86.5	5,100

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	900	32.4	29.5 - 35.4	1,900	67.6	64.6 - 70.5	2,700
\$15,001-\$25,000	600	27.1	23.8 - 30.7	1,500	72.9	69.3 - 76.2	2,000
\$25,001-\$40,000	700	22.7	20.1 - 25.6	2,300	77.3	74.4 - 79.9	3,000
>=\$40,001	900	15.5	13.8 - 17.3	4,800	84.5	82.7 - 86.2	5,700

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	29.9	24.4 - 36.0	600	70.1	64.0 - 75.6	900
Normal birth weight	3,000	22.5	21.2 - 23.8	10,200	77.5	76.2 - 78.8	13,200

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	39.7	34.6 - 45.0	600	60.3	55.0 - 65.4	900
No	2,800	21.6	20.3 - 22.9	10,100	78.4	77.1 - 79.7	12,900

PRAMS Health Indicator 3-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	34.7	31.5 - 37.9	1,700	65.3	62.1 - 68.5	2,600
No	2,200	19.9	18.6 - 21.3	8,900	80.1	78.7 - 81.4	11,100

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	30.3	28.2 - 32.4	4,000	69.7	67.6 - 71.8	5,700
No	1,500	18.0	16.5 - 19.6	6,800	82.0	80.4 - 83.5	8,300