

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration (grouped), among initiators	Breastfeeding (any) duration (grouped), among those who initiated breastfeeding

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/22/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- for
the State of Hawaii, for the Year(s)**

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013


Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	600	4.3	3.9 - 4.8	2,400	16.1	15.3 - 17.0	500	3.6	3.2 - 4.1	11,100	75.9	74.9 - 76.9	14,600
2001-2005	600	4.4	3.9 - 4.9	2,400	16.1	15.2 - 17.0	500	3.5	3.1 - 4.0	11,300	76.0	75.0 - 77.0	14,800
2002-2006	600	4.0	3.6 - 4.5	2,500	16.2	15.3 - 17.1	600	3.6	3.2 - 4.1	11,700	76.2	75.1 - 77.2	15,300
2003-2007	600	3.9	3.5 - 4.4	2,600	16.3	15.5 - 17.2	600	3.7	3.3 - 4.2	12,000	76.0	75.0 - 77.0	15,700
2004-2008	600	3.6	3.2 - 4.0	2,600	16.1	15.3 - 16.9	600	3.8	3.4 - 4.2	12,400	76.5	75.6 - 77.5	16,100
2005-2009	600	3.7	3.2 - 4.2	2,600	16.1	15.2 - 17.0	700	4.0	3.6 - 4.6	12,500	76.2	75.1 - 77.2	16,300
2006-2010	600	3.5	3.0 - 4.0	2,600	15.7	14.8 - 16.7	700	4.2	3.7 - 4.8	12,800	76.6	75.5 - 77.7	16,700
2007-2011	600	3.3	2.8 - 3.8	2,500	14.8	13.8 - 15.8	700	4.3	3.8 - 4.9	13,200	77.7	76.5 - 78.8	16,900
2008-2012	500	3.1	2.6 - 3.6	2,400	14.2	13.2 - 15.2	700	4.3	3.7 - 4.9	13,400	78.5	77.3 - 79.7	17,100
2009-2013	500	2.9	2.4 - 3.4	2,300	13.5	12.5 - 14.5	700	4.0	3.4 - 4.6	13,700	79.7	78.4 - 80.8	17,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	2.9	2.4 - 3.4	2,300	13.5	12.5 - 14.5	700	4.0	3.4 - 4.6	13,700	79.7	78.4 - 80.8	17,200

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.1	2.4 - 4.1	300	14.8	13.1 - 16.6	100	3.7	2.9 - 4.8	1,700	78.4	76.3 - 80.3	2,200
Honolulu	400	2.9	2.3 - 3.7	1,600	13.3	11.9 - 14.7	500	4.0	3.3 - 4.9	9,800	79.8	78.1 - 81.3	12,300
Kauai	<50	1.7	1.1 - 2.6	100	11.8	10.1 - 13.7	<50	3.3	2.5 - 4.5	700	83.2	81.1 - 85.2	800
Maui	100	2.9	2.2 - 3.8	300	14.1	12.6 - 15.8	100	4.0	3.2 - 4.9	1,500	79.0	77.1 - 80.8	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.7	1.8 - 4.1	400	9.8	8.0 - 11.9	200	4.2	3.1 - 5.7	3,600	83.3	80.7 - 85.5	4,300
Native Hawaiian	200	3.3	2.5 - 4.5	900	18.2	16.3 - 20.4	200	4.9	3.9 - 6.2	3,700	73.5	71.0 - 75.8	5,000
Chinese	<50	2.5	1.0 - 6.2	100	8.6	5.2 - 13.7	<50	2.2	0.8 - 5.7	700	86.7	80.8 - 91.0	800
Filipino	100	3.1	2.2 - 4.5	500	18.1	15.5 - 20.9	100	4.3	3.0 - 6.0	2,300	74.5	71.3 - 77.4	3,000
Japanese	<50	1.0	0.4 - 2.4	100	6.2	4.3 - 9.0	<50	1.7	0.8 - 3.3	1,500	91.1	88.1 - 93.5	1,700
Other	100	3.4	2.1 - 5.3	300	11.1	8.7 - 14.1	100	3.3	2.0 - 5.2	2,000	82.2	78.8 - 85.2	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	4.6	2.7 - 7.6	300	29.4	24.4 - 34.9	100	5.6	3.4 - 9.1	700	60.4	54.6 - 65.9	1,100
20-24	100	3.3	2.3 - 4.7	600	16.5	14.3 - 18.9	200	4.1	3.0 - 5.6	3,000	76.1	73.3 - 78.7	3,900
25-34	200	2.6	2.0 - 3.3	1,100	12.1	10.7 - 13.5	400	3.9	3.2 - 4.8	7,500	81.5	79.8 - 83.0	9,200
35+	100	2.7	1.8 - 4.1	300	8.3	6.6 - 10.4	100	3.4	2.3 - 4.8	2,600	85.6	83.0 - 87.8	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.1	2.5 - 4.0	1,200	13.2	11.9 - 14.7	300	4.0	3.3 - 4.9	6,900	79.6	77.9 - 81.3	8,700
Female	200	2.6	2.0 - 3.4	1,200	13.7	12.3 - 15.3	300	3.9	3.2 - 4.8	6,800	79.7	77.9 - 81.4	8,500

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.6	2.0 - 3.3	1,200	10.8	9.6 - 12.1	400	3.6	3.0 - 4.4	9,000	83.0	81.5 - 84.4	10,800
No	200	3.4	2.6 - 4.4	1,200	18.0	16.3 - 19.9	300	4.5	3.6 - 5.7	4,700	74.0	71.9 - 76.1	6,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	200	3.9	3.1 - 5.1	1,100	17.8	16.0 - 19.8	200	3.9	3.0 - 5.1	4,400	74.3	72.1 - 76.5	6,000
131-185%	100	3.3	2.1 - 5.2	400	14.4	11.8 - 17.4	100	4.5	3.2 - 6.4	2,100	77.8	74.3 - 80.9	2,600
186+%	200	2.1	1.5 - 2.9	600	8.7	7.4 - 10.2	300	3.8	3.0 - 4.8	6,200	85.4	83.6 - 87.0	7,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight													
Low birth weight	<50	3.0	2.2 - 4.0	200	16.4	14.6 - 18.3	100	5.1	4.1 - 6.4	900	75.5	73.3 - 77.7	1,200
Normal birth weight	500	2.9	2.4 - 3.5	2,100	13.3	12.2 - 14.4	600	3.9	3.3 - 4.6	12,800	80.0	78.6 - 81.2	16,000

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester													
Yes	100	7.6	4.8 - 11.7	200	25.3	20.4 - 30.8	100	6.0	3.6 - 9.7	600	61.2	55.1 - 66.9	1,000
No	400	2.7	2.2 - 3.2	2,000	12.7	11.7 - 13.8	600	3.8	3.3 - 4.5	12,800	80.8	79.5 - 82.0	15,900

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant													
Yes	200	5.3	3.9 - 7.2	700	20.7	18.1 - 23.6	200	5.6	4.2 - 7.5	2,200	68.4	65.1 - 71.5	3,200
No	300	2.4	1.9 - 3.0	1,600	11.8	10.7 - 12.9	500	3.6	3.0 - 4.2	11,300	82.3	80.9 - 83.5	13,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.2	3.3 - 5.2	1,300	18.2	16.5 - 20.0	300	4.2	3.4 - 5.2	5,400	73.5	71.4 - 75.4	7,300
No	200	1.9	1.4 - 2.6	1,000	9.9	8.8 - 11.2	400	3.7	3.0 - 4.5	8,200	84.5	82.9 - 85.9	9,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	500	3.1	2.6 - 3.6	2,400	14.2	13.2 - 15.2	700	4.3	3.7 - 4.9	13,400	78.5	77.3 - 79.7	17,100

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.8	2.9 - 5.0	300	15.8	14.0 - 17.8	100	3.6	2.8 - 4.7	1,700	76.7	74.5 - 78.9	2,200
Honolulu	400	3.0	2.4 - 3.7	1,700	14.1	12.8 - 15.4	500	4.3	3.6 - 5.2	9,600	78.6	77.0 - 80.1	12,200
Kauai	<50	1.8	1.1 - 2.9	100	11.9	9.8 - 14.3	<50	4.8	3.4 - 6.9	700	81.5	78.5 - 84.1	800
Maui	100	3.1	2.3 - 4.1	300	13.7	12.1 - 15.5	100	4.3	3.4 - 5.4	1,500	79.0	76.9 - 80.9	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.7	1.8 - 3.9	400	10.5	8.8 - 12.5	200	5.0	3.9 - 6.4	3,400	81.8	79.4 - 84.0	4,100
Native Hawaiian	200	3.6	2.7 - 4.7	900	18.4	16.5 - 20.5	300	5.4	4.3 - 6.7	3,600	72.6	70.3 - 74.9	4,900
Chinese	<50	2.7	1.1 - 6.6	<50	6.7	4.0 - 10.9	<50	3.0	1.4 - 6.4	600	87.6	82.3 - 91.5	700
Filipino	100	3.2	2.4 - 4.4	600	19.1	16.6 - 21.9	100	4.1	2.9 - 5.6	2,300	73.6	70.6 - 76.4	3,100
Japanese	<50	1.5	0.8 - 3.0	100	7.6	5.6 - 10.4	<50	2.2	1.3 - 3.8	1,500	88.6	85.5 - 91.1	1,700
Other	100	3.5	2.3 - 5.4	300	12.5	10.2 - 15.3	100	2.9	1.8 - 4.6	2,000	81.0	77.8 - 83.9	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.1	3.2 - 8.1	300	28.4	23.9 - 33.5	100	5.1	3.2 - 8.1	700	61.3	56.0 - 66.3	1,200
20-24	100	3.5	2.5 - 4.9	700	19.3	17.0 - 21.8	200	4.5	3.5 - 5.9	2,800	72.7	69.9 - 75.3	3,800
25-34	200	2.5	2.0 - 3.3	1,100	12.0	10.7 - 13.3	400	4.4	3.6 - 5.3	7,300	81.1	79.5 - 82.6	9,000
35+	100	3.2	2.2 - 4.5	300	8.7	7.0 - 10.7	100	3.2	2.3 - 4.5	2,600	84.9	82.5 - 87.1	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.3	2.6 - 4.1	1,300	14.3	13.0 - 15.7	400	4.5	3.8 - 5.4	6,800	77.9	76.2 - 79.5	8,800
Female	200	2.8	2.2 - 3.6	1,200	14.0	12.7 - 15.5	300	4.0	3.3 - 4.9	6,600	79.2	77.5 - 80.8	8,300

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.6	2.1 - 3.3	1,200	11.5	10.4 - 12.7	400	4.1	3.4 - 4.8	8,800	81.9	80.4 - 83.2	10,700
No	200	3.8	3.0 - 4.8	1,200	18.7	17.0 - 20.5	300	4.6	3.7 - 5.7	4,600	72.9	70.8 - 74.9	6,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	200	4.2	3.3 - 5.3	1,100	18.7	16.9 - 20.6	200	3.9	3.1 - 4.9	4,300	73.3	71.1 - 75.3	5,900
131-185%	100	3.2	2.0 - 5.0	400	14.3	11.8 - 17.1	100	4.7	3.4 - 6.5	2,100	77.8	74.6 - 80.8	2,700
186+%	200	2.1	1.6 - 2.9	700	9.6	8.4 - 11.0	300	4.5	3.6 - 5.5	6,100	83.8	82.1 - 85.3	7,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	3.6	2.6 - 5.0	200	16.4	14.2 - 18.7	100	5.0	3.9 - 6.4	900	75.0	72.3 - 77.5	1,200
Normal birth weight	500	3.0	2.5 - 3.6	2,200	14.0	13.0 - 15.1	700	4.2	3.7 - 4.9	12,500	78.8	77.5 - 80.0	15,900

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	6.3	3.9 - 9.9	300	26.5	21.9 - 31.6	100	7.7	5.2 - 11.4	600	59.5	53.9 - 64.9	1,000
No	400	2.9	2.4 - 3.4	2,100	13.3	12.3 - 14.3	600	4.0	3.5 - 4.7	12,500	79.8	78.6 - 81.0	15,700

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.6	3.4 - 6.4	700	21.7	19.2 - 24.5	200	6.9	5.4 - 8.8	2,100	66.7	63.6 - 69.8	3,100
No	400	2.7	2.2 - 3.3	1,700	12.4	11.3 - 13.4	500	3.7	3.1 - 4.3	11,100	81.3	80.0 - 82.5	13,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.4	3.6 - 5.4	1,400	18.8	17.2 - 20.6	300	4.2	3.4 - 5.2	5,200	72.5	70.6 - 74.4	7,200
No	200	2.1	1.6 - 2.7	1,000	10.7	9.6 - 12.0	400	4.2	3.5 - 5.0	8,000	83.0	81.6 - 84.4	9,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.3	2.8 - 3.8	2,500	14.8	13.8 - 15.8	700	4.3	3.8 - 4.9	13,200	77.7	76.5 - 78.8	16,900

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.4	3.4 - 5.8	300	14.8	13.0 - 16.9	100	3.1	2.3 - 4.1	1,700	77.6	75.3 - 79.8	2,200
Honolulu	400	3.1	2.5 - 3.8	1,800	15.0	13.8 - 16.3	500	4.5	3.8 - 5.3	9,300	77.4	75.9 - 78.9	12,000
Kauai	<50	1.4	0.7 - 2.7	100	12.5	10.1 - 15.4	<50	5.3	3.7 - 7.5	700	80.8	77.4 - 83.8	800
Maui	100	3.8	2.8 - 5.0	300	14.4	12.6 - 16.5	100	3.9	3.0 - 5.1	1,500	77.9	75.5 - 80.1	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.8	1.9 - 3.9	500	11.0	9.4 - 12.9	200	5.5	4.3 - 6.9	3,300	80.8	78.4 - 82.9	4,100
Native Hawaiian	200	4.4	3.4 - 5.6	900	19.4	17.5 - 21.5	200	4.8	3.8 - 6.0	3,500	71.4	69.0 - 73.6	4,800
Chinese	<50	2.1	0.9 - 4.9	<50	7.0	4.5 - 10.9	<50	2.7	1.3 - 5.2	600	88.2	83.6 - 91.5	700
Filipino	100	2.8	2.0 - 3.7	600	19.0	16.7 - 21.5	100	4.4	3.3 - 5.7	2,200	73.9	71.1 - 76.5	3,000
Japanese	<50	1.7	0.9 - 3.1	100	8.0	6.0 - 10.5	<50	2.4	1.4 - 4.0	1,600	87.9	85.0 - 90.4	1,800
Other	100	3.9	2.6 - 5.7	300	13.8	11.4 - 16.6	100	3.1	2.1 - 4.8	2,000	79.2	76.0 - 82.1	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.6	3.8 - 8.2	400	27.9	23.6 - 32.6	100	5.4	3.5 - 8.3	800	61.1	56.2 - 65.8	1,300
20-24	100	3.4	2.5 - 4.7	800	20.5	18.3 - 22.9	200	4.4	3.4 - 5.7	2,800	71.7	69.1 - 74.1	3,900
25-34	200	2.8	2.3 - 3.6	1,100	12.3	11.1 - 13.6	400	4.3	3.5 - 5.1	7,000	80.6	79.0 - 82.0	8,700
35+	100	3.2	2.3 - 4.6	300	8.9	7.3 - 10.8	100	3.8	2.8 - 5.1	2,600	84.1	81.7 - 86.2	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.1	2.5 - 3.9	1,300	15.1	13.8 - 16.5	400	4.7	3.9 - 5.5	6,800	77.1	75.5 - 78.7	8,800
Female	300	3.4	2.8 - 4.2	1,200	14.4	13.1 - 15.8	300	3.9	3.2 - 4.7	6,400	78.3	76.6 - 79.8	8,200

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.7	2.2 - 3.3	1,200	11.7	10.6 - 12.8	400	4.1	3.4 - 4.8	8,700	81.6	80.2 - 82.9	10,700
No	300	4.2	3.4 - 5.2	1,300	20.1	18.4 - 21.9	300	4.7	3.8 - 5.7	4,400	71.0	69.0 - 73.0	6,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.4	3.5 - 5.4	1,200	20.0	18.3 - 21.9	200	4.0	3.2 - 5.0	4,200	71.6	69.6 - 73.6	5,900
131-185%	100	3.6	2.4 - 5.4	400	14.3	12.0 - 17.0	100	4.4	3.1 - 6.0	2,000	77.7	74.6 - 80.6	2,600
186+%	200	2.4	1.9 - 3.1	700	9.9	8.7 - 11.2	300	4.5	3.7 - 5.4	6,100	83.2	81.6 - 84.7	7,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	4.9	3.6 - 6.5	600	21.9	19.3 - 24.8	100	3.9	2.7 - 5.4	2,000	69.4	66.2 - 72.3	2,800
\$10,000-\$24,999	100	4.2	3.1 - 5.6	600	17.2	15.1 - 19.5	200	4.6	3.5 - 6.0	2,400	74.0	71.3 - 76.6	3,300
\$25,000-\$49,999	100	3.3	2.4 - 4.7	600	15.6	13.6 - 17.7	200	4.6	3.5 - 6.1	3,000	76.5	73.9 - 78.8	3,900
>=\$50,000	100	2.1	1.6 - 2.9	500	8.7	7.5 - 10.1	200	4.0	3.2 - 5.1	5,000	85.1	83.4 - 86.7	5,900

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	<50	3.9	2.7 - 5.4	200	17.8	15.4 - 20.6	100	4.9	3.7 - 6.4	900	73.5	70.4 - 76.3	1,200
Normal birth weight	500	3.2	2.7 - 3.8	2,300	14.5	13.6 - 15.6	700	4.2	3.7 - 4.9	12,300	78.0	76.8 - 79.2	15,700

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.0	4.7 - 10.4	300	27.7	23.4 - 32.4	100	6.9	4.8 - 9.9	700	58.4	53.3 - 63.3	1,200
No	500	3.0	2.5 - 3.5	2,100	13.7	12.8 - 14.7	600	4.1	3.6 - 4.7	12,300	79.2	78.0 - 80.3	15,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	4.7	3.5 - 6.4	800	24.2	21.7 - 26.9	200	6.3	4.9 - 8.0	2,100	64.7	61.7 - 67.6	3,200
No	400	2.9	2.4 - 3.5	1,700	12.5	11.5 - 13.5	500	3.8	3.3 - 4.4	10,900	80.8	79.6 - 82.0	13,500

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.3	3.5 - 5.2	1,400	19.5	17.9 - 21.2	300	4.3	3.5 - 5.2	5,000	71.9	70.0 - 73.8	7,000
No	200	2.5	2.0 - 3.2	1,100	11.3	10.2 - 12.5	400	4.3	3.6 - 5.1	7,900	81.8	80.4 - 83.2	9,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.5	3.0 - 4.0	2,600	15.7	14.8 - 16.7	700	4.2	3.7 - 4.8	12,800	76.6	75.5 - 77.7	16,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.8	3.7 - 6.3	300	15.5	13.5 - 17.8	100	2.7	1.9 - 3.8	1,600	77.0	74.4 - 79.4	2,100
Honolulu	400	3.4	2.8 - 4.1	1,900	15.9	14.8 - 17.2	500	4.4	3.8 - 5.2	9,100	76.2	74.8 - 77.6	11,900
Kauai	<50	1.2	0.6 - 2.5	100	13.7	10.9 - 17.2	<50	5.2	3.5 - 7.6	700	79.9	76.0 - 83.2	800
Maui	100	3.5	2.5 - 4.8	300	15.3	13.3 - 17.7	100	3.9	2.9 - 5.3	1,500	77.2	74.6 - 79.7	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.0	2.2 - 4.2	500	11.4	9.8 - 13.2	200	5.9	4.7 - 7.3	3,200	79.7	77.4 - 81.8	4,000
Native Hawaiian	200	4.7	3.8 - 6.0	1,000	21.1	19.2 - 23.2	200	4.0	3.2 - 5.1	3,300	70.1	67.8 - 72.3	4,700
Chinese	<50	1.7	1.0 - 2.8	100	8.7	6.0 - 12.5	<50	2.9	1.5 - 5.5	600	86.7	82.6 - 90.0	700
Filipino	100	3.2	2.4 - 4.2	600	19.7	17.4 - 22.1	100	4.5	3.4 - 5.9	2,200	72.7	70.0 - 75.2	3,000
Japanese	<50	2.0	1.2 - 3.5	100	7.5	5.7 - 9.7	<50	2.1	1.3 - 3.6	1,600	88.3	85.7 - 90.6	1,800
Other	100	3.9	2.6 - 5.6	400	15.5	13.1 - 18.3	100	3.3	2.3 - 4.8	1,900	77.3	74.2 - 80.2	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.8	4.0 - 8.4	400	27.0	23.0 - 31.4	100	5.0	3.2 - 7.7	800	62.3	57.6 - 66.7	1,400
20-24	100	3.6	2.7 - 4.9	900	22.4	20.3 - 24.7	200	5.1	4.0 - 6.4	2,700	68.9	66.3 - 71.3	3,900
25-34	300	3.1	2.5 - 3.8	1,100	13.0	11.8 - 14.2	300	3.9	3.3 - 4.7	6,800	80.0	78.5 - 81.4	8,400
35+	100	3.5	2.5 - 4.8	300	9.6	8.0 - 11.6	100	3.4	2.5 - 4.6	2,500	83.5	81.1 - 85.6	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.2	2.6 - 4.0	1,400	16.5	15.2 - 17.9	400	4.8	4.1 - 5.7	6,500	75.4	73.8 - 76.9	8,700
Female	300	3.8	3.1 - 4.6	1,200	14.9	13.6 - 16.2	300	3.5	2.9 - 4.3	6,300	77.8	76.2 - 79.4	8,000

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.8	2.3 - 3.4	1,300	12.2	11.2 - 13.3	400	3.9	3.3 - 4.6	8,600	81.2	79.8 - 82.4	10,600
No	300	4.7	3.8 - 5.8	1,300	21.8	20.1 - 23.6	300	4.8	3.9 - 5.8	4,200	68.7	66.7 - 70.7	6,100

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.9	4.0 - 6.0	1,300	21.8	20.0 - 23.7	200	3.7	2.9 - 4.6	4,000	69.7	67.6 - 71.7	5,800
131-185%	100	3.8	2.5 - 5.6	400	14.4	12.1 - 17.0	100	4.8	3.5 - 6.5	1,900	77.0	73.9 - 79.8	2,500
186+%	200	2.5	1.9 - 3.2	800	10.6	9.5 - 11.9	300	4.4	3.7 - 5.3	6,000	82.5	81.0 - 83.9	7,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	100	5.3	4.0 - 7.0	700	24.4	21.7 - 27.3	100	3.9	2.8 - 5.4	1,800	66.5	63.3 - 69.5	2,800
\$10,000-\$24,999	200	4.9	3.7 - 6.3	600	18.1	16.0 - 20.3	100	4.2	3.2 - 5.5	2,400	72.9	70.3 - 75.3	3,300
\$25,000-\$49,999	100	3.5	2.5 - 4.8	600	16.6	14.7 - 18.7	200	4.6	3.6 - 5.9	2,900	75.4	72.9 - 77.6	3,800
>=\$50,000	100	2.1	1.6 - 2.8	500	8.9	7.7 - 10.3	200	4.1	3.3 - 5.1	4,900	84.9	83.2 - 86.4	5,700

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	4.3	3.0 - 6.1	200	17.1	14.5 - 20.1	100	4.3	3.1 - 5.8	900	74.3	70.9 - 77.3	1,200
Normal birth weight	500	3.4	2.9 - 4.0	2,400	15.6	14.6 - 16.6	700	4.2	3.7 - 4.8	11,900	76.8	75.6 - 77.9	15,500

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.0	5.6 - 11.3	300	26.0	22.1 - 30.4	100	6.2	4.3 - 8.9	700	59.8	55.0 - 64.4	1,200
No	500	3.1	2.7 - 3.7	2,300	14.8	13.9 - 15.8	600	4.0	3.5 - 4.6	11,900	78.0	76.8 - 79.1	15,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.6	4.2 - 7.3	800	24.1	21.6 - 26.7	200	6.4	5.1 - 8.0	2,000	63.9	61.0 - 66.8	3,200
No	400	3.0	2.5 - 3.6	1,800	13.7	12.7 - 14.7	500	3.7	3.1 - 4.3	10,600	79.7	78.5 - 80.8	13,300

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	5.0	4.1 - 6.0	1,400	20.8	19.2 - 22.5	300	4.2	3.5 - 5.2	4,800	70.0	68.1 - 71.8	6,800
No	200	2.4	2.0 - 3.0	1,200	12.2	11.1 - 13.3	400	4.1	3.5 - 4.9	7,800	81.3	79.9 - 82.6	9,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.7	3.2 - 4.2	2,600	16.1	15.2 - 17.0	700	4.0	3.6 - 4.6	12,500	76.2	75.1 - 77.2	16,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.6	3.4 - 6.1	300	15.4	13.3 - 17.8	100	2.5	1.7 - 3.6	1,600	77.5	74.8 - 80.0	2,100
Honolulu	400	3.6	3.1 - 4.2	1,900	16.2	15.2 - 17.4	500	4.3	3.7 - 4.9	8,900	75.9	74.6 - 77.2	11,700
Kauai	<50	1.3	0.6 - 2.6	100	14.3	11.2 - 18.1	<50	5.1	3.3 - 7.7	600	79.4	75.2 - 83.1	800
Maui	100	4.1	2.9 - 5.6	300	16.7	14.4 - 19.3	100	3.9	2.9 - 5.3	1,400	75.3	72.4 - 78.0	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.0	2.2 - 4.1	400	11.5	10.0 - 13.2	200	5.1	4.1 - 6.4	3,100	80.4	78.3 - 82.3	3,900
Native Hawaiian	200	5.3	4.3 - 6.5	1,000	21.6	19.7 - 23.7	200	4.2	3.4 - 5.3	3,100	68.8	66.5 - 71.0	4,500
Chinese	<50	1.6	1.0 - 2.7	100	9.3	6.9 - 12.4	<50	2.6	1.7 - 3.8	500	86.5	83.3 - 89.2	600
Filipino	100	3.6	2.8 - 4.6	600	19.8	17.8 - 22.1	100	4.6	3.6 - 6.0	2,100	71.9	69.5 - 74.3	3,000
Japanese	<50	1.9	1.1 - 3.3	200	8.8	7.0 - 11.0	<50	1.9	1.2 - 3.2	1,700	87.3	84.7 - 89.5	1,900
Other	100	3.7	2.6 - 5.3	400	16.0	13.7 - 18.6	100	3.2	2.2 - 4.7	1,800	77.0	74.1 - 79.7	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.4	4.5 - 8.9	400	26.9	23.1 - 31.0	100	4.0	2.6 - 6.2	900	62.8	58.4 - 67.0	1,400
20-24	200	4.3	3.3 - 5.5	900	23.5	21.4 - 25.7	200	5.2	4.2 - 6.5	2,600	67.0	64.6 - 69.4	3,800
25-34	300	3.2	2.6 - 3.9	1,100	13.0	11.9 - 14.2	300	3.8	3.2 - 4.6	6,500	80.0	78.5 - 81.3	8,200
35+	100	2.8	2.0 - 4.0	300	9.9	8.3 - 11.8	100	3.1	2.3 - 4.1	2,500	84.2	82.0 - 86.2	2,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.4	2.8 - 4.1	1,400	16.6	15.3 - 17.9	400	4.8	4.1 - 5.6	6,400	75.2	73.7 - 76.6	8,500
Female	300	3.9	3.3 - 4.7	1,200	15.6	14.3 - 16.9	300	3.2	2.6 - 3.9	6,100	77.3	75.8 - 78.7	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	2.4 - 3.5	1,300	12.4	11.4 - 13.4	400	3.7	3.2 - 4.3	8,400	81.0	79.8 - 82.2	10,400
No	300	5.0	4.1 - 6.1	1,300	22.6	20.9 - 24.4	300	4.6	3.8 - 5.6	4,000	67.8	65.8 - 69.7	6,000

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.3	4.4 - 6.4	1,300	22.5	20.8 - 24.3	200	3.6	2.9 - 4.4	3,900	68.6	66.6 - 70.5	5,700
131-185%	100	3.5	2.4 - 5.1	300	14.8	12.6 - 17.4	100	4.4	3.2 - 6.1	1,700	77.2	74.3 - 80.0	2,200
186+%	200	2.5	2.0 - 3.2	800	10.8	9.8 - 12.0	300	4.3	3.6 - 5.1	6,000	82.3	80.9 - 83.7	7,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	6.0	4.7 - 7.7	700	25.6	23.0 - 28.5	100	4.0	2.9 - 5.5	1,700	64.3	61.2 - 67.3	2,600
\$10,000-\$24,999	200	5.4	4.2 - 6.8	600	19.1	17.1 - 21.3	100	3.8	2.9 - 5.0	2,400	71.6	69.1 - 74.0	3,300
\$25,000-\$49,999	100	3.2	2.4 - 4.4	600	16.2	14.5 - 18.2	200	4.1	3.2 - 5.2	2,800	76.4	74.2 - 78.5	3,700
>=\$50,000	100	2.0	1.5 - 2.6	500	9.0	7.8 - 10.3	200	4.2	3.4 - 5.1	4,700	84.9	83.3 - 86.3	5,600

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	4.7	3.3 - 6.8	200	17.6	14.8 - 20.8	<50	3.8	2.6 - 5.5	900	73.9	70.3 - 77.2	1,200
Normal birth weight	500	3.6	3.1 - 4.1	2,400	16.0	15.1 - 16.9	600	4.1	3.6 - 4.6	11,600	76.4	75.3 - 77.5	15,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.5	6.2 - 11.5	300	27.3	23.5 - 31.6	100	6.8	4.8 - 9.6	700	57.4	52.8 - 61.8	1,200
No	500	3.2	2.8 - 3.7	2,200	15.1	14.2 - 16.0	600	3.8	3.3 - 4.3	11,500	77.9	76.8 - 78.9	14,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.4	4.2 - 7.0	800	24.6	22.3 - 27.2	200	6.6	5.3 - 8.2	2,000	63.4	60.6 - 66.1	3,100
No	400	3.2	2.7 - 3.8	1,800	14.0	13.0 - 14.9	400	3.4	3.0 - 4.0	10,300	79.4	78.3 - 80.5	12,900

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	5.3	4.4 - 6.3	1,400	21.4	19.8 - 23.0	300	3.9	3.2 - 4.8	4,600	69.4	67.6 - 71.2	6,600
No	200	2.5	2.0 - 3.0	1,200	12.4	11.4 - 13.5	400	4.0	3.4 - 4.7	7,600	81.1	79.8 - 82.3	9,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.6	3.2 - 4.0	2,600	16.1	15.3 - 16.9	600	3.8	3.4 - 4.2	12,400	76.5	75.6 - 77.5	16,100

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.3	3.1 - 5.8	300	14.8	12.6 - 17.3	<50	2.0	1.2 - 3.1	1,600	78.9	76.1 - 81.5	2,000
Honolulu	400	3.4	3.0 - 3.9	1,900	16.2	15.2 - 17.2	500	4.1	3.6 - 4.6	8,800	76.3	75.2 - 77.4	11,600
Kauai	<50	1.7	0.8 - 3.5	100	14.5	11.2 - 18.5	<50	4.5	2.8 - 7.2	600	79.3	74.9 - 83.2	800
Maui	100	4.6	3.4 - 6.2	300	17.7	15.2 - 20.5	100	3.7	2.6 - 5.2	1,300	74.1	71.0 - 77.0	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.5	1.9 - 3.4	500	12.4	10.9 - 14.0	200	4.7	3.8 - 5.8	3,000	80.4	78.5 - 82.2	3,800
Native Hawaiian	200	5.2	4.2 - 6.3	900	21.7	19.8 - 23.7	200	4.2	3.4 - 5.2	3,000	68.9	66.7 - 71.0	4,300
Chinese	<50	2.3	1.5 - 3.3	100	8.8	7.3 - 10.7	<50	2.9	2.0 - 4.0	500	86.0	83.9 - 88.0	600
Filipino	100	4.3	3.4 - 5.3	600	19.5	17.7 - 21.5	100	4.5	3.6 - 5.6	2,100	71.7	69.5 - 73.9	3,000
Japanese	<50	1.8	1.1 - 3.0	200	8.4	6.8 - 10.3	<50	2.0	1.3 - 3.1	1,800	87.7	85.5 - 89.7	2,000
Other	100	3.2	2.3 - 4.5	400	15.9	13.8 - 18.2	100	2.6	1.8 - 3.7	1,900	78.3	75.7 - 80.7	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.6	4.8 - 9.1	300	26.3	22.7 - 30.2	<50	3.5	2.2 - 5.3	800	63.6	59.4 - 67.6	1,300
20-24	200	4.1	3.3 - 5.2	1,000	24.8	22.8 - 26.9	200	4.6	3.7 - 5.7	2,600	66.5	64.2 - 68.7	3,900
25-34	300	3.2	2.7 - 3.8	1,000	12.7	11.7 - 13.8	300	3.8	3.2 - 4.4	6,500	80.4	79.1 - 81.6	8,000
35+	100	2.5	1.9 - 3.4	300	9.5	8.2 - 11.1	100	3.0	2.3 - 4.0	2,500	84.9	83.0 - 86.6	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.5	3.0 - 4.1	1,400	16.8	15.6 - 18.0	400	4.5	3.9 - 5.2	6,300	75.2	73.9 - 76.6	8,400
Female	300	3.7	3.1 - 4.3	1,200	15.3	14.2 - 16.6	200	3.1	2.6 - 3.6	6,000	78.0	76.6 - 79.3	7,700

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.8	2.4 - 3.3	1,300	12.7	11.8 - 13.6	400	3.6	3.1 - 4.1	8,600	80.9	79.8 - 82.0	10,600
No	300	5.0	4.2 - 6.0	1,300	22.5	20.9 - 24.3	200	4.2	3.5 - 5.1	3,800	68.2	66.3 - 70.0	5,600

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.1	4.3 - 6.1	1,300	22.8	21.2 - 24.5	200	3.2	2.6 - 3.9	3,900	68.9	67.1 - 70.7	5,600
131-185%	100	2.9	2.0 - 4.2	300	14.1	12.0 - 16.4	100	4.0	2.9 - 5.4	1,600	79.0	76.3 - 81.5	2,100
186+%	200	2.3	1.9 - 2.9	800	11.1	10.1 - 12.1	300	4.3	3.7 - 5.0	6,000	82.2	81.0 - 83.4	7,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	6.5	5.2 - 8.2	700	25.3	22.8 - 27.9	100	3.6	2.7 - 4.8	1,700	64.6	61.7 - 67.3	2,600
\$10,000-\$24,999	200	4.9	3.9 - 6.1	700	20.0	18.1 - 22.1	100	3.3	2.6 - 4.3	2,400	71.7	69.4 - 73.9	3,400
\$25,000-\$49,999	100	2.5	1.9 - 3.3	600	16.3	14.7 - 18.1	200	4.4	3.5 - 5.4	3,000	76.9	74.9 - 78.7	3,900
>=\$50,000	100	1.9	1.4 - 2.5	500	8.5	7.5 - 9.7	200	3.9	3.2 - 4.7	4,500	85.7	84.3 - 86.9	5,300

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.0	3.5 - 7.2	200	17.0	14.0 - 20.4	<50	3.5	2.3 - 5.3	900	74.5	70.7 - 78.0	1,200
Normal birth weight	500	3.5	3.1 - 3.9	2,400	16.0	15.2 - 16.9	600	3.8	3.4 - 4.3	11,500	76.7	75.7 - 77.7	15,000

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.5	5.5 - 10.1	300	26.2	22.5 - 30.2	100	6.4	4.6 - 8.9	700	59.9	55.5 - 64.1	1,200
No	500	3.2	2.8 - 3.6	2,200	15.2	14.4 - 16.1	500	3.6	3.2 - 4.1	11,500	77.9	76.9 - 78.9	14,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.6	3.6 - 5.9	700	23.9	21.6 - 26.2	200	6.4	5.3 - 7.8	2,000	65.1	62.5 - 67.6	3,000
No	400	3.3	2.8 - 3.7	1,800	14.2	13.3 - 15.1	400	3.2	2.8 - 3.7	10,200	79.3	78.3 - 80.3	12,900

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.6	3.9 - 5.4	1,400	21.1	19.6 - 22.6	200	3.5	2.9 - 4.3	4,700	70.8	69.1 - 72.4	6,600
No	200	2.7	2.3 - 3.2	1,200	12.5	11.6 - 13.5	400	4.0	3.5 - 4.6	7,400	80.8	79.6 - 81.9	9,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	3.9	3.5 - 4.4	2,600	16.3	15.5 - 17.2	600	3.7	3.3 - 4.2	12,000	76.0	75.0 - 77.0	15,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.9	2.8 - 5.4	300	15.3	13.1 - 17.7	<50	2.2	1.5 - 3.3	1,500	78.6	75.9 - 81.1	1,900
Honolulu	400	3.9	3.4 - 4.5	1,900	16.3	15.2 - 17.3	500	4.0	3.5 - 4.5	8,600	75.8	74.6 - 77.0	11,400
Kauai	<50	2.4	1.4 - 4.2	100	13.3	10.2 - 17.0	<50	3.8	2.4 - 6.0	600	80.5	76.4 - 84.1	800
Maui	100	4.6	3.4 - 6.2	300	19.2	16.7 - 22.0	100	3.9	2.8 - 5.3	1,200	72.3	69.2 - 75.1	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.6	1.9 - 3.5	400	12.3	10.7 - 14.0	200	4.3	3.4 - 5.4	2,900	80.9	78.8 - 82.8	3,600
Native Hawaiian	200	5.7	4.7 - 6.8	900	22.4	20.5 - 24.3	200	3.8	3.1 - 4.7	2,900	68.1	66.0 - 70.2	4,200
Chinese	<50	2.0	1.3 - 3.0	100	8.2	6.2 - 10.9	<50	2.5	1.7 - 3.6	500	87.3	84.5 - 89.7	600
Filipino	100	4.6	3.7 - 5.8	600	20.1	18.2 - 22.3	200	5.0	4.0 - 6.3	2,100	70.2	67.8 - 72.5	3,000
Japanese	<50	2.4	1.5 - 3.9	200	7.9	6.3 - 9.9	<50	2.0	1.3 - 3.2	1,800	87.6	85.2 - 89.7	2,000
Other	100	3.8	2.7 - 5.2	400	16.1	13.9 - 18.7	100	3.0	2.1 - 4.3	1,700	77.1	74.3 - 79.7	2,300

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.3	6.1 - 11.1	400	29.1	25.4 - 33.1	<50	3.6	2.4 - 5.3	800	59.0	54.8 - 63.2	1,300
20-24	200	4.3	3.5 - 5.3	1,000	25.1	23.1 - 27.2	100	3.9	3.1 - 5.0	2,500	66.6	64.3 - 68.8	3,800
25-34	300	3.7	3.1 - 4.4	1,000	12.4	11.4 - 13.6	300	3.9	3.3 - 4.6	6,200	79.9	78.6 - 81.2	7,800
35+	100	2.0	1.4 - 2.9	300	9.4	7.9 - 11.0	100	3.1	2.3 - 4.2	2,400	85.5	83.5 - 87.2	2,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.7	3.2 - 4.4	1,400	17.0	15.8 - 18.3	400	4.4	3.8 - 5.1	6,200	74.9	73.5 - 76.3	8,200
Female	300	4.2	3.5 - 4.9	1,200	15.5	14.4 - 16.8	200	3.1	2.6 - 3.7	5,800	77.2	75.8 - 78.6	7,500

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	2.4 - 3.4	1,300	12.4	11.5 - 13.3	400	3.6	3.1 - 4.2	8,500	81.2	80.0 - 82.2	10,500
No	300	6.0	5.1 - 7.1	1,300	24.1	22.4 - 25.9	200	4.0	3.3 - 4.8	3,500	65.8	63.9 - 67.7	5,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.7	4.9 - 6.7	1,400	23.7	22.0 - 25.4	200	3.1	2.5 - 3.9	3,900	67.5	65.6 - 69.3	5,700
131-185%	100	3.5	2.5 - 5.0	300	14.6	12.4 - 17.0	100	4.3	3.1 - 5.9	1,600	77.6	74.8 - 80.2	2,000
186+%	200	2.3	1.8 - 2.9	700	10.3	9.3 - 11.4	300	4.1	3.5 - 4.8	5,700	83.3	81.9 - 84.5	6,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight													
Low birth weight	100	5.1	3.5 - 7.3	200	18.3	15.1 - 22.0	<50	3.8	2.5 - 5.9	800	72.8	68.7 - 76.5	1,100
Normal birth weight	600	3.8	3.4 - 4.3	2,400	16.2	15.3 - 17.1	500	3.7	3.3 - 4.2	11,100	76.3	75.2 - 77.3	14,600

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester													
Yes	100	8.9	6.7 - 11.8	300	25.2	21.5 - 29.3	100	5.0	3.5 - 7.2	700	60.9	56.5 - 65.1	1,100
No	500	3.5	3.1 - 4.0	2,200	15.6	14.7 - 16.5	500	3.7	3.2 - 4.1	11,100	77.3	76.2 - 78.2	14,400

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant													
Yes	200	5.9	4.8 - 7.3	700	23.9	21.6 - 26.3	200	5.8	4.7 - 7.2	1,900	64.4	61.8 - 66.9	3,000
No	400	3.4	3.0 - 3.9	1,800	14.4	13.5 - 15.3	400	3.3	2.9 - 3.8	9,900	78.9	77.8 - 80.0	12,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	5.1	4.4 - 6.0	1,400	21.9	20.5 - 23.5	200	3.4	2.8 - 4.1	4,500	69.5	67.8 - 71.1	6,500
No	300	2.9	2.5 - 3.5	1,100	12.3	11.3 - 13.3	400	3.9	3.4 - 4.6	7,300	80.8	79.6 - 82.0	9,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.0	3.6 - 4.5	2,500	16.2	15.3 - 17.1	600	3.6	3.2 - 4.1	11,700	76.2	75.1 - 77.2	15,300

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.0	2.9 - 5.3	300	17.3	15.2 - 19.7	<50	2.4	1.7 - 3.5	1,400	76.3	73.7 - 78.8	1,800
Honolulu	500	4.0	3.5 - 4.7	1,700	15.6	14.6 - 16.7	400	3.8	3.3 - 4.4	8,600	76.5	75.3 - 77.8	11,200
Kauai	<50	2.5	1.4 - 4.3	100	14.3	11.3 - 17.9	<50	3.4	2.1 - 5.5	600	79.8	75.8 - 83.3	700
Maui	100	4.6	3.5 - 6.1	300	19.5	17.1 - 22.1	100	3.9	2.8 - 5.3	1,200	72.0	69.1 - 74.7	1,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.5	1.8 - 3.4	400	12.1	10.5 - 14.0	200	4.6	3.7 - 5.9	2,800	80.8	78.6 - 82.8	3,500
Native Hawaiian	200	5.4	4.5 - 6.4	900	21.3	19.6 - 23.1	200	4.1	3.3 - 5.0	2,800	69.2	67.2 - 71.1	4,000
Chinese	<50	2.4	1.3 - 4.5	100	9.1	6.4 - 12.7	<50	2.9	1.7 - 4.9	500	85.6	81.6 - 88.9	600
Filipino	200	5.4	4.3 - 6.8	600	20.4	18.3 - 22.6	100	4.1	3.2 - 5.3	2,100	70.1	67.6 - 72.6	3,000
Japanese	<50	2.2	1.3 - 3.6	200	7.9	6.3 - 10.0	<50	1.6	0.9 - 2.6	1,800	88.3	85.9 - 90.4	2,100
Other	100	4.3	3.0 - 6.1	400	17.3	14.7 - 20.1	100	2.6	1.7 - 3.8	1,600	75.9	72.7 - 78.8	2,100

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	7.8	5.7 - 10.4	400	30.7	26.9 - 34.7	<50	2.8	1.8 - 4.2	700	58.8	54.6 - 62.9	1,200
20-24	200	5.1	4.2 - 6.3	900	24.4	22.3 - 26.5	100	3.9	3.1 - 4.9	2,500	66.6	64.3 - 68.9	3,700
25-34	300	3.6	3.0 - 4.3	900	12.2	11.1 - 13.3	300	3.9	3.3 - 4.6	6,100	80.3	78.9 - 81.6	7,600
35+	100	2.0	1.3 - 2.9	300	9.7	8.2 - 11.5	100	2.8	2.0 - 3.9	2,300	85.5	83.4 - 87.3	2,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.2	3.5 - 4.9	1,300	16.7	15.5 - 18.0	300	4.2	3.6 - 4.9	5,900	74.9	73.4 - 76.3	7,900
Female	300	3.9	3.2 - 4.6	1,200	15.6	14.4 - 16.9	200	3.0	2.5 - 3.6	5,700	77.5	76.1 - 78.9	7,400

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.9	2.5 - 3.5	1,300	12.5	11.5 - 13.5	400	3.6	3.1 - 4.2	8,400	81.0	79.8 - 82.1	10,400
No	300	6.3	5.4 - 7.4	1,200	24.0	22.2 - 25.8	200	3.6	2.9 - 4.4	3,300	66.1	64.1 - 68.1	5,000

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.9	5.0 - 6.9	1,300	22.8	21.2 - 24.6	200	3.0	2.4 - 3.7	3,800	68.3	66.4 - 70.1	5,600
131-185%	100	3.9	2.8 - 5.4	300	15.0	12.8 - 17.4	100	4.0	2.8 - 5.5	1,700	77.2	74.3 - 79.8	2,200
186+%	100	2.0	1.5 - 2.6	700	10.2	9.2 - 11.4	300	4.2	3.5 - 5.0	5,400	83.6	82.2 - 84.9	6,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight													
Low birth weight	100	5.5	3.7 - 8.1	200	18.1	14.7 - 22.1	<50	3.8	2.4 - 6.1	800	72.5	68.1 - 76.6	1,100
Normal birth weight	600	3.9	3.5 - 4.4	2,300	16.0	15.2 - 17.0	500	3.6	3.2 - 4.1	10,900	76.4	75.4 - 77.5	14,200

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester													
Yes	100	8.5	6.3 - 11.4	300	24.3	20.6 - 28.5	<50	4.6	3.2 - 6.7	700	62.6	58.1 - 66.8	1,000
No	500	3.6	3.2 - 4.1	2,200	15.5	14.6 - 16.4	500	3.6	3.1 - 4.0	10,800	77.3	76.2 - 78.3	14,000

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant													
Yes	200	6.3	5.1 - 7.9	600	22.2	20.0 - 24.6	200	5.8	4.7 - 7.2	1,900	65.6	63.0 - 68.2	2,900
No	400	3.3	2.9 - 3.8	1,800	14.6	13.7 - 15.6	400	3.1	2.7 - 3.6	9,600	78.9	77.8 - 80.0	12,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	5.6	4.7 - 6.5	1,400	22.0	20.5 - 23.6	200	3.3	2.7 - 4.0	4,300	69.1	67.4 - 70.8	6,300
No	200	2.8	2.3 - 3.4	1,100	12.1	11.1 - 13.2	300	3.8	3.2 - 4.4	7,200	81.3	80.0 - 82.5	8,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.4	3.9 - 4.9	2,400	16.1	15.2 - 17.0	500	3.5	3.1 - 4.0	11,300	76.0	75.0 - 77.0	14,800

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.2	3.3 - 5.5	300	16.3	14.4 - 18.4	<50	2.7	2.0 - 3.6	1,300	76.8	74.4 - 79.0	1,700
Honolulu	500	4.3	3.7 - 4.9	1,700	15.7	14.6 - 16.8	400	3.6	3.1 - 4.2	8,300	76.4	75.1 - 77.6	10,900
Kauai	<50	3.4	2.2 - 5.3	100	14.0	11.3 - 17.3	<50	3.6	2.3 - 5.5	500	79.0	75.3 - 82.3	700
Maui	100	5.6	4.4 - 7.1	300	19.3	17.1 - 21.7	100	3.6	2.7 - 4.7	1,100	71.5	68.9 - 74.0	1,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.4	1.7 - 3.4	400	11.8	10.2 - 13.7	100	4.0	3.1 - 5.2	2,700	81.8	79.6 - 83.8	3,400
Native Hawaiian	200	6.2	5.3 - 7.2	800	20.6	19.1 - 22.1	200	4.3	3.6 - 5.1	2,700	69.0	67.2 - 70.7	3,900
Chinese	<50	2.6	1.4 - 4.8	100	10.0	6.8 - 14.4	<50	3.3	1.8 - 6.0	500	84.1	79.3 - 87.9	600
Filipino	200	5.9	4.8 - 7.3	600	20.6	18.4 - 22.9	100	3.9	3.0 - 5.1	2,100	69.6	67.1 - 72.1	3,000
Japanese	<50	2.0	1.2 - 3.3	200	8.7	7.0 - 10.9	<50	2.3	1.5 - 3.5	1,800	87.0	84.5 - 89.2	2,100
Other	100	4.9	3.5 - 6.8	300	17.2	14.6 - 20.2	<50	2.0	1.3 - 3.3	1,500	75.8	72.5 - 78.8	2,000

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.0	6.9 - 11.6	400	31.0	27.5 - 34.8	<50	2.6	1.8 - 4.0	700	57.3	53.4 - 61.2	1,200
20-24	200	5.8	4.8 - 7.1	900	23.6	21.7 - 25.7	100	3.5	2.8 - 4.5	2,400	67.0	64.7 - 69.2	3,600
25-34	300	3.7	3.1 - 4.3	900	12.1	11.0 - 13.2	300	3.8	3.3 - 4.5	5,900	80.4	79.1 - 81.7	7,400
35+	100	2.2	1.5 - 3.2	300	9.8	8.3 - 11.6	100	3.0	2.2 - 4.2	2,200	85.0	82.8 - 86.9	2,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.5	3.9 - 5.3	1,200	16.2	15.0 - 17.4	300	3.8	3.2 - 4.5	5,800	75.5	74.1 - 76.9	7,700
Female	300	4.2	3.6 - 4.9	1,100	16.0	14.8 - 17.3	200	3.2	2.7 - 3.9	5,500	76.6	75.1 - 78.0	7,200

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.0	2.6 - 3.6	1,300	12.8	11.8 - 13.8	400	3.6	3.1 - 4.2	8,200	80.5	79.3 - 81.7	10,200
No	300	7.3	6.3 - 8.4	1,100	23.3	21.6 - 25.0	200	3.2	2.7 - 3.9	3,100	66.2	64.3 - 68.1	4,700

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.2	5.3 - 7.2	1,200	21.8	20.3 - 23.5	200	3.0	2.4 - 3.6	3,800	69.0	67.2 - 70.7	5,500
131-185%	100	3.9	2.9 - 5.2	400	15.6	13.5 - 17.8	100	3.3	2.4 - 4.6	1,800	77.3	74.7 - 79.7	2,400
186+%	100	2.2	1.7 - 2.8	600	9.7	8.7 - 10.9	200	4.2	3.5 - 5.0	4,900	83.9	82.4 - 85.2	5,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight													
Low birth weight	100	5.5	3.6 - 8.2	200	19.7	16.1 - 23.8	<50	3.7	2.3 - 6.1	700	71.1	66.5 - 75.2	1,000
Normal birth weight	600	4.3	3.8 - 4.8	2,200	15.8	15.0 - 16.7	500	3.5	3.1 - 4.0	10,600	76.4	75.3 - 77.4	13,900

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester													
Yes	100	8.3	6.2 - 10.9	300	25.3	21.7 - 29.4	<50	4.1	2.8 - 5.9	600	62.3	58.0 - 66.4	1,000
No	500	4.0	3.6 - 4.5	2,100	15.3	14.5 - 16.3	500	3.5	3.1 - 4.0	10,500	77.2	76.1 - 78.2	13,600

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant													
Yes	200	6.5	5.3 - 8.0	600	22.9	20.7 - 25.2	200	5.4	4.4 - 6.7	1,800	65.2	62.6 - 67.7	2,800
No	400	3.7	3.2 - 4.2	1,700	14.4	13.5 - 15.4	400	3.1	2.7 - 3.6	9,300	78.8	77.7 - 79.9	11,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.9	5.1 - 6.9	1,300	21.7	20.2 - 23.2	200	3.2	2.7 - 3.9	4,200	69.1	67.4 - 70.8	6,100
No	300	3.2	2.7 - 3.8	1,000	12.2	11.2 - 13.2	300	3.7	3.1 - 4.3	6,900	80.9	79.6 - 82.1	8,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	600	4.3	3.9 - 4.8	2,400	16.1	15.3 - 17.0	500	3.6	3.2 - 4.1	11,100	75.9	74.9 - 76.9	14,600

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.8	3.8 - 6.0	300	17.9	16.1 - 19.8	100	3.1	2.4 - 4.0	1,200	74.3	72.1 - 76.4	1,700
Honolulu	400	4.2	3.6 - 4.8	1,700	15.6	14.6 - 16.8	400	3.7	3.2 - 4.2	8,200	76.5	75.3 - 77.8	10,700
Kauai	<50	4.1	2.7 - 6.1	100	14.3	11.8 - 17.3	<50	3.6	2.4 - 5.4	500	78.0	74.5 - 81.1	700
Maui	100	5.0	4.0 - 6.3	300	18.3	16.3 - 20.4	100	4.1	3.2 - 5.3	1,100	72.7	70.3 - 74.9	1,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	2.4	1.7 - 3.4	400	12.5	10.9 - 14.4	100	4.1	3.2 - 5.3	2,700	81.0	78.8 - 83.0	3,300
Native Hawaiian	200	5.8	5.1 - 6.6	800	21.5	20.2 - 22.9	200	4.1	3.5 - 4.8	2,600	68.5	67.0 - 70.0	3,800
Chinese	<50	2.8	1.5 - 5.3	100	11.5	7.9 - 16.5	<50	3.2	1.6 - 6.2	500	82.5	77.1 - 86.8	600
Filipino	200	5.8	4.6 - 7.2	600	19.4	17.3 - 21.7	100	3.8	2.9 - 5.0	2,100	70.9	68.3 - 73.4	3,000
Japanese	<50	2.1	1.3 - 3.5	100	7.5	5.8 - 9.7	100	3.0	2.0 - 4.4	1,700	87.4	84.8 - 89.6	2,000
Other	100	5.1	3.6 - 7.1	300	17.0	14.3 - 20.1	<50	2.4	1.5 - 3.8	1,400	75.5	72.1 - 78.7	1,900

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.6	7.6 - 12.0	400	31.2	27.9 - 34.7	<50	2.8	1.9 - 4.0	700	56.5	52.8 - 60.1	1,300
20-24	200	5.7	4.7 - 6.9	800	22.9	21.0 - 25.0	100	3.4	2.7 - 4.3	2,400	67.9	65.7 - 70.0	3,500
25-34	200	3.4	2.9 - 4.1	900	12.3	11.2 - 13.5	300	3.9	3.3 - 4.6	5,800	80.3	79.0 - 81.7	7,200
35+	100	2.2	1.5 - 3.2	300	9.8	8.2 - 11.7	100	3.6	2.6 - 4.9	2,100	84.4	82.2 - 86.4	2,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	4.4	3.8 - 5.2	1,200	16.5	15.3 - 17.7	300	3.6	3.0 - 4.2	5,700	75.5	74.1 - 76.9	7,600
Female	300	4.2	3.6 - 4.9	1,100	15.8	14.6 - 17.0	300	3.7	3.1 - 4.4	5,400	76.4	74.9 - 77.8	7,000

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	2.8	2.4 - 3.3	1,300	13.0	12.1 - 14.0	400	3.8	3.3 - 4.4	8,100	80.3	79.2 - 81.5	10,100
No	300	7.7	6.7 - 8.9	1,000	23.2	21.6 - 24.9	100	3.3	2.7 - 3.9	2,900	65.8	63.9 - 67.7	4,400

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	6.0	5.2 - 7.0	1,200	21.8	20.3 - 23.4	200	2.9	2.4 - 3.5	3,900	69.2	67.5 - 70.9	5,600
131-185%	100	3.7	2.8 - 5.0	400	14.9	13.0 - 17.1	100	3.2	2.4 - 4.4	1,900	78.1	75.6 - 80.4	2,400
186+%	100	2.2	1.7 - 2.8	500	9.8	8.7 - 11.0	300	4.6	3.8 - 5.5	4,600	83.5	82.0 - 84.9	5,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight													
Low birth weight	100	5.5	3.6 - 8.3	200	20.6	16.9 - 24.8	<50	3.6	2.2 - 5.9	700	70.3	65.7 - 74.6	900
Normal birth weight	600	4.2	3.8 - 4.8	2,200	15.8	14.9 - 16.7	500	3.6	3.2 - 4.1	10,400	76.4	75.3 - 77.4	13,600

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester													
Yes	100	7.7	5.8 - 10.1	300	25.7	22.1 - 29.7	<50	3.6	2.5 - 5.2	600	63.0	58.8 - 67.0	1,000
No	500	4.0	3.5 - 4.5	2,000	15.4	14.5 - 16.3	500	3.6	3.2 - 4.1	10,300	77.1	76.0 - 78.1	13,400

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant													
Yes	200	6.3	5.1 - 7.7	600	23.3	21.2 - 25.6	100	4.9	3.9 - 6.2	1,800	65.4	62.9 - 67.9	2,700
No	400	3.7	3.2 - 4.2	1,700	14.3	13.4 - 15.3	400	3.3	2.9 - 3.8	9,100	78.7	77.6 - 79.8	11,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.9	5.1 - 6.9	1,300	21.9	20.4 - 23.4	200	3.2	2.7 - 3.8	4,100	69.0	67.3 - 70.6	6,000
No	300	3.1	2.7 - 3.7	1,000	12.1	11.1 - 13.2	300	3.9	3.3 - 4.6	6,800	80.8	79.5 - 82.0	8,400