

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration 8+ weeks	Breastfeeding (any) duration 8+ weeks

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/22/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- for the State of Hawaii, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	5,100	31.3	30.3 - 32.4	11,200	68.7	67.6 - 69.7	16,300
2001-2005	5,100	31.1	30.1 - 32.1	11,400	68.9	67.9 - 69.9	16,500
2002-2006	5,200	30.6	29.6 - 31.7	11,800	69.4	68.3 - 70.4	16,900
2003-2007	5,200	30.3	29.3 - 31.3	12,100	69.7	68.7 - 70.7	17,300
2004-2008	5,200	29.3	28.4 - 30.3	12,500	70.7	69.7 - 71.6	17,600
2005-2009	5,200	29.2	28.2 - 30.3	12,600	70.8	69.7 - 71.8	17,800
2006-2010	5,100	28.3	27.2 - 29.4	12,900	71.7	70.6 - 72.8	18,000
2007-2011	4,900	26.8	25.6 - 28.0	13,300	73.2	72.0 - 74.4	18,100
2008-2012	4,600	25.4	24.2 - 26.7	13,500	74.6	73.3 - 75.8	18,200
2009-2013	4,300	23.9	22.7 - 25.2	13,800	76.1	74.8 - 77.3	18,100

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- for the State of Hawaii, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	4,300	23.9	22.7 - 25.2	13,800	76.1	74.8 - 77.3	18,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	25.1	23.1 - 27.2	1,700	74.9	72.8 - 76.9	2,300
Honolulu	3,100	24.2	22.5 - 25.9	9,900	75.8	74.1 - 77.5	13,000
Kauai	200	19.3	17.3 - 21.5	700	80.7	78.5 - 82.7	800
Maui	400	22.6	20.8 - 24.5	1,500	77.4	75.5 - 79.2	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	19.3	17.0 - 21.9	3,600	80.7	78.1 - 83.0	4,400
Native Hawaiian	1,700	30.9	28.6 - 33.4	3,700	69.1	66.6 - 71.4	5,400
Chinese	200	18.1	13.2 - 24.4	700	81.9	75.6 - 86.8	800
Filipino	900	27.9	24.9 - 31.0	2,300	72.1	69.0 - 75.1	3,200
Japanese	200	10.0	7.5 - 13.2	1,500	90.0	86.8 - 92.5	1,700
Other	600	23.3	20.0 - 27.0	2,000	76.7	73.0 - 80.0	2,600

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	42.5	37.0 - 48.2	700	57.5	51.8 - 63.0	1,100
20-24	1,100	27.9	25.2 - 30.8	3,000	72.1	69.2 - 74.8	4,100
25-34	2,100	21.9	20.2 - 23.6	7,500	78.1	76.4 - 79.8	9,700
35+	600	18.1	15.7 - 20.9	2,600	81.9	79.1 - 84.3	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,200	23.9	22.2 - 25.7	7,000	76.1	74.3 - 77.8	9,200
Female	2,100	23.9	22.1 - 25.7	6,800	76.1	74.3 - 77.9	9,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	19.5	18.0 - 21.1	9,100	80.5	78.9 - 82.0	11,200
No	2,100	31.0	28.9 - 33.2	4,800	69.0	66.8 - 71.1	6,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,000	30.4	28.2 - 32.7	4,500	69.6	67.3 - 71.8	6,400
131-185%	700	26.1	22.8 - 29.6	2,100	73.9	70.4 - 77.2	2,800
186+%	1,200	16.4	14.7 - 18.2	6,200	83.6	81.8 - 85.3	7,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	28.0	25.8 - 30.2	900	72.0	69.8 - 74.2	1,200
Normal birth weight	4,000	23.6	22.3 - 25.0	12,900	76.4	75.0 - 77.7	16,900

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	47.0	41.4 - 52.8	600	53.0	47.2 - 58.6	1,100
No	3,700	22.3	21.1 - 23.7	12,900	77.7	76.3 - 78.9	16,700

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	36.7	33.5 - 39.9	2,200	63.3	60.1 - 66.5	3,400
No	3,000	20.8	19.5 - 22.2	11,400	79.2	77.8 - 80.5	14,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	31.0	29.0 - 33.1	5,400	69.0	66.9 - 71.0	7,900
No	1,800	18.1	16.6 - 19.8	8,200	81.9	80.2 - 83.4	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	4,600	25.4	24.2 - 26.7	13,500	74.6	73.3 - 75.8	18,200

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	27.4	25.2 - 29.7	1,700	72.6	70.3 - 74.8	2,300
Honolulu	3,300	25.7	24.1 - 27.3	9,700	74.3	72.7 - 75.9	13,000
Kauai	200	20.3	17.6 - 23.2	700	79.7	76.8 - 82.4	800
Maui	500	23.4	21.4 - 25.6	1,500	76.6	74.4 - 78.6	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	20.7	18.5 - 23.2	3,400	79.3	76.8 - 81.5	4,300
Native Hawaiian	1,700	31.8	29.5 - 34.2	3,600	68.2	65.8 - 70.5	5,300
Chinese	100	17.1	12.5 - 23.1	700	82.9	76.9 - 87.5	800
Filipino	1,000	29.7	26.9 - 32.7	2,300	70.3	67.3 - 73.1	3,200
Japanese	200	12.8	10.1 - 15.9	1,500	87.2	84.1 - 89.9	1,700
Other	700	25.9	22.7 - 29.3	2,000	74.1	70.7 - 77.3	2,800

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	42.2	37.2 - 47.4	700	57.8	52.6 - 62.8	1,300
20-24	1,300	31.9	29.2 - 34.7	2,800	68.1	65.3 - 70.8	4,100
25-34	2,200	22.7	21.1 - 24.4	7,400	77.3	75.6 - 78.9	9,600
35+	600	18.6	16.3 - 21.2	2,600	81.4	78.8 - 83.7	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,400	25.8	24.2 - 27.6	6,900	74.2	72.4 - 75.8	9,300
Female	2,200	25.0	23.3 - 26.8	6,600	75.0	73.2 - 76.7	8,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	20.9	19.5 - 22.4	8,900	79.1	77.6 - 80.5	11,200
No	2,300	32.7	30.6 - 34.8	4,700	67.3	65.2 - 69.4	6,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,100	32.2	30.1 - 34.4	4,400	67.8	65.6 - 69.9	6,400
131-185%	800	26.6	23.5 - 29.9	2,100	73.4	70.1 - 76.5	2,800
186+%	1,400	17.9	16.3 - 19.7	6,200	82.1	80.3 - 83.7	7,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	28.9	26.4 - 31.6	900	71.1	68.4 - 73.6	1,200
Normal birth weight	4,300	25.2	23.9 - 26.5	12,700	74.8	73.5 - 76.1	16,900

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	48.4	43.3 - 53.6	600	51.6	46.4 - 56.7	1,200
No	3,900	23.7	22.5 - 25.0	12,700	76.3	75.0 - 77.5	16,600

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	38.9	35.9 - 42.0	2,100	61.1	58.0 - 64.1	3,500
No	3,200	22.2	20.9 - 23.5	11,200	77.8	76.5 - 79.1	14,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	32.7	30.7 - 34.6	5,300	67.3	65.4 - 69.3	7,900
No	2,000	19.7	18.3 - 21.3	8,100	80.3	78.7 - 81.7	10,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	4,900	26.8	25.6 - 28.0	13,300	73.2	72.0 - 74.4	18,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	28.0	25.7 - 30.5	1,700	72.0	69.5 - 74.3	2,300
Honolulu	3,500	27.1	25.6 - 28.6	9,400	72.9	71.4 - 74.4	12,900
Kauai	200	22.2	19.0 - 25.7	700	77.8	74.3 - 81.0	900
Maui	500	25.5	23.2 - 27.9	1,500	74.5	72.1 - 76.8	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,000	22.5	20.3 - 24.9	3,300	77.5	75.1 - 79.7	4,300
Native Hawaiian	1,800	33.5	31.2 - 35.8	3,500	66.5	64.2 - 68.8	5,300
Chinese	100	16.3	12.0 - 21.7	600	83.7	78.3 - 88.0	700
Filipino	1,000	30.5	27.8 - 33.3	2,300	69.5	66.7 - 72.2	3,300
Japanese	300	13.6	11.1 - 16.6	1,600	86.4	83.4 - 88.9	1,800
Other	800	27.9	24.8 - 31.2	2,000	72.1	68.8 - 75.2	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	43.4	38.7 - 48.1	800	56.6	51.9 - 61.3	1,400
20-24	1,400	33.7	31.2 - 36.4	2,800	66.3	63.6 - 68.8	4,300
25-34	2,200	23.7	22.2 - 25.3	7,100	76.3	74.7 - 77.8	9,300
35+	600	19.2	16.9 - 21.7	2,600	80.8	78.3 - 83.1	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,500	27.1	25.5 - 28.8	6,800	72.9	71.2 - 74.5	9,400
Female	2,300	26.4	24.8 - 28.1	6,400	73.6	71.9 - 75.2	8,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	21.8	20.4 - 23.2	8,800	78.2	76.8 - 79.6	11,200
No	2,400	35.0	33.0 - 37.0	4,500	65.0	63.0 - 67.0	6,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,200	34.4	32.3 - 36.5	4,300	65.6	63.5 - 67.7	6,500
131-185%	700	26.9	23.9 - 30.1	2,000	73.1	69.9 - 76.1	2,700
186+%	1,500	19.1	17.6 - 20.8	6,200	80.9	79.2 - 82.4	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,100	36.6	33.6 - 39.7	2,000	63.4	60.3 - 66.4	3,100
\$10,000-\$24,999	1,100	31.9	29.3 - 34.7	2,400	68.1	65.3 - 70.7	3,600
\$25,000-\$49,999	1,200	27.6	25.2 - 30.2	3,000	72.4	69.8 - 74.8	4,200
>=\$50,000	1,000	17.1	15.4 - 18.9	5,000	82.9	81.1 - 84.6	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	31.4	28.4 - 34.4	900	68.6	65.6 - 71.6	1,300
Normal birth weight	4,400	26.4	25.2 - 27.7	12,400	73.6	72.3 - 74.8	16,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	50.3	45.6 - 55.0	700	49.7	45.0 - 54.4	1,400
No	4,100	24.7	23.6 - 25.9	12,400	75.3	74.1 - 76.4	16,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	41.7	38.8 - 44.6	2,100	58.3	55.4 - 61.2	3,600
No	3,300	22.9	21.7 - 24.2	11,000	77.1	75.8 - 78.3	14,200

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	33.9	32.0 - 35.8	5,100	66.1	64.2 - 68.0	7,700
No	2,200	21.4	20.0 - 22.9	8,000	78.6	77.1 - 80.0	10,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,100	28.3	27.2 - 29.4	12,900	71.7	70.6 - 72.8	18,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	28.9	26.4 - 31.6	1,600	71.1	68.4 - 73.6	2,300
Honolulu	3,700	28.7	27.3 - 30.2	9,100	71.3	69.8 - 72.7	12,800
Kauai	200	24.0	20.4 - 27.9	700	76.0	72.1 - 79.6	900
Maui	500	26.9	24.4 - 29.6	1,500	73.1	70.4 - 75.6	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,000	24.2	22.0 - 26.4	3,200	75.8	73.6 - 78.0	4,300
Native Hawaiian	1,800	35.4	33.2 - 37.7	3,400	64.6	62.3 - 66.8	5,200
Chinese	100	16.4	12.5 - 21.2	600	83.6	78.8 - 87.5	700
Filipino	1,000	31.8	29.3 - 34.5	2,200	68.2	65.5 - 70.7	3,200
Japanese	300	13.8	11.4 - 16.6	1,600	86.2	83.4 - 88.6	1,900
Other	800	30.0	27.0 - 33.2	1,900	70.0	66.8 - 73.0	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	42.6	38.2 - 47.1	900	57.4	52.9 - 61.8	1,500
20-24	1,600	36.4	34.0 - 39.0	2,700	63.6	61.0 - 66.0	4,300
25-34	2,200	24.7	23.2 - 26.2	6,800	75.3	73.8 - 76.8	9,100
35+	700	21.1	18.8 - 23.6	2,500	78.9	76.4 - 81.2	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,700	29.0	27.5 - 30.6	6,600	71.0	69.4 - 72.5	9,300
Female	2,400	27.5	26.0 - 29.2	6,300	72.5	70.8 - 74.0	8,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	22.7	21.4 - 24.1	8,700	77.3	75.9 - 78.6	11,200
No	2,600	37.5	35.5 - 39.5	4,200	62.5	60.5 - 64.5	6,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,400	36.8	34.8 - 38.9	4,100	63.2	61.1 - 65.2	6,400
131-185%	800	28.2	25.3 - 31.3	1,900	71.8	68.7 - 74.7	2,700
186+%	1,500	20.0	18.6 - 21.6	6,100	80.0	78.4 - 81.4	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,200	40.0	37.0 - 43.0	1,800	60.0	57.0 - 63.0	3,100
\$10,000-\$24,999	1,200	33.4	30.8 - 36.0	2,400	66.6	64.0 - 69.2	3,700
\$25,000-\$49,999	1,200	28.9	26.6 - 31.4	2,900	71.1	68.6 - 73.4	4,100
>=\$50,000	1,100	17.8	16.2 - 19.5	4,900	82.2	80.5 - 83.8	6,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	31.7	28.5 - 35.0	900	68.3	65.0 - 71.5	1,300
Normal birth weight	4,700	28.0	26.9 - 29.2	12,000	72.0	70.8 - 73.1	16,700

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	50.6	46.2 - 54.9	800	49.4	45.1 - 53.8	1,500
No	4,200	26.2	25.1 - 27.4	12,000	73.8	72.6 - 74.9	16,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	43.3	40.5 - 46.1	2,100	56.7	53.9 - 59.5	3,600
No	3,400	24.4	23.2 - 25.7	10,600	75.6	74.3 - 76.8	14,100

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	36.3	34.4 - 38.1	4,800	63.7	61.9 - 65.6	7,500
No	2,300	22.5	21.1 - 23.9	7,900	77.5	76.1 - 78.9	10,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,200	29.2	28.2 - 30.3	12,600	70.8	69.7 - 71.8	17,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	29.8	27.2 - 32.6	1,600	70.2	67.4 - 72.8	2,300
Honolulu	3,700	29.4	28.1 - 30.8	9,000	70.6	69.2 - 71.9	12,700
Kauai	200	24.5	20.6 - 28.8	600	75.5	71.2 - 79.4	800
Maui	600	29.1	26.4 - 32.1	1,400	70.9	67.9 - 73.6	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,000	23.4	21.4 - 25.5	3,200	76.6	74.5 - 78.6	4,100
Native Hawaiian	1,900	37.5	35.2 - 39.7	3,200	62.5	60.3 - 64.8	5,100
Chinese	100	18.4	14.9 - 22.6	500	81.6	77.4 - 85.1	700
Filipino	1,100	32.7	30.4 - 35.2	2,200	67.3	64.8 - 69.6	3,200
Japanese	300	15.5	13.2 - 18.2	1,700	84.5	81.8 - 86.8	2,000
Other	800	31.2	28.3 - 34.1	1,800	68.8	65.9 - 71.7	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	42.8	38.7 - 47.1	900	57.2	52.9 - 61.3	1,600
20-24	1,600	38.6	36.3 - 41.0	2,600	61.4	59.0 - 63.7	4,200
25-34	2,200	25.4	24.0 - 26.8	6,600	74.6	73.2 - 76.0	8,900
35+	600	20.6	18.5 - 22.9	2,500	79.4	77.1 - 81.5	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,700	29.9	28.4 - 31.4	6,500	70.1	68.6 - 71.6	9,200
Female	2,400	28.5	27.0 - 30.1	6,100	71.5	69.9 - 73.0	8,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	23.4	22.2 - 24.6	8,500	76.6	75.4 - 77.8	11,100
No	2,600	38.9	37.0 - 40.9	4,100	61.1	59.1 - 63.0	6,700

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,500	39.0	37.1 - 41.0	3,900	61.0	59.0 - 62.9	6,400
131-185%	700	29.0	26.1 - 32.0	1,700	71.0	68.0 - 73.9	2,400
186+%	1,500	20.2	18.8 - 21.6	6,000	79.8	78.4 - 81.2	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,300	43.1	40.2 - 46.0	1,700	56.9	54.0 - 59.8	3,000
\$10,000-\$24,999	1,300	35.2	32.8 - 37.7	2,400	64.8	62.3 - 67.2	3,700
\$25,000-\$49,999	1,100	28.5	26.3 - 30.7	2,900	71.5	69.3 - 73.7	4,000
>=\$50,000	1,000	17.8	16.3 - 19.4	4,800	82.2	80.6 - 83.7	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	32.0	28.6 - 35.6	900	68.0	64.4 - 71.4	1,300
Normal birth weight	4,800	29.0	27.9 - 30.1	11,700	71.0	69.9 - 72.1	16,400

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	52.9	48.8 - 57.0	700	47.1	43.0 - 51.2	1,500
No	4,300	26.9	25.8 - 28.0	11,700	73.1	72.0 - 74.2	15,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	44.7	42.1 - 47.3	2,000	55.3	52.7 - 57.9	3,600
No	3,500	25.1	24.0 - 26.3	10,400	74.9	73.7 - 76.0	13,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	37.6	35.9 - 39.4	4,600	62.4	60.6 - 64.1	7,400
No	2,300	23.0	21.7 - 24.3	7,700	77.0	75.7 - 78.3	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,200	29.3	28.4 - 30.3	12,500	70.7	69.7 - 71.6	17,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	29.4	26.7 - 32.3	1,600	70.6	67.7 - 73.3	2,200
Honolulu	3,700	29.5	28.3 - 30.6	8,900	70.5	69.4 - 71.7	12,700
Kauai	200	24.4	20.3 - 29.0	600	75.6	71.0 - 79.7	800
Maui	600	30.7	27.7 - 33.8	1,300	69.3	66.2 - 72.3	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,000	24.0	22.2 - 26.0	3,100	76.0	74.0 - 77.8	4,100
Native Hawaiian	1,800	37.8	35.7 - 40.0	3,000	62.2	60.0 - 64.3	4,900
Chinese	100	17.5	15.4 - 19.8	500	82.5	80.2 - 84.6	600
Filipino	1,100	33.5	31.3 - 35.7	2,200	66.5	64.3 - 68.7	3,200
Japanese	300	15.7	13.6 - 18.1	1,800	84.3	81.9 - 86.4	2,100
Other	800	30.7	28.1 - 33.3	1,900	69.3	66.7 - 71.9	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	43.0	39.1 - 47.0	800	57.0	53.0 - 60.9	1,500
20-24	1,700	39.6	37.4 - 41.8	2,600	60.4	58.2 - 62.6	4,300
25-34	2,200	25.3	24.0 - 26.6	6,500	74.7	73.4 - 76.0	8,700
35+	700	20.5	18.6 - 22.6	2,500	79.5	77.4 - 81.4	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,800	30.2	28.9 - 31.6	6,400	69.8	68.4 - 71.1	9,200
Female	2,400	28.4	27.0 - 29.8	6,100	71.6	70.2 - 73.0	8,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	24.2	23.1 - 25.3	8,600	75.8	74.7 - 76.9	11,400
No	2,400	38.7	36.9 - 40.5	3,800	61.3	59.5 - 63.1	6,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,500	39.0	37.2 - 40.8	3,900	61.0	59.2 - 62.8	6,400
131-185%	700	28.2	25.6 - 31.0	1,700	71.8	69.0 - 74.4	2,300
186+%	1,600	20.8	19.6 - 22.1	6,000	79.2	77.9 - 80.4	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,300	42.8	40.1 - 45.5	1,700	57.2	54.5 - 59.9	2,900
\$10,000-\$24,999	1,400	35.8	33.5 - 38.0	2,500	64.2	62.0 - 66.5	3,800
\$25,000-\$49,999	1,200	28.1	26.1 - 30.1	3,000	71.9	69.9 - 73.9	4,200
≥\$50,000	1,000	17.9	16.5 - 19.4	4,600	82.1	80.6 - 83.5	5,600

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	31.7	28.1 - 35.5	900	68.3	64.5 - 71.9	1,300
Normal birth weight	4,800	29.2	28.2 - 30.2	11,600	70.8	69.8 - 71.8	16,400

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	51.0	47.1 - 54.9	700	49.0	45.1 - 52.9	1,500
No	4,300	27.3	26.3 - 28.3	11,600	72.7	71.7 - 73.7	15,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	43.0	40.6 - 45.5	2,000	57.0	54.5 - 59.4	3,500
No	3,600	25.8	24.8 - 26.9	10,300	74.2	73.1 - 75.2	13,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	36.6	35.0 - 38.3	4,700	63.4	61.7 - 65.0	7,500
No	2,300	23.7	22.5 - 24.9	7,500	76.3	75.1 - 77.5	9,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,200	30.3	29.3 - 31.3	12,100	69.7	68.7 - 70.7	17,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	30.5	27.9 - 33.3	1,500	69.5	66.7 - 72.1	2,100
Honolulu	3,800	30.4	29.2 - 31.6	8,700	69.6	68.4 - 70.8	12,500
Kauai	200	24.2	20.4 - 28.5	600	75.8	71.5 - 79.6	800
Maui	600	32.4	29.5 - 35.5	1,200	67.6	64.5 - 70.5	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	24.2	22.2 - 26.3	3,000	75.8	73.7 - 77.8	3,900
Native Hawaiian	1,900	39.5	37.5 - 41.5	2,900	60.5	58.5 - 62.5	4,800
Chinese	100	16.4	13.7 - 19.5	500	83.6	80.5 - 86.3	600
Filipino	1,200	35.3	33.0 - 37.7	2,100	64.7	62.3 - 67.0	3,300
Japanese	300	16.3	14.0 - 18.9	1,800	83.7	81.1 - 86.0	2,100
Other	800	31.3	28.6 - 34.2	1,800	68.7	65.8 - 71.4	2,600

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	47.8	43.9 - 51.8	800	52.2	48.2 - 56.1	1,500
20-24	1,700	40.2	38.0 - 42.4	2,500	59.8	57.6 - 62.0	4,300
25-34	2,200	26.1	24.8 - 27.5	6,300	73.9	72.5 - 75.2	8,500
35+	600	19.9	17.9 - 22.0	2,500	80.1	78.0 - 82.1	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,800	31.2	29.8 - 32.7	6,200	68.8	67.3 - 70.2	9,000
Female	2,400	29.3	27.9 - 30.8	5,800	70.7	69.2 - 72.1	8,300

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	24.2	23.1 - 25.4	8,600	75.8	74.6 - 76.9	11,300
No	2,500	41.8	39.9 - 43.7	3,500	58.2	56.3 - 60.1	6,000

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,700	40.9	39.1 - 42.7	3,900	59.1	57.3 - 60.9	6,600
131-185%	700	29.5	26.8 - 32.4	1,600	70.5	67.6 - 73.2	2,300
186+%	1,400	19.8	18.5 - 21.1	5,700	80.2	78.9 - 81.5	7,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	33.2	29.4 - 37.2	800	66.8	62.8 - 70.6	1,300
Normal birth weight	4,800	30.1	29.1 - 31.2	11,200	69.9	68.8 - 70.9	16,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	53.2	49.3 - 57.1	700	46.8	42.9 - 50.7	1,500
No	4,400	28.1	27.1 - 29.2	11,200	71.9	70.8 - 72.9	15,600

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	44.8	42.3 - 47.3	1,900	55.2	52.7 - 57.7	3,500
No	3,600	26.4	25.3 - 27.5	9,900	73.6	72.5 - 74.7	13,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	38.5	36.9 - 40.2	4,500	61.5	59.8 - 63.1	7,400
No	2,300	23.9	22.7 - 25.2	7,400	76.1	74.8 - 77.3	9,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,200	30.6	29.6 - 31.7	11,800	69.4	68.3 - 70.4	16,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	32.4	29.9 - 35.1	1,400	67.6	64.9 - 70.1	2,000
Honolulu	3,800	30.4	29.1 - 31.7	8,600	69.6	68.3 - 70.9	12,400
Kauai	200	24.8	21.1 - 28.9	600	75.2	71.1 - 78.9	800
Maui	600	32.6	29.9 - 35.5	1,200	67.4	64.5 - 70.1	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	24.0	21.9 - 26.2	2,900	76.0	73.8 - 78.1	3,800
Native Hawaiian	1,800	39.6	37.7 - 41.5	2,800	60.4	58.5 - 62.3	4,700
Chinese	100	20.2	16.2 - 24.8	500	79.8	75.2 - 83.8	600
Filipino	1,200	35.4	33.0 - 37.9	2,100	64.6	62.1 - 67.0	3,300
Japanese	300	15.3	13.0 - 17.8	1,800	84.7	82.2 - 87.0	2,200
Other	800	33.7	30.6 - 36.9	1,600	66.3	63.1 - 69.4	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	48.9	45.0 - 52.9	700	51.1	47.1 - 55.0	1,400
20-24	1,700	40.2	38.0 - 42.5	2,500	59.8	57.5 - 62.0	4,200
25-34	2,200	26.3	24.9 - 27.7	6,200	73.7	72.3 - 75.1	8,400
35+	600	20.6	18.5 - 22.9	2,400	79.4	77.1 - 81.5	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,800	32.0	30.6 - 33.5	6,000	68.0	66.5 - 69.4	8,800
Female	2,400	29.1	27.6 - 30.6	5,800	70.9	69.4 - 72.4	8,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	24.6	23.4 - 25.9	8,500	75.4	74.1 - 76.6	11,200
No	2,400	42.3	40.4 - 44.3	3,300	57.7	55.7 - 59.6	5,700

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,700	40.9	39.1 - 42.7	3,800	59.1	57.3 - 60.9	6,500
131-185%	700	30.2	27.4 - 33.2	1,700	69.8	66.8 - 72.6	2,400
186+%	1,300	19.1	17.7 - 20.5	5,400	80.9	79.5 - 82.3	6,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	33.1	29.1 - 37.4	800	66.9	62.6 - 70.9	1,200
Normal birth weight	4,800	30.4	29.4 - 31.5	11,000	69.6	68.5 - 70.6	15,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	53.4	49.5 - 57.3	700	46.6	42.7 - 50.5	1,400
No	4,300	28.4	27.4 - 29.5	10,900	71.6	70.5 - 72.6	15,300

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	44.3	41.8 - 46.8	1,900	55.7	53.2 - 58.2	3,400
No	3,500	26.8	25.7 - 28.0	9,700	73.2	72.0 - 74.3	13,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	39.5	37.8 - 41.3	4,400	60.5	58.7 - 62.2	7,200
No	2,300	23.8	22.5 - 25.1	7,200	76.2	74.9 - 77.5	9,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,100	31.1	30.1 - 32.1	11,400	68.9	67.9 - 69.9	16,500

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	33.2	30.8 - 35.6	1,300	66.8	64.4 - 69.2	2,000
Honolulu	3,700	30.9	29.6 - 32.2	8,400	69.1	67.8 - 70.4	12,100
Kauai	200	25.2	21.8 - 28.9	500	74.8	71.1 - 78.2	700
Maui	600	33.0	30.5 - 35.6	1,100	67.0	64.4 - 69.5	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	23.1	21.0 - 25.3	2,800	76.9	74.7 - 79.0	3,600
Native Hawaiian	1,800	40.4	38.7 - 42.1	2,700	59.6	57.9 - 61.3	4,500
Chinese	100	22.1	17.7 - 27.3	500	77.9	72.7 - 82.3	600
Filipino	1,200	36.1	33.6 - 38.6	2,100	63.9	61.4 - 66.4	3,300
Japanese	400	16.3	14.0 - 18.9	1,800	83.7	81.1 - 86.0	2,200
Other	800	34.7	31.5 - 38.0	1,500	65.3	62.0 - 68.5	2,300

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	51.1	47.4 - 54.7	700	48.9	45.3 - 52.6	1,400
20-24	1,700	40.4	38.3 - 42.6	2,400	59.6	57.4 - 61.7	4,100
25-34	2,200	26.5	25.1 - 27.9	6,000	73.5	72.1 - 74.9	8,200
35+	600	20.6	18.5 - 22.9	2,200	79.4	77.1 - 81.5	2,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,700	31.8	30.4 - 33.3	5,900	68.2	66.7 - 69.6	8,600
Female	2,400	30.3	28.9 - 31.8	5,500	69.7	68.2 - 71.1	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	25.0	23.8 - 26.2	8,300	75.0	73.8 - 76.2	11,000
No	2,400	43.5	41.6 - 45.3	3,100	56.5	54.7 - 58.4	5,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,600	40.7	39.0 - 42.5	3,800	59.3	57.5 - 61.0	6,400
131-185%	800	30.1	27.5 - 32.8	1,800	69.9	67.2 - 72.5	2,600
186+%	1,100	18.7	17.3 - 20.2	5,000	81.3	79.8 - 82.7	6,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	35.0	30.9 - 39.3	700	65.0	60.7 - 69.1	1,100
Normal birth weight	4,800	30.8	29.8 - 31.9	10,700	69.2	68.1 - 70.2	15,400

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	53.7	49.9 - 57.4	600	46.3	42.6 - 50.1	1,400
No	4,300	28.9	27.9 - 30.0	10,600	71.1	70.0 - 72.1	14,900

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	45.0	42.6 - 47.5	1,800	55.0	52.5 - 57.4	3,300
No	3,500	27.2	26.1 - 28.3	9,300	72.8	71.7 - 73.9	12,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd
Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	40.1	38.4 - 41.7	4,200	59.9	58.3 - 61.6	7,100
No	2,200	24.2	22.9 - 25.5	7,000	75.8	74.5 - 77.1	9,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,100	31.3	30.3 - 32.4	11,200	68.7	67.6 - 69.7	16,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	34.9	32.7 - 37.1	1,200	65.1	62.9 - 67.3	1,900
Honolulu	3,700	30.9	29.6 - 32.2	8,300	69.1	67.8 - 70.4	12,000
Kauai	200	26.3	23.1 - 29.8	500	73.7	70.2 - 76.9	700
Maui	500	32.4	30.1 - 34.7	1,100	67.6	65.3 - 69.9	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	24.7	22.5 - 26.9	2,700	75.3	73.1 - 77.5	3,600
Native Hawaiian	1,800	40.9	39.4 - 42.4	2,600	59.1	57.6 - 60.6	4,400
Chinese	100	22.9	18.1 - 28.6	500	77.1	71.4 - 81.9	600
Filipino	1,200	34.9	32.4 - 37.5	2,100	65.1	62.5 - 67.6	3,300
Japanese	300	16.2	13.8 - 18.9	1,800	83.8	81.1 - 86.2	2,100
Other	800	34.6	31.3 - 38.1	1,500	65.4	61.9 - 68.7	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	51.5	48.1 - 55.0	700	48.5	45.0 - 51.9	1,500
20-24	1,600	39.7	37.6 - 41.9	2,400	60.3	58.1 - 62.4	4,000
25-34	2,100	26.6	25.2 - 28.1	5,900	73.4	71.9 - 74.8	8,000
35+	600	21.6	19.4 - 24.0	2,200	78.4	76.0 - 80.6	2,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,700	32.2	30.8 - 33.7	5,700	67.8	66.3 - 69.2	8,500
Female	2,400	30.3	28.9 - 31.8	5,400	69.7	68.2 - 71.1	7,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	25.5	24.3 - 26.7	8,200	74.5	73.3 - 75.7	11,000
No	2,300	43.7	41.9 - 45.5	2,900	56.3	54.5 - 58.1	5,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,600	40.3	38.6 - 41.9	3,900	59.7	58.1 - 61.4	6,500
131-185%	800	29.1	26.7 - 31.7	1,900	70.9	68.3 - 73.3	2,700
186+%	1,100	19.2	17.8 - 20.8	4,700	80.8	79.2 - 82.2	5,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	36.6	32.4 - 40.9	700	63.4	59.1 - 67.6	1,000
Normal birth weight	4,700	30.9	29.9 - 32.0	10,500	69.1	68.0 - 70.1	15,200

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	54.0	50.3 - 57.6	600	46.0	42.4 - 49.7	1,300
No	4,200	29.0	28.0 - 30.1	10,400	71.0	69.9 - 72.0	14,600

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	44.8	42.5 - 47.2	1,800	55.2	52.8 - 57.5	3,200
No	3,500	27.4	26.3 - 28.6	9,100	72.6	71.4 - 73.7	12,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	40.4	38.8 - 42.0	4,200	59.6	58.0 - 61.2	7,000
No	2,200	24.4	23.1 - 25.7	6,900	75.6	74.3 - 76.9	9,100