

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

*The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.*

*Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.*

*Below are the indicators included in this report and the long description for each indicator.*

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (exclusive) duration (grouped)	Breastfeeding (exclusive) duration (grouped)

*Suggested citation:*

*Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/22/17*

*For documentation on other terms and coding in this report, please go to the following links:*

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact  
The Hawai'i Health Data Warehouse**

[profiles@hhdw.org](mailto:profiles@hhdw.org)

**(855) 946-5899 x15**

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**


*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

Years	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>2000-2004</b>	1,700	10.7	10.0 - 11.4	4,700	29.8	28.8 - 30.9	3,100	19.5	18.6 - 20.4	700	4.2	3.7 - 4.6	5,600	35.8	34.7 - 36.9	15,800
<b>2001-2005</b>	1,700	10.2	9.6 - 10.9	5,200	31.7	30.7 - 32.8	3,100	19.0	18.2 - 19.9	600	3.7	3.3 - 4.2	5,800	35.3	34.2 - 36.4	16,400
<b>2002-2006</b>	1,600	9.8	9.1 - 10.5	5,400	32.2	31.1 - 33.2	3,200	19.0	18.1 - 19.9	600	3.8	3.3 - 4.2	5,900	35.3	34.2 - 36.4	16,800
<b>2003-2007</b>	1,600	9.2	8.5 - 9.8	5,700	33.0	32.0 - 34.1	3,200	18.9	18.1 - 19.8	600	3.5	3.1 - 3.9	6,100	35.4	34.3 - 36.4	17,100
<b>2004-2008</b>	1,500	8.6	8.0 - 9.2	5,900	34.0	33.0 - 35.0	3,200	18.4	17.6 - 19.3	600	3.4	3.1 - 3.9	6,200	35.6	34.6 - 36.6	17,400
<b>2005-2009</b>	1,400	8.1	7.5 - 8.8	6,000	34.2	33.1 - 35.3	3,200	18.0	17.1 - 18.9	700	3.7	3.3 - 4.2	6,300	35.9	34.8 - 37.1	17,600
<b>2006-2010</b>	1,300	7.3	6.7 - 8.0	5,900	33.1	31.9 - 34.2	3,200	17.8	16.8 - 18.8	700	4.1	3.6 - 4.6	6,700	37.8	36.6 - 39.0	17,800
<b>2007-2011</b>	1,200	6.6	5.9 - 7.3	5,900	32.8	31.6 - 34.0	3,100	17.3	16.3 - 18.3	700	4.0	3.5 - 4.5	7,100	39.4	38.1 - 40.7	17,900
<b>2008-2012</b>	1,100	6.1	5.4 - 6.8	5,800	32.4	31.1 - 33.7	3,000	17.0	16.0 - 18.1	700	4.0	3.5 - 4.6	7,300	40.6	39.2 - 41.9	17,900
<b>2009-2013</b>	900	5.3	4.7 - 6.0	5,400	30.6	29.2 - 31.9	3,000	16.9	15.8 - 18.1	700	3.9	3.3 - 4.5	7,700	43.3	41.8 - 44.8	17,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	900	5.3	4.7 - 6.0	5,400	30.6	29.2 - 31.9	3,000	16.9	15.8 - 18.1	700	3.9	3.3 - 4.5	7,700	43.3	41.8 - 44.8	17,800

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	100	5.5	4.5 - 6.8	700	30.6	28.4 - 32.8	400	16.3	14.6 - 18.2	100	4.2	3.3 - 5.3	1,000	43.4	41.0 - 45.7	2,300
<b>Honolulu</b>	700	5.7	4.8 - 6.7	3,900	30.6	28.8 - 32.4	2,200	17.2	15.8 - 18.8	500	3.9	3.2 - 4.7	5,400	42.6	40.6 - 44.6	12,800
<b>Kauai</b>	<50	3.8	2.9 - 5.0	200	23.9	21.7 - 26.4	100	17.0	15.0 - 19.1	<50	4.2	3.2 - 5.4	400	51.1	48.3 - 53.8	800
<b>Maui</b>	100	3.0	2.3 - 3.9	600	33.1	31.0 - 35.3	300	15.8	14.2 - 17.5	100	3.3	2.6 - 4.2	900	44.8	42.6 - 47.0	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	3.7	2.7 - 5.1	800	19.2	17.0 - 21.7	700	16.6	14.4 - 19.0	200	4.1	3.1 - 5.5	2,400	56.4	53.4 - 59.3	4,300
Native Hawaiian	400	7.1	5.8 - 8.7	1,600	30.0	27.7 - 32.4	1,100	20.0	17.9 - 22.3	200	4.7	3.7 - 6.0	2,000	38.2	35.6 - 40.8	5,300
Chinese	100	6.8	3.8 - 11.8	300	38.0	31.3 - 45.3	100	11.6	7.8 - 17.0	<50	2.9	1.3 - 6.4	300	40.7	33.8 - 47.9	800
Filipino	100	4.5	3.3 - 6.1	1,300	40.7	37.3 - 44.2	600	18.7	16.1 - 21.6	100	4.0	2.8 - 5.6	1,000	32.2	28.9 - 35.6	3,100
Japanese	<50	1.3	0.6 - 3.1	700	40.1	35.4 - 45.0	200	13.2	10.2 - 17.1	<50	1.5	0.7 - 3.1	700	43.8	39.0 - 48.7	1,700
Other	200	7.4	5.4 - 10.0	800	30.1	26.5 - 34.0	300	13.3	10.7 - 16.4	100	3.6	2.3 - 5.4	1,200	45.7	41.6 - 49.8	2,600

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	5.0	2.9 - 8.7	400	36.8	31.5 - 42.4	200	20.9	16.5 - 26.1	<50	2.9	1.6 - 5.0	400	34.4	29.2 - 40.0	1,100
20-24	200	5.9	4.5 - 7.6	1,100	27.5	24.8 - 30.3	800	19.4	17.0 - 22.1	200	5.0	3.7 - 6.6	1,700	42.3	39.2 - 45.4	4,000
25-34	500	5.1	4.3 - 6.2	2,700	28.8	27.0 - 30.7	1,600	16.6	15.1 - 18.1	400	3.9	3.2 - 4.8	4,300	45.6	43.5 - 47.6	9,500
35+	200	5.2	3.8 - 7.0	1,200	37.6	34.2 - 41.0	400	13.4	11.3 - 15.9	100	2.8	1.9 - 4.2	1,300	41.0	37.7 - 44.5	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	500	5.4	4.5 - 6.5	2,700	30.2	28.3 - 32.1	1,500	16.8	15.3 - 18.4	300	3.8	3.1 - 4.6	3,900	43.8	41.8 - 45.9	9,000
<b>Female</b>	500	5.2	4.3 - 6.3	2,700	30.9	29.0 - 32.9	1,500	17.1	15.5 - 18.8	400	4.0	3.2 - 4.9	3,800	42.8	40.7 - 44.9	8,800

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	400	3.9	3.2 - 4.8	3,200	28.7	27.0 - 30.4	1,800	16.4	15.0 - 17.9	400	3.8	3.1 - 4.6	5,200	47.1	45.2 - 49.1	11,000
<b>No</b>	500	7.5	6.3 - 8.9	2,300	33.6	31.4 - 35.8	1,200	17.8	16.1 - 19.7	300	4.0	3.2 - 5.0	2,500	37.1	34.8 - 39.4	6,800

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	400	7.1	5.9 - 8.5	2,100	33.1	30.8 - 35.4	1,100	16.9	15.2 - 18.8	200	3.2	2.5 - 4.2	2,500	39.7	37.3 - 42.1	6,300
<b>131-185%</b>	200	5.7	4.1 - 8.0	800	29.8	26.5 - 33.5	500	18.7	15.8 - 21.8	200	6.0	4.4 - 8.1	1,100	39.8	36.1 - 43.6	2,800
<b>186+%</b>	200	2.9	2.2 - 3.9	2,200	29.5	27.5 - 31.7	1,200	15.9	14.3 - 17.7	300	3.7	2.9 - 4.7	3,500	47.9	45.6 - 50.2	7,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.8	4.7 - 7.1	500	42.8	40.4 - 45.3	200	16.3	14.5 - 18.2	<50	3.4	2.6 - 4.4	400	31.8	29.5 - 34.1	1,200
Normal birth weight	900	5.3	4.6 - 6.1	4,900	29.7	28.3 - 31.2	2,800	17.0	15.8 - 18.2	600	3.9	3.3 - 4.6	7,300	44.2	42.6 - 45.7	16,600

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	14.0	10.4 - 18.6	400	33.8	28.5 - 39.5	200	15.8	12.1 - 20.3	<50	3.3	1.8 - 5.8	400	33.2	28.0 - 38.8	1,100
No	800	4.7	4.1 - 5.5	5,000	30.4	29.0 - 31.8	2,800	17.0	15.9 - 18.2	700	4.0	3.4 - 4.6	7,200	43.9	42.3 - 45.4	16,300

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	8.5	6.8 - 10.6	1,100	31.4	28.4 - 34.6	600	16.8	14.4 - 19.4	100	4.4	3.2 - 6.0	1,300	38.9	35.6 - 42.2	3,400
No	600	4.6	3.9 - 5.4	4,300	30.4	28.9 - 32.0	2,400	17.0	15.7 - 18.3	500	3.8	3.2 - 4.5	6,200	44.2	42.5 - 45.9	14,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	500	7.0	5.9 - 8.2	2,600	33.3	31.2 - 35.5	1,300	16.9	15.3 - 18.6	300	4.1	3.3 - 5.1	3,000	38.7	36.6 - 40.9	7,700
<b>No</b>	400	4.0	3.2 - 4.9	2,800	28.5	26.7 - 30.3	1,700	17.0	15.5 - 18.6	400	3.7	3.0 - 4.6	4,600	46.8	44.8 - 48.9	9,800



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	1,100	6.1	5.4 - 6.8	5,800	32.4	31.1 - 33.7	3,000	17.0	16.0 - 18.1	700	4.0	3.5 - 4.6	7,300	40.6	39.2 - 41.9	17,900

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	100	6.5	5.3 - 8.0	700	30.8	28.5 - 33.2	400	16.8	15.0 - 18.8	100	4.1	3.3 - 5.2	1,000	41.8	39.4 - 44.3	2,300
<b>Honolulu</b>	800	6.5	5.6 - 7.5	4,200	32.9	31.3 - 34.7	2,200	17.0	15.6 - 18.4	500	4.0	3.4 - 4.8	5,100	39.6	37.8 - 41.4	12,800
<b>Kauai</b>	<50	3.3	2.3 - 4.7	200	26.4	23.4 - 29.5	100	18.3	15.7 - 21.2	<50	4.4	3.2 - 5.9	400	47.7	44.3 - 51.1	800
<b>Maui</b>	100	4.3	3.3 - 5.6	600	32.8	30.6 - 35.2	300	16.9	15.1 - 18.8	100	3.5	2.7 - 4.5	800	42.5	40.1 - 44.9	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	4.3	3.2 - 5.7	800	19.8	17.7 - 22.1	700	17.4	15.3 - 19.7	200	4.8	3.7 - 6.2	2,300	53.7	50.8 - 56.4	4,300
Native Hawaiian	400	7.7	6.4 - 9.3	1,700	32.5	30.2 - 34.9	1,000	19.0	17.0 - 21.1	300	5.2	4.1 - 6.4	1,900	35.7	33.3 - 38.1	5,200
Chinese	100	6.7	3.7 - 11.7	300	36.2	30.0 - 42.9	100	12.1	8.7 - 16.6	<50	3.2	1.4 - 6.8	300	41.8	35.3 - 48.7	800
Filipino	200	5.7	4.4 - 7.4	1,300	42.2	39.0 - 45.4	600	18.9	16.4 - 21.6	100	3.3	2.3 - 4.7	1,000	30.0	27.0 - 33.0	3,200
Japanese	<50	1.6	0.9 - 3.1	800	45.1	40.7 - 49.6	200	13.3	10.6 - 16.7	<50	1.7	1.0 - 3.0	700	38.2	33.9 - 42.6	1,700
Other	200	8.9	6.9 - 11.4	800	31.1	27.7 - 34.6	400	14.0	11.6 - 16.9	100	3.1	2.0 - 4.7	1,200	42.9	39.2 - 46.7	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	6.7	4.4 - 10.0	400	35.5	30.7 - 40.5	300	21.0	17.0 - 25.7	<50	3.0	1.8 - 5.1	400	33.8	29.1 - 38.8	1,300
20-24	300	7.4	5.9 - 9.2	1,200	30.4	27.7 - 33.1	800	20.0	17.7 - 22.6	200	5.1	3.9 - 6.6	1,500	37.2	34.3 - 40.1	4,100
25-34	600	5.9	5.0 - 6.9	2,900	30.5	28.8 - 32.3	1,500	16.2	14.8 - 17.6	400	4.1	3.4 - 4.9	4,100	43.4	41.5 - 45.3	9,400
35+	200	4.9	3.7 - 6.6	1,200	39.1	36.0 - 42.3	400	14.0	12.0 - 16.3	100	2.8	2.0 - 4.1	1,200	39.1	36.0 - 42.3	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	500	6.0	5.1 - 7.0	3,000	32.1	30.3 - 33.9	1,600	17.6	16.2 - 19.2	400	4.0	3.3 - 4.8	3,700	40.3	38.5 - 42.3	9,200
<b>Female</b>	500	6.2	5.3 - 7.3	2,800	32.6	30.8 - 34.5	1,400	16.3	14.9 - 17.9	400	4.0	3.3 - 4.9	3,600	40.8	38.8 - 42.7	8,700

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	500	4.5	3.8 - 5.4	3,400	31.1	29.5 - 32.8	1,800	16.1	14.9 - 17.5	400	3.9	3.3 - 4.7	4,900	44.3	42.5 - 46.1	11,100
<b>No</b>	600	8.6	7.4 - 10.0	2,400	34.3	32.2 - 36.5	1,300	18.4	16.7 - 20.2	300	4.2	3.4 - 5.2	2,400	34.5	32.5 - 36.7	6,900

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	500	8.6	7.3 - 10.0	2,200	34.6	32.4 - 36.8	1,100	17.4	15.7 - 19.2	200	3.3	2.6 - 4.2	2,300	36.2	34.0 - 38.4	6,300
<b>131-185%</b>	200	6.3	4.7 - 8.4	800	29.7	26.5 - 33.0	500	19.0	16.3 - 22.0	200	5.8	4.3 - 7.7	1,100	39.3	35.8 - 42.8	2,800
<b>186+%</b>	200	3.2	2.5 - 4.1	2,400	31.9	29.9 - 33.9	1,200	15.7	14.2 - 17.3	300	4.0	3.2 - 5.0	3,400	45.3	43.1 - 47.4	7,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	6.5	5.2 - 8.2	500	44.4	41.5 - 47.3	200	15.4	13.4 - 17.7	<50	3.6	2.7 - 4.8	400	30.1	27.6 - 32.8	1,200
Normal birth weight	1,000	6.1	5.4 - 6.8	5,300	31.5	30.1 - 32.9	2,900	17.1	16.0 - 18.2	700	4.0	3.5 - 4.7	6,900	41.3	39.9 - 42.8	16,700

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	15.3	11.8 - 19.5	400	34.9	30.0 - 40.0	200	17.6	14.0 - 21.8	100	4.5	2.8 - 7.0	300	27.9	23.4 - 32.8	1,200
No	900	5.4	4.8 - 6.2	5,300	32.2	30.9 - 33.6	2,800	17.0	15.9 - 18.1	700	4.0	3.5 - 4.7	6,800	41.4	39.9 - 42.8	16,300

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	10.0	8.2 - 12.1	1,100	32.9	30.0 - 35.9	600	17.7	15.4 - 20.2	200	5.0	3.8 - 6.5	1,200	34.4	31.5 - 37.5	3,400
No	700	5.2	4.5 - 6.0	4,600	32.3	30.8 - 33.7	2,400	16.8	15.7 - 18.0	500	3.9	3.3 - 4.5	5,900	41.9	40.3 - 43.4	14,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	600	8.3	7.1 - 9.5	2,600	34.0	32.0 - 36.0	1,300	17.2	15.7 - 18.9	300	4.1	3.4 - 5.0	2,800	36.4	34.4 - 38.4	7,700
<b>No</b>	400	4.4	3.6 - 5.3	3,100	31.2	29.6 - 33.0	1,700	16.8	15.4 - 18.3	400	3.9	3.3 - 4.8	4,300	43.6	41.8 - 45.5	9,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	1,200	6.6	5.9 - 7.3	5,900	32.8	31.6 - 34.0	3,100	17.3	16.3 - 18.3	700	4.0	3.5 - 4.5	7,100	39.4	38.1 - 40.7	17,900

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	200	7.8	6.4 - 9.5	700	30.2	27.9 - 32.7	400	15.8	14.0 - 17.8	100	3.7	2.9 - 4.8	1,000	42.4	39.8 - 45.0	2,300
<b>Honolulu</b>	800	6.6	5.8 - 7.6	4,300	33.8	32.2 - 35.4	2,200	17.5	16.2 - 18.9	500	4.0	3.4 - 4.7	4,800	38.1	36.4 - 39.7	12,700
<b>Kauai</b>	<50	4.9	3.3 - 7.3	200	23.1	20.0 - 26.4	200	20.9	17.8 - 24.4	<50	5.0	3.6 - 6.9	400	46.1	42.3 - 50.0	800
<b>Maui</b>	100	5.4	4.2 - 6.9	700	33.4	30.9 - 36.0	300	16.2	14.3 - 18.3	100	3.6	2.7 - 4.7	800	41.4	38.8 - 44.0	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.1	4.1 - 6.5	900	20.4	18.4 - 22.6	700	17.0	15.0 - 19.2	200	5.3	4.2 - 6.7	2,200	52.2	49.5 - 54.8	4,300
Native Hawaiian	400	8.2	6.9 - 9.7	1,700	32.8	30.6 - 35.2	1,000	19.3	17.4 - 21.3	300	4.8	3.8 - 6.0	1,800	34.9	32.6 - 37.3	5,200
Chinese	<50	5.4	2.8 - 10.1	300	35.6	29.9 - 41.8	100	14.9	11.2 - 19.6	<50	3.1	1.5 - 6.2	300	41.0	34.8 - 47.4	700
Filipino	200	6.6	5.3 - 8.2	1,400	43.0	40.0 - 46.0	600	18.5	16.3 - 21.0	100	2.9	2.1 - 4.1	900	29.0	26.3 - 31.9	3,200
Japanese	<50	1.9	1.1 - 3.3	900	46.8	42.8 - 50.9	300	14.0	11.4 - 17.1	<50	2.0	1.2 - 3.3	600	35.2	31.5 - 39.2	1,800
Other	200	9.3	7.4 - 11.6	800	29.8	26.7 - 33.1	400	15.3	12.9 - 18.2	100	3.1	2.0 - 4.7	1,100	42.5	39.1 - 46.1	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.1	5.7 - 11.2	400	32.2	28.0 - 36.8	300	22.5	18.7 - 26.8	<50	3.0	1.8 - 5.1	500	34.2	29.8 - 38.9	1,400
20-24	400	8.4	6.9 - 10.0	1,300	30.8	28.4 - 33.4	900	20.4	18.2 - 22.8	200	4.4	3.4 - 5.8	1,500	36.0	33.4 - 38.7	4,200
25-34	600	6.2	5.3 - 7.2	2,900	31.7	30.0 - 33.4	1,500	15.9	14.6 - 17.3	400	4.1	3.4 - 4.9	3,900	42.1	40.3 - 43.9	9,200
35+	100	4.7	3.5 - 6.2	1,200	38.8	35.8 - 41.8	500	15.0	13.0 - 17.3	100	3.3	2.4 - 4.6	1,200	38.2	35.3 - 41.2	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	600	6.5	5.6 - 7.5	3,000	32.3	30.6 - 34.0	1,700	18.0	16.6 - 19.4	400	3.9	3.3 - 4.7	3,600	39.3	37.6 - 41.1	9,300
<b>Female</b>	600	6.7	5.8 - 7.7	2,900	33.3	31.6 - 35.1	1,400	16.6	15.2 - 18.1	300	4.0	3.3 - 4.8	3,400	39.5	37.6 - 41.3	8,600

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	600	5.0	4.3 - 5.8	3,500	32.0	30.5 - 33.5	1,800	16.1	15.0 - 17.4	400	3.8	3.2 - 4.5	4,800	43.1	41.5 - 44.8	11,100
<b>No</b>	600	9.2	8.0 - 10.5	2,300	34.1	32.1 - 36.2	1,300	19.2	17.5 - 21.0	300	4.2	3.4 - 5.2	2,300	33.3	31.3 - 35.3	6,800

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	600	9.3	8.1 - 10.7	2,200	34.1	32.0 - 36.2	1,200	18.3	16.6 - 20.0	200	3.2	2.6 - 4.1	2,200	35.1	33.0 - 37.2	6,400
<b>131-185%</b>	200	6.6	5.1 - 8.5	800	30.2	27.2 - 33.4	500	18.2	15.7 - 21.0	100	4.9	3.6 - 6.7	1,100	40.1	36.8 - 43.5	2,700
<b>186+%</b>	300	3.8	3.1 - 4.6	2,500	32.9	31.1 - 34.8	1,200	16.0	14.5 - 17.5	300	4.3	3.6 - 5.3	3,200	43.0	41.0 - 45.0	7,600



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	9.2	7.5 - 11.1	1,100	34.3	31.3 - 37.5	600	19.6	17.1 - 22.3	100	2.7	1.8 - 3.9	1,000	34.2	31.3 - 37.3	3,100
\$10,000-\$24,999	300	8.9	7.3 - 10.8	1,200	33.0	30.4 - 35.8	700	18.7	16.6 - 21.1	100	4.0	3.1 - 5.3	1,300	35.3	32.6 - 38.2	3,600
\$25,000-\$49,999	300	6.4	5.1 - 7.9	1,300	30.8	28.4 - 33.4	700	17.6	15.6 - 19.7	200	4.5	3.5 - 5.8	1,700	40.8	38.1 - 43.5	4,200
>=\$50,000	200	3.5	2.8 - 4.4	2,000	33.7	31.6 - 35.8	900	15.1	13.5 - 16.8	300	4.3	3.4 - 5.4	2,600	43.4	41.2 - 45.7	6,000

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.6	6.0 - 9.6	500	43.5	40.3 - 46.7	200	16.4	14.0 - 19.0	100	4.3	3.2 - 5.7	400	28.3	25.5 - 31.2	1,300
Normal birth weight	1,100	6.5	5.8 - 7.2	5,300	32.0	30.7 - 33.3	2,900	17.4	16.3 - 18.5	700	3.9	3.4 - 4.5	6,700	40.2	38.9 - 41.6	16,700

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	16.6	13.3 - 20.6	500	35.4	31.0 - 40.1	200	15.9	12.9 - 19.4	100	4.9	3.3 - 7.3	400	27.2	23.2 - 31.6	1,400
No	900	5.7	5.1 - 6.4	5,300	32.6	31.3 - 33.9	2,800	17.4	16.4 - 18.5	600	3.9	3.4 - 4.5	6,600	40.3	39.0 - 41.7	16,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	400	11.1	9.3 - 13.1	1,200	33.8	31.1 - 36.7	700	18.5	16.4 - 20.8	200	4.9	3.8 - 6.3	1,100	31.7	29.0 - 34.5	3,600
<b>No</b>	800	5.5	4.8 - 6.2	4,600	32.6	31.2 - 34.0	2,400	17.0	15.9 - 18.1	500	3.8	3.2 - 4.4	5,800	41.2	39.8 - 42.7	14,100

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	700	9.0	7.9 - 10.2	2,500	33.0	31.1 - 34.9	1,300	17.6	16.1 - 19.2	300	4.0	3.2 - 4.8	2,800	36.4	34.5 - 38.4	7,600
<b>No</b>	500	4.8	4.0 - 5.6	3,300	32.7	31.1 - 34.3	1,700	17.1	15.8 - 18.5	400	4.0	3.4 - 4.8	4,200	41.5	39.8 - 43.2	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	1,300	7.3	6.7 - 8.0	5,900	33.1	31.9 - 34.2	3,200	17.8	16.8 - 18.8	700	4.1	3.6 - 4.6	6,700	37.8	36.6 - 39.0	17,800

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	200	8.4	6.9 - 10.2	700	29.1	26.6 - 31.8	400	16.5	14.5 - 18.7	100	3.9	3.0 - 5.0	1,000	42.1	39.3 - 44.9	2,300
<b>Honolulu</b>	900	7.4	6.6 - 8.3	4,400	34.7	33.3 - 36.2	2,200	17.7	16.5 - 18.9	500	4.0	3.4 - 4.7	4,600	36.2	34.7 - 37.7	12,700
<b>Kauai</b>	<50	5.9	4.0 - 8.6	200	20.6	17.4 - 24.2	200	21.9	18.5 - 25.8	<50	5.6	3.9 - 7.8	400	46.0	41.8 - 50.4	800
<b>Maui</b>	100	6.3	5.0 - 8.1	600	32.3	29.6 - 35.1	400	18.1	16.0 - 20.5	100	3.7	2.7 - 5.0	800	39.5	36.7 - 42.4	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.1	5.0 - 7.5	900	20.8	18.8 - 22.9	700	16.3	14.5 - 18.3	300	6.1	4.9 - 7.4	2,100	50.8	48.2 - 53.3	4,200
Native Hawaiian	500	9.3	8.0 - 10.8	1,700	32.7	30.5 - 35.0	1,000	20.0	18.2 - 22.0	200	4.2	3.4 - 5.3	1,700	33.7	31.5 - 36.0	5,100
Chinese	<50	4.1	2.0 - 8.1	200	36.8	31.6 - 42.3	100	16.9	13.3 - 21.3	<50	3.5	1.8 - 6.7	300	38.7	33.2 - 44.6	700
Filipino	200	7.0	5.7 - 8.5	1,400	42.6	39.9 - 45.5	600	18.6	16.4 - 20.9	100	2.9	2.1 - 4.0	900	28.9	26.4 - 31.5	3,200
Japanese	<50	2.5	1.6 - 4.0	900	47.8	44.1 - 51.6	300	14.7	12.2 - 17.5	<50	2.1	1.3 - 3.2	600	32.9	29.4 - 36.5	1,900
Other	300	9.9	8.0 - 12.2	800	30.4	27.5 - 33.5	500	17.3	14.9 - 20.0	100	3.4	2.3 - 5.0	1,100	39.0	35.8 - 42.2	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.4	6.2 - 11.3	500	31.5	27.5 - 35.8	300	23.4	19.7 - 27.5	<50	2.5	1.4 - 4.4	500	34.1	30.0 - 38.6	1,500
20-24	400	8.8	7.4 - 10.4	1,300	31.3	29.0 - 33.8	900	20.3	18.2 - 22.5	200	4.5	3.5 - 5.7	1,500	35.1	32.7 - 37.7	4,200
25-34	600	6.8	6.0 - 7.8	2,900	32.1	30.5 - 33.7	1,500	16.5	15.2 - 17.8	400	4.3	3.6 - 5.1	3,600	40.3	38.6 - 42.0	8,900
35+	200	6.2	4.9 - 7.8	1,200	38.9	36.1 - 41.7	500	15.4	13.5 - 17.6	100	3.6	2.7 - 4.8	1,100	35.9	33.2 - 38.7	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	600	7.1	6.2 - 8.0	3,000	33.1	31.5 - 34.7	1,700	18.7	17.4 - 20.1	400	4.3	3.6 - 5.0	3,400	37.0	35.3 - 38.6	9,200
<b>Female</b>	700	7.6	6.7 - 8.6	2,800	33.1	31.4 - 34.8	1,400	16.8	15.5 - 18.2	300	3.8	3.2 - 4.6	3,300	38.7	37.0 - 40.5	8,600

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	600	5.7	5.0 - 6.5	3,600	32.8	31.4 - 34.3	1,800	16.3	15.2 - 17.5	400	4.0	3.4 - 4.7	4,500	41.1	39.6 - 42.6	11,000
<b>No</b>	700	9.9	8.7 - 11.3	2,300	33.5	31.5 - 35.4	1,400	20.2	18.5 - 21.9	300	4.1	3.4 - 5.0	2,200	32.4	30.5 - 34.3	6,700

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	600	10.2	9.0 - 11.5	2,200	33.9	31.9 - 35.9	1,200	18.7	17.1 - 20.4	200	3.3	2.7 - 4.1	2,200	33.9	31.9 - 35.9	6,400
<b>131-185%</b>	200	7.7	6.1 - 9.7	800	31.0	28.0 - 34.1	500	18.4	16.0 - 21.1	100	5.1	3.8 - 6.8	1,000	37.8	34.7 - 41.1	2,600
<b>186+%</b>	300	4.1	3.4 - 5.0	2,500	33.4	31.7 - 35.2	1,200	16.6	15.2 - 18.0	300	4.4	3.7 - 5.3	3,100	41.5	39.7 - 43.3	7,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	10.2	8.5 - 12.3	1,000	33.2	30.4 - 36.2	600	19.6	17.2 - 22.3	100	2.9	2.0 - 4.1	1,000	34.0	31.2 - 37.0	3,000
\$10,000-\$24,999	300	9.4	7.8 - 11.2	1,200	34.4	31.8 - 37.1	700	18.7	16.7 - 20.9	100	4.0	3.1 - 5.2	1,200	33.5	31.0 - 36.2	3,600
\$25,000-\$49,999	300	6.8	5.6 - 8.3	1,300	31.5	29.2 - 34.0	700	18.2	16.3 - 20.3	200	4.9	3.9 - 6.2	1,600	38.5	36.0 - 41.0	4,100
>=\$50,000	200	4.1	3.4 - 5.1	2,000	33.7	31.8 - 35.7	900	15.8	14.3 - 17.5	200	4.1	3.3 - 5.1	2,500	42.2	40.1 - 44.3	5,900

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.2	7.3 - 11.6	500	41.5	38.1 - 45.0	200	16.9	14.3 - 19.8	100	3.9	2.8 - 5.3	400	28.4	25.4 - 31.7	1,300
Normal birth weight	1,200	7.2	6.5 - 7.9	5,300	32.4	31.2 - 33.6	2,900	17.8	16.9 - 18.9	700	4.1	3.6 - 4.6	6,400	38.5	37.3 - 39.8	16,500

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	18.7	15.4 - 22.4	500	33.6	29.5 - 37.9	200	16.3	13.5 - 19.7	100	4.0	2.7 - 5.8	400	27.4	23.6 - 31.5	1,500
No	1,000	6.3	5.7 - 7.0	5,300	33.0	31.8 - 34.2	2,900	18.0	17.0 - 19.0	700	4.1	3.6 - 4.7	6,200	38.7	37.4 - 39.9	16,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	500	12.5	10.7 - 14.5	1,200	32.7	30.1 - 35.3	700	19.6	17.5 - 21.9	200	5.0	4.0 - 6.3	1,100	30.3	27.8 - 32.9	3,600
<b>No</b>	800	6.0	5.4 - 6.8	4,600	33.1	31.8 - 34.4	2,400	17.4	16.3 - 18.5	500	3.9	3.3 - 4.5	5,500	39.6	38.3 - 41.0	13,900

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	700	9.9	8.8 - 11.1	2,500	33.1	31.3 - 34.9	1,300	17.9	16.5 - 19.5	300	4.1	3.4 - 5.0	2,600	35.0	33.2 - 36.9	7,400
<b>No</b>	500	5.5	4.7 - 6.3	3,300	33.0	31.5 - 34.6	1,800	17.6	16.4 - 18.9	400	4.1	3.4 - 4.8	4,000	39.8	38.2 - 41.4	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	1,400	8.1	7.5 - 8.8	6,000	34.2	33.1 - 35.3	3,200	18.0	17.1 - 18.9	700	3.7	3.3 - 4.2	6,300	35.9	34.8 - 37.1	17,600

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	200	10.0	8.3 - 12.0	600	28.5	25.9 - 31.3	400	17.7	15.5 - 20.1	100	3.3	2.5 - 4.5	900	40.5	37.6 - 43.5	2,300
<b>Honolulu</b>	1,000	8.1	7.3 - 8.9	4,600	36.4	35.1 - 37.8	2,200	17.9	16.8 - 19.0	500	3.7	3.1 - 4.3	4,300	34.0	32.7 - 35.4	12,600
<b>Kauai</b>	100	6.3	4.3 - 9.3	200	18.9	15.5 - 22.7	200	23.2	19.4 - 27.4	<50	5.4	3.7 - 7.9	400	46.2	41.6 - 51.0	800
<b>Maui</b>	100	6.9	5.4 - 8.8	600	33.0	30.1 - 36.0	300	17.4	15.1 - 19.9	100	4.0	3.0 - 5.5	700	38.8	35.8 - 41.9	1,900



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	6.0	5.0 - 7.3	900	22.2	20.3 - 24.3	700	16.2	14.6 - 18.1	200	5.7	4.6 - 6.9	2,000	49.8	47.4 - 52.2	4,100
Native Hawaiian	500	10.7	9.3 - 12.3	1,700	33.3	31.2 - 35.5	1,000	20.4	18.6 - 22.4	200	3.7	2.9 - 4.7	1,600	31.9	29.7 - 34.1	5,000
Chinese	<50	6.0	3.6 - 9.9	300	40.4	35.8 - 45.1	100	18.1	15.0 - 21.7	<50	2.5	1.7 - 3.7	200	33.0	28.7 - 37.6	600
Filipino	200	7.5	6.3 - 9.0	1,400	43.0	40.5 - 45.6	600	18.4	16.4 - 20.5	100	2.9	2.2 - 3.7	900	28.2	25.9 - 30.6	3,200
Japanese	100	3.4	2.4 - 4.9	1,000	49.1	45.7 - 52.4	300	14.8	12.6 - 17.4	<50	2.3	1.5 - 3.5	600	30.4	27.4 - 33.6	2,000
Other	300	11.0	9.2 - 13.1	800	30.9	28.1 - 33.8	500	18.2	16.0 - 20.7	100	3.3	2.2 - 4.7	1,000	36.6	33.7 - 39.6	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.6	7.4 - 12.3	500	32.1	28.3 - 36.1	300	22.6	19.2 - 26.3	<50	2.5	1.5 - 4.3	500	33.3	29.3 - 37.4	1,500
20-24	400	9.5	8.2 - 11.1	1,300	32.2	30.0 - 34.6	900	21.3	19.3 - 23.4	200	4.4	3.5 - 5.5	1,400	32.6	30.3 - 34.9	4,200
25-34	700	7.8	6.9 - 8.7	2,900	33.5	31.9 - 35.0	1,400	16.5	15.3 - 17.7	300	3.9	3.3 - 4.6	3,400	38.4	36.8 - 40.0	8,800
35+	200	6.4	5.2 - 7.8	1,200	40.0	37.4 - 42.6	500	15.8	14.0 - 17.8	100	3.0	2.2 - 4.1	1,100	34.8	32.4 - 37.4	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	700	7.8	7.0 - 8.8	3,100	34.0	32.5 - 35.6	1,700	19.2	17.9 - 20.5	300	3.7	3.1 - 4.4	3,200	35.2	33.7 - 36.8	9,100
<b>Female</b>	700	8.4	7.5 - 9.4	2,900	34.4	32.8 - 36.0	1,400	16.8	15.6 - 18.1	300	3.8	3.2 - 4.5	3,100	36.7	35.1 - 38.3	8,500

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	700	6.5	5.8 - 7.2	3,700	34.0	32.7 - 35.4	1,800	16.4	15.4 - 17.5	400	3.8	3.3 - 4.4	4,300	39.3	37.9 - 40.7	11,000
<b>No</b>	700	10.8	9.6 - 12.1	2,300	34.4	32.6 - 36.3	1,400	20.7	19.2 - 22.4	200	3.7	3.0 - 4.5	2,000	30.4	28.7 - 32.3	6,600

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	800	11.8	10.6 - 13.2	2,200	34.5	32.6 - 36.4	1,200	18.7	17.2 - 20.3	200	3.1	2.5 - 3.9	2,000	31.8	30.0 - 33.7	6,300
<b>131-185%</b>	200	8.9	7.2 - 10.9	700	30.7	27.9 - 33.7	500	19.4	17.0 - 22.1	100	4.6	3.3 - 6.3	900	36.4	33.4 - 39.5	2,400
<b>186+%</b>	300	4.3	3.6 - 5.0	2,700	35.4	33.8 - 37.0	1,200	16.5	15.3 - 17.8	300	4.1	3.5 - 4.9	3,000	39.7	38.1 - 41.4	7,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	400	12.2	10.3 - 14.3	1,000	34.4	31.6 - 37.3	600	18.8	16.6 - 21.2	100	2.8	2.0 - 4.0	900	31.7	29.1 - 34.6	3,000
\$10,000-\$24,999	400	10.3	8.9 - 12.1	1,300	34.9	32.5 - 37.4	700	19.5	17.6 - 21.6	100	3.6	2.7 - 4.7	1,200	31.7	29.3 - 34.1	3,700
\$25,000-\$49,999	300	7.4	6.2 - 8.8	1,300	32.0	29.8 - 34.3	700	18.2	16.4 - 20.1	200	4.7	3.8 - 5.9	1,500	37.7	35.4 - 40.1	4,000
>=\$50,000	300	4.5	3.7 - 5.4	2,100	35.8	33.9 - 37.6	900	16.1	14.7 - 17.6	200	3.7	3.0 - 4.6	2,300	39.9	38.0 - 41.9	5,700

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.2	7.2 - 11.7	600	43.3	39.6 - 47.0	200	16.4	13.8 - 19.5	<50	3.8	2.6 - 5.3	400	27.3	24.1 - 30.8	1,300
Normal birth weight	1,300	8.0	7.3 - 8.7	5,400	33.5	32.3 - 34.6	3,000	18.2	17.2 - 19.1	600	3.7	3.3 - 4.2	6,000	36.6	35.5 - 37.8	16,300

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	19.5	16.4 - 23.1	500	34.9	31.1 - 39.0	300	16.4	13.7 - 19.6	100	3.8	2.6 - 5.6	400	25.3	21.9 - 29.1	1,500
No	1,100	7.1	6.4 - 7.7	5,400	34.0	32.9 - 35.2	2,900	18.2	17.2 - 19.1	600	3.8	3.3 - 4.3	5,800	37.0	35.8 - 38.2	15,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	500	13.8	12.0 - 15.8	1,200	33.0	30.6 - 35.6	700	19.6	17.7 - 21.7	200	5.2	4.1 - 6.5	1,000	28.3	26.0 - 30.8	3,600
<b>No</b>	900	6.7	6.0 - 7.4	4,700	34.4	33.2 - 35.6	2,400	17.6	16.6 - 18.6	500	3.4	3.0 - 3.9	5,200	37.9	36.7 - 39.2	13,700

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	800	11.0	9.9 - 12.2	2,400	33.1	31.4 - 34.8	1,400	18.7	17.4 - 20.2	300	3.7	3.1 - 4.5	2,400	33.4	31.7 - 35.2	7,300
<b>No</b>	600	6.0	5.3 - 6.8	3,500	34.9	33.4 - 36.3	1,700	17.4	16.2 - 18.6	400	3.8	3.3 - 4.5	3,800	37.9	36.5 - 39.4	9,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	1,500	8.6	8.0 - 9.2	5,900	34.0	33.0 - 35.0	3,200	18.4	17.6 - 19.3	600	3.4	3.1 - 3.9	6,200	35.6	34.6 - 36.6	17,400

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	200	11.0	9.2 - 13.2	600	27.6	24.9 - 30.4	400	17.6	15.4 - 20.1	100	2.6	1.8 - 3.8	900	41.2	38.1 - 44.2	2,200
<b>Honolulu</b>	1,100	8.5	7.8 - 9.3	4,600	36.5	35.3 - 37.7	2,300	18.2	17.3 - 19.2	400	3.3	2.9 - 3.8	4,200	33.4	32.3 - 34.6	12,500
<b>Kauai</b>	<50	5.9	3.8 - 8.9	100	17.6	14.2 - 21.7	200	23.9	19.9 - 28.4	<50	4.7	3.0 - 7.3	400	47.9	43.0 - 52.9	800
<b>Maui</b>	100	7.2	5.6 - 9.1	600	31.9	28.9 - 35.0	400	18.6	16.2 - 21.3	100	4.5	3.3 - 6.1	700	37.9	34.7 - 41.1	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.6	5.6 - 7.8	900	21.4	19.7 - 23.3	700	16.8	15.2 - 18.5	200	5.1	4.2 - 6.2	2,000	50.0	47.8 - 52.2	4,000
Native Hawaiian	500	11.0	9.7 - 12.5	1,600	33.8	31.7 - 35.9	1,000	20.6	18.9 - 22.5	200	3.5	2.7 - 4.4	1,500	31.1	29.1 - 33.2	4,800
Chinese	<50	4.6	3.5 - 5.9	300	40.2	37.4 - 43.0	100	19.5	17.3 - 21.9	<50	2.6	1.9 - 3.7	200	33.2	30.5 - 35.9	600
Filipino	300	8.2	7.0 - 9.6	1,300	41.6	39.3 - 43.8	600	18.7	17.0 - 20.6	100	3.3	2.5 - 4.2	900	28.2	26.2 - 30.4	3,200
Japanese	100	4.2	3.0 - 5.7	1,000	48.3	45.3 - 51.4	300	16.2	14.1 - 18.6	<50	2.2	1.5 - 3.2	600	29.1	26.4 - 31.9	2,100
Other	300	11.8	10.1 - 13.8	800	31.4	28.9 - 34.0	500	18.1	16.0 - 20.4	100	2.3	1.6 - 3.3	1,000	36.4	33.7 - 39.1	2,700

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	11.1	8.8 - 14.0	400	30.7	27.1 - 34.5	300	23.0	19.8 - 26.6	<50	2.8	1.8 - 4.5	500	32.4	28.7 - 36.3	1,400
20-24	400	10.0	8.7 - 11.4	1,300	31.7	29.7 - 33.8	900	21.5	19.7 - 23.4	200	4.0	3.2 - 5.0	1,400	32.8	30.7 - 34.9	4,200
25-34	700	8.0	7.2 - 8.9	2,900	34.2	32.8 - 35.6	1,400	16.7	15.6 - 17.8	300	3.5	3.0 - 4.1	3,200	37.6	36.2 - 39.1	8,600
35+	200	6.9	5.7 - 8.2	1,200	38.1	35.8 - 40.4	500	17.2	15.5 - 19.1	100	2.8	2.1 - 3.7	1,100	35.1	32.9 - 37.4	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	800	8.3	7.5 - 9.2	3,100	34.2	32.8 - 35.6	1,700	19.1	18.0 - 20.3	300	3.5	3.0 - 4.0	3,200	34.9	33.5 - 36.3	9,100
<b>Female</b>	700	8.8	7.9 - 9.8	2,800	33.8	32.4 - 35.3	1,500	17.7	16.5 - 18.9	300	3.4	2.9 - 4.0	3,000	36.3	34.8 - 37.8	8,400

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	800	7.2	6.5 - 7.9	3,900	34.2	33.0 - 35.4	1,900	17.2	16.3 - 18.2	400	3.4	3.0 - 3.9	4,300	38.0	36.7 - 39.2	11,300
<b>No</b>	700	11.1	9.9 - 12.3	2,100	33.6	31.8 - 35.4	1,300	20.7	19.2 - 22.3	200	3.5	2.9 - 4.3	1,900	31.2	29.5 - 33.0	6,200

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	800	12.2	11.0 - 13.5	2,100	33.0	31.3 - 34.7	1,300	20.0	18.6 - 21.5	200	2.8	2.3 - 3.5	2,000	31.9	30.3 - 33.7	6,400
<b>131-185%</b>	200	10.0	8.3 - 11.9	700	30.0	27.3 - 32.8	400	18.7	16.5 - 21.1	100	3.9	2.9 - 5.2	900	37.5	34.7 - 40.4	2,300
<b>186+%</b>	400	4.7	4.1 - 5.5	2,700	36.2	34.7 - 37.6	1,300	16.7	15.5 - 17.9	300	3.9	3.3 - 4.5	2,900	38.6	37.1 - 40.1	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Income (2004-2011)	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	300	12.1	10.3 - 14.0	1,000	34.0	31.5 - 36.7	600	20.0	17.8 - 22.3	100	2.8	2.0 - 3.9	900	31.2	28.6 - 33.8	2,900
\$10,000-\$24,999	400	11.3	9.9 - 12.9	1,200	32.3	30.1 - 34.5	800	20.2	18.4 - 22.2	100	3.2	2.5 - 4.1	1,200	33.0	30.9 - 35.2	3,800
\$25,000-\$49,999	300	7.4	6.3 - 8.6	1,300	32.0	30.0 - 34.0	800	19.0	17.3 - 20.8	200	4.3	3.5 - 5.3	1,500	37.4	35.3 - 39.5	4,100
>=\$50,000	300	5.1	4.4 - 6.0	2,000	37.0	35.3 - 38.7	900	15.7	14.4 - 17.1	200	3.3	2.7 - 4.1	2,100	38.8	37.1 - 40.6	5,500

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.3	7.2 - 12.0	600	44.2	40.3 - 48.2	200	16.3	13.5 - 19.6	<50	3.8	2.6 - 5.6	300	26.3	22.9 - 30.0	1,200
Normal birth weight	1,400	8.5	7.9 - 9.2	5,400	33.2	32.2 - 34.2	3,000	18.6	17.8 - 19.5	600	3.4	3.0 - 3.8	5,900	36.3	35.2 - 37.3	16,200

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	19.6	16.6 - 22.9	500	32.0	28.5 - 35.7	300	18.7	15.9 - 22.0	100	4.7	3.3 - 6.7	400	25.0	21.8 - 28.5	1,500
No	1,200	7.6	7.0 - 8.2	5,400	34.1	33.1 - 35.2	2,900	18.4	17.6 - 19.3	500	3.4	3.0 - 3.8	5,700	36.5	35.5 - 37.6	15,700



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	500	13.5	11.8 - 15.3	1,100	31.4	29.2 - 33.8	800	21.7	19.8 - 23.9	200	5.4	4.4 - 6.6	1,000	28.0	25.8 - 30.3	3,500
<b>No</b>	1,000	7.3	6.7 - 8.0	4,700	34.6	33.4 - 35.7	2,400	17.6	16.7 - 18.6	400	3.0	2.6 - 3.4	5,100	37.5	36.3 - 38.6	13,700

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	800	11.4	10.3 - 12.5	2,400	32.2	30.6 - 33.8	1,400	19.5	18.2 - 20.9	200	3.2	2.7 - 3.9	2,500	33.7	32.1 - 35.3	7,400
<b>No</b>	600	6.4	5.7 - 7.1	3,400	35.3	34.0 - 36.6	1,700	17.5	16.5 - 18.6	400	3.6	3.2 - 4.2	3,600	37.2	35.9 - 38.5	9,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	1,600	9.2	8.5 - 9.8	5,700	33.0	32.0 - 34.1	3,200	18.9	18.1 - 19.8	600	3.5	3.1 - 3.9	6,100	35.4	34.3 - 36.4	17,100

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	300	12.1	10.3 - 14.2	500	25.6	23.1 - 28.3	400	17.9	15.7 - 20.3	100	3.1	2.2 - 4.3	900	41.4	38.5 - 44.3	2,100
<b>Honolulu</b>	1,100	9.1	8.4 - 9.9	4,400	35.8	34.5 - 37.0	2,300	18.9	17.9 - 20.0	400	3.3	2.9 - 3.8	4,100	32.9	31.7 - 34.2	12,400
<b>Kauai</b>	100	7.3	5.1 - 10.3	100	15.6	12.5 - 19.2	200	22.9	19.2 - 27.1	<50	4.8	3.1 - 7.3	400	49.4	44.7 - 54.1	800
<b>Maui</b>	100	6.9	5.4 - 8.7	600	31.0	28.1 - 34.0	300	18.7	16.4 - 21.3	100	4.6	3.5 - 6.2	700	38.8	35.7 - 41.9	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	7.0	5.8 - 8.3	800	21.5	19.6 - 23.5	600	16.7	15.0 - 18.5	200	5.5	4.5 - 6.7	1,900	49.4	47.0 - 51.7	3,900
Native Hawaiian	600	12.0	10.7 - 13.4	1,500	32.1	30.1 - 34.0	1,000	21.2	19.5 - 23.0	200	3.2	2.5 - 4.0	1,500	31.6	29.7 - 33.5	4,700
Chinese	<50	4.8	3.3 - 6.8	300	41.3	37.2 - 45.5	100	17.5	14.5 - 20.8	<50	2.4	1.7 - 3.5	200	34.1	30.2 - 38.1	600
Filipino	300	9.4	8.0 - 11.0	1,300	38.5	36.1 - 40.9	700	20.1	18.1 - 22.1	100	3.4	2.6 - 4.4	900	28.6	26.5 - 30.9	3,300
Japanese	100	5.0	3.7 - 6.7	1,000	47.9	44.7 - 51.1	300	15.8	13.6 - 18.3	<50	1.8	1.2 - 2.9	600	29.4	26.6 - 32.5	2,100
Other	300	11.5	9.6 - 13.6	800	31.0	28.3 - 33.9	500	19.7	17.4 - 22.2	100	2.8	1.9 - 4.1	900	35.0	32.2 - 37.9	2,500

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	11.9	9.6 - 14.7	400	30.8	27.2 - 34.6	400	24.4	21.1 - 28.0	<50	2.4	1.5 - 3.9	400	30.5	26.9 - 34.2	1,400
20-24	500	10.8	9.5 - 12.3	1,300	30.2	28.2 - 32.3	900	22.2	20.4 - 24.2	200	3.8	3.0 - 4.8	1,400	32.9	30.8 - 35.0	4,200
25-34	700	8.6	7.7 - 9.5	2,800	33.4	32.0 - 34.9	1,400	17.2	16.1 - 18.4	300	3.8	3.2 - 4.5	3,100	37.0	35.5 - 38.5	8,400
35+	200	7.2	5.9 - 8.6	1,100	36.9	34.5 - 39.4	500	16.5	14.7 - 18.5	100	2.7	2.0 - 3.7	1,100	36.7	34.3 - 39.2	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	800	9.1	8.2 - 10.0	3,000	33.3	31.9 - 34.8	1,800	19.6	18.4 - 20.9	300	3.6	3.0 - 4.2	3,100	34.5	33.0 - 35.9	8,900
<b>Female</b>	800	9.3	8.4 - 10.3	2,700	32.7	31.2 - 34.2	1,500	18.2	17.0 - 19.4	300	3.4	2.9 - 4.1	3,000	36.4	34.9 - 37.9	8,200

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	800	7.5	6.8 - 8.2	3,700	33.3	32.1 - 34.6	2,000	17.7	16.7 - 18.8	400	3.6	3.1 - 4.1	4,200	37.9	36.6 - 39.2	11,100
<b>No</b>	700	12.3	11.1 - 13.6	1,900	32.5	30.7 - 34.3	1,300	21.2	19.7 - 22.8	200	3.4	2.8 - 4.2	1,800	30.6	28.9 - 32.4	6,000

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	900	13.2	11.9 - 14.5	2,100	31.8	30.1 - 33.5	1,400	20.9	19.4 - 22.4	200	2.8	2.2 - 3.5	2,000	31.4	29.8 - 33.1	6,500
<b>131-185%</b>	200	9.8	8.1 - 11.9	700	29.8	27.0 - 32.7	400	19.2	16.9 - 21.7	100	4.2	3.1 - 5.6	800	37.0	34.1 - 40.0	2,200
<b>186+%</b>	300	4.6	4.0 - 5.3	2,500	35.5	34.0 - 37.1	1,200	16.9	15.7 - 18.2	300	4.0	3.4 - 4.7	2,800	38.9	37.3 - 40.6	7,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.8	7.6 - 12.5	500	41.5	37.5 - 45.6	200	17.9	14.9 - 21.5	<50	3.9	2.6 - 5.9	300	26.9	23.4 - 30.7	1,200
Normal birth weight	1,400	9.1	8.5 - 9.8	5,100	32.4	31.3 - 33.4	3,000	19.0	18.1 - 19.9	600	3.5	3.1 - 3.9	5,700	36.0	34.9 - 37.1	15,900

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	23.6	20.4 - 27.2	400	30.5	27.1 - 34.2	200	16.9	14.2 - 20.0	100	3.6	2.4 - 5.3	400	25.3	22.1 - 28.8	1,500
No	1,200	7.8	7.2 - 8.5	5,100	33.2	32.1 - 34.3	3,000	19.2	18.3 - 20.1	500	3.5	3.1 - 4.0	5,600	36.3	35.2 - 37.4	15,400

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	15.1	13.4 - 17.1	1,100	31.0	28.7 - 33.3	700	21.2	19.2 - 23.3	200	4.9	3.9 - 6.1	1,000	27.8	25.7 - 30.1	3,500
No	1,000	7.5	6.9 - 8.2	4,500	33.5	32.3 - 34.6	2,500	18.4	17.4 - 19.4	400	3.2	2.8 - 3.6	5,000	37.4	36.2 - 38.6	13,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	900	12.3	11.2 - 13.4	2,300	31.1	29.5 - 32.7	1,500	20.0	18.7 - 21.4	200	3.2	2.7 - 3.9	2,400	33.4	31.8 - 35.1	7,300
<b>No</b>	600	6.7	6.0 - 7.5	3,300	34.3	33.0 - 35.7	1,700	18.0	16.9 - 19.2	400	3.7	3.2 - 4.3	3,600	37.2	35.8 - 38.6	9,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	1,600	9.8	9.1 - 10.5	5,400	32.2	31.1 - 33.2	3,200	19.0	18.1 - 19.9	600	3.8	3.3 - 4.2	5,900	35.3	34.2 - 36.4	16,800

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	200	12.1	10.4 - 14.1	500	25.8	23.4 - 28.3	400	19.5	17.3 - 21.8	100	3.2	2.4 - 4.4	800	39.4	36.7 - 42.2	2,000
<b>Honolulu</b>	1,200	10.0	9.1 - 10.9	4,300	34.7	33.4 - 36.0	2,300	18.6	17.6 - 19.7	400	3.6	3.2 - 4.2	4,100	33.1	31.8 - 34.4	12,300
<b>Kauai</b>	100	7.4	5.3 - 10.1	100	15.5	12.6 - 18.9	200	21.9	18.4 - 26.0	<50	4.3	2.8 - 6.7	400	50.8	46.3 - 55.4	700
<b>Maui</b>	100	6.9	5.5 - 8.6	500	28.9	26.3 - 31.7	300	19.8	17.5 - 22.3	100	4.9	3.7 - 6.4	700	39.4	36.6 - 42.4	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.8	5.6 - 8.3	800	20.5	18.5 - 22.6	600	16.5	14.7 - 18.4	200	6.5	5.4 - 8.0	1,900	49.7	47.2 - 52.2	3,700
Native Hawaiian	600	13.6	12.3 - 15.0	1,400	31.2	29.4 - 33.0	1,000	20.7	19.1 - 22.3	200	3.4	2.7 - 4.1	1,400	31.2	29.5 - 33.0	4,600
Chinese	<50	7.3	4.7 - 11.1	300	41.2	36.1 - 46.4	100	18.6	14.9 - 23.1	<50	2.6	1.5 - 4.5	200	30.3	25.9 - 35.1	600
Filipino	300	9.4	7.9 - 11.0	1,200	37.6	35.1 - 40.2	700	20.1	18.1 - 22.3	100	3.2	2.5 - 4.2	1,000	29.7	27.4 - 32.2	3,300
Japanese	100	4.5	3.3 - 6.1	1,000	45.5	42.1 - 48.9	400	17.6	15.2 - 20.4	<50	2.0	1.2 - 3.2	700	30.4	27.4 - 33.6	2,100
Other	300	13.2	11.0 - 15.7	700	30.5	27.5 - 33.6	500	19.6	17.1 - 22.3	100	2.7	1.9 - 4.0	800	34.0	30.9 - 37.2	2,400

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	13.4	10.9 - 16.4	500	32.5	28.9 - 36.3	300	23.5	20.3 - 27.1	<50	2.5	1.5 - 4.2	400	28.1	24.8 - 31.7	1,400
20-24	500	10.9	9.6 - 12.5	1,200	29.9	27.8 - 32.0	900	22.4	20.5 - 24.4	200	3.9	3.1 - 4.9	1,400	32.9	30.8 - 35.1	4,100
25-34	800	9.3	8.3 - 10.3	2,700	32.0	30.5 - 33.5	1,400	17.3	16.1 - 18.6	300	4.2	3.6 - 4.9	3,100	37.3	35.7 - 38.8	8,300
35+	200	8.0	6.6 - 9.6	1,000	35.7	33.1 - 38.3	500	16.9	15.0 - 19.0	100	2.9	2.1 - 3.9	1,100	36.6	34.1 - 39.2	2,900



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	900	10.2	9.2 - 11.2	2,800	32.5	31.0 - 34.0	1,700	19.3	18.1 - 20.6	300	3.9	3.3 - 4.6	3,000	34.1	32.7 - 35.6	8,700
<b>Female</b>	800	9.4	8.5 - 10.4	2,600	31.8	30.3 - 33.4	1,500	18.7	17.4 - 20.0	300	3.6	3.0 - 4.3	2,900	36.5	35.0 - 38.1	8,100

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	900	7.9	7.1 - 8.7	3,600	32.3	31.0 - 33.6	2,000	17.7	16.6 - 18.8	400	4.0	3.4 - 4.6	4,200	38.2	36.9 - 39.6	11,100
<b>No</b>	800	13.6	12.3 - 15.0	1,800	31.9	30.1 - 33.8	1,200	21.6	20.0 - 23.2	200	3.3	2.7 - 4.1	1,700	29.6	27.9 - 31.3	5,700

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	900	14.2	12.9 - 15.6	2,000	31.0	29.3 - 32.7	1,300	20.6	19.1 - 22.1	200	2.7	2.2 - 3.4	2,000	31.5	29.8 - 33.2	6,400
<b>131-185%</b>	200	10.1	8.3 - 12.2	700	28.9	26.1 - 31.9	400	18.3	16.0 - 20.8	100	5.0	3.8 - 6.6	900	37.7	34.7 - 40.8	2,400
<b>186+%</b>	300	4.3	3.6 - 5.0	2,300	34.6	32.9 - 36.3	1,200	17.5	16.2 - 18.9	300	4.3	3.7 - 5.2	2,600	39.3	37.6 - 41.1	6,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.3	7.2 - 12.0	500	41.5	37.3 - 45.9	200	17.8	14.5 - 21.6	<50	3.5	2.2 - 5.5	300	27.9	24.1 - 32.0	1,100
Normal birth weight	1,500	9.8	9.1 - 10.6	4,900	31.5	30.4 - 32.6	3,000	19.1	18.2 - 20.1	600	3.8	3.3 - 4.3	5,600	35.8	34.7 - 37.0	15,600

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	26.1	22.8 - 29.8	400	29.1	25.7 - 32.8	200	16.7	14.0 - 19.9	<50	3.1	2.1 - 4.7	400	24.9	21.7 - 28.4	1,400
No	1,300	8.3	7.6 - 9.0	4,900	32.4	31.3 - 33.5	2,900	19.3	18.3 - 20.2	600	3.9	3.4 - 4.4	5,500	36.2	35.1 - 37.4	15,100

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	16.1	14.3 - 18.1	1,000	30.4	28.1 - 32.8	700	20.2	18.3 - 22.3	200	4.9	4.0 - 6.1	1,000	28.4	26.2 - 30.7	3,400
No	1,100	8.1	7.4 - 8.8	4,200	32.5	31.3 - 33.7	2,400	18.7	17.7 - 19.8	500	3.5	3.1 - 4.0	4,800	37.2	35.9 - 38.4	13,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,000	13.3	12.2 - 14.6	2,200	30.9	29.3 - 32.5	1,400	20.2	18.8 - 21.6	200	3.3	2.8 - 4.0	2,300	32.3	30.7 - 34.0	7,200
<b>No</b>	700	7.1	6.3 - 7.9	3,100	33.0	31.6 - 34.5	1,700	18.1	16.9 - 19.3	400	4.1	3.5 - 4.7	3,500	37.8	36.3 - 39.2	9,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	1,700	10.2	9.6 - 10.9	5,200	31.7	30.7 - 32.8	3,100	19.0	18.2 - 19.9	600	3.7	3.3 - 4.2	5,800	35.3	34.2 - 36.4	16,400

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	300	13.5	11.8 - 15.4	500	25.1	22.9 - 27.4	400	19.2	17.2 - 21.3	100	3.0	2.3 - 4.0	800	39.2	36.7 - 41.8	2,000
<b>Honolulu</b>	1,200	10.4	9.5 - 11.3	4,100	34.0	32.7 - 35.4	2,300	18.8	17.7 - 19.9	400	3.7	3.2 - 4.3	4,000	33.2	31.8 - 34.5	12,000
<b>Kauai</b>	100	6.9	5.1 - 9.3	100	16.4	13.6 - 19.6	200	22.1	18.8 - 25.7	<50	4.3	2.9 - 6.3	400	50.3	46.2 - 54.5	700
<b>Maui</b>	100	7.0	5.8 - 8.5	500	29.8	27.3 - 32.3	300	19.3	17.2 - 21.5	100	4.4	3.3 - 5.7	700	39.6	37.0 - 42.3	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	6.9	5.7 - 8.3	700	20.1	18.1 - 22.2	600	16.8	15.0 - 18.8	200	6.1	4.9 - 7.5	1,800	50.1	47.6 - 52.6	3,600
Native Hawaiian	700	14.6	13.4 - 15.8	1,400	31.0	29.4 - 32.7	900	19.5	18.2 - 20.9	200	3.6	2.9 - 4.3	1,400	31.4	29.8 - 33.0	4,500
Chinese	<50	7.7	5.0 - 11.6	300	39.8	34.4 - 45.5	100	19.3	15.1 - 24.3	<50	2.8	1.5 - 5.3	200	30.3	25.5 - 35.7	600
Filipino	300	9.5	8.0 - 11.2	1,200	36.6	34.1 - 39.2	700	20.9	18.8 - 23.1	100	3.1	2.3 - 4.1	1,000	29.9	27.5 - 32.4	3,300
Japanese	100	4.4	3.2 - 5.9	1,000	44.6	41.3 - 48.1	400	18.6	16.1 - 21.4	100	2.6	1.7 - 4.0	600	29.7	26.7 - 32.9	2,100
Other	300	14.3	12.0 - 17.0	700	29.9	26.9 - 33.1	400	19.1	16.6 - 21.9	100	2.5	1.6 - 3.7	800	34.2	31.1 - 37.5	2,300

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	15.0	12.5 - 17.8	500	33.0	29.6 - 36.6	300	23.1	20.2 - 26.4	<50	3.0	1.9 - 4.6	400	25.9	22.9 - 29.1	1,400
20-24	500	11.5	10.2 - 13.0	1,200	28.7	26.7 - 30.8	900	22.8	21.0 - 24.7	100	3.6	2.9 - 4.5	1,400	33.4	31.3 - 35.5	4,100
25-34	800	9.7	8.7 - 10.7	2,600	31.8	30.3 - 33.4	1,400	16.9	15.7 - 18.1	300	4.2	3.5 - 4.9	3,000	37.5	35.9 - 39.0	8,100
35+	200	7.6	6.3 - 9.3	1,000	35.2	32.6 - 37.9	500	17.6	15.6 - 19.8	100	2.9	2.1 - 3.9	1,000	36.7	34.1 - 39.3	2,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	900	10.6	9.6 - 11.6	2,700	32.0	30.6 - 33.5	1,600	19.1	17.9 - 20.4	300	3.7	3.1 - 4.3	3,000	34.6	33.2 - 36.1	8,500
<b>Female</b>	800	9.9	9.0 - 10.9	2,500	31.4	29.9 - 33.0	1,500	18.9	17.7 - 20.2	300	3.8	3.2 - 4.4	2,800	36.0	34.5 - 37.6	7,900

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	900	7.8	7.1 - 8.6	3,500	31.7	30.4 - 33.1	2,000	18.1	17.1 - 19.2	400	4.0	3.5 - 4.6	4,200	38.3	37.0 - 39.7	11,000
<b>No</b>	800	15.2	13.9 - 16.6	1,700	31.7	30.0 - 33.5	1,100	20.8	19.4 - 22.4	200	3.1	2.5 - 3.7	1,600	29.2	27.6 - 30.9	5,500

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	1,000	14.9	13.7 - 16.2	1,900	30.5	28.8 - 32.2	1,300	20.4	19.0 - 21.8	200	2.6	2.1 - 3.2	2,000	31.7	30.0 - 33.3	6,400
<b>131-185%</b>	300	10.0	8.3 - 11.9	700	28.3	25.8 - 31.1	500	18.9	16.7 - 21.2	100	5.1	3.9 - 6.5	1,000	37.8	35.0 - 40.6	2,600
<b>186+%</b>	300	4.2	3.5 - 4.9	2,100	34.5	32.7 - 36.2	1,100	17.3	16.0 - 18.8	300	4.5	3.8 - 5.4	2,400	39.5	37.7 - 41.4	6,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.7	7.6 - 12.2	400	41.6	37.2 - 46.0	200	18.3	15.0 - 22.3	<50	3.5	2.2 - 5.6	300	26.8	23.1 - 31.0	1,100
Normal birth weight	1,600	10.3	9.6 - 11.0	4,800	31.0	30.0 - 32.1	2,900	19.1	18.2 - 20.0	600	3.7	3.3 - 4.2	5,500	35.9	34.8 - 37.0	15,300

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	26.5	23.2 - 30.0	400	29.2	25.9 - 32.8	200	17.0	14.4 - 20.0	<50	2.8	1.9 - 4.1	300	24.5	21.4 - 27.9	1,400
No	1,300	8.7	8.1 - 9.4	4,700	31.9	30.8 - 33.1	2,800	19.2	18.3 - 20.2	600	3.8	3.4 - 4.3	5,400	36.3	35.1 - 37.4	14,800

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	16.5	14.7 - 18.4	1,000	30.3	28.1 - 32.6	700	20.0	18.1 - 22.0	100	4.2	3.4 - 5.3	1,000	29.0	26.8 - 31.3	3,300
No	1,100	8.5	7.8 - 9.3	4,100	32.0	30.8 - 33.2	2,400	18.8	17.8 - 19.9	500	3.6	3.2 - 4.2	4,700	37.0	35.8 - 38.3	12,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,000	14.0	12.8 - 15.2	2,100	30.4	28.9 - 32.0	1,400	19.9	18.6 - 21.3	200	3.2	2.6 - 3.8	2,300	32.6	31.0 - 34.2	7,000
<b>No</b>	700	7.3	6.5 - 8.1	3,000	32.7	31.2 - 34.1	1,700	18.4	17.2 - 19.6	400	4.2	3.6 - 4.8	3,500	37.5	36.1 - 39.0	9,200



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

State	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	1,700	10.7	10.0 - 11.4	4,700	29.8	28.8 - 30.9	3,100	19.5	18.6 - 20.4	700	4.2	3.7 - 4.6	5,600	35.8	34.7 - 36.9	15,800

County	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	200	13.4	11.9 - 15.1	500	25.2	23.2 - 27.3	300	18.3	16.5 - 20.2	100	3.5	2.7 - 4.5	700	39.6	37.3 - 42.0	1,800
<b>Honolulu</b>	1,300	10.9	10.0 - 11.8	3,700	31.6	30.3 - 33.0	2,200	19.2	18.1 - 20.4	500	4.2	3.7 - 4.8	4,000	34.1	32.7 - 35.4	11,600
<b>Kauai</b>	<50	7.0	5.3 - 9.1	100	16.4	13.7 - 19.5	200	21.9	18.9 - 25.3	<50	4.5	3.2 - 6.5	300	50.1	46.2 - 54.0	700
<b>Maui</b>	100	7.8	6.6 - 9.2	500	27.8	25.6 - 30.1	400	21.9	19.9 - 24.1	100	4.4	3.4 - 5.7	600	38.1	35.6 - 40.6	1,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

DOH Race-Ethnicity	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	8.1	6.8 - 9.6	600	18.5	16.6 - 20.6	600	17.2	15.4 - 19.2	200	6.9	5.7 - 8.4	1,700	49.3	46.8 - 51.8	3,500
Native Hawaiian	600	14.9	13.8 - 16.0	1,300	29.5	28.1 - 30.9	900	20.9	19.7 - 22.1	200	3.7	3.2 - 4.4	1,300	31.1	29.7 - 32.5	4,300
Chinese	<50	7.6	4.7 - 12.0	200	38.4	32.5 - 44.6	100	19.0	14.5 - 24.4	<50	2.6	1.2 - 5.6	200	32.4	27.0 - 38.4	600
Filipino	300	9.5	8.0 - 11.2	1,100	34.5	31.9 - 37.2	700	21.3	19.1 - 23.6	100	3.4	2.6 - 4.5	1,000	31.3	28.9 - 33.9	3,200
Japanese	100	5.0	3.7 - 6.8	800	41.2	37.7 - 44.8	400	19.3	16.6 - 22.4	100	3.2	2.1 - 4.7	600	31.3	28.0 - 34.7	2,000
Other	300	14.4	12.0 - 17.3	600	28.8	25.6 - 32.1	400	18.3	15.7 - 21.2	100	3.0	2.1 - 4.5	800	35.5	32.1 - 38.9	2,200

PRAMS Age Group	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	14.9	12.6 - 17.6	500	32.0	28.8 - 35.4	300	23.4	20.6 - 26.4	<50	3.0	2.0 - 4.5	400	26.7	23.8 - 29.7	1,500
20-24	500	12.0	10.7 - 13.4	1,100	27.0	25.0 - 29.0	900	23.2	21.4 - 25.1	100	3.6	2.8 - 4.5	1,400	34.3	32.3 - 36.4	3,900
25-34	800	10.0	9.0 - 11.0	2,300	29.7	28.1 - 31.2	1,300	17.4	16.2 - 18.7	400	4.8	4.1 - 5.6	2,900	38.2	36.6 - 39.8	7,700
35+	200	8.5	7.0 - 10.3	900	33.3	30.6 - 36.0	500	17.9	15.9 - 20.2	100	3.9	3.0 - 5.0	1,000	36.4	33.7 - 39.2	2,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

Infant Gender	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	900	11.3	10.3 - 12.3	2,500	29.9	28.5 - 31.4	1,600	19.5	18.3 - 20.8	300	4.1	3.5 - 4.8	2,900	35.2	33.7 - 36.7	8,200
<b>Female</b>	800	10.0	9.1 - 11.0	2,200	29.7	28.2 - 31.3	1,500	19.5	18.2 - 20.8	300	4.3	3.6 - 5.0	2,700	36.5	34.9 - 38.1	7,500

Married	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	900	8.5	7.7 - 9.3	3,100	29.5	28.2 - 30.8	2,000	18.7	17.6 - 19.8	500	4.5	4.0 - 5.2	4,100	38.8	37.4 - 40.2	10,600
<b>No</b>	800	15.2	13.9 - 16.7	1,600	30.5	28.8 - 32.3	1,100	21.2	19.7 - 22.8	200	3.4	2.8 - 4.1	1,500	29.7	28.0 - 31.4	5,100

Poverty Level	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	900	14.8	13.7 - 16.1	1,800	28.5	26.9 - 30.1	1,300	20.9	19.6 - 22.4	200	3.0	2.5 - 3.6	2,100	32.8	31.2 - 34.4	6,400
<b>131-185%</b>	300	10.1	8.5 - 12.0	700	27.0	24.5 - 29.7	500	19.4	17.3 - 21.7	100	5.7	4.5 - 7.1	1,000	37.8	35.0 - 40.6	2,600
<b>186+%</b>	200	4.5	3.8 - 5.4	1,800	32.3	30.5 - 34.2	1,000	17.9	16.4 - 19.4	300	5.1	4.3 - 6.1	2,200	40.2	38.3 - 42.1	5,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

Birthweight	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	11.1	8.9 - 13.7	400	37.7	33.4 - 42.3	200	20.6	17.1 - 24.7	<50	3.6	2.3 - 5.7	300	26.9	23.1 - 31.1	1,000
Normal birth weight	1,600	10.6	9.9 - 11.4	4,300	29.2	28.2 - 30.3	2,900	19.4	18.5 - 20.4	600	4.2	3.8 - 4.7	5,400	36.4	35.3 - 37.6	14,700

Smoke 3rd Trimester	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	27.8	24.6 - 31.3	400	26.8	23.6 - 30.2	300	19.2	16.5 - 22.2	<50	3.2	2.2 - 4.6	300	23.0	20.1 - 26.2	1,300
No	1,300	9.0	8.3 - 9.7	4,300	30.1	29.0 - 31.3	2,800	19.6	18.7 - 20.6	600	4.3	3.8 - 4.8	5,200	37.0	35.8 - 38.1	14,100

Smoke Before Pregnant	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	17.2	15.4 - 19.1	900	27.9	25.7 - 30.1	700	21.0	19.1 - 23.1	100	4.2	3.4 - 5.3	900	29.7	27.5 - 31.9	3,200
No	1,100	8.8	8.1 - 9.6	3,700	30.2	29.0 - 31.4	2,300	19.2	18.2 - 20.3	500	4.2	3.7 - 4.7	4,600	37.5	36.3 - 38.8	12,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped)- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

WIC During Pregnancy	Never breastfed			Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,000	14.6	13.4 - 15.8	2,000	29.0	27.5 - 30.6	1,400	20.0	18.7 - 21.3	200	3.6	3.0 - 4.3	2,200	32.8	31.2 - 34.4	6,800
<b>No</b>	700	7.5	6.8 - 8.4	2,700	30.4	29.0 - 31.9	1,700	19.2	18.0 - 20.4	400	4.6	4.0 - 5.3	3,400	38.3	36.8 - 39.8	8,800