

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

*The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.*

*Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.*

*Below are the indicators included in this report and the long description for each indicator.*

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (exclusive) duration 8+ weeks	Breastfeeding (exclusive) duration 8+ weeks

*Suggested citation:*

*Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/22/17*

*For documentation on other terms and coding in this report, please go to the following links:*

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact  
The Hawai'i Health Data Warehouse**

[profiles@hhdw.org](mailto:profiles@hhdw.org)

**(855) 946-5899 x15**

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- for the State of Hawaii,  
for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**


*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

Years	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	10,000	63.2	62.1 - 64.3	5,800	36.8	35.7 - 37.9	15,800
2001-2005	10,500	63.9	62.8 - 65.0	5,900	36.1	35.0 - 37.2	16,400
2002-2006	10,700	64.0	62.9 - 65.0	6,000	36.0	35.0 - 37.1	16,800
2003-2007	11,000	64.0	63.0 - 65.1	6,200	36.0	34.9 - 37.0	17,100
2004-2008	11,100	63.9	62.9 - 64.9	6,300	36.1	35.1 - 37.1	17,400
2005-2009	11,100	63.4	62.3 - 64.5	6,400	36.6	35.5 - 37.7	17,600
2006-2010	10,900	61.4	60.2 - 62.6	6,900	38.6	37.4 - 39.8	17,800
2007-2011	10,700	59.9	58.6 - 61.2	7,200	40.1	38.8 - 41.4	17,900
2008-2012	10,500	58.6	57.2 - 59.9	7,400	41.4	40.1 - 42.8	17,900
2009-2013	9,900	55.8	54.3 - 57.3	7,900	44.2	42.7 - 45.7	17,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	9,900	55.8	54.3 - 57.3	7,900	44.2	42.7 - 45.7	17,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,300	55.5	53.1 - 57.8	1,000	44.5	42.2 - 46.9	2,300
<b>Honolulu</b>	7,200	56.6	54.6 - 58.5	5,600	43.4	41.5 - 45.4	12,800
<b>Kauai</b>	400	47.8	45.0 - 50.5	400	52.2	49.5 - 55.0	800
<b>Maui</b>	1,000	54.3	52.1 - 56.6	900	45.7	43.4 - 47.9	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	42.9	40.0 - 45.9	2,500	57.1	54.1 - 60.0	4,300
Native Hawaiian	3,200	60.8	58.2 - 63.4	2,100	39.2	36.6 - 41.8	5,300
Chinese	500	57.8	50.5 - 64.7	400	42.2	35.3 - 49.5	800
Filipino	2,000	66.5	63.1 - 69.8	1,000	33.5	30.2 - 36.9	3,100
Japanese	900	56.1	51.1 - 60.9	700	43.9	39.1 - 48.9	1,700
Other	1,400	53.5	49.4 - 57.6	1,200	46.5	42.4 - 50.6	2,600

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	65.0	59.4 - 70.3	400	35.0	29.7 - 40.6	1,100
20-24	2,300	56.7	53.6 - 59.9	1,800	43.3	40.1 - 46.4	4,000
25-34	5,100	53.4	51.4 - 55.5	4,400	46.6	44.5 - 48.6	9,500
35+	1,800	58.4	54.9 - 61.7	1,300	41.6	38.3 - 45.1	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,000	55.3	53.2 - 57.3	4,000	44.7	42.7 - 46.8	9,000
Female	4,900	56.3	54.2 - 58.4	3,800	43.7	41.6 - 45.8	8,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,700	52.0	50.0 - 53.9	5,300	48.0	46.1 - 50.0	11,000
No	4,200	62.0	59.7 - 64.3	2,600	38.0	35.7 - 40.3	6,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,700	59.7	57.3 - 62.1	2,500	40.3	37.9 - 42.7	6,300
131-185%	1,600	59.2	55.4 - 62.9	1,100	40.8	37.1 - 44.6	2,800
186+%	3,700	51.0	48.7 - 53.3	3,600	49.0	46.7 - 51.3	7,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	800	67.5	65.1 - 69.7	400	32.5	30.3 - 34.9	1,200
<b>Normal birth weight</b>	9,100	54.9	53.4 - 56.5	7,500	45.1	43.5 - 46.6	16,600

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	700	66.0	60.4 - 71.2	400	34.0	28.8 - 39.6	1,100
<b>No</b>	9,000	55.2	53.7 - 56.8	7,300	44.8	43.2 - 46.3	16,300

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	2,000	60.3	56.9 - 63.5	1,300	39.7	36.5 - 43.1	3,400
<b>No</b>	7,700	54.9	53.2 - 56.5	6,300	45.1	43.5 - 46.8	14,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,700	60.3	58.1 - 62.5	3,100	39.7	37.5 - 41.9	7,700
<b>No</b>	5,200	52.3	50.3 - 54.4	4,700	47.7	45.6 - 49.7	9,800



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,500	58.6	57.2 - 59.9	7,400	41.4	40.1 - 42.8	17,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,300	57.2	54.7 - 59.6	1,000	42.8	40.4 - 45.3	2,300
<b>Honolulu</b>	7,700	59.6	57.8 - 61.4	5,200	40.4	38.6 - 42.2	12,800
<b>Kauai</b>	400	51.3	47.9 - 54.6	400	48.7	45.4 - 52.1	800
<b>Maui</b>	1,100	56.8	54.4 - 59.2	900	43.2	40.8 - 45.6	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	45.5	42.7 - 48.3	2,300	54.5	51.7 - 57.3	4,300
Native Hawaiian	3,300	63.3	60.8 - 65.7	1,900	36.7	34.3 - 39.2	5,200
Chinese	400	56.5	49.7 - 63.1	300	43.5	36.9 - 50.3	800
Filipino	2,200	69.0	65.9 - 71.9	1,000	31.0	28.1 - 34.1	3,200
Japanese	1,100	61.8	57.4 - 66.0	700	38.2	34.0 - 42.6	1,700
Other	1,500	56.5	52.8 - 60.2	1,200	43.5	39.8 - 47.2	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	65.7	60.7 - 70.4	400	34.3	29.6 - 39.3	1,300
20-24	2,500	61.8	58.9 - 64.7	1,500	38.2	35.3 - 41.1	4,100
25-34	5,300	55.7	53.8 - 57.6	4,200	44.3	42.4 - 46.2	9,400
35+	1,900	60.2	57.0 - 63.3	1,300	39.8	36.7 - 43.0	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,400	58.8	56.9 - 60.7	3,800	41.2	39.3 - 43.1	9,200
Female	5,100	58.3	56.3 - 60.3	3,600	41.7	39.7 - 43.7	8,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,100	54.8	53.0 - 56.5	5,000	45.2	43.5 - 47.0	11,100
No	4,400	64.7	62.6 - 66.8	2,400	35.3	33.2 - 37.4	6,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,000	63.2	60.9 - 65.4	2,300	36.8	34.6 - 39.1	6,300
131-185%	1,700	59.9	56.4 - 63.4	1,100	40.1	36.6 - 43.6	2,800
186+%	4,000	53.6	51.5 - 55.8	3,400	46.4	44.2 - 48.5	7,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	800	69.2	66.5 - 71.7	400	30.8	28.3 - 33.5	1,200
<b>Normal birth weight</b>	9,700	57.8	56.3 - 59.2	7,000	42.2	40.8 - 43.7	16,700

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	900	71.3	66.4 - 75.8	400	28.7	24.2 - 33.6	1,200
<b>No</b>	9,400	57.8	56.3 - 59.2	6,900	42.2	40.8 - 43.7	16,300

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	2,200	64.7	61.7 - 67.7	1,200	35.3	32.3 - 38.3	3,400
<b>No</b>	8,100	57.2	55.7 - 58.8	6,000	42.8	41.2 - 44.3	14,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,900	62.7	60.7 - 64.7	2,900	37.3	35.3 - 39.3	7,700
<b>No</b>	5,500	55.6	53.7 - 57.4	4,400	44.4	42.6 - 46.3	9,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,700	59.9	58.6 - 61.2	7,200	40.1	38.8 - 41.4	17,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,300	56.8	54.2 - 59.4	1,000	43.2	40.6 - 45.8	2,300
<b>Honolulu</b>	7,800	61.2	59.5 - 62.9	4,900	38.8	37.1 - 40.5	12,700
<b>Kauai</b>	400	52.9	49.0 - 56.7	400	47.1	43.3 - 51.0	800
<b>Maui</b>	1,200	58.0	55.4 - 60.6	800	42.0	39.4 - 44.6	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,000	46.9	44.3 - 49.6	2,300	53.1	50.4 - 55.7	4,300
Native Hawaiian	3,400	64.2	61.8 - 66.5	1,900	35.8	33.5 - 38.2	5,200
Chinese	400	58.1	51.6 - 64.3	300	41.9	35.7 - 48.4	700
Filipino	2,200	70.3	67.4 - 73.0	900	29.7	27.0 - 32.6	3,200
Japanese	1,200	64.6	60.6 - 68.4	600	35.4	31.6 - 39.4	1,800
Other	1,500	57.0	53.5 - 60.5	1,200	43.0	39.5 - 46.5	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	65.3	60.6 - 69.7	500	34.7	30.3 - 39.4	1,400
20-24	2,700	63.3	60.6 - 65.9	1,500	36.7	34.1 - 39.4	4,200
25-34	5,200	57.0	55.2 - 58.8	3,900	43.0	41.2 - 44.8	9,200
35+	1,900	61.2	58.2 - 64.1	1,200	38.8	35.9 - 41.8	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,600	60.0	58.2 - 61.7	3,700	40.0	38.3 - 41.8	9,300
Female	5,200	59.8	57.9 - 61.6	3,500	40.2	38.4 - 42.1	8,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,200	56.1	54.5 - 57.8	4,900	43.9	42.2 - 45.5	11,100
No	4,500	66.0	63.9 - 68.0	2,300	34.0	32.0 - 36.1	6,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,100	64.3	62.2 - 66.4	2,300	35.7	33.6 - 37.8	6,400
131-185%	1,600	59.5	56.1 - 62.8	1,100	40.5	37.2 - 43.9	2,700
186+%	4,200	56.0	54.0 - 58.0	3,300	44.0	42.0 - 46.0	7,600



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	65.5	62.4 - 68.5	1,100	34.5	31.5 - 37.6	3,100
\$10,000-\$24,999	2,300	63.7	60.9 - 66.5	1,300	36.3	33.5 - 39.1	3,600
\$25,000-\$49,999	2,400	58.7	56.0 - 61.3	1,700	41.3	38.7 - 44.0	4,200
>=\$50,000	3,300	55.5	53.3 - 57.7	2,700	44.5	42.3 - 46.7	6,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	70.7	67.7 - 73.5	400	29.3	26.5 - 32.3	1,300
Normal birth weight	9,800	59.1	57.7 - 60.4	6,800	40.9	39.6 - 42.3	16,700

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	71.9	67.4 - 75.9	400	28.1	24.1 - 32.6	1,400
No	9,600	59.0	57.6 - 60.3	6,700	41.0	39.7 - 42.4	16,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,400	67.5	64.7 - 70.2	1,200	32.5	29.8 - 35.3	3,600
<b>No</b>	8,200	58.1	56.6 - 59.5	5,900	41.9	40.5 - 43.4	14,100

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,800	62.8	60.8 - 64.8	2,800	37.2	35.2 - 39.2	7,600
<b>No</b>	5,800	57.8	56.1 - 59.6	4,200	42.2	40.4 - 43.9	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,900	61.4	60.2 - 62.6	6,900	38.6	37.4 - 39.8	17,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,300	57.1	54.2 - 59.8	1,000	42.9	40.2 - 45.8	2,300
<b>Honolulu</b>	8,000	63.0	61.5 - 64.5	4,700	37.0	35.5 - 38.5	12,700
<b>Kauai</b>	400	52.9	48.6 - 57.2	400	47.1	42.8 - 51.4	800
<b>Maui</b>	1,200	59.9	57.0 - 62.7	800	40.1	37.3 - 43.0	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,000	48.1	45.6 - 50.6	2,200	51.9	49.4 - 54.4	4,200
Native Hawaiian	3,400	65.5	63.2 - 67.8	1,800	34.5	32.2 - 36.8	5,100
Chinese	400	60.2	54.3 - 65.8	300	39.8	34.2 - 45.7	700
Filipino	2,200	70.4	67.7 - 72.9	900	29.6	27.1 - 32.3	3,200
Japanese	1,200	66.9	63.2 - 70.3	600	33.1	29.7 - 36.8	1,900
Other	1,600	60.4	57.2 - 63.6	1,100	39.6	36.4 - 42.8	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	65.4	61.0 - 69.5	500	34.6	30.5 - 39.0	1,500
20-24	2,700	64.2	61.7 - 66.7	1,500	35.8	33.3 - 38.3	4,200
25-34	5,300	58.8	57.1 - 60.5	3,700	41.2	39.5 - 42.9	9,000
35+	2,000	63.3	60.6 - 66.1	1,200	36.7	33.9 - 39.4	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,700	62.2	60.5 - 63.8	3,500	37.8	36.2 - 39.5	9,200
Female	5,200	60.6	58.9 - 62.4	3,400	39.4	37.6 - 41.1	8,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,400	58.0	56.5 - 59.5	4,600	42.0	40.5 - 43.5	11,000
No	4,500	67.1	65.1 - 69.0	2,200	32.9	31.0 - 34.9	6,700

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,200	65.5	63.5 - 67.5	2,200	34.5	32.5 - 36.5	6,400
131-185%	1,600	61.5	58.3 - 64.6	1,000	38.5	35.4 - 41.7	2,600
186+%	4,300	57.4	55.6 - 59.3	3,200	42.6	40.7 - 44.4	7,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	65.7	62.8 - 68.6	1,000	34.3	31.4 - 37.2	3,000
\$10,000-\$24,999	2,400	65.6	62.9 - 68.2	1,200	34.4	31.8 - 37.1	3,600
\$25,000-\$49,999	2,500	61.0	58.4 - 63.5	1,600	39.0	36.5 - 41.6	4,100
>=\$50,000	3,300	56.6	54.5 - 58.7	2,600	43.4	41.3 - 45.5	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	70.4	67.1 - 73.5	400	29.6	26.5 - 32.9	1,300
Normal birth weight	10,000	60.7	59.5 - 62.0	6,500	39.3	38.0 - 40.5	16,500

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	71.8	67.6 - 75.6	400	28.2	24.4 - 32.4	1,500
No	9,700	60.6	59.3 - 61.8	6,300	39.4	38.2 - 40.7	16,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,500	69.1	66.4 - 71.6	1,100	30.9	28.4 - 33.6	3,600
<b>No</b>	8,300	59.6	58.2 - 60.9	5,600	40.4	39.1 - 41.8	13,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,800	64.2	62.3 - 66.0	2,700	35.8	34.0 - 37.7	7,400
<b>No</b>	6,000	59.5	57.8 - 61.1	4,100	40.5	38.9 - 42.2	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	11,100	63.4	62.3 - 64.5	6,400	36.6	35.5 - 37.7	17,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,300	58.9	55.9 - 61.8	900	41.1	38.2 - 44.1	2,300
<b>Honolulu</b>	8,200	65.4	64.0 - 66.7	4,300	34.6	33.3 - 36.0	12,600
<b>Kauai</b>	400	52.8	48.1 - 57.5	400	47.2	42.5 - 51.9	800
<b>Maui</b>	1,200	60.6	57.5 - 63.6	800	39.4	36.4 - 42.5	1,900



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,000	49.2	46.8 - 51.6	2,100	50.8	48.4 - 53.2	4,100
Native Hawaiian	3,400	67.6	65.3 - 69.7	1,600	32.4	30.3 - 34.7	5,000
Chinese	400	66.5	61.9 - 70.8	200	33.5	29.2 - 38.1	600
Filipino	2,300	71.2	68.8 - 73.5	900	28.8	26.5 - 31.2	3,200
Japanese	1,400	69.3	66.1 - 72.3	600	30.7	27.7 - 33.9	2,000
Other	1,700	62.8	59.8 - 65.7	1,000	37.2	34.3 - 40.2	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	66.3	62.1 - 70.2	500	33.7	29.8 - 37.9	1,500
20-24	2,800	66.9	64.5 - 69.1	1,400	33.1	30.9 - 35.5	4,200
25-34	5,300	60.8	59.2 - 62.4	3,400	39.2	37.6 - 40.8	8,800
35+	2,000	64.5	62.0 - 67.0	1,100	35.5	33.0 - 38.0	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,800	64.1	62.5 - 65.7	3,300	35.9	34.3 - 37.5	9,100
Female	5,300	62.6	61.0 - 64.2	3,200	37.4	35.8 - 39.0	8,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,600	60.0	58.5 - 61.4	4,400	40.0	38.6 - 41.5	11,000
No	4,600	69.1	67.3 - 70.9	2,000	30.9	29.1 - 32.7	6,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,300	67.6	65.7 - 69.5	2,100	32.4	30.5 - 34.3	6,300
131-185%	1,500	63.1	60.0 - 66.1	900	36.9	33.9 - 40.0	2,400
186+%	4,500	59.4	57.7 - 61.1	3,000	40.6	38.9 - 42.3	7,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	68.0	65.2 - 70.7	900	32.0	29.3 - 34.8	3,000
\$10,000-\$24,999	2,500	67.6	65.2 - 70.0	1,200	32.4	30.0 - 34.8	3,700
\$25,000-\$49,999	2,500	61.7	59.4 - 64.1	1,500	38.3	35.9 - 40.6	4,000
>=\$50,000	3,400	59.1	57.2 - 61.0	2,300	40.9	39.0 - 42.8	5,700

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	71.7	68.2 - 75.0	400	28.3	25.0 - 31.8	1,300
Normal birth weight	10,200	62.7	61.6 - 63.9	6,100	37.3	36.1 - 38.4	16,300

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	73.8	70.0 - 77.3	400	26.2	22.7 - 30.0	1,500
No	9,800	62.4	61.2 - 63.6	5,900	37.6	36.4 - 38.8	15,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,500	70.9	68.5 - 73.3	1,000	29.1	26.7 - 31.5	3,600
<b>No</b>	8,400	61.4	60.2 - 62.7	5,300	38.6	37.3 - 39.8	13,700

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,800	65.9	64.1 - 67.6	2,500	34.1	32.4 - 35.9	7,300
<b>No</b>	6,100	61.4	60.0 - 62.9	3,800	38.6	37.1 - 40.0	9,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	11,100	63.9	62.9 - 64.9	6,300	36.1	35.1 - 37.1	17,400

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,300	58.6	55.5 - 61.6	900	41.4	38.4 - 44.5	2,200
<b>Honolulu</b>	8,300	66.0	64.8 - 67.1	4,300	34.0	32.9 - 35.2	12,500
<b>Kauai</b>	400	51.6	46.6 - 56.6	400	48.4	43.4 - 53.4	800
<b>Maui</b>	1,200	61.6	58.3 - 64.7	700	38.4	35.3 - 41.7	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,000	49.0	46.8 - 51.2	2,000	51.0	48.8 - 53.2	4,000
Native Hawaiian	3,300	68.5	66.4 - 70.5	1,500	31.5	29.5 - 33.6	4,800
Chinese	400	66.2	63.4 - 68.8	200	33.8	31.2 - 36.6	600
Filipino	2,300	71.3	69.1 - 73.3	900	28.7	26.7 - 30.9	3,200
Japanese	1,500	70.6	67.8 - 73.3	600	29.4	26.7 - 32.2	2,100
Other	1,700	63.3	60.5 - 65.9	1,000	36.7	34.1 - 39.5	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	67.0	63.1 - 70.7	500	33.0	29.3 - 36.9	1,400
20-24	2,800	66.8	64.7 - 68.9	1,400	33.2	31.1 - 35.3	4,200
25-34	5,300	61.7	60.2 - 63.1	3,300	38.3	36.9 - 39.8	8,600
35+	2,000	64.4	62.1 - 66.6	1,100	35.6	33.4 - 37.9	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,900	64.6	63.2 - 66.0	3,200	35.4	34.0 - 36.8	9,100
Female	5,300	63.1	61.6 - 64.5	3,100	36.9	35.5 - 38.4	8,400

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,900	61.4	60.1 - 62.6	4,300	38.6	37.4 - 39.9	11,300
No	4,200	68.4	66.6 - 70.1	2,000	31.6	29.9 - 33.4	6,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,300	67.7	66.0 - 69.4	2,100	32.3	30.6 - 34.0	6,400
131-185%	1,400	62.1	59.2 - 65.0	900	37.9	35.0 - 40.8	2,300
186+%	4,600	60.6	59.1 - 62.1	3,000	39.4	37.9 - 40.9	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	68.6	66.0 - 71.2	900	31.4	28.8 - 34.0	2,900
\$10,000-\$24,999	2,500	66.6	64.4 - 68.8	1,300	33.4	31.2 - 35.6	3,800
\$25,000-\$49,999	2,600	62.0	59.9 - 64.1	1,600	38.0	35.9 - 40.1	4,100
>=\$50,000	3,300	60.4	58.6 - 62.1	2,200	39.6	37.9 - 41.4	5,500

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	72.8	69.1 - 76.2	300	27.2	23.8 - 30.9	1,200
Normal birth weight	10,200	63.2	62.1 - 64.2	6,000	36.8	35.8 - 37.9	16,200

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	74.0	70.5 - 77.3	400	26.0	22.7 - 29.5	1,500
No	9,900	62.9	61.9 - 64.0	5,800	37.1	36.0 - 38.1	15,700



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,500	71.2	68.9 - 73.4	1,000	28.8	26.6 - 31.1	3,500
<b>No</b>	8,500	62.0	60.9 - 63.2	5,200	38.0	36.8 - 39.1	13,700

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,900	65.8	64.2 - 67.4	2,500	34.2	32.6 - 35.8	7,400
<b>No</b>	6,100	62.2	60.9 - 63.5	3,700	37.8	36.5 - 39.1	9,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	11,000	64.0	63.0 - 65.1	6,200	36.0	34.9 - 37.0	17,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,200	58.2	55.3 - 61.1	900	41.8	38.9 - 44.7	2,100
<b>Honolulu</b>	8,200	66.5	65.2 - 67.7	4,100	33.5	32.3 - 34.8	12,400
<b>Kauai</b>	400	49.6	44.9 - 54.3	400	50.4	45.7 - 55.1	800
<b>Maui</b>	1,100	60.5	57.4 - 63.6	700	39.5	36.4 - 42.6	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	49.5	47.2 - 51.9	2,000	50.5	48.1 - 52.8	3,900
Native Hawaiian	3,200	68.1	66.1 - 70.0	1,500	31.9	30.0 - 33.9	4,700
Chinese	400	65.2	61.1 - 69.1	200	34.8	30.9 - 38.9	600
Filipino	2,300	70.8	68.5 - 73.0	1,000	29.2	27.0 - 31.5	3,300
Japanese	1,500	70.3	67.2 - 73.1	600	29.7	26.9 - 32.8	2,100
Other	1,600	64.5	61.6 - 67.3	900	35.5	32.7 - 38.4	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	69.1	65.3 - 72.6	400	30.9	27.4 - 34.7	1,400
20-24	2,800	66.7	64.6 - 68.8	1,400	33.3	31.2 - 35.4	4,200
25-34	5,200	62.2	60.7 - 63.7	3,200	37.8	36.3 - 39.3	8,400
35+	1,900	62.8	60.4 - 65.3	1,100	37.2	34.7 - 39.6	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,800	65.0	63.6 - 66.4	3,100	35.0	33.6 - 36.4	8,900
Female	5,100	63.0	61.4 - 64.5	3,000	37.0	35.5 - 38.6	8,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,800	61.4	60.1 - 62.7	4,300	38.6	37.3 - 39.9	11,100
No	4,100	68.9	67.2 - 70.7	1,900	31.1	29.3 - 32.8	6,000

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,400	68.2	66.4 - 69.8	2,100	31.8	30.2 - 33.6	6,500
131-185%	1,400	62.6	59.6 - 65.5	800	37.4	34.5 - 40.4	2,200
186+%	4,300	60.2	58.6 - 61.8	2,800	39.8	38.2 - 41.4	7,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	900	72.3	68.5 - 75.8	300	27.7	24.2 - 31.5	1,200
<b>Normal birth weight</b>	10,100	63.4	62.3 - 64.5	5,800	36.6	35.5 - 37.7	15,900

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	1,100	73.8	70.3 - 77.1	400	26.2	22.9 - 29.7	1,500
<b>No</b>	9,700	63.1	62.0 - 64.2	5,700	36.9	35.8 - 38.0	15,400

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	2,500	71.3	69.1 - 73.5	1,000	28.7	26.5 - 30.9	3,500
<b>No</b>	8,300	62.0	60.8 - 63.2	5,100	38.0	36.8 - 39.2	13,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,800	66.0	64.4 - 67.6	2,500	34.0	32.4 - 35.6	7,300
<b>No</b>	6,000	62.2	60.8 - 63.6	3,600	37.8	36.4 - 39.2	9,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,700	64.0	62.9 - 65.0	6,000	36.0	35.0 - 37.1	16,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,200	60.1	57.4 - 62.9	800	39.9	37.1 - 42.6	2,000
<b>Honolulu</b>	8,100	66.1	64.8 - 67.4	4,200	33.9	32.6 - 35.2	12,300
<b>Kauai</b>	400	48.3	43.8 - 52.9	400	51.7	47.1 - 56.2	700
<b>Maui</b>	1,000	59.7	56.8 - 62.6	700	40.3	37.4 - 43.2	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	48.6	46.1 - 51.1	1,900	51.4	48.9 - 53.9	3,700
Native Hawaiian	3,200	68.4	66.5 - 70.1	1,500	31.6	29.9 - 33.5	4,600
Chinese	400	68.9	64.1 - 73.4	200	31.1	26.6 - 35.9	600
Filipino	2,300	69.6	67.1 - 72.0	1,000	30.4	28.0 - 32.9	3,300
Japanese	1,500	69.4	66.2 - 72.4	700	30.6	27.6 - 33.8	2,100
Other	1,600	65.5	62.3 - 68.6	800	34.5	31.4 - 37.7	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	71.1	67.5 - 74.5	400	28.9	25.5 - 32.5	1,400
20-24	2,700	66.6	64.4 - 68.8	1,400	33.4	31.2 - 35.6	4,100
25-34	5,100	61.9	60.3 - 63.5	3,200	38.1	36.5 - 39.7	8,300
35+	1,800	62.4	59.8 - 65.0	1,100	37.6	35.0 - 40.2	2,900



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,700	65.2	63.7 - 66.7	3,000	34.8	33.3 - 36.3	8,700
Female	5,000	62.6	61.0 - 64.2	3,000	37.4	35.8 - 39.0	8,100

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,800	60.9	59.5 - 62.2	4,300	39.1	37.8 - 40.5	11,100
No	4,000	70.0	68.2 - 71.7	1,700	30.0	28.3 - 31.8	5,700

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,400	68.0	66.3 - 69.7	2,100	32.0	30.3 - 33.7	6,400
131-185%	1,500	61.4	58.3 - 64.4	900	38.6	35.6 - 41.7	2,400
186+%	4,000	59.7	57.9 - 61.4	2,700	40.3	38.6 - 42.1	6,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	800	71.7	67.6 - 75.5	300	28.3	24.5 - 32.4	1,100
<b>Normal birth weight</b>	9,900	63.4	62.3 - 64.5	5,700	36.6	35.5 - 37.7	15,600

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	1,100	74.4	70.9 - 77.7	400	25.6	22.3 - 29.1	1,400
<b>No</b>	9,500	63.0	61.9 - 64.2	5,600	37.0	35.8 - 38.1	15,100

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	2,400	70.7	68.4 - 73.0	1,000	29.3	27.0 - 31.6	3,400
<b>No</b>	8,100	62.1	60.8 - 63.3	4,900	37.9	36.7 - 39.2	13,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,800	67.1	65.4 - 68.7	2,400	32.9	31.3 - 34.6	7,200
<b>No</b>	5,800	61.4	59.9 - 62.8	3,600	38.6	37.2 - 40.1	9,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,500	63.9	62.8 - 65.0	5,900	36.1	35.0 - 37.2	16,400

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,200	60.4	57.8 - 62.9	800	39.6	37.1 - 42.2	2,000
<b>Honolulu</b>	7,900	66.0	64.7 - 67.3	4,100	34.0	32.7 - 35.3	12,000
<b>Kauai</b>	400	48.4	44.3 - 52.6	400	51.6	47.4 - 55.7	700
<b>Maui</b>	1,000	59.8	57.1 - 62.4	700	40.2	37.6 - 42.9	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,700	48.2	45.7 - 50.7	1,800	51.8	49.3 - 54.3	3,600
Native Hawaiian	3,100	68.1	66.5 - 69.7	1,400	31.9	30.3 - 33.5	4,500
Chinese	400	69.0	63.7 - 73.9	200	31.0	26.1 - 36.3	600
Filipino	2,300	69.3	66.8 - 71.7	1,000	30.7	28.3 - 33.2	3,300
Japanese	1,500	70.0	66.8 - 73.0	600	30.0	27.0 - 33.2	2,100
Other	1,500	65.3	62.0 - 68.5	800	34.7	31.5 - 38.0	2,300

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	73.3	70.0 - 76.4	400	26.7	23.6 - 30.0	1,400
20-24	2,700	66.1	64.0 - 68.2	1,400	33.9	31.8 - 36.0	4,100
25-34	5,000	61.7	60.1 - 63.3	3,100	38.3	36.7 - 39.9	8,100
35+	1,700	62.3	59.6 - 64.9	1,100	37.7	35.1 - 40.4	2,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,500	64.7	63.2 - 66.1	3,000	35.3	33.9 - 36.8	8,500
Female	5,000	63.1	61.5 - 64.7	2,900	36.9	35.3 - 38.5	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,700	60.8	59.4 - 62.1	4,300	39.2	37.9 - 40.6	11,000
No	3,800	70.3	68.6 - 71.9	1,600	29.7	28.1 - 31.4	5,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,300	67.8	66.2 - 69.5	2,100	32.2	30.5 - 33.8	6,400
131-185%	1,600	60.9	58.1 - 63.7	1,000	39.1	36.3 - 41.9	2,600
186+%	3,600	59.5	57.7 - 61.3	2,500	40.5	38.7 - 42.3	6,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	800	72.7	68.6 - 76.5	300	27.3	23.5 - 31.4	1,100
<b>Normal birth weight</b>	9,700	63.3	62.2 - 64.4	5,600	36.7	35.6 - 37.8	15,300

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	1,000	75.0	71.7 - 78.1	300	25.0	21.9 - 28.3	1,400
<b>No</b>	9,300	62.9	61.8 - 64.1	5,500	37.1	35.9 - 38.2	14,800

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	2,300	70.1	67.8 - 72.3	1,000	29.9	27.7 - 32.2	3,300
<b>No</b>	7,900	62.2	61.0 - 63.4	4,800	37.8	36.6 - 39.0	12,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,700	66.8	65.2 - 68.4	2,300	33.2	31.6 - 34.8	7,000
<b>No</b>	5,700	61.5	60.1 - 63.0	3,500	38.5	37.0 - 39.9	9,200



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,000	63.2	62.1 - 64.3	5,800	36.8	35.7 - 37.9	15,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,100	59.7	57.4 - 62.1	700	40.3	37.9 - 42.6	1,800
<b>Honolulu</b>	7,500	64.9	63.5 - 66.3	4,100	35.1	33.7 - 36.5	11,600
<b>Kauai</b>	300	48.6	44.7 - 52.5	400	51.4	47.5 - 55.3	700
<b>Maui</b>	1,000	61.2	58.7 - 63.7	600	38.8	36.3 - 41.3	1,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,700	48.9	46.4 - 51.4	1,800	51.1	48.6 - 53.6	3,500
Native Hawaiian	2,900	68.2	66.8 - 69.6	1,400	31.8	30.4 - 33.2	4,300
Chinese	400	67.3	61.3 - 72.7	200	32.7	27.3 - 38.7	600
Filipino	2,200	67.7	65.1 - 70.2	1,000	32.3	29.8 - 34.9	3,200
Japanese	1,400	68.1	64.6 - 71.3	600	31.9	28.7 - 35.4	2,000
Other	1,400	64.0	60.5 - 67.3	800	36.0	32.7 - 39.5	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	72.5	69.4 - 75.4	400	27.5	24.6 - 30.6	1,500
20-24	2,600	65.1	63.0 - 67.2	1,400	34.9	32.8 - 37.0	3,900
25-34	4,700	60.8	59.2 - 62.4	3,000	39.2	37.6 - 40.8	7,700
35+	1,700	62.2	59.4 - 64.9	1,000	37.8	35.1 - 40.6	2,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,300	63.9	62.4 - 65.4	3,000	36.1	34.6 - 37.6	8,200
Female	4,700	62.5	60.9 - 64.1	2,800	37.5	35.9 - 39.1	7,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,400	60.1	58.7 - 61.5	4,200	39.9	38.5 - 41.3	10,600
No	3,600	69.7	68.0 - 71.4	1,500	30.3	28.6 - 32.0	5,100

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,200	66.6	64.9 - 68.2	2,100	33.4	31.8 - 35.1	6,400
131-185%	1,600	60.6	57.8 - 63.4	1,000	39.4	36.6 - 42.2	2,600
186+%	3,200	58.7	56.7 - 60.6	2,300	41.3	39.4 - 43.3	5,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	700	72.2	68.0 - 76.1	300	27.8	23.9 - 32.0	1,000
<b>Normal birth weight</b>	9,200	62.6	61.5 - 63.7	5,500	37.4	36.3 - 38.5	14,700

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	1,000	76.5	73.4 - 79.5	300	23.5	20.5 - 26.6	1,300
<b>No</b>	8,800	62.1	60.9 - 63.2	5,400	37.9	36.8 - 39.1	14,100

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	2,200	69.4	67.2 - 71.6	1,000	30.6	28.4 - 32.8	3,200
<b>No</b>	7,500	61.5	60.2 - 62.7	4,700	38.5	37.3 - 39.8	12,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,500	66.4	64.8 - 68.0	2,300	33.6	32.0 - 35.2	6,800
<b>No</b>	5,300	60.7	59.2 - 62.2	3,400	39.3	37.8 - 40.8	8,800