

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

*The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.*

*Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.*

*Below are the indicators included in this report and the long description for each indicator.*

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration 8+ weeks, among initiators	Breastfeeding (any) duration 8+ weeks, among those who initiated breastfeeding

*Suggested citation:*

*Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/22/17*

*For documentation on other terms and coding in this report, please go to the following links:*

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact  
The Hawai'i Health Data Warehouse**

[profiles@hhdw.org](mailto:profiles@hhdw.org)

**(855) 946-5899 x15**

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**


*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

Years	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	3,400	23.4	22.4 - 24.4	11,200	76.6	75.6 - 77.6	14,600
2001-2005	3,500	23.3	22.3 - 24.3	11,400	76.7	75.7 - 77.7	14,800
2002-2006	3,500	23.2	22.2 - 24.2	11,800	76.8	75.8 - 77.8	15,300
2003-2007	3,700	23.4	22.4 - 24.4	12,100	76.6	75.6 - 77.6	15,700
2004-2008	3,700	22.8	21.9 - 23.8	12,500	77.2	76.2 - 78.1	16,100
2005-2009	3,800	23.1	22.0 - 24.1	12,600	76.9	75.9 - 78.0	16,300
2006-2010	3,800	22.7	21.7 - 23.8	12,900	77.3	76.2 - 78.3	16,700
2007-2011	3,700	21.7	20.6 - 22.8	13,300	78.3	77.2 - 79.4	16,900
2008-2012	3,500	20.7	19.5 - 21.8	13,500	79.3	78.2 - 80.5	17,100
2009-2013	3,400	19.7	18.6 - 21.0	13,800	80.3	79.0 - 81.4	17,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,400	19.7	18.6 - 21.0	13,800	80.3	79.0 - 81.4	17,200

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	500	20.8	18.9 - 22.9	1,700	79.2	77.1 - 81.1	2,200
<b>Honolulu</b>	2,400	19.7	18.1 - 21.3	9,900	80.3	78.7 - 81.9	12,300
<b>Kauai</b>	100	16.2	14.3 - 18.3	700	83.8	81.7 - 85.7	800
<b>Maui</b>	400	20.3	18.5 - 22.2	1,500	79.7	77.8 - 81.5	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	16.3	14.1 - 18.8	3,600	83.7	81.2 - 85.9	4,300
Native Hawaiian	1,300	25.7	23.4 - 28.1	3,700	74.3	71.9 - 76.6	5,000
Chinese	100	12.3	8.2 - 18.0	700	87.7	82.0 - 91.8	800
Filipino	700	24.6	21.7 - 27.7	2,300	75.4	72.3 - 78.3	3,000
Japanese	100	8.8	6.5 - 11.9	1,500	91.2	88.1 - 93.5	1,700
Other	400	17.3	14.3 - 20.8	2,000	82.7	79.2 - 85.7	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	39.5	34.0 - 45.2	700	60.5	54.8 - 66.0	1,100
20-24	900	23.5	20.9 - 26.3	3,000	76.5	73.7 - 79.1	3,900
25-34	1,600	17.7	16.2 - 19.4	7,500	82.3	80.6 - 83.8	9,200
35+	400	13.8	11.6 - 16.4	2,600	86.2	83.6 - 88.4	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,700	19.7	18.1 - 21.4	7,000	80.3	78.6 - 81.9	8,700
Female	1,700	19.8	18.1 - 21.5	6,800	80.2	78.5 - 81.9	8,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	16.3	14.9 - 17.8	9,100	83.7	82.2 - 85.1	10,800
No	1,600	25.5	23.5 - 27.7	4,800	74.5	72.3 - 76.5	6,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,500	25.2	23.1 - 27.4	4,500	74.8	72.6 - 76.9	6,000
131-185%	600	21.6	18.6 - 25.1	2,100	78.4	74.9 - 81.4	2,600
186+%	1,000	13.9	12.3 - 15.6	6,200	86.1	84.4 - 87.7	7,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	300	23.8	21.7 - 26.0	900	76.2	74.0 - 78.3	1,200
<b>Normal birth weight</b>	3,100	19.4	18.2 - 20.7	12,900	80.6	79.3 - 81.8	16,000

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	400	38.4	32.7 - 44.5	600	61.6	55.5 - 67.3	1,000
<b>No</b>	3,000	18.6	17.4 - 19.8	12,900	81.4	80.2 - 82.6	15,900

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	1,000	30.9	27.8 - 34.2	2,200	69.1	65.8 - 72.2	3,200
<b>No</b>	2,300	17.1	15.9 - 18.4	11,400	82.9	81.6 - 84.1	13,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,900	26.0	24.0 - 28.0	5,400	74.0	72.0 - 76.0	7,300
<b>No</b>	1,400	14.8	13.4 - 16.3	8,200	85.2	83.7 - 86.6	9,700



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,500	20.7	19.5 - 21.8	13,500	79.3	78.2 - 80.5	17,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	500	22.5	20.4 - 24.7	1,700	77.5	75.3 - 79.6	2,200
<b>Honolulu</b>	2,500	20.6	19.2 - 22.2	9,700	79.4	77.8 - 80.8	12,200
<b>Kauai</b>	100	17.7	15.1 - 20.6	700	82.3	79.4 - 84.9	800
<b>Maui</b>	400	20.0	18.1 - 22.1	1,500	80.0	77.9 - 81.9	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	17.3	15.1 - 19.6	3,400	82.7	80.4 - 84.9	4,100
Native Hawaiian	1,300	26.2	23.9 - 28.5	3,600	73.8	71.5 - 76.1	4,900
Chinese	100	11.2	7.6 - 16.2	700	88.8	83.8 - 92.4	700
Filipino	800	25.5	22.8 - 28.5	2,300	74.5	71.5 - 77.2	3,100
Japanese	200	11.3	8.8 - 14.4	1,500	88.7	85.6 - 91.2	1,700
Other	500	18.8	15.9 - 22.0	2,000	81.2	78.0 - 84.1	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	38.1	33.1 - 43.4	700	61.9	56.6 - 66.9	1,200
20-24	1,000	26.5	23.9 - 29.3	2,800	73.5	70.7 - 76.1	3,800
25-34	1,600	18.0	16.5 - 19.6	7,400	82.0	80.4 - 83.5	9,000
35+	400	14.5	12.4 - 16.9	2,600	85.5	83.1 - 87.6	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,900	21.2	19.6 - 22.9	6,900	78.8	77.1 - 80.4	8,800
Female	1,700	20.1	18.5 - 21.8	6,600	79.9	78.2 - 81.5	8,300

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	17.2	15.9 - 18.6	8,900	82.8	81.4 - 84.1	10,700
No	1,700	26.5	24.5 - 28.6	4,700	73.5	71.4 - 75.5	6,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,500	25.9	23.9 - 28.1	4,400	74.1	71.9 - 76.1	5,900
131-185%	600	21.7	18.7 - 24.9	2,100	78.3	75.1 - 81.3	2,700
186+%	1,100	15.3	13.8 - 16.9	6,200	84.7	83.1 - 86.2	7,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	300	24.1	21.6 - 26.7	900	75.9	73.3 - 78.4	1,200
<b>Normal birth weight</b>	3,200	20.4	19.2 - 21.7	12,700	79.6	78.3 - 80.8	15,900

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	400	39.1	33.8 - 44.7	600	60.9	55.3 - 66.2	1,000
<b>No</b>	3,000	19.4	18.2 - 20.6	12,700	80.6	79.4 - 81.8	15,700

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	1,000	32.2	29.2 - 35.3	2,100	67.8	64.7 - 70.8	3,100
<b>No</b>	2,400	18.0	16.8 - 19.2	11,200	82.0	80.8 - 83.2	13,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,900	26.7	24.8 - 28.6	5,300	73.3	71.4 - 75.2	7,200
<b>No</b>	1,600	16.1	14.7 - 17.6	8,100	83.9	82.4 - 85.3	9,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,700	21.7	20.6 - 22.8	13,300	78.3	77.2 - 79.4	16,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	500	21.9	19.7 - 24.3	1,700	78.1	75.7 - 80.3	2,200
<b>Honolulu</b>	2,600	22.0	20.5 - 23.4	9,400	78.0	76.6 - 79.5	12,000
<b>Kauai</b>	200	18.3	15.3 - 21.6	700	81.7	78.4 - 84.7	800
<b>Maui</b>	400	21.3	19.1 - 23.6	1,500	78.7	76.4 - 80.9	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	18.4	16.3 - 20.7	3,300	81.6	79.3 - 83.7	4,100
Native Hawaiian	1,300	27.6	25.4 - 29.9	3,500	72.4	70.1 - 74.6	4,800
Chinese	100	11.5	8.1 - 16.0	600	88.5	84.0 - 91.9	700
Filipino	800	25.7	23.1 - 28.4	2,300	74.3	71.6 - 76.9	3,000
Japanese	200	11.9	9.5 - 14.9	1,600	88.1	85.1 - 90.5	1,800
Other	500	20.7	17.8 - 23.9	2,000	79.3	76.1 - 82.2	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	38.5	33.8 - 43.4	800	61.5	56.6 - 66.2	1,300
20-24	1,100	27.8	25.3 - 30.4	2,800	72.2	69.6 - 74.7	3,900
25-34	1,600	18.7	17.3 - 20.2	7,100	81.3	79.8 - 82.7	8,700
35+	500	15.3	13.2 - 17.7	2,600	84.7	82.3 - 86.8	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,900	22.1	20.6 - 23.7	6,800	77.9	76.3 - 79.4	8,800
Female	1,700	21.2	19.7 - 22.9	6,400	78.8	77.1 - 80.3	8,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	17.7	16.4 - 19.1	8,800	82.3	80.9 - 83.6	10,700
No	1,800	28.5	26.5 - 30.5	4,500	71.5	69.5 - 73.5	6,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,600	27.7	25.8 - 29.8	4,300	72.3	70.2 - 74.2	5,900
131-185%	600	21.8	18.9 - 24.9	2,000	78.2	75.1 - 81.1	2,600
186+%	1,200	16.0	14.5 - 17.6	6,200	84.0	82.4 - 85.5	7,300



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	30.4	27.4 - 33.5	2,000	69.6	66.5 - 72.6	2,800
\$10,000-\$24,999	800	25.4	22.9 - 28.0	2,400	74.6	72.0 - 77.1	3,300
\$25,000-\$49,999	900	22.7	20.4 - 25.3	3,000	77.3	74.7 - 79.6	3,900
>=\$50,000	800	14.1	12.6 - 15.8	5,000	85.9	84.2 - 87.4	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	25.9	23.1 - 28.9	900	74.1	71.1 - 76.9	1,200
Normal birth weight	3,400	21.4	20.2 - 22.6	12,400	78.6	77.4 - 79.8	15,700

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	40.4	35.5 - 45.5	700	59.6	54.5 - 64.5	1,200
No	3,100	20.2	19.1 - 21.4	12,400	79.8	78.6 - 80.9	15,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,100	34.5	31.6 - 37.5	2,100	65.5	62.5 - 68.4	3,200
<b>No</b>	2,500	18.6	17.4 - 19.8	11,000	81.4	80.2 - 82.6	13,500

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,900	27.4	25.6 - 29.3	5,100	72.6	70.7 - 74.4	7,000
<b>No</b>	1,700	17.5	16.2 - 18.9	8,000	82.5	81.1 - 83.8	9,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,800	22.7	21.7 - 23.8	12,900	77.3	76.2 - 78.3	16,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	500	22.5	20.1 - 25.0	1,600	77.5	75.0 - 79.9	2,100
<b>Honolulu</b>	2,700	23.1	21.7 - 24.5	9,100	76.9	75.5 - 78.3	11,900
<b>Kauai</b>	200	19.4	16.0 - 23.2	700	80.6	76.8 - 84.0	800
<b>Maui</b>	400	22.0	19.6 - 24.6	1,500	78.0	75.4 - 80.4	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	19.3	17.3 - 21.5	3,200	80.7	78.5 - 82.7	4,000
Native Hawaiian	1,400	28.9	26.7 - 31.2	3,400	71.1	68.8 - 73.3	4,700
Chinese	100	12.9	9.6 - 17.1	600	87.1	82.9 - 90.4	700
Filipino	800	26.8	24.3 - 29.5	2,200	73.2	70.5 - 75.7	3,000
Japanese	200	11.5	9.3 - 14.2	1,600	88.5	85.8 - 90.7	1,800
Other	600	22.4	19.6 - 25.5	1,900	77.6	74.5 - 80.4	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	37.4	32.9 - 42.0	900	62.6	58.0 - 67.1	1,400
20-24	1,200	30.4	28.0 - 32.9	2,700	69.6	67.1 - 72.0	3,900
25-34	1,600	19.2	17.8 - 20.7	6,800	80.8	79.3 - 82.2	8,400
35+	500	16.0	13.9 - 18.3	2,500	84.0	81.7 - 86.1	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,100	23.7	22.2 - 25.3	6,600	76.3	74.7 - 77.8	8,700
Female	1,700	21.7	20.2 - 23.3	6,300	78.3	76.7 - 79.8	8,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.1	16.9 - 19.4	8,700	81.9	80.6 - 83.1	10,600
No	1,900	30.7	28.8 - 32.7	4,200	69.3	67.3 - 71.2	6,100

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	29.8	27.8 - 31.8	4,100	70.2	68.2 - 72.2	5,800
131-185%	600	22.3	19.5 - 25.4	1,900	77.7	74.6 - 80.5	2,500
186+%	1,200	16.6	15.2 - 18.2	6,100	83.4	81.8 - 84.8	7,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	33.2	30.2 - 36.4	1,800	66.8	63.6 - 69.8	2,800
\$10,000-\$24,999	900	26.6	24.1 - 29.1	2,400	73.4	70.9 - 75.9	3,300
\$25,000-\$49,999	900	23.8	21.5 - 26.2	2,900	76.2	73.8 - 78.5	3,800
>=\$50,000	800	14.3	12.8 - 15.9	4,900	85.7	84.1 - 87.2	5,700

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	24.9	21.9 - 28.2	900	75.1	71.8 - 78.1	1,200
Normal birth weight	3,500	22.6	21.4 - 23.7	12,000	77.4	76.3 - 78.6	15,500

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	39.2	34.6 - 44.0	800	60.8	56.0 - 65.4	1,200
No	3,200	21.3	20.2 - 22.5	12,000	78.7	77.5 - 79.8	15,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,100	35.3	32.5 - 38.2	2,100	64.7	61.8 - 67.5	3,200
<b>No</b>	2,600	19.7	18.5 - 20.8	10,600	80.3	79.2 - 81.5	13,300

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,000	29.4	27.5 - 31.2	4,800	70.6	68.8 - 72.5	6,800
<b>No</b>	1,700	18.0	16.7 - 19.4	7,900	82.0	80.6 - 83.3	9,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,800	23.1	22.0 - 24.1	12,600	76.9	75.9 - 78.0	16,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	500	22.1	19.6 - 24.8	1,600	77.9	75.2 - 80.4	2,100
<b>Honolulu</b>	2,700	23.3	22.1 - 24.6	9,000	76.7	75.4 - 77.9	11,700
<b>Kauai</b>	200	19.5	15.9 - 23.6	600	80.5	76.4 - 84.1	800
<b>Maui</b>	400	24.0	21.3 - 26.9	1,400	76.0	73.1 - 78.7	1,800



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	18.6	16.7 - 20.6	3,200	81.4	79.4 - 83.3	3,900
Native Hawaiian	1,400	30.1	27.9 - 32.4	3,200	69.9	67.6 - 72.1	4,500
Chinese	100	13.3	10.6 - 16.5	500	86.7	83.5 - 89.4	600
Filipino	800	27.3	25.0 - 29.8	2,200	72.7	70.2 - 75.0	3,000
Japanese	200	12.6	10.4 - 15.1	1,700	87.4	84.9 - 89.6	1,900
Other	500	22.7	20.0 - 25.6	1,800	77.3	74.4 - 80.0	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	36.8	32.6 - 41.2	900	63.2	58.8 - 67.4	1,400
20-24	1,200	32.3	29.9 - 34.7	2,600	67.7	65.3 - 70.1	3,800
25-34	1,600	19.2	17.8 - 20.5	6,600	80.8	79.5 - 82.2	8,200
35+	400	15.3	13.3 - 17.4	2,500	84.7	82.6 - 86.7	2,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	2,000	23.9	22.5 - 25.4	6,500	76.1	74.6 - 77.5	8,500
<b>Female</b>	1,700	22.1	20.7 - 23.6	6,100	77.9	76.4 - 79.3	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,900	18.1	17.0 - 19.3	8,500	81.9	80.7 - 83.0	10,400
<b>No</b>	1,900	31.6	29.7 - 33.6	4,100	68.4	66.4 - 70.3	6,000

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	1,700	30.9	29.0 - 32.9	3,900	69.1	67.1 - 71.0	5,700
<b>131-185%</b>	500	22.2	19.5 - 25.1	1,700	77.8	74.9 - 80.5	2,200
<b>186+%</b>	1,200	16.6	15.3 - 18.0	6,000	83.4	82.0 - 84.7	7,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	35.3	32.3 - 38.4	1,700	64.7	61.6 - 67.7	2,600
\$10,000-\$24,999	900	27.7	25.4 - 30.2	2,400	72.3	69.8 - 74.6	3,300
\$25,000-\$49,999	800	22.8	20.8 - 25.0	2,900	77.2	75.0 - 79.2	3,700
>=\$50,000	800	14.0	12.6 - 15.5	4,800	86.0	84.5 - 87.4	5,600

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	25.2	21.9 - 28.8	900	74.8	71.2 - 78.1	1,200
Normal birth weight	3,500	22.9	21.8 - 24.0	11,700	77.1	76.0 - 78.2	15,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	41.5	37.1 - 46.1	700	58.5	53.9 - 62.9	1,200
No	3,200	21.4	20.4 - 22.5	11,700	78.6	77.5 - 79.6	14,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,100	35.9	33.2 - 38.7	2,000	64.1	61.3 - 66.8	3,100
<b>No</b>	2,600	19.8	18.8 - 21.0	10,400	80.2	79.0 - 81.2	12,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,000	30.0	28.2 - 31.8	4,600	70.0	68.2 - 71.8	6,600
<b>No</b>	1,700	18.1	16.9 - 19.3	7,700	81.9	80.7 - 83.1	9,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,700	22.8	21.9 - 23.8	12,500	77.2	76.2 - 78.1	16,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	400	20.8	18.2 - 23.6	1,600	79.2	76.4 - 81.8	2,000
<b>Honolulu</b>	2,700	23.0	21.9 - 24.1	8,900	77.0	75.9 - 78.1	11,600
<b>Kauai</b>	200	19.7	16.0 - 24.1	600	80.3	75.9 - 84.0	800
<b>Maui</b>	400	25.4	22.5 - 28.5	1,300	74.6	71.5 - 77.5	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	18.7	17.0 - 20.5	3,100	81.3	79.5 - 83.0	3,800
Native Hawaiian	1,300	30.2	28.1 - 32.4	3,000	69.8	67.6 - 71.9	4,300
Chinese	100	13.6	11.7 - 15.8	500	86.4	84.2 - 88.3	600
Filipino	800	27.6	25.5 - 29.8	2,200	72.4	70.2 - 74.5	3,000
Japanese	200	12.1	10.2 - 14.3	1,800	87.9	85.7 - 89.8	2,000
Other	500	21.5	19.1 - 24.0	1,900	78.5	76.0 - 80.9	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	35.9	32.0 - 40.1	800	64.1	59.9 - 68.0	1,300
20-24	1,300	32.9	30.7 - 35.2	2,600	67.1	64.8 - 69.3	3,900
25-34	1,500	18.8	17.6 - 20.1	6,500	81.2	79.9 - 82.4	8,000
35+	400	14.8	13.1 - 16.6	2,500	85.2	83.4 - 86.9	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	24.0	22.7 - 25.3	6,400	76.0	74.7 - 77.3	8,400
Female	1,700	21.6	20.2 - 22.9	6,100	78.4	77.1 - 79.8	7,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.4	17.4 - 19.5	8,600	81.6	80.5 - 82.6	10,600
No	1,700	31.2	29.3 - 33.0	3,800	68.8	67.0 - 70.7	5,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	30.6	28.8 - 32.4	3,900	69.4	67.6 - 71.2	5,600
131-185%	400	20.4	18.0 - 23.0	1,700	79.6	77.0 - 82.0	2,100
186+%	1,200	16.9	15.7 - 18.2	6,000	83.1	81.8 - 84.3	7,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	35.1	32.3 - 37.9	1,700	64.9	62.1 - 67.7	2,600
\$10,000-\$24,999	900	27.7	25.5 - 29.9	2,500	72.3	70.1 - 74.5	3,400
\$25,000-\$49,999	900	22.4	20.6 - 24.4	3,000	77.6	75.6 - 79.4	3,900
>=\$50,000	700	13.5	12.3 - 14.8	4,600	86.5	85.2 - 87.7	5,300

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	24.8	21.4 - 28.6	900	75.2	71.4 - 78.6	1,200
Normal birth weight	3,400	22.7	21.7 - 23.7	11,600	77.3	76.3 - 78.3	15,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	39.0	34.9 - 43.4	700	61.0	56.6 - 65.1	1,200
No	3,200	21.4	20.5 - 22.4	11,600	78.6	77.6 - 79.5	14,700



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,000	34.2	31.7 - 36.7	2,000	65.8	63.3 - 68.3	3,000
<b>No</b>	2,600	20.0	19.1 - 21.0	10,300	80.0	79.0 - 80.9	12,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,900	28.6	27.0 - 30.3	4,700	71.4	69.7 - 73.0	6,600
<b>No</b>	1,700	18.5	17.4 - 19.7	7,500	81.5	80.3 - 82.6	9,200

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,700	23.4	22.4 - 24.4	12,100	76.6	75.6 - 77.6	15,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	400	21.0	18.6 - 23.7	1,500	79.0	76.3 - 81.4	1,900
<b>Honolulu</b>	2,700	23.5	22.3 - 24.7	8,700	76.5	75.3 - 77.7	11,400
<b>Kauai</b>	100	18.3	14.8 - 22.4	600	81.7	77.6 - 85.2	800
<b>Maui</b>	500	27.4	24.6 - 30.5	1,200	72.6	69.5 - 75.4	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	18.6	16.7 - 20.6	3,000	81.4	79.4 - 83.3	3,600
Native Hawaiian	1,300	31.3	29.3 - 33.4	2,900	68.7	66.6 - 70.7	4,200
Chinese	100	12.3	9.9 - 15.1	500	87.7	84.9 - 90.1	600
Filipino	900	28.7	26.4 - 31.1	2,100	71.3	68.9 - 73.6	3,000
Japanese	200	11.9	9.9 - 14.3	1,800	88.1	85.7 - 90.1	2,000
Other	500	22.5	19.9 - 25.3	1,800	77.5	74.7 - 80.1	2,300

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	40.9	36.8 - 45.1	800	59.1	54.9 - 63.2	1,300
20-24	1,300	33.0	30.8 - 35.3	2,500	67.0	64.7 - 69.2	3,800
25-34	1,500	19.3	18.0 - 20.6	6,300	80.7	79.4 - 82.0	7,800
35+	400	13.8	12.1 - 15.7	2,500	86.2	84.3 - 87.9	2,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	24.5	23.1 - 25.9	6,200	75.5	74.1 - 76.9	8,200
Female	1,700	22.2	20.8 - 23.6	5,800	77.8	76.4 - 79.2	7,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.2	17.1 - 19.3	8,600	81.8	80.7 - 82.9	10,500
No	1,800	33.7	31.8 - 35.7	3,500	66.3	64.3 - 68.2	5,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,800	32.0	30.2 - 33.9	3,900	68.0	66.1 - 69.8	5,700
131-185%	400	21.9	19.4 - 24.7	1,600	78.1	75.3 - 80.6	2,000
186+%	1,100	15.9	14.7 - 17.2	5,700	84.1	82.8 - 85.3	6,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	300	26.2	22.5 - 30.2	800	73.8	69.8 - 77.5	1,100
<b>Normal birth weight</b>	3,400	23.2	22.2 - 24.2	11,200	76.8	75.8 - 77.8	14,600

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	400	38.7	34.4 - 43.1	700	61.3	56.9 - 65.6	1,100
<b>No</b>	3,200	22.1	21.2 - 23.2	11,200	77.9	76.8 - 78.8	14,400

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	1,000	35.0	32.5 - 37.6	1,900	65.0	62.4 - 67.5	3,000
<b>No</b>	2,600	20.5	19.5 - 21.5	9,900	79.5	78.5 - 80.5	12,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,900	30.1	28.4 - 31.8	4,500	69.9	68.2 - 71.6	6,500
<b>No</b>	1,700	18.5	17.3 - 19.7	7,400	81.5	80.3 - 82.7	9,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,500	23.2	22.2 - 24.2	11,800	76.8	75.8 - 77.8	15,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	400	23.2	20.8 - 25.8	1,400	76.8	74.2 - 79.2	1,800
<b>Honolulu</b>	2,500	22.8	21.6 - 24.0	8,600	77.2	76.0 - 78.4	11,200
<b>Kauai</b>	100	18.9	15.4 - 22.8	600	81.1	77.2 - 84.6	700
<b>Maui</b>	400	27.6	24.9 - 30.5	1,200	72.4	69.5 - 75.1	1,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	18.5	16.5 - 20.6	2,900	81.5	79.4 - 83.5	3,500
Native Hawaiian	1,200	30.2	28.3 - 32.2	2,800	69.8	67.8 - 71.7	4,000
Chinese	100	14.0	10.7 - 18.0	500	86.0	82.0 - 89.3	600
Filipino	900	28.8	26.4 - 31.4	2,100	71.2	68.6 - 73.6	3,000
Japanese	200	11.3	9.3 - 13.7	1,800	88.7	86.3 - 90.7	2,100
Other	500	23.7	20.8 - 26.9	1,600	76.3	73.1 - 79.2	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	41.1	37.0 - 45.3	700	58.9	54.7 - 63.0	1,200
20-24	1,200	32.9	30.7 - 35.3	2,500	67.1	64.7 - 69.3	3,700
25-34	1,400	18.9	17.6 - 20.2	6,200	81.1	79.8 - 82.4	7,600
35+	400	13.8	12.0 - 15.8	2,400	86.2	84.2 - 88.0	2,700



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,900	24.4	23.0 - 25.9	6,000	75.6	74.1 - 77.0	7,900
Female	1,600	21.9	20.5 - 23.3	5,800	78.1	76.7 - 79.5	7,400

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.3	17.2 - 19.5	8,500	81.7	80.5 - 82.8	10,400
No	1,700	33.4	31.4 - 35.3	3,300	66.6	64.7 - 68.6	5,000

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	31.2	29.4 - 33.0	3,800	68.8	67.0 - 70.6	5,600
131-185%	500	22.4	19.8 - 25.3	1,700	77.6	74.7 - 80.2	2,200
186+%	1,000	15.5	14.2 - 16.9	5,400	84.5	83.1 - 85.8	6,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	300	26.4	22.4 - 30.8	800	73.6	69.2 - 77.6	1,100
<b>Normal birth weight</b>	3,300	22.9	21.9 - 24.0	11,000	77.1	76.0 - 78.1	14,200

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	400	36.8	32.6 - 41.3	700	63.2	58.7 - 67.4	1,000
<b>No</b>	3,100	22.1	21.0 - 23.1	10,900	77.9	76.9 - 79.0	14,000

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	1,000	33.6	31.1 - 36.3	1,900	66.4	63.7 - 68.9	2,900
<b>No</b>	2,500	20.5	19.4 - 21.6	9,700	79.5	78.4 - 80.6	12,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,900	30.3	28.6 - 32.1	4,400	69.7	67.9 - 71.4	6,300
<b>No</b>	1,600	18.0	16.8 - 19.3	7,200	82.0	80.7 - 83.2	8,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,500	23.3	22.3 - 24.3	11,400	76.7	75.7 - 77.7	14,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	400	22.8	20.6 - 25.2	1,300	77.2	74.8 - 79.4	1,700
<b>Honolulu</b>	2,500	22.9	21.7 - 24.2	8,400	77.1	75.8 - 78.3	10,900
<b>Kauai</b>	100	19.6	16.5 - 23.2	500	80.4	76.8 - 83.5	700
<b>Maui</b>	400	28.0	25.5 - 30.6	1,100	72.0	69.4 - 74.5	1,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	17.5	15.5 - 19.6	2,800	82.5	80.4 - 84.5	3,400
Native Hawaiian	1,200	30.3	28.6 - 32.0	2,700	69.7	68.0 - 71.4	3,900
Chinese	100	15.7	11.8 - 20.5	500	84.3	79.5 - 88.2	600
Filipino	900	29.5	27.0 - 32.0	2,100	70.5	68.0 - 73.0	3,000
Japanese	300	12.5	10.4 - 15.0	1,800	87.5	85.0 - 89.6	2,100
Other	500	23.9	20.8 - 27.1	1,500	76.1	72.9 - 79.2	2,000

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	42.5	38.7 - 46.5	700	57.5	53.5 - 61.3	1,200
20-24	1,200	32.7	30.5 - 35.0	2,400	67.3	65.0 - 69.5	3,600
25-34	1,400	18.7	17.4 - 20.0	6,000	81.3	80.0 - 82.6	7,400
35+	400	14.1	12.3 - 16.2	2,200	85.9	83.8 - 87.7	2,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	1,800	23.8	22.4 - 25.3	5,900	76.2	74.7 - 77.6	7,700
<b>Female</b>	1,600	22.7	21.4 - 24.2	5,500	77.3	75.8 - 78.6	7,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,900	18.7	17.6 - 19.8	8,300	81.3	80.2 - 82.4	10,200
<b>No</b>	1,600	33.4	31.6 - 35.4	3,100	66.6	64.6 - 68.4	4,700

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	1,700	30.4	28.7 - 32.2	3,800	69.6	67.8 - 71.3	5,500
<b>131-185%</b>	500	22.3	20.0 - 24.9	1,800	77.7	75.1 - 80.0	2,400
<b>186+%</b>	900	15.2	13.9 - 16.6	5,000	84.8	83.4 - 86.1	5,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	300	28.1	24.0 - 32.6	700	71.9	67.4 - 76.0	1,000
<b>Normal birth weight</b>	3,200	23.0	22.0 - 24.0	10,700	77.0	76.0 - 78.0	13,900

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	400	36.9	32.8 - 41.2	600	63.1	58.8 - 67.2	1,000
<b>No</b>	3,000	22.2	21.2 - 23.2	10,600	77.8	76.8 - 78.8	13,600

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	900	34.2	31.7 - 36.8	1,800	65.8	63.2 - 68.3	2,800
<b>No</b>	2,400	20.5	19.4 - 21.6	9,300	79.5	78.4 - 80.6	11,700

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,800	30.4	28.7 - 32.1	4,200	69.6	67.9 - 71.3	6,100
<b>No</b>	1,600	18.3	17.1 - 19.5	7,000	81.7	80.5 - 82.9	8,600



**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	3,400	23.4	22.4 - 24.4	11,200	76.6	75.6 - 77.6	14,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	400	25.2	23.1 - 27.4	1,200	74.8	72.6 - 76.9	1,700
<b>Honolulu</b>	2,400	22.8	21.5 - 24.0	8,300	77.2	76.0 - 78.5	10,700
<b>Kauai</b>	100	21.0	17.9 - 24.4	500	79.0	75.6 - 82.1	700
<b>Maui</b>	400	26.8	24.6 - 29.2	1,100	73.2	70.8 - 75.4	1,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	18.3	16.3 - 20.5	2,700	81.7	79.5 - 83.7	3,300
Native Hawaiian	1,200	30.8	29.3 - 32.3	2,600	69.2	67.7 - 70.7	3,800
Chinese	100	16.8	12.5 - 22.1	500	83.2	77.9 - 87.5	600
Filipino	800	28.3	25.9 - 30.9	2,100	71.7	69.1 - 74.1	3,000
Japanese	200	12.0	9.8 - 14.5	1,800	88.0	85.5 - 90.2	2,000
Other	500	23.9	20.8 - 27.4	1,500	76.1	72.6 - 79.2	1,900

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	43.3	39.7 - 47.0	700	56.7	53.0 - 60.3	1,300
20-24	1,100	31.7	29.5 - 33.9	2,400	68.3	66.1 - 70.5	3,500
25-34	1,400	18.9	17.6 - 20.2	5,900	81.1	79.8 - 82.4	7,200
35+	400	14.7	12.7 - 16.8	2,200	85.3	83.2 - 87.3	2,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,800	23.9	22.5 - 25.3	5,700	76.1	74.7 - 77.5	7,600
Female	1,600	22.9	21.5 - 24.3	5,400	77.1	75.7 - 78.5	7,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.8	17.7 - 20.0	8,200	81.2	80.0 - 82.3	10,100
No	1,500	33.8	32.0 - 35.7	2,900	66.2	64.3 - 68.0	4,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	30.1	28.5 - 31.9	3,900	69.9	68.1 - 71.5	5,600
131-185%	500	21.4	19.2 - 23.9	1,900	78.6	76.1 - 80.8	2,400
186+%	900	15.6	14.2 - 17.0	4,700	84.4	83.0 - 85.8	5,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Birthweight</b>							
<b>Low birth weight</b>	300	28.9	24.7 - 33.5	700	71.1	66.5 - 75.3	900
<b>Normal birth weight</b>	3,100	23.0	22.0 - 24.0	10,500	77.0	76.0 - 78.0	13,600

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke 3rd Trimester</b>							
<b>Yes</b>	400	36.4	32.4 - 40.6	600	63.6	59.4 - 67.6	1,000
<b>No</b>	3,000	22.3	21.3 - 23.3	10,400	77.7	76.7 - 78.7	13,400

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Smoke Before Pregnant</b>							
<b>Yes</b>	900	33.6	31.2 - 36.2	1,800	66.4	63.8 - 68.8	2,700
<b>No</b>	2,400	20.7	19.6 - 21.8	9,100	79.3	78.2 - 80.4	11,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,800	30.5	28.9 - 32.1	4,200	69.5	67.9 - 71.1	6,000
<b>No</b>	1,600	18.4	17.2 - 19.7	6,900	81.6	80.3 - 82.8	8,400