

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (exclusive) duration (grouped), among initiators	Breastfeeding (exclusive) duration (grouped), among those who initiated breastfeeding

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 9/29/15

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among
 initiators- for the State of Hawaii, for the Year(s)
 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**


Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	4,700	33.4	32.2 - 34.5	3,100	21.8	20.9 - 22.9	700	4.7	4.2 - 5.2	5,600	40.1	38.9 - 41.3	14,100
2001-2005	5,200	35.4	34.2 - 36.5	3,100	21.2	20.2 - 22.2	600	4.1	3.7 - 4.6	5,800	39.3	38.2 - 40.5	14,700
2002-2006	5,400	35.6	34.5 - 36.8	3,200	21.1	20.1 - 22.1	600	4.2	3.7 - 4.7	5,900	39.1	38.0 - 40.3	15,100
2003-2007	5,700	36.4	35.3 - 37.5	3,200	20.8	19.9 - 21.8	600	3.9	3.4 - 4.3	6,100	38.9	37.8 - 40.1	15,500
2004-2008	5,900	37.2	36.1 - 38.2	3,200	20.2	19.3 - 21.1	600	3.8	3.4 - 4.2	6,200	38.9	37.8 - 40.0	16,000
2005-2009	6,000	37.2	36.0 - 38.4	3,200	19.6	18.7 - 20.6	700	4.1	3.6 - 4.6	6,300	39.1	37.9 - 40.3	16,100
2006-2010	5,900	35.7	34.5 - 36.9	3,200	19.2	18.2 - 20.2	700	4.4	3.9 - 4.9	6,700	40.8	39.5 - 42.0	16,500
2007-2011	5,900	35.1	33.8 - 36.4	3,100	18.5	17.5 - 19.6	700	4.2	3.7 - 4.8	7,100	42.2	40.8 - 43.5	16,700
2008-2012	5,800	34.5	33.1 - 35.8	3,000	18.1	17.0 - 19.2	700	4.3	3.7 - 4.9	7,300	43.2	41.8 - 44.6	16,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,800	34.5	33.1 - 35.8	3,000	18.1	17.0 - 19.2	700	4.3	3.7 - 4.9	7,300	43.2	41.8 - 44.6	16,800

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	32.9	30.5 - 35.4	400	18.0	16.0 - 20.0	100	4.4	3.5 - 5.5	1,000	44.7	42.2 - 47.3	2,100
Honolulu	4,200	35.2	33.4 - 37.0	2,200	18.1	16.7 - 19.7	500	4.3	3.6 - 5.2	5,100	42.3	40.5 - 44.2	12,000
Kauai	200	27.3	24.3 - 30.5	100	18.9	16.2 - 21.9	<50	4.5	3.3 - 6.1	400	49.3	45.9 - 52.8	800
Maui	600	34.3	32.0 - 36.7	300	17.6	15.8 - 19.6	100	3.7	2.8 - 4.8	800	44.4	42.0 - 46.9	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	20.7	18.5 - 23.1	700	18.2	16.0 - 20.6	200	5.0	3.9 - 6.5	2,300	56.0	53.2 - 58.9	4,100
Native Hawaiian	1,700	35.2	32.7 - 37.7	1,000	20.6	18.5 - 22.8	300	5.6	4.5 - 7.0	1,900	38.7	36.1 - 41.3	4,800
Chinese	300	38.8	32.2 - 45.7	100	13.0	9.4 - 17.8	<50	3.4	1.6 - 7.3	300	44.8	38.0 - 51.8	700
Filipino	1,300	44.7	41.4 - 48.1	600	20.0	17.5 - 22.9	100	3.5	2.4 - 5.0	1,000	31.8	28.7 - 35.0	3,000
Japanese	800	45.9	41.4 - 50.4	200	13.6	10.7 - 17.0	<50	1.7	1.0 - 3.0	700	38.8	34.5 - 43.3	1,700
Other	800	34.1	30.5 - 37.9	400	15.4	12.7 - 18.5	100	3.4	2.2 - 5.2	1,200	47.1	43.2 - 51.1	2,500

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	38.0	33.0 - 43.3	300	22.5	18.2 - 27.5	<50	3.3	1.9 - 5.5	400	36.2	31.3 - 41.4	1,200
20-24	1,200	32.8	30.0 - 35.7	800	21.6	19.1 - 24.4	200	5.5	4.2 - 7.1	1,500	40.1	37.1 - 43.2	3,800
25-34	2,900	32.4	30.6 - 34.3	1,500	17.2	15.7 - 18.7	400	4.3	3.6 - 5.2	4,100	46.1	44.1 - 48.0	8,900
35+	1,200	41.1	37.9 - 44.4	400	14.7	12.6 - 17.2	100	3.0	2.1 - 4.3	1,200	41.2	37.9 - 44.5	3,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,000	34.1	32.3 - 36.0	1,600	18.7	17.2 - 20.4	400	4.2	3.5 - 5.1	3,700	42.9	40.9 - 44.9	8,600
Female	2,800	34.8	32.9 - 36.8	1,400	17.4	15.9 - 19.1	400	4.3	3.5 - 5.2	3,600	43.5	41.5 - 45.5	8,200

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,400	32.6	31.0 - 34.3	1,800	16.9	15.6 - 18.3	400	4.1	3.4 - 4.9	4,900	46.4	44.6 - 48.2	10,600
No	2,400	37.5	35.3 - 39.8	1,300	20.1	18.3 - 22.1	300	4.6	3.7 - 5.6	2,400	37.8	35.6 - 40.0	6,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,200	37.8	35.5 - 40.2	1,100	19.0	17.2 - 21.0	200	3.6	2.8 - 4.6	2,300	39.6	37.2 - 41.9	5,800
131-185%	800	31.7	28.4 - 35.2	500	20.2	17.4 - 23.4	200	6.2	4.6 - 8.3	1,100	41.9	38.3 - 45.6	2,600
186+%	2,400	32.9	30.9 - 35.0	1,200	16.2	14.7 - 17.9	300	4.1	3.3 - 5.1	3,400	46.7	44.6 - 48.9	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	47.5	44.5 - 50.5	200	16.5	14.3 - 18.9	<50	3.9	2.9 - 5.2	400	32.2	29.5 - 35.0	1,100
Normal birth weight	5,300	33.5	32.1 - 35.0	2,900	18.2	17.0 - 19.4	700	4.3	3.7 - 5.0	6,900	44.0	42.5 - 45.5	15,700

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	41.1	35.7 - 46.8	200	20.7	16.6 - 25.5	100	5.3	3.3 - 8.2	300	32.9	27.8 - 38.4	1,000
No	5,300	34.1	32.7 - 35.5	2,800	17.9	16.8 - 19.1	700	4.3	3.7 - 4.9	6,800	43.7	42.3 - 45.2	15,500

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	36.5	33.4 - 39.8	600	19.7	17.2 - 22.4	200	5.5	4.2 - 7.2	1,200	38.2	35.1 - 41.5	3,100
No	4,600	34.0	32.6 - 35.6	2,400	17.7	16.5 - 19.0	500	4.1	3.5 - 4.8	5,900	44.2	42.6 - 45.7	13,400

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,600	37.1	35.0 - 39.2	1,300	18.8	17.1 - 20.5	300	4.5	3.7 - 5.5	2,800	39.7	37.6 - 41.8	7,100
No	3,100	32.7	30.9 - 34.5	1,700	17.6	16.1 - 19.1	400	4.1	3.4 - 5.0	4,300	45.6	43.7 - 47.5	9,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,900	35.1	33.8 - 36.4	3,100	18.5	17.5 - 19.6	700	4.2	3.7 - 4.8	7,100	42.2	40.8 - 43.5	16,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	32.8	30.3 - 35.4	400	17.1	15.2 - 19.3	100	4.0	3.2 - 5.2	1,000	46.0	43.3 - 48.8	2,100
Honolulu	4,300	36.2	34.5 - 37.9	2,200	18.8	17.4 - 20.2	500	4.3	3.6 - 5.1	4,800	40.8	39.0 - 42.5	11,900
Kauai	200	24.3	21.0 - 27.8	200	22.0	18.8 - 25.6	<50	5.2	3.8 - 7.2	400	48.5	44.6 - 52.4	800
Maui	700	35.3	32.7 - 38.0	300	17.1	15.2 - 19.3	100	3.8	2.9 - 5.0	800	43.8	41.1 - 46.5	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	21.5	19.4 - 23.8	700	17.9	15.8 - 20.2	200	5.6	4.4 - 7.0	2,200	55.0	52.3 - 57.7	4,000
Native Hawaiian	1,700	35.8	33.4 - 38.2	1,000	21.0	19.0 - 23.2	300	5.2	4.2 - 6.5	1,800	38.0	35.6 - 40.5	4,800
Chinese	300	37.7	31.7 - 44.0	100	15.8	11.9 - 20.6	<50	3.3	1.6 - 6.5	300	43.3	36.9 - 49.9	700
Filipino	1,400	46.0	42.9 - 49.1	600	19.8	17.4 - 22.5	100	3.1	2.2 - 4.4	900	31.0	28.2 - 34.0	3,000
Japanese	900	47.7	43.7 - 51.8	300	14.3	11.6 - 17.5	<50	2.0	1.2 - 3.3	600	35.9	32.1 - 40.0	1,800
Other	800	32.8	29.5 - 36.3	400	16.9	14.2 - 20.0	100	3.4	2.2 - 5.1	1,100	46.9	43.2 - 50.6	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	35.1	30.5 - 39.9	300	24.4	20.3 - 29.0	<50	3.3	2.0 - 5.5	500	37.2	32.5 - 42.1	1,300
20-24	1,300	33.7	31.0 - 36.4	900	22.3	19.9 - 24.8	200	4.8	3.7 - 6.3	1,500	39.2	36.5 - 42.1	3,900
25-34	2,900	33.8	32.1 - 35.6	1,500	16.9	15.6 - 18.4	400	4.4	3.6 - 5.2	3,900	44.9	43.0 - 46.8	8,600
35+	1,200	40.7	37.6 - 43.8	500	15.8	13.6 - 18.1	100	3.5	2.5 - 4.9	1,200	40.1	37.0 - 43.2	3,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,000	34.5	32.8 - 36.3	1,700	19.2	17.8 - 20.8	400	4.2	3.5 - 5.0	3,600	42.1	40.2 - 43.9	8,700
Female	2,900	35.7	33.8 - 37.5	1,400	17.8	16.3 - 19.3	300	4.3	3.5 - 5.2	3,400	42.3	40.4 - 44.2	8,100

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,500	33.6	32.1 - 35.2	1,800	17.0	15.7 - 18.3	400	4.0	3.4 - 4.7	4,800	45.4	43.7 - 47.1	10,500
No	2,300	37.6	35.4 - 39.7	1,300	21.1	19.3 - 23.1	300	4.7	3.8 - 5.7	2,300	36.7	34.6 - 38.8	6,200

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,200	37.6	35.4 - 39.8	1,200	20.1	18.3 - 22.1	200	3.6	2.8 - 4.5	2,200	38.7	36.5 - 41.0	5,800
131-185%	800	32.3	29.1 - 35.7	500	19.5	16.9 - 22.5	100	5.2	3.8 - 7.1	1,100	42.9	39.4 - 46.4	2,500
186+%	2,500	34.2	32.3 - 36.1	1,200	16.6	15.1 - 18.2	300	4.5	3.7 - 5.5	3,200	44.7	42.7 - 46.7	7,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,100	37.8	34.6 - 41.1	600	21.6	18.8 - 24.5	100	3.0	2.0 - 4.3	1,000	37.7	34.5 - 41.0	2,800
\$10,000-\$24,999	1,200	36.2	33.4 - 39.2	700	20.5	18.2 - 23.1	100	4.4	3.4 - 5.8	1,300	38.8	35.9 - 41.8	3,200
\$25,000-\$49,999	1,300	32.9	30.3 - 35.6	700	18.7	16.6 - 21.0	200	4.8	3.7 - 6.2	1,700	43.6	40.8 - 46.4	3,900
>=\$50,000	2,000	34.9	32.8 - 37.1	900	15.6	14.0 - 17.4	300	4.5	3.5 - 5.6	2,600	45.0	42.7 - 47.3	5,800

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	47.1	43.8 - 50.4	200	17.7	15.2 - 20.5	100	4.6	3.4 - 6.2	400	30.6	27.7 - 33.7	1,200
Normal birth weight	5,300	34.2	32.9 - 35.5	2,900	18.6	17.5 - 19.7	700	4.2	3.7 - 4.8	6,700	43.0	41.6 - 44.4	15,600

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	42.4	37.4 - 47.6	200	19.1	15.6 - 23.1	100	5.9	3.9 - 8.7	400	32.6	28.0 - 37.6	1,200
No	5,300	34.6	33.3 - 35.9	2,800	18.5	17.4 - 19.6	600	4.2	3.6 - 4.8	6,600	42.8	41.4 - 44.2	15,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	38.0	35.1 - 41.1	700	20.8	18.4 - 23.4	200	5.5	4.2 - 7.1	1,100	35.7	32.7 - 38.7	3,200
No	4,600	34.5	33.0 - 35.9	2,400	18.0	16.8 - 19.2	500	4.0	3.4 - 4.7	5,800	43.6	42.1 - 45.1	13,300

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	36.3	34.2 - 38.3	1,300	19.4	17.7 - 21.1	300	4.3	3.6 - 5.3	2,800	40.0	38.0 - 42.1	6,900
No	3,300	34.3	32.6 - 36.0	1,700	17.9	16.6 - 19.4	400	4.2	3.5 - 5.0	4,200	43.6	41.8 - 45.3	9,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,900	35.7	34.5 - 36.9	3,200	19.2	18.2 - 20.2	700	4.4	3.9 - 4.9	6,700	40.8	39.5 - 42.0	16,500

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	31.8	29.1 - 34.6	400	18.0	15.8 - 20.4	100	4.2	3.3 - 5.5	1,000	45.9	43.0 - 48.9	2,100
Honolulu	4,400	37.5	35.9 - 39.0	2,200	19.1	17.8 - 20.4	500	4.4	3.7 - 5.1	4,600	39.1	37.5 - 40.7	11,700
Kauai	200	21.9	18.5 - 25.7	200	23.3	19.6 - 27.4	<50	5.9	4.2 - 8.3	400	48.9	44.5 - 53.3	800
Maui	600	34.5	31.7 - 37.4	400	19.4	17.1 - 21.9	100	3.9	2.9 - 5.3	800	42.2	39.3 - 45.2	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	22.1	20.1 - 24.3	700	17.3	15.4 - 19.5	300	6.5	5.2 - 7.9	2,100	54.1	51.5 - 56.6	4,000
Native Hawaiian	1,700	36.1	33.7 - 38.5	1,000	22.1	20.1 - 24.2	200	4.7	3.7 - 5.9	1,700	37.2	34.8 - 39.7	4,700
Chinese	200	38.3	33.0 - 43.9	100	17.7	13.9 - 22.2	<50	3.6	1.9 - 6.9	300	40.4	34.7 - 46.3	600
Filipino	1,400	45.8	42.9 - 48.8	600	20.0	17.7 - 22.5	100	3.1	2.3 - 4.3	900	31.1	28.4 - 33.9	3,000
Japanese	900	49.1	45.3 - 52.9	300	15.0	12.6 - 17.9	<50	2.1	1.4 - 3.3	600	33.7	30.2 - 37.4	1,800
Other	800	33.7	30.6 - 37.1	500	19.2	16.5 - 22.2	100	3.8	2.6 - 5.5	1,100	43.3	39.9 - 46.7	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	34.4	30.1 - 39.0	300	25.5	21.6 - 29.9	<50	2.7	1.6 - 4.8	500	37.3	32.8 - 41.9	1,400
20-24	1,300	34.3	31.8 - 37.0	900	22.2	20.0 - 24.7	200	4.9	3.8 - 6.2	1,500	38.5	35.9 - 41.2	3,800
25-34	2,900	34.4	32.8 - 36.1	1,500	17.7	16.4 - 19.1	400	4.6	3.9 - 5.4	3,600	43.3	41.5 - 45.1	8,300
35+	1,200	41.4	38.6 - 44.4	500	16.5	14.4 - 18.7	100	3.8	2.9 - 5.1	1,100	38.2	35.4 - 41.1	3,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,000	35.6	33.9 - 37.3	1,700	20.1	18.7 - 21.6	400	4.6	3.9 - 5.4	3,400	39.8	38.0 - 41.5	8,600
Female	2,800	35.8	34.0 - 37.6	1,400	18.2	16.8 - 19.7	300	4.1	3.5 - 5.0	3,300	41.9	40.1 - 43.7	7,900

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,600	34.8	33.3 - 36.3	1,800	17.3	16.1 - 18.6	400	4.3	3.6 - 5.0	4,500	43.6	42.0 - 45.2	10,400
No	2,300	37.1	35.1 - 39.2	1,400	22.4	20.6 - 24.2	300	4.6	3.7 - 5.6	2,200	35.9	33.9 - 38.0	6,100

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,200	37.7	35.6 - 39.9	1,200	20.8	19.0 - 22.7	200	3.7	3.0 - 4.6	2,200	37.8	35.6 - 39.9	5,700
131-185%	800	33.5	30.4 - 36.8	500	20.0	17.4 - 22.8	100	5.5	4.1 - 7.4	1,000	41.0	37.7 - 44.4	2,400
186+%	2,500	34.9	33.1 - 36.7	1,200	17.3	15.9 - 18.8	300	4.6	3.8 - 5.5	3,100	43.3	41.4 - 45.2	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,000	37.0	33.9 - 40.3	600	21.9	19.2 - 24.8	100	3.2	2.2 - 4.6	1,000	37.9	34.8 - 41.1	2,700
\$10,000-\$24,999	1,200	38.0	35.2 - 40.8	700	20.6	18.4 - 23.0	100	4.4	3.4 - 5.8	1,200	37.0	34.2 - 39.8	3,300
\$25,000-\$49,999	1,300	33.9	31.4 - 36.4	700	19.6	17.5 - 21.8	200	5.3	4.2 - 6.7	1,600	41.3	38.7 - 43.9	3,800
>=\$50,000	2,000	35.2	33.2 - 37.2	900	16.5	14.9 - 18.2	200	4.3	3.5 - 5.4	2,500	44.0	41.9 - 46.1	5,700

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	45.8	42.1 - 49.4	200	18.6	15.8 - 21.8	100	4.3	3.1 - 5.9	400	31.3	28.0 - 34.8	1,200
Normal birth weight	5,300	34.9	33.6 - 36.2	2,900	19.2	18.2 - 20.3	700	4.4	3.8 - 5.0	6,400	41.5	40.2 - 42.8	15,300

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	41.3	36.6 - 46.2	200	20.1	16.6 - 24.1	100	4.9	3.4 - 7.1	400	33.7	29.2 - 38.5	1,200
No	5,300	35.2	33.9 - 36.5	2,900	19.2	18.1 - 20.3	700	4.4	3.8 - 5.0	6,200	41.3	39.9 - 42.6	15,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	37.3	34.5 - 40.2	700	22.4	20.0 - 24.9	200	5.7	4.5 - 7.2	1,100	34.6	31.8 - 37.5	3,200
No	4,600	35.3	33.9 - 36.6	2,400	18.5	17.4 - 19.6	500	4.1	3.5 - 4.8	5,500	42.2	40.7 - 43.6	13,000

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	36.7	34.8 - 38.7	1,300	19.9	18.3 - 21.6	300	4.6	3.8 - 5.5	2,600	38.8	36.9 - 40.8	6,700
No	3,300	34.9	33.4 - 36.5	1,800	18.7	17.3 - 20.0	400	4.3	3.6 - 5.1	4,000	42.1	40.5 - 43.8	9,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	6,000	37.2	36.0 - 38.4	3,200	19.6	18.7 - 20.6	700	4.1	3.6 - 4.6	6,300	39.1	37.9 - 40.3	16,100

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	31.7	28.8 - 34.6	400	19.6	17.2 - 22.3	100	3.7	2.7 - 4.9	900	45.0	41.9 - 48.1	2,000
Honolulu	4,600	39.6	38.2 - 41.1	2,200	19.4	18.3 - 20.6	500	4.0	3.4 - 4.6	4,300	37.0	35.6 - 38.4	11,500
Kauai	200	20.1	16.6 - 24.2	200	24.7	20.8 - 29.2	<50	5.8	3.9 - 8.4	400	49.4	44.5 - 54.2	800
Maui	600	35.4	32.4 - 38.5	300	18.7	16.2 - 21.3	100	4.3	3.2 - 5.9	700	41.6	38.5 - 44.8	1,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	23.7	21.6 - 25.8	700	17.3	15.5 - 19.2	200	6.0	4.9 - 7.4	2,000	53.0	50.6 - 55.5	3,800
Native Hawaiian	1,700	37.3	35.0 - 39.7	1,000	22.9	20.9 - 25.0	200	4.2	3.3 - 5.3	1,600	35.7	33.4 - 38.1	4,500
Chinese	300	42.9	38.3 - 47.8	100	19.3	16.0 - 23.1	<50	2.7	1.8 - 3.9	200	35.1	30.6 - 39.8	600
Filipino	1,400	46.6	43.9 - 49.3	600	19.9	17.8 - 22.1	100	3.1	2.4 - 4.0	900	30.5	28.0 - 33.1	2,900
Japanese	1,000	50.8	47.4 - 54.2	300	15.3	13.1 - 18.0	<50	2.4	1.6 - 3.6	600	31.5	28.4 - 34.7	1,900
Other	800	34.7	31.7 - 37.9	500	20.5	18.0 - 23.2	100	3.7	2.5 - 5.3	1,000	41.1	38.0 - 44.3	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	35.5	31.3 - 39.8	300	24.9	21.3 - 29.0	<50	2.8	1.7 - 4.7	500	36.8	32.5 - 41.2	1,400
20-24	1,300	35.6	33.2 - 38.1	900	23.5	21.4 - 25.8	200	4.9	3.8 - 6.1	1,400	36.0	33.6 - 38.5	3,800
25-34	2,900	36.3	34.7 - 37.9	1,400	17.8	16.6 - 19.2	300	4.2	3.6 - 5.0	3,400	41.6	40.0 - 43.3	8,100
35+	1,200	42.7	40.0 - 45.5	500	16.9	15.0 - 19.0	100	3.2	2.3 - 4.4	1,100	37.2	34.6 - 39.9	2,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,100	36.9	35.3 - 38.5	1,700	20.8	19.5 - 22.2	300	4.0	3.4 - 4.7	3,200	38.2	36.6 - 39.9	8,400
Female	2,900	37.5	35.9 - 39.2	1,400	18.3	17.0 - 19.7	300	4.1	3.5 - 4.9	3,100	40.1	38.4 - 41.8	7,700

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,700	36.4	35.0 - 37.8	1,800	17.5	16.4 - 18.7	400	4.1	3.5 - 4.7	4,300	42.0	40.5 - 43.5	10,200
No	2,300	38.6	36.6 - 40.6	1,400	23.2	21.5 - 25.1	200	4.1	3.4 - 5.0	2,000	34.1	32.2 - 36.1	5,900

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,200	39.1	37.0 - 41.2	1,200	21.2	19.6 - 23.0	200	3.5	2.8 - 4.4	2,000	36.1	34.1 - 38.2	5,600
131-185%	700	33.7	30.6 - 36.9	500	21.3	18.7 - 24.2	100	5.0	3.6 - 6.9	900	40.0	36.8 - 43.3	2,200
186+%	2,700	37.0	35.3 - 38.7	1,200	17.2	15.9 - 18.6	300	4.3	3.6 - 5.1	3,000	41.5	39.8 - 43.2	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,000	39.2	36.1 - 42.3	600	21.4	19.0 - 24.1	100	3.2	2.3 - 4.5	900	36.1	33.2 - 39.2	2,600
\$10,000-\$24,999	1,300	38.9	36.3 - 41.6	700	21.8	19.7 - 24.1	100	4.0	3.0 - 5.2	1,200	35.3	32.8 - 38.0	3,300
\$25,000-\$49,999	1,300	34.6	32.2 - 37.0	700	19.6	17.7 - 21.7	200	5.1	4.1 - 6.4	1,500	40.7	38.3 - 43.2	3,700
>=\$50,000	2,100	37.4	35.5 - 39.4	900	16.8	15.4 - 18.4	200	3.9	3.2 - 4.8	2,300	41.8	39.8 - 43.8	5,500

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	47.7	43.8 - 51.6	200	18.1	15.2 - 21.4	<50	4.1	2.9 - 5.9	400	30.1	26.6 - 33.8	1,200
Normal birth weight	5,400	36.4	35.2 - 37.6	3,000	19.7	18.7 - 20.8	600	4.1	3.6 - 4.6	6,000	39.8	38.6 - 41.1	15,000

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	43.4	38.9 - 48.0	300	20.4	17.0 - 24.3	100	4.8	3.2 - 7.0	400	31.4	27.3 - 35.8	1,200
No	5,400	36.6	35.4 - 37.8	2,900	19.5	18.6 - 20.6	600	4.1	3.6 - 4.6	5,800	39.8	38.6 - 41.0	14,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	38.3	35.6 - 41.1	700	22.8	20.5 - 25.2	200	6.0	4.8 - 7.5	1,000	32.9	30.3 - 35.6	3,100
No	4,700	36.8	35.6 - 38.1	2,400	18.9	17.8 - 20.0	500	3.7	3.2 - 4.2	5,200	40.6	39.3 - 42.0	12,800

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	37.2	35.3 - 39.1	1,400	21.1	19.5 - 22.7	300	4.2	3.5 - 5.0	2,400	37.6	35.7 - 39.5	6,500
No	3,500	37.1	35.6 - 38.6	1,700	18.5	17.3 - 19.7	400	4.1	3.5 - 4.8	3,800	40.3	38.8 - 41.9	9,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,900	37.2	36.1 - 38.2	3,200	20.2	19.3 - 21.1	600	3.8	3.4 - 4.2	6,200	38.9	37.8 - 40.0	16,000

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	31.0	28.0 - 34.1	400	19.8	17.3 - 22.6	100	3.0	2.1 - 4.3	900	46.2	43.0 - 49.5	2,000
Honolulu	4,600	39.9	38.7 - 41.2	2,300	19.9	18.9 - 21.0	400	3.6	3.2 - 4.2	4,200	36.5	35.3 - 37.8	11,400
Kauai	100	18.7	15.1 - 23.0	200	25.4	21.2 - 30.1	<50	5.0	3.2 - 7.8	400	50.9	45.8 - 56.0	800
Maui	600	34.3	31.1 - 37.6	400	20.1	17.4 - 23.0	100	4.8	3.6 - 6.6	700	40.8	37.5 - 44.2	1,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	900	23.0	21.1 - 25.0	700	18.0	16.3 - 19.9	200	5.5	4.5 - 6.6	2,000	53.5	51.2 - 55.8	3,700
Native Hawaiian	1,600	37.9	35.7 - 40.2	1,000	23.2	21.3 - 25.2	200	3.9	3.1 - 4.9	1,500	35.0	32.8 - 37.2	4,300
Chinese	300	42.1	39.2 - 45.0	100	20.4	18.1 - 22.9	<50	2.8	1.9 - 3.9	200	34.7	32.0 - 37.6	600
Filipino	1,300	45.3	42.9 - 47.7	600	20.4	18.5 - 22.4	100	3.5	2.8 - 4.6	900	30.8	28.6 - 33.1	2,900
Japanese	1,000	50.4	47.4 - 53.5	300	16.9	14.7 - 19.4	<50	2.3	1.6 - 3.4	600	30.4	27.6 - 33.3	2,000
Other	800	35.6	32.8 - 38.5	500	20.5	18.2 - 23.1	100	2.6	1.8 - 3.7	1,000	41.3	38.4 - 44.2	2,400

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	34.5	30.6 - 38.7	300	25.9	22.3 - 29.8	<50	3.2	2.0 - 5.0	500	36.4	32.4 - 40.7	1,300
20-24	1,300	35.2	33.0 - 37.5	900	23.8	21.9 - 25.9	200	4.5	3.6 - 5.6	1,400	36.4	34.2 - 38.7	3,800
25-34	2,900	37.2	35.7 - 38.7	1,400	18.1	16.9 - 19.3	300	3.8	3.2 - 4.4	3,200	40.9	39.4 - 42.5	7,900
35+	1,200	40.9	38.5 - 43.3	500	18.5	16.6 - 20.5	100	3.0	2.2 - 3.9	1,100	37.7	35.3 - 40.1	2,900

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,100	37.3	35.8 - 38.8	1,700	20.9	19.6 - 22.2	300	3.8	3.2 - 4.4	3,200	38.1	36.6 - 39.6	8,300
Female	2,800	37.1	35.5 - 38.6	1,500	19.4	18.1 - 20.7	300	3.8	3.2 - 4.4	3,000	39.8	38.2 - 41.4	7,600

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,900	36.9	35.6 - 38.2	1,900	18.5	17.5 - 19.6	400	3.7	3.2 - 4.2	4,300	40.9	39.6 - 42.2	10,400
No	2,100	37.7	35.8 - 39.7	1,300	23.2	21.6 - 25.0	200	3.9	3.2 - 4.8	1,900	35.1	33.2 - 37.0	5,500

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,100	37.6	35.7 - 39.5	1,300	22.8	21.2 - 24.5	200	3.2	2.6 - 4.0	2,000	36.4	34.5 - 38.3	5,600
131-185%	700	33.3	30.4 - 36.3	400	20.8	18.3 - 23.4	100	4.3	3.2 - 5.8	900	41.6	38.6 - 44.7	2,000
186+%	2,700	38.0	36.4 - 39.5	1,300	17.5	16.3 - 18.8	300	4.1	3.5 - 4.7	2,900	40.5	38.9 - 42.1	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Income (2004-2011)	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,000	38.7	35.9 - 41.6	600	22.7	20.3 - 25.3	100	3.2	2.3 - 4.4	900	35.4	32.7 - 38.3	2,500
\$10,000-\$24,999	1,200	36.4	34.0 - 38.8	800	22.8	20.8 - 24.9	100	3.6	2.8 - 4.6	1,200	37.2	34.9 - 39.6	3,400
\$25,000-\$49,999	1,300	34.5	32.4 - 36.7	800	20.5	18.7 - 22.4	200	4.7	3.8 - 5.7	1,500	40.3	38.1 - 42.6	3,800
>=\$50,000	2,000	39.0	37.2 - 40.8	900	16.6	15.2 - 18.0	200	3.5	2.9 - 4.3	2,100	40.9	39.1 - 42.8	5,200

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	48.8	44.6 - 52.9	200	18.0	14.9 - 21.5	<50	4.2	2.9 - 6.1	300	29.0	25.3 - 33.0	1,100
Normal birth weight	5,400	36.3	35.2 - 37.4	3,000	20.3	19.4 - 21.3	600	3.7	3.3 - 4.2	5,900	39.6	38.5 - 40.8	14,800

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	39.8	35.6 - 44.1	300	23.3	19.8 - 27.2	100	5.9	4.1 - 8.3	400	31.1	27.2 - 35.2	1,200
No	5,400	36.9	35.8 - 38.0	2,900	19.9	19.0 - 20.9	500	3.6	3.2 - 4.1	5,700	39.5	38.4 - 40.7	14,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	36.3	33.8 - 38.9	800	25.1	22.9 - 27.5	200	6.2	5.0 - 7.6	1,000	32.4	29.9 - 34.9	3,000
No	4,700	37.3	36.1 - 38.5	2,400	19.0	18.1 - 20.0	400	3.2	2.8 - 3.7	5,100	40.5	39.3 - 41.7	12,700

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	36.3	34.6 - 38.1	1,400	22.0	20.6 - 23.6	200	3.7	3.0 - 4.4	2,500	38.0	36.3 - 39.7	6,600
No	3,400	37.7	36.3 - 39.0	1,700	18.7	17.6 - 19.9	400	3.9	3.4 - 4.5	3,600	39.7	38.3 - 41.1	9,100

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,700	36.4	35.3 - 37.5	3,200	20.8	19.9 - 21.8	600	3.9	3.4 - 4.3	6,100	38.9	37.8 - 40.1	15,500

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	29.1	26.3 - 32.1	400	20.3	17.9 - 23.0	100	3.5	2.5 - 4.9	900	47.1	43.9 - 50.2	1,900
Honolulu	4,400	39.3	38.0 - 40.7	2,300	20.8	19.7 - 21.9	400	3.7	3.2 - 4.2	4,100	36.2	34.9 - 37.6	11,200
Kauai	100	16.8	13.6 - 20.6	200	24.7	20.7 - 29.2	<50	5.2	3.4 - 7.9	400	53.3	48.4 - 58.1	800
Maui	600	33.3	30.3 - 36.5	300	20.1	17.6 - 22.9	100	5.0	3.7 - 6.6	700	41.6	38.4 - 44.9	1,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	23.1	21.1 - 25.3	600	18.0	16.1 - 19.9	200	5.9	4.8 - 7.2	1,900	53.1	50.6 - 55.5	3,600
Native Hawaiian	1,500	36.4	34.3 - 38.6	1,000	24.1	22.2 - 26.0	200	3.6	2.9 - 4.6	1,500	35.8	33.7 - 38.0	4,200
Chinese	300	43.4	39.1 - 47.7	100	18.3	15.3 - 21.9	<50	2.5	1.7 - 3.7	200	35.8	31.7 - 40.0	600
Filipino	1,300	42.5	40.0 - 45.1	700	22.1	20.1 - 24.4	100	3.7	2.9 - 4.8	900	31.6	29.3 - 34.1	2,900
Japanese	1,000	50.5	47.2 - 53.8	300	16.6	14.3 - 19.2	<50	1.9	1.2 - 3.0	600	31.0	28.0 - 34.1	2,000
Other	800	35.0	32.1 - 38.2	500	22.2	19.7 - 25.1	100	3.2	2.2 - 4.6	900	39.5	36.5 - 42.7	2,200

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	35.0	31.0 - 39.2	400	27.7	24.0 - 31.6	<50	2.8	1.7 - 4.5	400	34.6	30.7 - 38.7	1,300
20-24	1,300	33.9	31.7 - 36.2	900	24.9	22.9 - 27.1	200	4.3	3.4 - 5.3	1,400	36.9	34.6 - 39.2	3,800
25-34	2,800	36.5	35.0 - 38.1	1,400	18.8	17.6 - 20.1	300	4.2	3.6 - 4.9	3,100	40.5	38.9 - 42.1	7,700
35+	1,100	39.8	37.2 - 42.4	500	17.8	15.9 - 19.9	100	2.9	2.2 - 3.9	1,100	39.5	37.0 - 42.1	2,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,000	36.6	35.1 - 38.2	1,800	21.6	20.3 - 22.9	300	3.9	3.3 - 4.6	3,100	37.9	36.4 - 39.4	8,100
Female	2,700	36.1	34.5 - 37.7	1,500	20.0	18.7 - 21.4	300	3.8	3.2 - 4.5	3,000	40.1	38.5 - 41.8	7,400

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,700	36.0	34.7 - 37.4	2,000	19.2	18.1 - 20.3	400	3.8	3.3 - 4.4	4,200	41.0	39.6 - 42.4	10,300
No	1,900	37.0	35.1 - 39.0	1,300	24.2	22.5 - 26.0	200	3.9	3.2 - 4.7	1,800	34.9	33.0 - 36.8	5,200

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,100	36.6	34.7 - 38.5	1,400	24.0	22.4 - 25.8	200	3.2	2.6 - 4.0	2,000	36.2	34.3 - 38.1	5,600
131-185%	700	33.0	30.1 - 36.2	400	21.3	18.7 - 24.1	100	4.6	3.4 - 6.2	800	41.1	37.9 - 44.3	2,000
186+%	2,500	37.2	35.6 - 38.9	1,200	17.7	16.5 - 19.1	300	4.2	3.6 - 5.0	2,800	40.8	39.2 - 42.5	6,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight													
Low birth weight	500	46.0	41.7 - 50.3	200	19.9	16.5 - 23.7	<50	4.4	2.9 - 6.5	300	29.8	26.0 - 33.9	1,100
Normal birth weight	5,100	35.6	34.5 - 36.8	3,000	20.9	20.0 - 21.9	600	3.8	3.4 - 4.3	5,700	39.6	38.5 - 40.8	14,400

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester													
Yes	400	40.0	35.7 - 44.4	200	22.2	18.7 - 26.1	100	4.7	3.2 - 6.9	400	33.2	29.2 - 37.5	1,100
No	5,100	36.0	34.9 - 37.2	3,000	20.8	19.8 - 21.8	500	3.8	3.4 - 4.3	5,600	39.4	38.3 - 40.6	14,200

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant													
Yes	1,100	36.5	34.0 - 39.1	700	25.0	22.7 - 27.4	200	5.8	4.6 - 7.2	1,000	32.8	30.3 - 35.3	3,000
No	4,500	36.2	35.0 - 37.4	2,500	19.9	18.9 - 20.9	400	3.4	3.0 - 3.9	5,000	40.5	39.2 - 41.7	12,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	35.4	33.7 - 37.2	1,500	22.8	21.3 - 24.4	200	3.7	3.1 - 4.5	2,400	38.1	36.3 - 39.9	6,400
No	3,300	36.8	35.4 - 38.3	1,700	19.3	18.2 - 20.6	400	4.0	3.4 - 4.6	3,600	39.9	38.4 - 41.3	8,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,400	35.6	34.5 - 36.8	3,200	21.1	20.1 - 22.1	600	4.2	3.7 - 4.7	5,900	39.1	38.0 - 40.3	15,100

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	29.3	26.7 - 32.1	400	22.2	19.8 - 24.8	100	3.7	2.7 - 5.0	800	44.9	41.9 - 47.9	1,800
Honolulu	4,300	38.5	37.1 - 40.0	2,300	20.7	19.5 - 21.9	400	4.1	3.5 - 4.7	4,100	36.7	35.3 - 38.2	11,000
Kauai	100	16.7	13.6 - 20.4	200	23.7	19.9 - 28.0	<50	4.7	3.0 - 7.2	400	54.9	50.2 - 59.5	700
Maui	500	31.1	28.3 - 34.0	300	21.3	18.8 - 24.0	100	5.2	4.0 - 6.9	700	42.4	39.3 - 45.5	1,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	22.0	19.9 - 24.3	600	17.7	15.8 - 19.8	200	7.0	5.8 - 8.5	1,900	53.3	50.7 - 55.9	3,500
Native Hawaiian	1,400	36.1	34.1 - 38.1	1,000	23.9	22.2 - 25.8	200	3.9	3.1 - 4.8	1,400	36.1	34.1 - 38.1	4,000
Chinese	300	44.4	39.2 - 49.8	100	20.1	16.1 - 24.8	<50	2.8	1.6 - 4.8	200	32.7	28.0 - 37.8	600
Filipino	1,200	41.5	38.8 - 44.3	700	22.2	20.0 - 24.5	100	3.6	2.7 - 4.7	1,000	32.8	30.2 - 35.4	3,000
Japanese	1,000	47.6	44.2 - 51.1	400	18.5	15.9 - 21.3	<50	2.1	1.3 - 3.3	700	31.8	28.7 - 35.2	2,000
Other	700	35.1	31.8 - 38.5	500	22.6	19.8 - 25.7	100	3.2	2.1 - 4.7	800	39.1	35.8 - 42.6	2,100

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	37.5	33.5 - 41.7	300	27.2	23.6 - 31.1	<50	2.9	1.8 - 4.8	400	32.4	28.7 - 36.4	1,200
20-24	1,200	33.5	31.3 - 35.9	900	25.2	23.1 - 27.3	200	4.3	3.5 - 5.4	1,400	37.0	34.6 - 39.3	3,700
25-34	2,700	35.3	33.6 - 36.9	1,400	19.1	17.7 - 20.4	300	4.6	4.0 - 5.4	3,100	41.1	39.4 - 42.7	7,500
35+	1,000	38.7	36.1 - 41.5	500	18.3	16.3 - 20.6	100	3.1	2.3 - 4.3	1,100	39.8	37.1 - 42.5	2,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,800	36.2	34.6 - 37.8	1,700	21.5	20.2 - 22.9	300	4.3	3.7 - 5.1	3,000	38.0	36.4 - 39.6	7,800
Female	2,600	35.1	33.4 - 36.8	1,500	20.6	19.2 - 22.0	300	4.0	3.4 - 4.7	2,900	40.3	38.6 - 42.0	7,300

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,600	35.0	33.6 - 36.4	2,000	19.2	18.1 - 20.4	400	4.3	3.7 - 5.0	4,200	41.5	40.0 - 42.9	10,200
No	1,800	36.9	34.9 - 39.0	1,200	25.0	23.2 - 26.8	200	3.8	3.1 - 4.7	1,700	34.2	32.3 - 36.2	4,900

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,000	36.1	34.2 - 38.1	1,300	24.0	22.3 - 25.7	200	3.2	2.5 - 3.9	2,000	36.7	34.9 - 38.7	5,500
131-185%	700	32.2	29.1 - 35.4	400	20.3	17.8 - 23.1	100	5.6	4.2 - 7.3	900	41.9	38.7 - 45.2	2,100
186+%	2,300	36.1	34.4 - 37.9	1,200	18.3	16.9 - 19.7	300	4.5	3.8 - 5.4	2,600	41.1	39.3 - 42.9	6,300

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	45.8	41.2 - 50.5	200	19.6	16.0 - 23.8	<50	3.8	2.4 - 6.0	300	30.8	26.6 - 35.2	1,000
Normal birth weight	4,900	34.9	33.7 - 36.1	3,000	21.2	20.2 - 22.2	600	4.2	3.7 - 4.7	5,600	39.7	38.5 - 40.9	14,100

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	39.4	35.1 - 43.9	200	22.6	19.1 - 26.7	<50	4.2	2.8 - 6.3	400	33.7	29.6 - 38.1	1,000
No	4,900	35.3	34.1 - 36.5	2,900	21.0	20.0 - 22.0	600	4.2	3.7 - 4.7	5,500	39.5	38.3 - 40.7	13,900

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	36.2	33.6 - 38.9	700	24.1	21.8 - 26.6	200	5.9	4.7 - 7.3	1,000	33.8	31.3 - 36.5	2,900
No	4,200	35.3	34.1 - 36.6	2,400	20.4	19.3 - 21.5	500	3.8	3.3 - 4.4	4,800	40.4	39.1 - 41.8	12,000

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	35.6	33.8 - 37.4	1,400	23.2	21.7 - 24.9	200	3.8	3.2 - 4.6	2,300	37.3	35.5 - 39.1	6,200
No	3,100	35.5	34.1 - 37.1	1,700	19.4	18.2 - 20.7	400	4.4	3.8 - 5.1	3,500	40.6	39.1 - 42.2	8,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	5,200	35.4	34.2 - 36.5	3,100	21.2	20.2 - 22.2	600	4.1	3.7 - 4.6	5,800	39.3	38.2 - 40.5	14,700

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	29.0	26.6 - 31.6	400	22.2	19.9 - 24.6	100	3.5	2.6 - 4.7	800	45.3	42.6 - 48.1	1,700
Honolulu	4,100	37.9	36.5 - 39.4	2,300	20.9	19.8 - 22.2	400	4.1	3.6 - 4.8	4,000	37.0	35.6 - 38.4	10,800
Kauai	100	17.6	14.6 - 21.0	200	23.7	20.3 - 27.6	<50	4.6	3.1 - 6.8	400	54.1	49.8 - 58.3	700
Maui	500	32.0	29.4 - 34.7	300	20.7	18.5 - 23.1	100	4.7	3.5 - 6.2	700	42.6	39.8 - 45.4	1,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	21.6	19.5 - 23.8	600	18.1	16.2 - 20.2	200	6.5	5.3 - 8.0	1,800	53.8	51.2 - 56.4	3,300
Native Hawaiian	1,400	36.3	34.5 - 38.1	900	22.8	21.3 - 24.4	200	4.2	3.5 - 5.0	1,400	36.7	35.0 - 38.5	3,900
Chinese	300	43.2	37.4 - 49.1	100	20.9	16.4 - 26.3	<50	3.1	1.6 - 5.7	200	32.9	27.7 - 38.5	600
Filipino	1,200	40.5	37.8 - 43.3	700	23.1	20.8 - 25.5	100	3.4	2.6 - 4.5	1,000	33.0	30.5 - 35.7	3,000
Japanese	1,000	46.7	43.2 - 50.2	400	19.5	16.8 - 22.4	100	2.8	1.8 - 4.2	600	31.1	28.0 - 34.4	2,100
Other	700	34.8	31.5 - 38.4	400	22.3	19.5 - 25.5	100	2.9	1.9 - 4.3	800	39.9	36.4 - 43.6	2,000

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	38.8	35.0 - 42.8	300	27.2	23.8 - 30.9	<50	3.5	2.3 - 5.4	400	30.4	27.0 - 34.1	1,200
20-24	1,200	32.4	30.2 - 34.7	900	25.8	23.7 - 27.9	100	4.1	3.3 - 5.1	1,400	37.7	35.5 - 40.0	3,600
25-34	2,600	35.3	33.6 - 36.9	1,400	18.7	17.4 - 20.0	300	4.6	3.9 - 5.4	3,000	41.5	39.8 - 43.2	7,300
35+	1,000	38.1	35.3 - 40.9	500	19.1	16.9 - 21.4	100	3.1	2.3 - 4.3	1,000	39.7	36.9 - 42.5	2,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,700	35.8	34.2 - 37.4	1,600	21.4	20.1 - 22.8	300	4.1	3.5 - 4.8	3,000	38.7	37.2 - 40.3	7,600
Female	2,500	34.9	33.3 - 36.6	1,500	21.0	19.6 - 22.4	300	4.2	3.5 - 4.9	2,800	40.0	38.3 - 41.7	7,100

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,500	34.4	33.0 - 35.8	2,000	19.6	18.5 - 20.8	400	4.4	3.8 - 5.0	4,200	41.6	40.1 - 43.0	10,100
No	1,700	37.4	35.4 - 39.4	1,100	24.6	22.9 - 26.4	200	3.6	3.0 - 4.4	1,600	34.4	32.6 - 36.3	4,600

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,900	35.8	34.0 - 37.7	1,300	24.0	22.3 - 25.6	200	3.0	2.4 - 3.7	2,000	37.2	35.4 - 39.1	5,400
131-185%	700	31.5	28.7 - 34.4	500	20.9	18.6 - 23.5	100	5.6	4.3 - 7.2	1,000	42.0	39.0 - 45.0	2,300
186+%	2,100	36.0	34.2 - 37.8	1,100	18.1	16.7 - 19.6	300	4.7	3.9 - 5.6	2,400	41.3	39.4 - 43.1	5,800

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight													
Low birth weight	400	46.0	41.3 - 50.8	200	20.3	16.6 - 24.6	<50	3.9	2.4 - 6.2	300	29.7	25.6 - 34.2	1,000
Normal birth weight	4,800	34.6	33.4 - 35.8	2,900	21.3	20.3 - 22.3	600	4.2	3.7 - 4.7	5,500	40.0	38.8 - 41.2	13,800

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester													
Yes	400	39.7	35.5 - 44.1	200	23.1	19.7 - 27.0	<50	3.8	2.6 - 5.6	300	33.3	29.4 - 37.5	1,000
No	4,700	35.0	33.8 - 36.2	2,800	21.1	20.1 - 22.1	600	4.2	3.7 - 4.8	5,400	39.7	38.5 - 41.0	13,500

	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant													
Yes	1,000	36.3	33.8 - 38.9	700	23.9	21.7 - 26.3	100	5.0	4.0 - 6.3	1,000	34.7	32.2 - 37.3	2,800
No	4,100	34.9	33.7 - 36.2	2,400	20.6	19.5 - 21.7	500	4.0	3.5 - 4.6	4,700	40.5	39.2 - 41.8	11,600

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	35.4	33.6 - 37.2	1,400	23.1	21.6 - 24.7	200	3.7	3.0 - 4.4	2,300	37.9	36.1 - 39.6	6,000
No	3,000	35.2	33.7 - 36.8	1,700	19.8	18.6 - 21.1	400	4.5	3.9 - 5.2	3,500	40.5	38.9 - 42.0	8,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

State	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	4,700	33.4	32.2 - 34.5	3,100	21.8	20.9 - 22.9	700	4.7	4.2 - 5.2	5,600	40.1	38.9 - 41.3	14,100

County	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	29.1	26.8 - 31.5	300	21.1	19.1 - 23.3	100	4.1	3.1 - 5.2	700	45.7	43.2 - 48.3	1,600
Honolulu	3,700	35.5	34.0 - 37.0	2,200	21.6	20.3 - 22.8	500	4.7	4.1 - 5.4	4,000	38.2	36.8 - 39.7	10,300
Kauai	100	17.7	14.8 - 21.0	200	23.6	20.3 - 27.2	<50	4.9	3.4 - 6.9	300	53.9	49.8 - 57.9	600
Maui	500	30.1	27.8 - 32.6	400	23.8	21.6 - 26.1	100	4.8	3.7 - 6.2	600	41.3	38.7 - 43.9	1,500

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

DOH Race-Ethnicity	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	20.1	18.0 - 22.4	600	18.7	16.7 - 20.8	200	7.5	6.2 - 9.1	1,700	53.6	51.0 - 56.2	3,200
Native Hawaiian	1,300	34.6	33.0 - 36.2	900	24.5	23.1 - 26.0	200	4.4	3.7 - 5.1	1,300	36.5	35.0 - 38.1	3,700
Chinese	200	41.5	35.3 - 48.0	100	20.5	15.7 - 26.3	<50	2.9	1.3 - 6.0	200	35.1	29.3 - 41.4	600
Filipino	1,100	38.1	35.3 - 41.0	700	23.5	21.2 - 26.0	100	3.8	2.9 - 5.0	1,000	34.6	31.9 - 37.4	2,900
Japanese	800	43.4	39.7 - 47.1	400	20.3	17.5 - 23.5	100	3.4	2.3 - 5.0	600	32.9	29.6 - 36.5	1,900
Other	600	33.6	30.1 - 37.3	400	21.4	18.4 - 24.6	100	3.6	2.4 - 5.2	800	41.5	37.8 - 45.3	1,900

PRAMS Age Group	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	37.6	34.0 - 41.4	300	27.5	24.3 - 30.9	<50	3.5	2.3 - 5.3	400	31.3	28.1 - 34.8	1,300
20-24	1,100	30.6	28.5 - 32.9	900	26.3	24.3 - 28.5	100	4.1	3.2 - 5.1	1,400	39.0	36.7 - 41.3	3,500
25-34	2,300	32.9	31.3 - 34.6	1,300	19.4	18.0 - 20.8	400	5.3	4.6 - 6.2	2,900	42.4	40.7 - 44.1	6,900
35+	900	36.4	33.5 - 39.3	500	19.6	17.4 - 22.1	100	4.2	3.2 - 5.5	1,000	39.8	36.9 - 42.7	2,400

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Infant Gender	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,500	33.7	32.1 - 35.3	1,600	22.0	20.7 - 23.4	300	4.6	3.9 - 5.4	2,900	39.7	38.1 - 41.3	7,300
Female	2,200	33.0	31.4 - 34.7	1,500	21.7	20.3 - 23.1	300	4.7	4.1 - 5.5	2,700	40.6	38.9 - 42.3	6,800

Married	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,100	32.2	30.8 - 33.6	2,000	20.4	19.3 - 21.7	500	5.0	4.3 - 5.7	4,100	42.4	40.9 - 43.9	9,700
No	1,600	36.0	34.1 - 38.0	1,100	25.0	23.3 - 26.8	200	4.0	3.3 - 4.8	1,500	35.0	33.1 - 36.9	4,300

Poverty Level	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,800	33.4	31.7 - 35.3	1,300	24.6	23.0 - 26.2	200	3.5	2.9 - 4.2	2,100	38.5	36.7 - 40.3	5,400
131-185%	700	30.1	27.3 - 32.9	500	21.6	19.3 - 24.1	100	6.3	5.0 - 7.9	1,000	42.0	39.1 - 45.0	2,400
186+%	1,800	33.8	31.9 - 35.8	1,000	18.7	17.2 - 20.3	300	5.4	4.5 - 6.4	2,200	42.1	40.2 - 44.1	5,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Birthweight	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	42.4	37.7 - 47.4	200	23.2	19.3 - 27.7	<50	4.0	2.5 - 6.4	300	30.3	26.0 - 34.9	900
Normal birth weight	4,300	32.7	31.6 - 33.9	2,900	21.8	20.8 - 22.8	600	4.7	4.2 - 5.3	5,400	40.8	39.6 - 42.0	13,200

Smoke 3rd Trimester	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	37.1	33.0 - 41.4	300	26.6	23.0 - 30.5	<50	4.4	3.1 - 6.3	300	31.9	28.0 - 36.0	1,000
No	4,300	33.1	31.9 - 34.3	2,800	21.6	20.6 - 22.6	600	4.7	4.2 - 5.3	5,200	40.6	39.4 - 41.9	12,900

Smoke Before Pregnant	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	33.7	31.2 - 36.2	700	25.4	23.2 - 27.8	100	5.1	4.1 - 6.4	900	35.8	33.3 - 38.4	2,600
No	3,700	33.1	31.8 - 34.5	2,300	21.1	20.0 - 22.2	500	4.6	4.0 - 5.2	4,600	41.2	39.8 - 42.5	11,100

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration (grouped), among initiators- by
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

WIC During Pregnancy	Breastfed <1 week			Breastfed 1-4 weeks			Breastfed 5-8 weeks			Breastfed 9+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	34.0	32.3 - 35.8	1,400	23.4	21.9 - 25.0	200	4.2	3.5 - 5.0	2,200	38.4	36.6 - 40.1	5,800
No	2,700	32.9	31.4 - 34.5	1,700	20.7	19.5 - 22.1	400	5.0	4.3 - 5.7	3,400	41.4	39.8 - 43.0	8,100