

**PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (exclusive) duration 8+ weeks, among initiators	Breastfeeding (exclusive) duration 8+ weeks, among those who initiated breastfeeding

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/22/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013


Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	8,300	58.8	57.6 - 60.0	5,800	41.2	40.0 - 42.4	14,100
2001-2005	8,800	59.8	58.6 - 61.0	5,900	40.2	39.0 - 41.4	14,700
2002-2006	9,100	60.0	58.9 - 61.2	6,000	40.0	38.8 - 41.1	15,100
2003-2007	9,400	60.4	59.3 - 61.5	6,200	39.6	38.5 - 40.7	15,500
2004-2008	9,600	60.5	59.4 - 61.6	6,300	39.5	38.4 - 40.6	16,000
2005-2009	9,700	60.2	59.0 - 61.3	6,400	39.8	38.7 - 41.0	16,100
2006-2010	9,600	58.4	57.1 - 59.6	6,900	41.6	40.4 - 42.9	16,500
2007-2011	9,500	57.1	55.7 - 58.4	7,200	42.9	41.6 - 44.3	16,700
2008-2012	9,400	55.9	54.5 - 57.3	7,400	44.1	42.7 - 45.5	16,800
2009-2013	9,000	53.3	51.8 - 54.8	7,900	46.7	45.2 - 48.2	16,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,000	53.3	51.8 - 54.8	7,900	46.7	45.2 - 48.2	16,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	52.8	50.4 - 55.3	1,000	47.2	44.7 - 49.6	2,100
Honolulu	6,500	53.9	51.9 - 56.0	5,600	46.1	44.0 - 48.1	12,100
Kauai	400	45.7	42.9 - 48.5	400	54.3	51.5 - 57.1	800
Maui	1,000	52.9	50.6 - 55.2	900	47.1	44.8 - 49.4	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,700	40.7	37.8 - 43.8	2,500	59.3	56.2 - 62.2	4,200
Native Hawaiian	2,800	57.8	55.1 - 60.5	2,100	42.2	39.5 - 44.9	4,900
Chinese	400	54.7	47.2 - 62.0	400	45.3	38.0 - 52.8	800
Filipino	1,900	65.0	61.4 - 68.4	1,000	35.0	31.6 - 38.6	2,900
Japanese	900	55.5	50.5 - 60.4	700	44.5	39.6 - 49.5	1,700
Other	1,200	49.8	45.5 - 54.0	1,200	50.2	46.0 - 54.5	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	63.2	57.4 - 68.6	400	36.8	31.4 - 42.6	1,100
20-24	2,100	54.0	50.8 - 57.3	1,800	46.0	42.7 - 49.2	3,800
25-34	4,600	50.9	48.8 - 53.0	4,400	49.1	47.0 - 51.2	9,000
35+	1,700	56.1	52.5 - 59.6	1,300	43.9	40.4 - 47.5	3,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,500	52.7	50.6 - 54.9	4,000	47.3	45.1 - 49.4	8,500
Female	4,500	53.9	51.7 - 56.1	3,800	46.1	43.9 - 48.3	8,300

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,300	50.0	48.0 - 51.9	5,300	50.0	48.1 - 52.0	10,600
No	3,700	58.9	56.5 - 61.3	2,600	41.1	38.7 - 43.5	6,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,300	56.6	54.1 - 59.1	2,500	43.4	40.9 - 45.9	5,800
131-185%	1,500	56.7	52.8 - 60.6	1,100	43.3	39.4 - 47.2	2,600
186+%	3,500	49.5	47.2 - 51.9	3,600	50.5	48.1 - 52.8	7,100

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight							
Low birth weight	700	65.5	63.0 - 67.8	400	34.5	32.2 - 37.0	1,100
Normal birth weight	8,200	52.4	50.8 - 54.0	7,500	47.6	46.0 - 49.2	15,700

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester							
Yes	600	60.5	54.3 - 66.3	400	39.5	33.7 - 45.7	1,000
No	8,200	53.0	51.4 - 54.6	7,300	47.0	45.4 - 48.6	15,500

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant							
Yes	1,700	56.6	53.0 - 60.0	1,300	43.4	40.0 - 47.0	3,100
No	7,100	52.7	51.0 - 54.4	6,300	47.3	45.6 - 49.0	13,400

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,100	57.3	55.0 - 59.6	3,100	42.7	40.4 - 45.0	7,200
No	4,800	50.4	48.3 - 52.4	4,700	49.6	47.6 - 51.7	9,400

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,400	55.9	54.5 - 57.3	7,400	44.1	42.7 - 45.5	16,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	54.2	51.6 - 56.7	1,000	45.8	43.3 - 48.4	2,100
Honolulu	6,800	56.8	54.9 - 58.6	5,200	43.2	41.4 - 45.1	12,000
Kauai	400	49.6	46.1 - 53.0	400	50.4	47.0 - 53.9	800
Maui	1,000	54.8	52.4 - 57.3	900	45.2	42.7 - 47.6	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	43.0	40.2 - 45.9	2,300	57.0	54.1 - 59.8	4,100
Native Hawaiian	2,900	60.2	57.6 - 62.7	1,900	39.8	37.3 - 42.4	4,800
Chinese	400	53.4	46.4 - 60.2	300	46.6	39.8 - 53.6	700
Filipino	2,000	67.1	63.9 - 70.2	1,000	32.9	29.8 - 36.1	3,000
Japanese	1,000	61.2	56.7 - 65.5	700	38.8	34.5 - 43.3	1,700
Other	1,300	52.3	48.4 - 56.2	1,200	47.7	43.8 - 51.6	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	63.2	58.0 - 68.2	400	36.8	31.8 - 42.0	1,200
20-24	2,200	58.8	55.7 - 61.8	1,500	41.2	38.2 - 44.3	3,800
25-34	4,700	52.9	51.0 - 54.9	4,200	47.1	45.1 - 49.0	8,900
35+	1,700	58.1	54.8 - 61.4	1,300	41.9	38.6 - 45.2	3,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,900	56.2	54.2 - 58.2	3,800	43.8	41.8 - 45.8	8,600
Female	4,500	55.5	53.5 - 57.6	3,600	44.5	42.4 - 46.5	8,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,600	52.6	50.8 - 54.4	5,000	47.4	45.6 - 49.2	10,600
No	3,900	61.4	59.2 - 63.7	2,400	38.6	36.3 - 40.8	6,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	59.8	57.4 - 62.1	2,300	40.2	37.9 - 42.6	5,800
131-185%	1,500	57.2	53.5 - 60.8	1,100	42.8	39.2 - 46.5	2,600
186+%	3,800	52.1	49.9 - 54.3	3,400	47.9	45.7 - 50.1	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight							
Low birth weight	800	67.0	64.2 - 69.7	400	33.0	30.3 - 35.8	1,100
Normal birth weight	8,600	55.1	53.6 - 56.6	7,000	44.9	43.4 - 46.4	15,700

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester							
Yes	700	66.1	60.6 - 71.2	400	33.9	28.8 - 39.4	1,000
No	8,600	55.3	53.8 - 56.8	6,900	44.7	43.2 - 46.2	15,500

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant							
Yes	1,900	60.8	57.5 - 64.0	1,200	39.2	36.0 - 42.5	3,100
No	7,400	54.9	53.3 - 56.5	6,000	45.1	43.5 - 46.7	13,400

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,200	59.3	57.2 - 61.5	2,900	40.7	38.5 - 42.8	7,100
No	5,100	53.5	51.6 - 55.4	4,400	46.5	44.6 - 48.4	9,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,500	57.1	55.7 - 58.4	7,200	42.9	41.6 - 44.3	16,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	53.2	50.4 - 55.9	1,000	46.8	44.1 - 49.6	2,100
Honolulu	6,900	58.4	56.7 - 60.2	4,900	41.6	39.8 - 43.3	11,900
Kauai	400	50.4	46.5 - 54.3	400	49.6	45.7 - 53.5	800
Maui	1,100	55.6	52.9 - 58.3	800	44.4	41.7 - 47.1	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	44.0	41.3 - 46.8	2,300	56.0	53.2 - 58.7	4,000
Native Hawaiian	2,900	61.0	58.5 - 63.4	1,900	39.0	36.6 - 41.5	4,800
Chinese	400	55.7	49.2 - 62.1	300	44.3	37.9 - 50.8	700
Filipino	2,000	68.2	65.2 - 71.0	900	31.8	29.0 - 34.8	3,000
Japanese	1,200	63.9	59.9 - 67.8	600	36.1	32.2 - 40.1	1,800
Other	1,300	52.6	48.9 - 56.3	1,200	47.4	43.7 - 51.1	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	62.3	57.3 - 67.0	500	37.7	33.0 - 42.7	1,300
20-24	2,300	60.0	57.1 - 62.8	1,500	40.0	37.2 - 42.9	3,900
25-34	4,700	54.2	52.3 - 56.1	3,900	45.8	43.9 - 47.7	8,600
35+	1,800	59.3	56.2 - 62.3	1,200	40.7	37.7 - 43.8	3,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,000	57.2	55.3 - 59.0	3,700	42.8	41.0 - 44.7	8,700
Female	4,600	56.9	55.0 - 58.8	3,500	43.1	41.2 - 45.0	8,100

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,700	53.8	52.1 - 55.5	4,900	46.2	44.5 - 47.9	10,500
No	3,900	62.5	60.4 - 64.7	2,300	37.5	35.3 - 39.6	6,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	60.7	58.4 - 62.9	2,300	39.3	37.1 - 41.6	5,800
131-185%	1,400	56.6	53.1 - 60.1	1,100	43.4	39.9 - 46.9	2,500
186+%	3,900	54.2	52.2 - 56.3	3,300	45.8	43.7 - 47.8	7,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,700	62.0	58.7 - 65.2	1,100	38.0	34.8 - 41.3	2,800
\$10,000-\$24,999	2,000	60.2	57.2 - 63.1	1,300	39.8	36.9 - 42.8	3,200
\$25,000-\$49,999	2,200	55.9	53.0 - 58.6	1,700	44.1	41.4 - 47.0	3,900
>=\$50,000	3,100	53.9	51.6 - 56.2	2,700	46.1	43.8 - 48.4	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	68.3	65.1 - 71.2	400	31.7	28.8 - 34.9	1,200
Normal birth weight	8,800	56.2	54.8 - 57.6	6,800	43.8	42.4 - 45.2	15,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	66.3	61.2 - 71.0	400	33.7	29.0 - 38.8	1,200
No	8,700	56.5	55.0 - 57.8	6,700	43.5	42.2 - 45.0	15,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	63.4	60.4 - 66.4	1,200	36.6	33.6 - 39.6	3,200
No	7,400	55.6	54.1 - 57.1	5,900	44.4	42.9 - 45.9	13,300

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,100	59.2	57.1 - 61.2	2,800	40.8	38.8 - 42.9	6,900
No	5,300	55.7	54.0 - 57.5	4,200	44.3	42.5 - 46.0	9,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,600	58.4	57.1 - 59.6	6,900	41.6	40.4 - 42.9	16,500

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	53.1	50.2 - 56.1	1,000	46.9	43.9 - 49.8	2,100
Honolulu	7,100	60.1	58.5 - 61.7	4,700	39.9	38.3 - 41.5	11,700
Kauai	400	50.0	45.6 - 54.4	400	50.0	45.6 - 54.4	800
Maui	1,100	57.2	54.2 - 60.1	800	42.8	39.9 - 45.8	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	44.7	42.1 - 47.3	2,200	55.3	52.7 - 57.9	4,000
Native Hawaiian	2,900	62.0	59.5 - 64.4	1,800	38.0	35.6 - 40.5	4,700
Chinese	400	58.5	52.6 - 64.2	300	41.5	35.8 - 47.4	600
Filipino	2,000	68.1	65.3 - 70.8	900	31.9	29.2 - 34.7	3,000
Japanese	1,200	66.0	62.3 - 69.6	600	34.0	30.4 - 37.7	1,800
Other	1,400	56.1	52.6 - 59.5	1,100	43.9	40.5 - 47.4	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	62.2	57.6 - 66.7	500	37.8	33.3 - 42.4	1,400
20-24	2,300	60.8	58.1 - 63.4	1,500	39.2	36.6 - 41.9	3,800
25-34	4,700	55.8	54.0 - 57.5	3,700	44.2	42.5 - 46.0	8,300
35+	1,800	60.9	58.0 - 63.8	1,200	39.1	36.2 - 42.0	3,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,100	59.3	57.6 - 61.1	3,500	40.7	38.9 - 42.4	8,600
Female	4,500	57.4	55.5 - 59.2	3,400	42.6	40.8 - 44.5	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,800	55.4	53.8 - 57.0	4,600	44.6	43.0 - 46.2	10,400
No	3,900	63.4	61.4 - 65.5	2,200	36.6	34.5 - 38.6	6,100

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	61.6	59.4 - 63.8	2,200	38.4	36.2 - 40.6	5,700
131-185%	1,400	58.3	54.9 - 61.6	1,000	41.7	38.4 - 45.1	2,400
186+%	4,000	55.6	53.7 - 57.5	3,200	44.4	42.5 - 46.3	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,700	61.8	58.6 - 65.0	1,000	38.2	35.0 - 41.4	2,700
\$10,000-\$24,999	2,000	62.1	59.2 - 64.8	1,200	37.9	35.2 - 40.8	3,300
\$25,000-\$49,999	2,200	58.1	55.5 - 60.7	1,600	41.9	39.3 - 44.5	3,800
>=\$50,000	3,100	54.7	52.6 - 56.9	2,600	45.3	43.1 - 47.4	5,700

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	67.4	63.9 - 70.7	400	32.6	29.3 - 36.1	1,200
Normal birth weight	8,800	57.7	56.4 - 59.0	6,500	42.3	41.0 - 43.6	15,300

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	65.3	60.5 - 69.8	400	34.7	30.2 - 39.5	1,200
No	8,700	57.9	56.6 - 59.2	6,300	42.1	40.8 - 43.4	15,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	64.7	61.8 - 67.5	1,100	35.3	32.5 - 38.2	3,200
No	7,400	57.0	55.5 - 58.4	5,600	43.0	41.6 - 44.5	13,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,000	60.3	58.2 - 62.2	2,700	39.7	37.8 - 41.8	6,700
No	5,400	57.1	55.4 - 58.8	4,100	42.9	41.2 - 44.6	9,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,700	60.2	59.0 - 61.3	6,400	39.8	38.7 - 41.0	16,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	54.3	51.2 - 57.4	900	45.7	42.6 - 48.8	2,000
Honolulu	7,200	62.3	60.9 - 63.7	4,300	37.7	36.3 - 39.1	11,500
Kauai	400	49.6	44.8 - 54.4	400	50.4	45.6 - 55.2	800
Maui	1,000	57.7	54.4 - 60.8	800	42.3	39.2 - 45.6	1,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	45.9	43.5 - 48.4	2,100	54.1	51.6 - 56.5	3,800
Native Hawaiian	2,900	63.7	61.3 - 66.0	1,600	36.3	34.0 - 38.7	4,500
Chinese	400	64.3	59.6 - 68.8	200	35.7	31.2 - 40.4	600
Filipino	2,000	68.9	66.3 - 71.3	900	31.1	28.7 - 33.7	2,900
Japanese	1,300	68.2	64.9 - 71.3	600	31.8	28.7 - 35.1	1,900
Other	1,400	58.2	55.0 - 61.4	1,000	41.8	38.6 - 45.0	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	62.7	58.3 - 67.0	500	37.3	33.0 - 41.7	1,400
20-24	2,400	63.4	60.9 - 65.8	1,400	36.6	34.2 - 39.1	3,800
25-34	4,700	57.5	55.9 - 59.2	3,400	42.5	40.8 - 44.1	8,100
35+	1,800	62.1	59.4 - 64.7	1,100	37.9	35.3 - 40.6	2,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,100	61.1	59.4 - 62.7	3,300	38.9	37.3 - 40.6	8,400
Female	4,600	59.2	57.5 - 60.9	3,200	40.8	39.1 - 42.5	7,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,900	57.2	55.7 - 58.6	4,400	42.8	41.4 - 44.3	10,200
No	3,900	65.4	63.4 - 67.3	2,000	34.6	32.7 - 36.6	5,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	63.3	61.2 - 65.3	2,100	36.7	34.7 - 38.8	5,600
131-185%	1,300	59.6	56.3 - 62.8	900	40.4	37.2 - 43.7	2,200
186+%	4,100	57.6	55.8 - 59.3	3,000	42.4	40.7 - 44.2	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,700	63.6	60.5 - 66.6	900	36.4	33.4 - 39.5	2,600
\$10,000-\$24,999	2,100	63.9	61.2 - 66.4	1,200	36.1	33.6 - 38.8	3,300
\$25,000-\$49,999	2,200	58.7	56.2 - 61.1	1,500	41.3	38.9 - 43.8	3,700
>=\$50,000	3,100	57.2	55.2 - 59.2	2,300	42.8	40.8 - 44.8	5,500

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	68.9	65.1 - 72.4	400	31.1	27.6 - 34.9	1,200
Normal birth weight	8,900	59.5	58.3 - 60.7	6,100	40.5	39.3 - 41.7	15,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	67.5	63.0 - 71.6	400	32.5	28.4 - 37.0	1,200
No	8,700	59.5	58.3 - 60.8	5,900	40.5	39.2 - 41.7	14,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	66.3	63.6 - 68.9	1,000	33.7	31.1 - 36.4	3,100
No	7,500	58.7	57.3 - 60.0	5,300	41.3	40.0 - 42.7	12,800

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,000	61.6	59.7 - 63.5	2,500	38.4	36.5 - 40.3	6,500
No	5,500	59.0	57.4 - 60.5	3,800	41.0	39.5 - 42.6	9,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,600	60.5	59.4 - 61.6	6,300	39.5	38.4 - 40.6	16,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	53.4	50.2 - 56.7	900	46.6	43.3 - 49.8	2,000
Honolulu	7,200	62.8	61.5 - 64.0	4,300	37.2	36.0 - 38.5	11,400
Kauai	400	48.6	43.5 - 53.7	400	51.4	46.3 - 56.5	800
Maui	1,000	58.6	55.2 - 61.9	700	41.4	38.1 - 44.8	1,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,700	45.4	43.1 - 47.7	2,000	54.6	52.3 - 56.9	3,700
Native Hawaiian	2,800	64.6	62.3 - 66.8	1,500	35.4	33.2 - 37.7	4,300
Chinese	400	64.5	61.7 - 67.3	200	35.5	32.7 - 38.3	600
Filipino	2,000	68.7	66.4 - 70.9	900	31.3	29.1 - 33.6	2,900
Japanese	1,400	69.3	66.4 - 72.1	600	30.7	27.9 - 33.6	2,000
Other	1,400	58.3	55.4 - 61.2	1,000	41.7	38.8 - 44.6	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	62.9	58.7 - 66.9	500	37.1	33.1 - 41.3	1,300
20-24	2,400	63.2	60.8 - 65.4	1,400	36.8	34.6 - 39.2	3,800
25-34	4,600	58.4	56.8 - 59.9	3,300	41.6	40.1 - 43.2	7,900
35+	1,800	61.8	59.4 - 64.1	1,100	38.2	35.9 - 40.6	2,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,100	61.4	59.9 - 62.9	3,200	38.6	37.1 - 40.1	8,300
Female	4,500	59.5	57.9 - 61.1	3,100	40.5	38.9 - 42.1	7,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,100	58.4	57.1 - 59.7	4,300	41.6	40.3 - 42.9	10,400
No	3,600	64.5	62.5 - 66.4	2,000	35.5	33.6 - 37.5	5,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	63.3	61.4 - 65.1	2,100	36.7	34.9 - 38.6	5,600
131-185%	1,200	57.9	54.8 - 61.0	900	42.1	39.0 - 45.2	2,000
186+%	4,200	58.7	57.1 - 60.2	3,000	41.3	39.8 - 42.9	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,600	64.3	61.5 - 67.1	900	35.7	32.9 - 38.5	2,500
\$10,000-\$24,999	2,100	62.4	60.0 - 64.7	1,300	37.6	35.3 - 40.0	3,400
\$25,000-\$49,999	2,300	59.0	56.8 - 61.2	1,600	41.0	38.8 - 43.2	3,800
>=\$50,000	3,000	58.2	56.4 - 60.0	2,200	41.8	40.0 - 43.6	5,200

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	70.0	66.0 - 73.7	300	30.0	26.3 - 34.0	1,100
Normal birth weight	8,900	59.8	58.6 - 60.9	6,000	40.2	39.1 - 41.4	14,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	67.7	63.5 - 71.6	400	32.3	28.4 - 36.5	1,200
No	8,700	59.9	58.8 - 61.0	5,800	40.1	39.0 - 41.2	14,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	66.7	64.2 - 69.1	1,000	33.3	30.9 - 35.8	3,000
No	7,500	59.0	57.8 - 60.2	5,200	41.0	39.8 - 42.2	12,700

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,000	61.5	59.7 - 63.2	2,500	38.5	36.8 - 40.3	6,600
No	5,400	59.6	58.2 - 61.0	3,700	40.4	39.0 - 41.8	9,100

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,400	60.4	59.3 - 61.5	6,200	39.6	38.5 - 40.7	15,500

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	52.5	49.3 - 55.6	900	47.5	44.4 - 50.7	1,900
Honolulu	7,100	63.1	61.8 - 64.5	4,100	36.9	35.5 - 38.2	11,200
Kauai	300	45.6	40.8 - 50.5	400	54.4	49.5 - 59.2	800
Maui	1,000	57.6	54.3 - 60.8	700	42.4	39.2 - 45.7	1,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	45.7	43.3 - 48.2	2,000	54.3	51.8 - 56.7	3,600
Native Hawaiian	2,600	63.7	61.6 - 65.8	1,500	36.3	34.2 - 38.4	4,200
Chinese	400	63.5	59.2 - 67.5	200	36.5	32.5 - 40.8	600
Filipino	2,000	67.8	65.3 - 70.1	1,000	32.2	29.9 - 34.7	2,900
Japanese	1,400	68.7	65.6 - 71.7	600	31.3	28.3 - 34.4	2,000
Other	1,300	59.9	56.7 - 63.0	900	40.1	37.0 - 43.3	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	64.9	60.8 - 68.8	400	35.1	31.2 - 39.2	1,300
20-24	2,400	62.7	60.3 - 65.0	1,400	37.3	35.0 - 39.7	3,800
25-34	4,500	58.7	57.1 - 60.3	3,200	41.3	39.7 - 42.9	7,700
35+	1,700	60.0	57.4 - 62.5	1,100	40.0	37.5 - 42.6	2,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,000	61.5	60.0 - 63.0	3,100	38.5	37.0 - 40.0	8,100
Female	4,400	59.2	57.5 - 60.8	3,000	40.8	39.2 - 42.5	7,400

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,000	58.3	56.9 - 59.6	4,300	41.7	40.4 - 43.1	10,300
No	3,400	64.6	62.6 - 66.5	1,900	35.4	33.5 - 37.4	5,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,600	63.3	61.4 - 65.2	2,100	36.7	34.8 - 38.6	5,600
131-185%	1,200	58.6	55.3 - 61.7	800	41.4	38.3 - 44.7	2,000
186+%	3,900	58.3	56.6 - 60.0	2,800	41.7	40.0 - 43.4	6,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight							
Low birth weight	800	69.3	65.2 - 73.1	300	30.7	26.9 - 34.8	1,100
Normal birth weight	8,600	59.7	58.6 - 60.9	5,800	40.3	39.1 - 41.4	14,400

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester							
Yes	700	65.7	61.4 - 69.8	400	34.3	30.2 - 38.6	1,100
No	8,500	60.0	58.8 - 61.1	5,700	40.0	38.9 - 41.2	14,200

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant							
Yes	2,000	66.2	63.7 - 68.7	1,000	33.8	31.3 - 36.3	3,000
No	7,300	58.9	57.7 - 60.2	5,100	41.1	39.8 - 42.3	12,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,900	61.3	59.5 - 63.0	2,500	38.7	37.0 - 40.5	6,400
No	5,300	59.5	58.0 - 60.9	3,600	40.5	39.1 - 42.0	8,900

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,100	60.0	58.9 - 61.2	6,000	40.0	38.8 - 41.1	15,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	54.6	51.6 - 57.6	800	45.4	42.4 - 48.4	1,800
Honolulu	6,900	62.4	60.9 - 63.8	4,200	37.6	36.2 - 39.1	11,000
Kauai	300	44.2	39.6 - 48.9	400	55.8	51.1 - 60.4	700
Maui	900	56.8	53.7 - 59.8	700	43.2	40.2 - 46.3	1,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	44.9	42.3 - 47.4	1,900	55.1	52.6 - 57.7	3,500
Native Hawaiian	2,500	63.4	61.4 - 65.4	1,500	36.6	34.6 - 38.6	4,000
Chinese	400	66.5	61.4 - 71.2	200	33.5	28.8 - 38.6	600
Filipino	2,000	66.5	63.8 - 69.0	1,000	33.5	31.0 - 36.2	3,000
Japanese	1,400	68.0	64.6 - 71.1	700	32.0	28.9 - 35.4	2,000
Other	1,300	60.3	56.8 - 63.7	800	39.7	36.3 - 43.2	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	66.7	62.6 - 70.5	400	33.3	29.5 - 37.4	1,200
20-24	2,300	62.5	60.2 - 64.9	1,400	37.5	35.1 - 39.8	3,700
25-34	4,400	58.0	56.3 - 59.7	3,200	42.0	40.3 - 43.7	7,500
35+	1,600	59.2	56.4 - 61.9	1,100	40.8	38.1 - 43.6	2,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,800	61.3	59.6 - 62.8	3,000	38.7	37.2 - 40.4	7,800
Female	4,300	58.7	57.0 - 60.4	3,000	41.3	39.6 - 43.0	7,300

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,900	57.5	56.1 - 59.0	4,300	42.5	41.0 - 43.9	10,200
No	3,200	65.2	63.3 - 67.2	1,700	34.8	32.8 - 36.7	4,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	62.7	60.8 - 64.6	2,100	37.3	35.4 - 39.2	5,500
131-185%	1,200	57.0	53.7 - 60.3	900	43.0	39.7 - 46.3	2,100
186+%	3,700	57.9	56.1 - 59.6	2,700	42.1	40.4 - 43.9	6,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight							
Low birth weight	700	68.8	64.4 - 72.9	300	31.2	27.1 - 35.6	1,000
Normal birth weight	8,400	59.4	58.2 - 60.6	5,700	40.6	39.4 - 41.8	14,100

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester							
Yes	700	65.4	61.0 - 69.6	400	34.6	30.4 - 39.0	1,000
No	8,300	59.7	58.4 - 60.9	5,600	40.3	39.1 - 41.6	13,900

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant							
Yes	1,900	65.1	62.5 - 67.7	1,000	34.9	32.3 - 37.5	2,900
No	7,000	58.8	57.4 - 60.1	4,900	41.2	39.9 - 42.6	12,000

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,800	62.0	60.2 - 63.8	2,400	38.0	36.2 - 39.8	6,200
No	5,100	58.4	56.8 - 60.0	3,600	41.6	40.0 - 43.2	8,700

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,800	59.8	58.6 - 61.0	5,900	40.2	39.0 - 41.4	14,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	900	54.2	51.4 - 57.0	800	45.8	43.0 - 48.6	1,700
Honolulu	6,700	62.1	60.6 - 63.5	4,100	37.9	36.5 - 39.4	10,800
Kauai	300	44.6	40.4 - 48.9	400	55.4	51.1 - 59.6	700
Maui	900	56.7	53.9 - 59.5	700	43.3	40.5 - 46.1	1,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,500	44.4	41.8 - 47.0	1,800	55.6	53.0 - 58.2	3,300
Native Hawaiian	2,400	62.7	60.9 - 64.5	1,400	37.3	35.5 - 39.1	3,900
Chinese	400	66.5	60.8 - 71.7	200	33.5	28.3 - 39.2	600
Filipino	2,000	66.0	63.3 - 68.6	1,000	34.0	31.4 - 36.7	3,000
Japanese	1,400	68.7	65.3 - 71.8	600	31.3	28.2 - 34.7	2,100
Other	1,200	59.5	55.9 - 63.1	800	40.5	36.9 - 44.1	2,000

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	68.6	64.9 - 72.1	400	31.4	27.9 - 35.1	1,200
20-24	2,200	61.7	59.4 - 64.0	1,400	38.3	36.0 - 40.6	3,600
25-34	4,200	57.6	55.9 - 59.3	3,100	42.4	40.7 - 44.1	7,300
35+	1,500	59.2	56.4 - 62.0	1,100	40.8	38.0 - 43.6	2,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,600	60.5	58.9 - 62.1	3,000	39.5	37.9 - 41.1	7,600
Female	4,200	59.1	57.4 - 60.7	2,900	40.9	39.3 - 42.6	7,100

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,800	57.4	56.0 - 58.9	4,300	42.6	41.1 - 44.0	10,100
No	3,000	65.0	63.1 - 66.9	1,600	35.0	33.1 - 36.9	4,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,400	62.2	60.3 - 64.0	2,100	37.8	36.0 - 39.7	5,400
131-185%	1,300	56.6	53.5 - 59.6	1,000	43.4	40.4 - 46.5	2,300
186+%	3,400	57.8	55.9 - 59.6	2,500	42.2	40.4 - 44.1	5,800

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight							
Low birth weight	700	69.8	65.3 - 73.9	300	30.2	26.1 - 34.7	1,000
Normal birth weight	8,100	59.1	57.9 - 60.3	5,600	40.9	39.7 - 42.1	13,800

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester							
Yes	700	66.0	61.8 - 70.0	300	34.0	30.0 - 38.2	1,000
No	8,000	59.4	58.2 - 60.6	5,500	40.6	39.4 - 41.8	13,500

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant							
Yes	1,800	64.2	61.6 - 66.7	1,000	35.8	33.3 - 38.4	2,800
No	6,800	58.7	57.4 - 60.0	4,800	41.3	40.0 - 42.6	11,600

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,700	61.4	59.7 - 63.2	2,300	38.6	36.8 - 40.3	6,000
No	5,000	58.5	57.0 - 60.1	3,500	41.5	39.9 - 43.0	8,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,300	58.8	57.6 - 60.0	5,800	41.2	40.0 - 42.4	14,100

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	800	53.5	50.9 - 56.1	700	46.5	43.9 - 49.1	1,600
Honolulu	6,300	60.6	59.1 - 62.1	4,100	39.4	37.9 - 40.9	10,300
Kauai	300	44.7	40.7 - 48.8	400	55.3	51.2 - 59.3	600
Maui	900	58.0	55.3 - 60.6	600	42.0	39.4 - 44.7	1,500

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,400	44.4	41.8 - 47.1	1,800	55.6	52.9 - 58.2	3,200
Native Hawaiian	2,300	62.7	61.1 - 64.2	1,400	37.3	35.8 - 38.9	3,700
Chinese	400	64.6	58.3 - 70.4	200	35.4	29.6 - 41.7	600
Filipino	1,900	64.3	61.5 - 67.0	1,000	35.7	33.0 - 38.5	2,900
Japanese	1,300	66.4	62.8 - 69.8	600	33.6	30.2 - 37.2	1,900
Other	1,100	57.9	54.1 - 61.6	800	42.1	38.4 - 45.9	1,900

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	67.7	64.2 - 71.1	400	32.3	28.9 - 35.8	1,300
20-24	2,100	60.4	58.1 - 62.6	1,400	39.6	37.4 - 41.9	3,500
25-34	3,900	56.5	54.7 - 58.2	3,000	43.5	41.8 - 45.3	6,900
35+	1,400	58.7	55.7 - 61.6	1,000	41.3	38.4 - 44.3	2,400

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,300	59.3	57.7 - 60.9	3,000	40.7	39.1 - 42.3	7,300
Female	3,900	58.3	56.6 - 60.0	2,800	41.7	40.0 - 43.4	6,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,500	56.4	54.9 - 57.9	4,200	43.6	42.1 - 45.1	9,700
No	2,800	64.3	62.3 - 66.1	1,500	35.7	33.9 - 37.7	4,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,300	60.8	58.9 - 62.6	2,100	39.2	37.4 - 41.1	5,400
131-185%	1,300	56.2	53.2 - 59.1	1,000	43.8	40.9 - 46.8	2,400
186+%	3,000	56.7	54.7 - 58.7	2,300	43.3	41.3 - 45.3	5,300

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Birthweight							
Low birth weight	600	68.7	64.1 - 73.0	300	31.3	27.0 - 35.9	900
Normal birth weight	7,600	58.1	56.9 - 59.4	5,500	41.9	40.6 - 43.1	13,200

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke 3rd Trimester							
Yes	700	67.5	63.4 - 71.4	300	32.5	28.6 - 36.6	1,000
No	7,500	58.3	57.1 - 59.5	5,400	41.7	40.5 - 42.9	12,900

	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Smoke Before Pregnant							
Yes	1,700	63.1	60.5 - 65.6	1,000	36.9	34.4 - 39.5	2,600
No	6,400	57.8	56.4 - 59.1	4,700	42.2	40.9 - 43.6	11,100

PRAMS Health Indicator 5-Year Aggregate Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,500	60.7	58.9 - 62.4	2,300	39.3	37.6 - 41.1	5,800
No	4,700	57.5	55.9 - 59.1	3,400	42.5	40.9 - 44.1	8,100