

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (any) duration 8+ weeks, among initiators	Breastfeeding (any) duration 8+ weeks, among initiators

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/13/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


**PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	3,300	23.3	21.4 - 25.4	10,800	76.7	74.6 - 78.6	14,000
2001	3,400	23.9	21.9 - 26.0	10,700	76.1	74.0 - 78.1	14,000
2002	3,200	21.9	19.6 - 24.5	11,200	78.1	75.5 - 80.4	14,400
2003	3,600	24.3	21.8 - 26.8	11,200	75.7	73.2 - 78.2	14,800
2004	3,700	23.5	21.6 - 25.5	12,000	76.5	74.5 - 78.4	15,700
2005	3,500	22.9	20.8 - 25.0	11,800	77.1	75.0 - 79.2	15,300
2006	3,800	23.2	21.0 - 25.5	12,500	76.8	74.5 - 79.0	16,300
2007	3,800	23.1	21.1 - 25.3	12,700	76.9	74.7 - 78.9	16,600
2008	3,600	21.4	19.4 - 23.5	13,200	78.6	76.5 - 80.6	16,700
2009	4,100	24.7	21.9 - 27.7	12,600	75.3	72.3 - 78.1	16,700
2010	3,700	21.3	18.7 - 24.2	13,500	78.7	75.8 - 81.3	17,200
2011	3,200	18.1	15.8 - 20.8	14,300	81.9	79.2 - 84.2	17,500
2012	3,100	18.0	15.5 - 20.7	14,200	82.0	79.3 - 84.5	17,300
2013	2,900	16.7	14.2 - 19.6	14,500	83.3	80.4 - 85.8	17,400

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	2,900	16.7	14.2 - 19.6	14,500	83.3	80.4 - 85.8	17,400

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	14.7	11.0 - 19.3	1,900	85.3	80.7 - 89.0	2,200
Honolulu	2,100	16.9	13.6 - 20.8	10,400	83.1	79.2 - 86.4	12,500
Kauai	100	16.4	12.8 - 20.9	700	83.6	79.1 - 87.2	800
Maui	300	17.9	14.0 - 22.7	1,500	82.1	77.3 - 86.0	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	14.4	10.0 - 20.4	4,100	85.6	79.6 - 90.0	4,800
Native Hawaiian	1,200	23.6	18.6 - 29.6	3,800	76.4	70.4 - 81.4	5,000
Chinese	200	17.9	8.5 - 33.8	700	82.1	66.2 - 91.5	900
Filipino	600	19.9	14.0 - 27.5	2,200	80.1	72.5 - 86.0	2,800
Japanese	100	3.0	0.9 - 9.3	1,700	97.0	90.7 - 99.1	1,800
Other	300	12.5	7.1 - 21.1	1,900	87.5	78.9 - 92.9	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	29.5	17.6 - 45.0	600	70.5	55.0 - 82.4	800
20-24	700	16.9	11.9 - 23.3	3,300	83.1	76.7 - 88.1	3,900
25-34	1,600	16.6	13.2 - 20.7	7,900	83.4	79.3 - 86.8	9,400
35+	400	13.7	9.1 - 20.2	2,800	86.3	79.8 - 90.9	3,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,300	14.8	11.6 - 18.7	7,200	85.2	81.3 - 88.4	8,500
Female	1,600	18.5	14.9 - 22.8	7,300	81.5	77.2 - 85.1	8,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	15.3	12.2 - 19.0	9,500	84.7	81.0 - 87.8	11,200
No	1,200	19.4	15.3 - 24.1	5,000	80.6	75.9 - 84.7	6,100

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,400	21.9	17.4 - 27.2	4,900	78.1	72.8 - 82.6	6,300
131-185%	400	15.4	9.5 - 24.0	1,900	84.6	76.0 - 90.5	2,300
186+%	800	11.6	8.4 - 15.8	6,300	88.4	84.2 - 91.6	7,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Income (2012+)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	900	23.4	17.6 - 30.3	3,000	76.6	69.7 - 82.4	3,900
\$17,001-\$26,000	400	17.5	11.1 - 26.3	1,900	82.5	73.7 - 88.9	2,300
\$26,001-\$51,000	600	16.9	11.8 - 23.6	3,100	83.1	76.4 - 88.2	3,700
>=\$51,001	600	10.5	7.3 - 14.9	5,400	89.5	85.1 - 92.7	6,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	20.3	16.6 - 24.6	1,000	79.7	75.4 - 83.4	1,200
Normal birth weight	2,700	16.4	13.8 - 19.5	13,500	83.6	80.5 - 86.2	16,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	30.1	18.4 - 45.1	600	69.9	54.9 - 81.6	900
No	2,600	16.1	13.5 - 19.0	13,500	83.9	81.0 - 86.5	16,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	25.4	18.7 - 33.6	2,300	74.6	66.4 - 81.3	3,100
No	2,100	14.9	12.3 - 18.0	11,800	85.1	82.0 - 87.7	13,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	23.0	18.7 - 28.0	5,700	77.0	72.0 - 81.3	7,400
No	1,200	11.8	9.0 - 15.2	8,800	88.2	84.8 - 91.0	9,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,100	18.0	15.5 - 20.7	14,200	82.0	79.3 - 84.5	17,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	19.4	15.4 - 24.2	1,800	80.6	75.8 - 84.6	2,200
Honolulu	2,200	17.7	14.5 - 21.4	10,200	82.3	78.6 - 85.5	12,400
Kauai	100	16.9	12.5 - 22.4	700	83.1	77.6 - 87.5	800
Maui	400	18.8	15.1 - 23.3	1,500	81.2	76.7 - 84.9	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	500	12.9	8.7 - 18.7	3,400	87.1	81.3 - 91.3	3,900
Native Hawaiian	1,300	25.7	20.7 - 31.4	3,600	74.3	68.6 - 79.3	4,900
Chinese	100	10.8	4.4 - 24.3	900	89.2	75.7 - 95.6	1,000
Filipino	800	26.1	19.6 - 33.8	2,300	73.9	66.2 - 80.4	3,100
Japanese	100	6.9	3.3 - 13.8	1,400	93.1	86.2 - 96.7	1,500
Other	300	11.1	6.7 - 17.9	2,500	88.9	82.1 - 93.3	2,800

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	36.3	24.2 - 50.5	600	63.7	49.5 - 75.8	1,000
20-24	800	23.6	17.8 - 30.7	2,700	76.4	69.3 - 82.2	3,600
25-34	1,600	16.5	13.4 - 20.1	8,100	83.5	79.9 - 86.6	9,700
35+	300	10.3	6.7 - 15.3	2,700	89.7	84.7 - 93.3	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,600	18.7	15.3 - 22.8	7,000	81.3	77.2 - 84.7	8,700
Female	1,500	17.2	13.9 - 21.1	7,100	82.8	78.9 - 86.1	8,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	14.5	11.7 - 17.9	9,200	85.5	82.1 - 88.3	10,700
No	1,500	23.7	19.4 - 28.5	5,000	76.3	71.5 - 80.6	6,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,400	23.6	19.2 - 28.7	4,600	76.4	71.3 - 80.8	6,000
131-185%	600	20.6	14.5 - 28.4	2,300	79.4	71.6 - 85.5	2,900
186+%	800	12.3	9.1 - 16.3	6,000	87.7	83.7 - 90.9	6,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Income (2012+)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	900	23.4	17.9 - 29.9	2,900	76.6	70.1 - 82.1	3,800
\$17,001-\$26,000	500	22.5	15.8 - 31.0	1,700	77.5	69.0 - 84.2	2,200
\$26,001-\$51,000	900	21.3	15.9 - 27.8	3,200	78.7	72.2 - 84.1	4,000
>=\$51,001	700	11.2	8.0 - 15.4	5,200	88.8	84.6 - 92.0	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	20.2	15.8 - 25.4	900	79.8	74.6 - 84.2	1,100
Normal birth weight	2,900	17.9	15.3 - 20.8	13,200	82.1	79.2 - 84.7	16,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	44.0	28.9 - 60.4	400	56.0	39.6 - 71.1	700
No	2,800	17.3	14.8 - 20.1	13,200	82.7	79.9 - 85.2	16,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race- Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	28.2	21.2 - 36.4	1,900	71.8	63.6 - 78.8	2,700
No	2,300	16.4	13.8 - 19.4	11,700	83.6	80.6 - 86.2	14,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	25.5	21.4 - 30.2	5,700	74.5	69.8 - 78.6	7,600
No	1,100	11.6	9.0 - 14.9	8,400	88.4	85.1 - 91.0	9,500

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,200	18.1	15.8 - 20.8	14,300	81.9	79.2 - 84.2	17,500

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	21.9	17.8 - 26.7	1,800	78.1	73.3 - 82.2	2,300
Honolulu	2,200	17.4	14.3 - 21.0	10,300	82.6	79.0 - 85.7	12,500
Kauai	100	14.0	10.1 - 19.0	700	86.0	81.0 - 89.9	800
Maui	400	20.4	16.8 - 24.5	1,500	79.6	75.5 - 83.2	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	16.4	11.9 - 22.3	3,600	83.6	77.7 - 88.1	4,400
Native Hawaiian	1,200	23.5	18.9 - 28.9	3,800	76.5	71.1 - 81.1	4,900
Chinese	100	8.0	2.5 - 22.4	600	92.0	77.6 - 97.5	600
Filipino	600	19.9	14.8 - 26.3	2,500	80.1	73.7 - 85.2	3,100
Japanese	200	12.4	6.8 - 21.5	1,600	87.6	78.5 - 93.2	1,800
Other	400	15.2	9.7 - 22.9	2,200	84.8	77.1 - 90.3	2,600

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	45.6	33.4 - 58.3	500	54.4	41.7 - 66.6	1,000
20-24	800	20.5	15.4 - 26.6	3,200	79.5	73.4 - 84.6	4,000
25-34	1,600	17.1	13.8 - 20.8	7,800	82.9	79.2 - 86.2	9,400
35+	300	9.8	6.4 - 14.7	2,800	90.2	85.3 - 93.6	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,700	18.9	15.5 - 22.9	7,200	81.1	77.1 - 84.5	8,900
Female	1,500	17.4	14.2 - 21.1	7,100	82.6	78.9 - 85.8	8,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	15.6	12.8 - 19.0	9,200	84.4	81.0 - 87.2	10,900
No	1,500	22.3	18.3 - 27.0	5,100	77.7	73.0 - 81.7	6,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,400	22.9	18.7 - 27.6	4,700	77.1	72.4 - 81.3	6,100
131-185%	500	18.5	12.6 - 26.3	2,100	81.5	73.7 - 87.4	2,600
186+%	1,100	13.8	10.6 - 17.7	6,700	86.2	82.3 - 89.4	7,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	700	23.6	17.9 - 30.5	2,400	76.4	69.5 - 82.1	3,100
\$10,000-\$24,999	700	22.0	16.4 - 28.8	2,500	78.0	71.2 - 83.6	3,200
\$25,000-\$49,999	900	20.6	15.5 - 26.8	3,400	79.4	73.2 - 84.5	4,200
>=\$50,000	700	11.2	8.1 - 15.3	5,300	88.8	84.7 - 91.9	6,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	26.3	21.5 - 31.7	900	73.7	68.3 - 78.5	1,200
Normal birth weight	2,900	17.5	15.0 - 20.4	13,400	82.5	79.6 - 85.0	16,300

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	42.9	30.2 - 56.6	500	57.1	43.4 - 69.8	800
No	2,700	16.6	14.2 - 19.3	13,700	83.4	80.7 - 85.8	16,500

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race- Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	27.5	21.5 - 34.5	2,400	72.5	65.5 - 78.5	3,300
No	2,200	15.6	13.1 - 18.5	11,800	84.4	81.5 - 86.9	14,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	22.0	18.3 - 26.2	6,000	78.0	73.8 - 81.7	7,800
No	1,400	14.9	11.9 - 18.4	8,100	85.1	81.6 - 88.1	9,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,700	21.3	18.7 - 24.2	13,500	78.7	75.8 - 81.3	17,200

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	22.0	17.9 - 26.7	1,700	78.0	73.3 - 82.1	2,200
Honolulu	2,700	21.7	18.2 - 25.6	9,700	78.3	74.4 - 81.8	12,300
Kauai	100	14.1	10.4 - 18.8	700	85.9	81.2 - 89.6	800
Maui	400	20.7	17.0 - 24.9	1,500	79.3	75.1 - 83.0	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	20.1	14.9 - 26.5	3,300	79.9	73.5 - 85.1	4,200
Native Hawaiian	1,200	24.4	19.8 - 29.8	3,800	75.6	70.2 - 80.2	5,100
Chinese	100	13.0	5.2 - 28.8	600	87.0	71.2 - 94.8	700
Filipino	900	28.1	21.6 - 35.5	2,200	71.9	64.5 - 78.4	3,000
Japanese	100	6.3	3.1 - 12.7	1,600	93.7	87.3 - 96.9	1,700
Other	500	21.0	14.4 - 29.7	1,900	79.0	70.3 - 85.6	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	39.6	28.0 - 52.6	600	60.4	47.4 - 72.0	1,100
20-24	1,100	25.9	20.3 - 32.4	3,000	74.1	67.6 - 79.7	4,100
25-34	1,700	18.8	15.4 - 22.7	7,200	81.2	77.3 - 84.6	8,900
35+	500	16.1	11.2 - 22.5	2,600	83.9	77.5 - 88.8	3,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,900	22.0	18.3 - 26.1	6,800	78.0	73.9 - 81.7	8,700
Female	1,800	20.6	17.0 - 24.7	6,700	79.4	75.3 - 83.0	8,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	17.7	14.6 - 21.2	9,000	82.3	78.8 - 85.4	10,900
No	1,700	27.7	23.1 - 32.7	4,500	72.3	67.3 - 76.9	6,200

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,500	26.0	21.5 - 31.1	4,300	74.0	68.9 - 78.5	5,900
131-185%	700	24.2	17.5 - 32.4	2,100	75.8	67.6 - 82.5	2,800
186+%	1,100	15.0	11.7 - 19.2	6,300	85.0	80.8 - 88.3	7,400

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	900	28.4	22.2 - 35.6	2,300	71.6	64.4 - 77.8	3,100
\$10,000-\$24,999	700	22.7	17.0 - 29.7	2,300	77.3	70.3 - 83.0	2,900
\$25,000-\$49,999	1,000	24.2	18.7 - 30.7	3,200	75.8	69.3 - 81.3	4,300
>=\$50,000	800	14.0	10.4 - 18.7	5,100	86.0	81.3 - 89.6	6,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	24.4	19.7 - 29.7	900	75.6	70.3 - 80.3	1,200
Normal birth weight	3,400	21.1	18.3 - 24.1	12,600	78.9	75.9 - 81.7	16,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	31.2	20.6 - 44.2	800	68.8	55.8 - 79.4	1,100
No	3,300	20.8	18.1 - 23.7	12,600	79.2	76.3 - 81.9	15,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	35.2	28.5 - 42.6	2,200	64.8	57.4 - 71.5	3,400
No	2,400	18.0	15.3 - 21.1	11,100	82.0	78.9 - 84.7	13,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	26.0	21.9 - 30.6	5,500	74.0	69.4 - 78.1	7,400
No	1,700	17.9	14.6 - 21.7	7,900	82.1	78.3 - 85.4	9,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	4,100	24.7	21.9 - 27.7	12,600	75.3	72.3 - 78.1	16,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	600	26.3	21.8 - 31.3	1,600	73.7	68.7 - 78.2	2,200
Honolulu	2,900	25.0	21.3 - 29.1	8,800	75.0	70.9 - 78.7	11,800
Kauai	200	19.7	15.2 - 25.2	600	80.3	74.8 - 84.8	800
Maui	500	23.3	19.5 - 27.6	1,500	76.7	72.4 - 80.5	2,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	17.8	13.2 - 23.6	3,400	82.2	76.4 - 86.8	4,100
Native Hawaiian	1,600	31.2	26.0 - 36.9	3,500	68.8	63.1 - 74.0	5,100
Chinese	100	10.5	3.9 - 25.6	600	89.5	74.4 - 96.1	700
Filipino	900	28.6	22.2 - 35.8	2,200	71.4	64.2 - 77.8	3,000
Japanese	200	15.6	9.1 - 25.4	1,300	84.4	74.6 - 90.9	1,600
Other	600	27.9	20.0 - 37.4	1,600	72.1	62.6 - 80.0	2,300

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	42.4	32.2 - 53.4	900	57.6	46.6 - 67.8	1,600
20-24	1,200	30.8	24.8 - 37.5	2,600	69.2	62.5 - 75.2	3,800
25-34	1,700	20.1	16.6 - 24.1	6,700	79.9	75.9 - 83.4	8,400
35+	600	19.8	14.0 - 27.2	2,300	80.2	72.8 - 86.0	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,100	23.9	20.1 - 28.0	6,700	76.1	72.0 - 79.9	8,800
Female	2,000	25.6	21.7 - 30.0	5,900	74.4	70.0 - 78.3	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.4	15.3 - 22.0	8,300	81.6	78.0 - 84.7	10,200
No	2,300	34.4	29.6 - 39.6	4,300	65.6	60.4 - 70.4	6,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,800	32.2	27.2 - 37.8	3,800	67.8	62.2 - 72.8	5,600
131-185%	800	28.6	21.7 - 36.7	1,900	71.4	63.3 - 78.3	2,600
186+%	1,200	16.7	13.2 - 21.0	5,900	83.3	79.0 - 86.8	7,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,000	36.0	28.2 - 44.5	1,700	64.0	55.5 - 71.8	2,700
\$10,000-\$24,999	1,000	29.7	23.8 - 36.4	2,400	70.3	63.6 - 76.2	3,500
\$25,000-\$49,999	800	23.7	17.9 - 30.6	2,600	76.3	69.4 - 82.1	3,400
>=\$50,000	1,000	16.4	12.6 - 21.1	4,900	83.6	78.9 - 87.4	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	28.0	23.2 - 33.4	800	72.0	66.6 - 76.8	1,100
Normal birth weight	3,800	24.4	21.5 - 27.6	11,800	75.6	72.4 - 78.5	15,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	44.2	33.2 - 55.8	800	55.8	44.2 - 66.8	1,400
No	3,400	22.5	19.7 - 25.6	11,600	77.5	74.4 - 80.3	15,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	37.0	30.3 - 44.3	2,100	63.0	55.7 - 69.7	3,400
No	2,700	21.0	18.1 - 24.3	10,200	79.0	75.7 - 81.9	13,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	34.5	29.8 - 39.5	4,300	65.5	60.5 - 70.2	6,500
No	1,700	18.0	14.8 - 21.8	7,900	82.0	78.2 - 85.2	9,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,600	21.4	19.4 - 23.5	13,200	78.6	76.5 - 80.6	16,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	22.7	17.2 - 29.5	1,600	77.3	70.5 - 82.8	2,100
Honolulu	2,600	21.7	19.4 - 24.3	9,300	78.3	75.7 - 80.6	11,900
Kauai	200	23.5	15.3 - 34.3	600	76.5	65.7 - 84.7	800
Maui	300	16.8	11.9 - 23.2	1,600	83.2	76.8 - 88.1	1,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	18.8	15.2 - 23.0	3,400	81.2	77.0 - 84.8	4,100
Native Hawaiian	1,200	25.8	21.5 - 30.7	3,400	74.2	69.3 - 78.5	4,500
Chinese	100	13.6	9.6 - 19.1	500	86.4	80.9 - 90.4	600
Filipino	800	25.0	20.5 - 30.1	2,300	75.0	69.9 - 79.5	3,000
Japanese	300	14.6	10.1 - 20.7	1,700	85.4	79.3 - 89.9	2,000
Other	500	20.5	15.6 - 26.5	2,000	79.5	73.5 - 84.4	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	27.2	19.6 - 36.5	900	72.8	63.5 - 80.4	1,300
20-24	1,200	32.3	27.3 - 37.7	2,400	67.7	62.3 - 72.7	3,600
25-34	1,500	17.7	15.2 - 20.5	7,100	82.3	79.5 - 84.8	8,700
35+	500	16.8	13.0 - 21.5	2,700	83.2	78.5 - 87.0	3,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	22.5	19.7 - 25.6	6,800	77.5	74.4 - 80.3	8,800
Female	1,600	20.1	17.4 - 23.2	6,300	79.9	76.8 - 82.6	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	19.9	17.6 - 22.5	8,600	80.1	77.5 - 82.4	10,800
No	1,400	24.1	20.5 - 28.0	4,500	75.9	72.0 - 79.5	6,000

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,500	25.3	21.7 - 29.4	4,400	74.7	70.6 - 78.3	5,800
131-185%	400	15.7	11.4 - 21.3	2,000	84.3	78.7 - 88.6	2,400
186+%	1,400	18.5	15.8 - 21.5	6,100	81.5	78.5 - 84.2	7,400

**PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-
Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income
(2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	700	28.0	22.5 - 34.4	1,900	72.0	65.6 - 77.5	2,600
\$10,000-\$24,999	800	24.3	19.8 - 29.5	2,600	75.7	70.5 - 80.2	3,500
\$25,000-\$49,999	900	22.3	18.3 - 26.9	3,000	77.7	73.1 - 81.7	3,800
>=\$50,000	900	14.7	12.0 - 17.9	5,000	85.3	82.1 - 88.0	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	21.5	14.5 - 30.6	900	78.5	69.4 - 85.5	1,100
Normal birth weight	3,300	21.4	19.3 - 23.6	12,300	78.6	76.4 - 80.7	15,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	35.6	27.1 - 45.2	800	64.4	54.8 - 72.9	1,300
No	3,100	20.2	18.2 - 22.4	12,100	79.8	77.6 - 81.8	15,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	31.9	26.4 - 37.9	2,000	68.1	62.1 - 73.6	2,900
No	2,600	19.1	17.0 - 21.4	10,900	80.9	78.6 - 83.0	13,500

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	26.5	23.1 - 30.3	5,000	73.5	69.7 - 76.9	6,700
No	1,800	18.0	15.6 - 20.6	8,000	82.0	79.4 - 84.4	9,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,800	23.1	21.1 - 25.3	12,700	76.9	74.7 - 78.9	16,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	16.6	11.8 - 22.9	1,800	83.4	77.1 - 88.2	2,100
Honolulu	2,800	24.3	21.8 - 26.9	8,800	75.7	73.1 - 78.2	11,600
Kauai	200	19.6	12.4 - 29.6	700	80.4	70.4 - 87.6	900
Maui	500	25.0	19.0 - 32.3	1,500	75.0	67.7 - 81.0	2,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	18.8	15.0 - 23.3	3,000	81.2	76.7 - 85.0	3,700
Native Hawaiian	1,500	33.2	28.5 - 38.3	3,000	66.8	61.7 - 71.5	4,600
Chinese	100	12.2	8.3 - 17.7	600	87.8	82.3 - 91.7	700
Filipino	800	27.0	22.4 - 32.1	2,200	73.0	67.9 - 77.6	3,000
Japanese	200	10.8	7.0 - 16.2	1,800	89.2	83.8 - 93.0	2,000
Other	500	19.6	14.9 - 25.4	2,100	80.4	74.6 - 85.1	2,600

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	38.5	29.6 - 48.3	900	61.5	51.7 - 70.4	1,500
20-24	1,200	30.1	25.6 - 35.1	2,800	69.9	64.9 - 74.4	4,100
25-34	1,600	20.1	17.4 - 23.1	6,500	79.9	76.9 - 82.6	8,100
35+	400	14.1	10.6 - 18.6	2,500	85.9	81.4 - 89.4	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	23.4	20.6 - 26.5	6,600	76.6	73.5 - 79.4	8,600
Female	1,800	22.8	19.9 - 26.0	6,100	77.2	74.0 - 80.1	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	16.9	14.8 - 19.4	8,700	83.1	80.6 - 85.2	10,500
No	2,000	33.9	29.9 - 38.2	4,000	66.1	61.8 - 70.1	6,000

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,900	32.5	28.5 - 36.7	4,000	67.5	63.3 - 71.5	6,000
131-185%	500	21.1	16.2 - 26.9	1,900	78.9	73.1 - 83.8	2,500
186+%	1,100	16.1	13.5 - 19.0	5,800	83.9	81.0 - 86.5	7,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,000	37.0	30.8 - 43.6	1,700	63.0	56.4 - 69.2	2,700
\$10,000-\$24,999	900	27.6	22.8 - 32.9	2,400	72.4	67.1 - 77.2	3,300
\$25,000-\$49,999	900	23.0	18.9 - 27.7	2,900	77.0	72.3 - 81.1	3,800
>=\$50,000	800	14.2	11.6 - 17.4	4,800	85.8	82.6 - 88.4	5,600

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	400	28.7	21.3 - 37.4	1,000	71.3	62.6 - 78.7	1,300
Normal birth weight	3,400	22.6	20.5 - 24.9	11,800	77.4	75.1 - 79.5	15,200

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	47.5	38.1 - 57.1	700	52.5	42.9 - 61.9	1,300
No	3,200	21.3	19.3 - 23.6	11,900	78.7	76.4 - 80.7	15,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	40.7	35.0 - 46.7	1,800	59.3	53.3 - 65.0	3,100
No	2,600	19.3	17.2 - 21.6	10,700	80.7	78.4 - 82.8	13,300

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	29.3	25.7 - 33.2	4,600	70.7	66.8 - 74.3	6,600
No	1,800	18.7	16.4 - 21.4	7,900	81.3	78.6 - 83.6	9,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,800	23.2	21.0 - 25.5	12,500	76.8	74.5 - 79.0	16,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	24.7	18.7 - 32.0	1,500	75.3	68.0 - 81.3	2,000
Honolulu	2,700	23.0	20.5 - 25.7	9,200	77.0	74.3 - 79.5	11,900
Kauai	100	19.5	11.4 - 31.3	600	80.5	68.7 - 88.6	800
Maui	400	24.5	18.1 - 32.4	1,300	75.5	67.6 - 81.9	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	800	20.9	16.9 - 25.6	3,100	79.1	74.4 - 83.1	3,900
Native Hawaiian	1,300	30.0	25.3 - 35.3	3,100	70.0	64.7 - 74.7	4,400
Chinese	100	15.6	11.1 - 21.5	500	84.4	78.5 - 88.9	600
Filipino	800	25.4	20.7 - 30.7	2,200	74.6	69.3 - 79.3	3,000
Japanese	200	10.5	6.7 - 16.0	1,700	89.5	84.0 - 93.3	1,900
Other	600	23.7	18.3 - 30.1	1,900	76.3	69.9 - 81.7	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	37.9	29.0 - 47.7	800	62.1	52.3 - 71.0	1,300
20-24	1,300	33.2	28.2 - 38.6	2,700	66.8	61.4 - 71.8	4,000
25-34	1,600	19.5	16.7 - 22.5	6,500	80.5	77.5 - 83.3	8,100
35+	400	13.0	9.6 - 17.4	2,500	87.0	82.6 - 90.4	2,900

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,200	26.8	23.7 - 30.2	6,100	73.2	69.8 - 76.3	8,400
Female	1,500	19.3	16.5 - 22.5	6,400	80.7	77.5 - 83.5	8,000

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	17.5	15.2 - 20.0	8,600	82.5	80.0 - 84.8	10,500
No	2,000	33.3	29.1 - 37.8	3,900	66.7	62.2 - 70.9	5,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,800	32.9	28.7 - 37.5	3,700	67.1	62.5 - 71.3	5,500
131-185%	400	20.9	15.5 - 27.6	1,700	79.1	72.4 - 84.5	2,100
186+%	1,300	16.8	14.3 - 19.8	6,300	83.2	80.2 - 85.7	7,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,000	37.4	31.0 - 44.2	1,700	62.6	55.8 - 69.0	2,700
\$10,000-\$24,999	900	28.0	23.0 - 33.6	2,400	72.0	66.4 - 77.0	3,400
\$25,000-\$49,999	1,000	25.5	21.1 - 30.4	2,900	74.5	69.6 - 78.9	3,900
>=\$50,000	600	11.7	9.2 - 14.7	4,800	88.3	85.3 - 90.8	5,400

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	21.7	14.8 - 30.5	1,000	78.3	69.5 - 85.2	1,300
Normal birth weight	3,500	23.3	21.1 - 25.7	11,500	76.7	74.3 - 78.9	15,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	35.8	26.8 - 45.9	800	64.2	54.1 - 73.2	1,200
No	3,300	22.0	19.8 - 24.3	11,600	78.0	75.7 - 80.2	14,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	31.3	25.9 - 37.3	2,200	68.7	62.7 - 74.1	3,200
No	2,700	21.0	18.7 - 23.5	10,200	79.0	76.5 - 81.3	12,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	30.9	27.1 - 35.0	4,600	69.1	65.0 - 72.9	6,700
No	1,600	17.5	15.1 - 20.2	7,700	82.5	79.8 - 84.9	9,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,500	22.9	20.8 - 25.0	11,800	77.1	75.0 - 79.2	15,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	20.0	14.7 - 26.7	1,500	80.0	73.3 - 85.3	1,900
Honolulu	2,500	22.6	20.3 - 25.2	8,700	77.4	74.8 - 79.7	11,200
Kauai	100	13.9	7.5 - 24.4	600	86.1	75.6 - 92.5	700
Maui	500	31.9	25.0 - 39.7	1,000	68.1	60.3 - 75.0	1,500

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	16.2	12.6 - 20.5	2,900	83.8	79.5 - 87.4	3,500
Native Hawaiian	1,200	30.0	25.3 - 35.1	2,900	70.0	64.9 - 74.7	4,100
Chinese	100	14.9	10.7 - 20.4	500	85.1	79.6 - 89.3	600
Filipino	900	30.8	26.1 - 36.0	2,000	69.2	64.0 - 73.9	2,900
Japanese	300	12.2	8.5 - 17.2	1,900	87.8	82.8 - 91.5	2,200
Other	500	22.4	17.3 - 28.6	1,600	77.6	71.4 - 82.7	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	36.4	28.2 - 45.6	800	63.6	54.4 - 71.8	1,300
20-24	1,300	35.1	30.3 - 40.2	2,400	64.9	59.8 - 69.7	3,700
25-34	1,400	18.4	15.8 - 21.3	6,200	81.6	78.7 - 84.2	7,600
35+	300	12.2	9.0 - 16.4	2,400	87.8	83.6 - 91.0	2,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,800	23.1	20.3 - 26.3	6,000	76.9	73.7 - 79.7	7,800
Female	1,700	22.6	19.7 - 25.7	5,800	77.4	74.3 - 80.3	7,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	17.8	15.6 - 20.3	8,200	82.2	79.7 - 84.4	10,000
No	1,700	32.3	28.3 - 36.5	3,600	67.7	63.5 - 71.7	5,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	31.6	27.7 - 35.8	3,600	68.4	64.2 - 72.3	5,300
131-185%	400	24.3	18.2 - 31.7	1,200	75.7	68.3 - 81.8	1,600
186+%	1,100	14.9	12.5 - 17.7	6,100	85.1	82.3 - 87.5	7,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,000	38.0	32.1 - 44.4	1,600	62.0	55.6 - 67.9	2,600
\$10,000-\$24,999	900	29.4	24.5 - 34.8	2,100	70.6	65.2 - 75.5	3,000
\$25,000-\$49,999	700	19.5	15.6 - 23.9	2,900	80.5	76.1 - 84.4	3,600
>=\$50,000	600	12.6	10.0 - 15.8	4,400	87.4	84.2 - 90.0	5,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	25.9	18.5 - 35.0	800	74.1	65.0 - 81.5	1,100
Normal birth weight	3,200	22.7	20.6 - 24.9	11,000	77.3	75.1 - 79.4	14,200

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	44.5	35.1 - 54.3	600	55.5	45.7 - 64.9	1,100
No	3,000	21.0	19.0 - 23.3	11,100	79.0	76.7 - 81.0	14,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	38.4	32.9 - 44.2	1,900	61.6	55.8 - 67.1	3,000
No	2,300	18.8	16.7 - 21.0	9,800	81.2	79.0 - 83.3	12,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	28.7	25.2 - 32.4	4,500	71.3	67.6 - 74.8	6,300
No	1,600	18.2	15.8 - 20.9	7,100	81.8	79.1 - 84.2	8,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,700	23.5	21.6 - 25.5	12,000	76.5	74.5 - 78.4	15,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	19.8	14.9 - 25.8	1,500	80.2	74.2 - 85.1	1,900
Honolulu	2,600	23.3	21.1 - 25.7	8,700	76.7	74.3 - 78.9	11,400
Kauai	200	21.1	13.7 - 31.0	600	78.9	69.0 - 86.3	700
Maui	500	30.3	24.4 - 37.0	1,200	69.7	63.0 - 75.6	1,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	18.4	15.0 - 22.3	3,000	81.6	77.7 - 85.0	3,600
Native Hawaiian	1,300	32.1	27.7 - 36.9	2,800	67.9	63.1 - 72.3	4,100
Chinese	100	12.0	9.3 - 15.5	500	88.0	84.5 - 90.7	600
Filipino	900	30.0	25.8 - 34.5	2,100	70.0	65.5 - 74.2	3,000
Japanese	300	12.3	8.6 - 17.3	1,900	87.7	82.7 - 91.4	2,100
Other	500	21.3	16.5 - 27.0	1,800	78.7	73.0 - 83.5	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	40.0	31.5 - 49.2	600	60.0	50.8 - 68.5	1,100
20-24	1,300	34.0	29.6 - 38.7	2,600	66.0	61.3 - 70.4	3,900
25-34	1,400	18.4	16.0 - 21.0	6,300	81.6	79.0 - 84.0	7,700
35+	500	17.1	13.5 - 21.5	2,500	82.9	78.5 - 86.5	3,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	23.9	21.4 - 26.7	6,500	76.1	73.3 - 78.6	8,500
Female	1,700	23.1	20.3 - 26.1	5,600	76.9	73.9 - 79.7	7,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	19.8	17.7 - 22.0	9,000	80.2	78.0 - 82.3	11,200
No	1,500	32.7	28.7 - 37.0	3,100	67.3	63.0 - 71.3	4,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,700	30.6	27.1 - 34.4	3,900	69.4	65.6 - 72.9	5,600
131-185%	400	21.4	16.3 - 27.5	1,500	78.6	72.5 - 83.7	1,900
186+%	1,300	18.2	15.8 - 20.9	5,900	81.8	79.1 - 84.2	7,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	800	34.8	29.1 - 40.9	1,500	65.2	59.1 - 70.9	2,400
\$10,000-\$24,999	1,100	29.2	25.0 - 33.8	2,600	70.8	66.2 - 75.0	3,700
\$25,000-\$49,999	900	21.6	18.1 - 25.5	3,200	78.4	74.5 - 81.9	4,100
>=\$50,000	600	14.2	11.5 - 17.3	3,900	85.8	82.7 - 88.5	4,500

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	26.1	18.9 - 35.0	700	73.9	65.0 - 81.1	1,000
Normal birth weight	3,500	23.4	21.4 - 25.4	11,300	76.6	74.6 - 78.6	14,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	31.4	23.5 - 40.6	700	68.6	59.4 - 76.5	1,100
No	3,300	22.7	20.8 - 24.8	11,200	77.3	75.2 - 79.2	14,500

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	28.6	23.9 - 33.7	2,200	71.4	66.3 - 76.1	3,000
No	2,800	22.0	20.0 - 24.2	9,800	78.0	75.8 - 80.0	12,500

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	27.6	24.5 - 31.0	5,000	72.4	69.0 - 75.5	6,900
No	1,800	20.4	18.1 - 23.0	6,800	79.6	77.0 - 81.9	8,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,600	24.3	21.8 - 26.8	11,200	75.7	73.2 - 78.2	14,800

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	400	25.2	21.0 - 29.8	1,200	74.8	70.2 - 79.0	1,500
Honolulu	2,700	24.4	21.3 - 27.8	8,200	75.6	72.2 - 78.7	10,900
Kauai	100	16.7	11.8 - 23.1	600	83.3	76.9 - 88.2	800
Maui	400	26.0	21.6 - 30.9	1,200	74.0	69.1 - 78.4	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	18.4	13.5 - 24.5	2,800	81.6	75.5 - 86.5	3,400
Native Hawaiian	1,200	31.2	28.2 - 34.4	2,600	68.8	65.6 - 71.8	3,800
Chinese	<50	7.4	2.3 - 21.7	600	92.6	78.3 - 97.7	700
Filipino	1,000	30.3	24.3 - 37.0	2,200	69.7	63.0 - 75.7	3,100
Japanese	200	13.9	8.3 - 22.4	1,500	86.1	77.6 - 91.7	1,800
Other	500	26.4	18.9 - 35.6	1,400	73.6	64.4 - 81.1	1,900

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	51.7	42.6 - 60.7	600	48.3	39.3 - 57.4	1,300
20-24	1,100	32.7	27.4 - 38.6	2,200	67.3	61.4 - 72.6	3,300
25-34	1,500	20.1	16.9 - 23.6	6,000	79.9	76.4 - 83.1	7,500
35+	300	12.3	8.2 - 17.9	2,400	87.7	82.1 - 91.8	2,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	2,000	25.0	21.6 - 28.7	5,900	75.0	71.3 - 78.4	7,800
Female	1,600	23.4	20.0 - 27.1	5,300	76.6	72.9 - 80.0	6,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	18.6	15.9 - 21.7	8,300	81.4	78.3 - 84.1	10,100
No	1,700	36.6	32.1 - 41.3	2,900	63.4	58.7 - 67.9	4,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,000	32.4	28.4 - 36.7	4,200	67.6	63.3 - 71.6	6,200
131-185%	500	22.6	16.8 - 29.6	1,700	77.4	70.4 - 83.2	2,200
186+%	600	12.7	9.5 - 16.7	4,500	87.3	83.3 - 90.5	5,100

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,100	36.1	30.5 - 42.1	2,000	63.9	57.9 - 69.5	3,100
\$15,001-\$25,000	500	26.0	19.4 - 33.8	1,400	74.0	66.2 - 80.6	1,900
\$25,001-\$40,000	800	27.3	21.6 - 33.8	2,100	72.7	66.2 - 78.4	2,900
>=\$40,001	900	14.4	11.4 - 18.1	5,200	85.6	81.9 - 88.6	6,100

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	28.8	19.3 - 40.7	700	71.2	59.3 - 80.7	1,000
Normal birth weight	3,300	23.9	21.4 - 26.6	10,500	76.1	73.4 - 78.6	13,700

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	32.4	23.5 - 42.8	700	67.6	57.2 - 76.5	1,000
No	3,200	23.8	21.3 - 26.5	10,300	76.2	73.5 - 78.7	13,500

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race- Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	36.3	30.2 - 42.9	1,700	63.7	57.1 - 69.8	2,600
No	2,500	21.4	18.8 - 24.2	9,300	78.6	75.8 - 81.2	11,800

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	34.3	30.3 - 38.7	3,800	65.7	61.3 - 69.7	5,800
No	1,600	17.7	14.8 - 20.9	7,200	82.3	79.1 - 85.2	8,800

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,200	21.9	19.6 - 24.5	11,200	78.1	75.5 - 80.4	14,400

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	27.1	23.0 - 31.6	1,200	72.9	68.4 - 77.0	1,700
Honolulu	2,200	20.5	17.5 - 23.9	8,400	79.5	76.1 - 82.5	10,500
Kauai	100	23.8	17.3 - 31.9	400	76.2	68.1 - 82.7	600
Maui	400	25.4	21.0 - 30.3	1,200	74.6	69.7 - 79.0	1,600

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	600	18.1	13.2 - 24.4	2,500	81.9	75.6 - 86.8	3,100
Native Hawaiian	1,000	27.4	24.5 - 30.5	2,700	72.6	69.5 - 75.5	3,700
Chinese	100	21.8	10.5 - 39.9	400	78.2	60.1 - 89.5	500
Filipino	800	27.7	21.8 - 34.6	2,100	72.3	65.4 - 78.2	3,000
Japanese	200	8.3	4.8 - 14.0	2,100	91.7	86.0 - 95.2	2,300
Other	500	25.5	17.4 - 35.6	1,300	74.5	64.4 - 82.6	1,800

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	500	39.1	30.3 - 48.7	700	60.9	51.3 - 69.7	1,200
20-24	1,000	29.3	24.1 - 34.9	2,500	70.7	65.1 - 75.9	3,600
25-34	1,300	18.0	14.9 - 21.5	5,900	82.0	78.5 - 85.1	7,200
35+	300	14.2	9.9 - 20.1	2,000	85.8	79.9 - 90.1	2,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,600	22.8	19.4 - 26.5	5,400	77.2	73.5 - 80.6	7,000
Female	1,500	21.2	18.0 - 24.7	5,800	78.8	75.3 - 82.0	7,300

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	17.6	14.9 - 20.7	8,200	82.4	79.3 - 85.1	10,000
No	1,400	32.0	27.6 - 36.7	3,000	68.0	63.3 - 72.4	4,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,500	28.0	24.0 - 32.4	3,800	72.0	67.6 - 76.0	5,300
131-185%	700	22.9	17.7 - 29.1	2,300	77.1	70.9 - 82.3	3,000
186+%	700	13.3	10.2 - 17.2	4,300	86.7	82.8 - 89.8	5,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	800	31.2	25.4 - 37.6	1,700	68.8	62.4 - 74.6	2,500
\$15,001-\$25,000	500	24.3	18.4 - 31.3	1,600	75.7	68.7 - 81.6	2,100
\$25,001-\$40,000	700	24.1	18.8 - 30.3	2,300	75.9	69.7 - 81.2	3,000
>=\$40,001	900	14.7	11.7 - 18.4	5,100	85.3	81.6 - 88.3	6,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	31.0	20.4 - 43.9	600	69.0	56.1 - 79.6	900
Normal birth weight	2,900	21.3	18.9 - 23.9	10,600	78.7	76.1 - 81.1	13,400

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	40.5	30.4 - 51.4	500	59.5	48.6 - 69.6	900
No	2,800	20.7	18.3 - 23.4	10,500	79.3	76.6 - 81.7	13,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	34.1	28.0 - 40.8	1,700	65.9	59.2 - 72.0	2,600
No	2,200	19.0	16.5 - 21.7	9,300	81.0	78.3 - 83.5	11,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	30.7	26.5 - 35.2	3,900	69.3	64.8 - 73.5	5,600
No	1,400	16.4	13.8 - 19.4	7,200	83.6	80.6 - 86.2	8,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,400	23.9	21.9 - 26.0	10,700	76.1	74.0 - 78.1	14,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	300	22.9	19.0 - 27.4	1,200	77.1	72.6 - 81.0	1,500
Honolulu	2,500	23.7	21.3 - 26.4	7,900	76.3	73.6 - 78.7	10,400
Kauai	200	23.4	17.1 - 31.2	500	76.6	68.8 - 82.9	700
Maui	400	25.9	21.4 - 30.9	1,100	74.1	69.1 - 78.6	1,500

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	500	16.1	12.6 - 20.4	2,600	83.9	79.6 - 87.4	3,100
Native Hawaiian	1,100	30.5	28.1 - 33.1	2,500	69.5	66.9 - 71.9	3,600
Chinese	100	23.9	13.6 - 38.4	500	76.1	61.6 - 86.4	600
Filipino	900	28.4	23.3 - 34.1	2,100	71.6	65.9 - 76.7	3,000
Japanese	300	16.8	11.9 - 23.2	1,600	83.2	76.8 - 88.1	1,900
Other	400	24.3	18.1 - 31.8	1,400	75.7	68.2 - 81.9	1,800

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	44.7	38.0 - 51.6	700	55.3	48.4 - 62.0	1,300
20-24	1,200	32.1	27.9 - 36.7	2,500	67.9	63.3 - 72.1	3,600
25-34	1,300	18.7	16.1 - 21.6	5,600	81.3	78.4 - 83.9	6,800
35+	300	14.6	10.7 - 19.7	2,000	85.4	80.3 - 89.3	2,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,800	24.2	21.3 - 27.2	5,500	75.8	72.8 - 78.7	7,300
Female	1,600	23.6	20.9 - 26.6	5,100	76.4	73.4 - 79.1	6,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	19.4	17.1 - 22.0	7,800	80.6	78.0 - 82.9	9,600
No	1,500	33.7	30.0 - 37.6	2,900	66.3	62.4 - 70.0	4,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,400	28.9	25.6 - 32.5	3,600	71.1	67.5 - 74.4	5,000
131-185%	700	21.2	17.5 - 25.5	2,500	78.8	74.5 - 82.5	3,100
186+%	800	15.8	12.9 - 19.2	4,200	84.2	80.8 - 87.1	5,000

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	800	30.0	25.4 - 35.1	1,800	70.0	64.9 - 74.6	2,600
\$15,001-\$25,000	600	31.5	25.8 - 37.9	1,300	68.5	62.1 - 74.2	1,800
\$25,001-\$40,000	700	23.3	19.4 - 27.7	2,400	76.7	72.3 - 80.6	3,100
>=\$40,001	900	16.2	13.5 - 19.3	4,800	83.8	80.7 - 86.5	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	29.2	21.1 - 38.9	600	70.8	61.1 - 78.9	900
Normal birth weight	3,100	23.6	21.5 - 25.8	10,000	76.4	74.2 - 78.5	13,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	36.2	28.5 - 44.6	700	63.8	55.4 - 71.5	1,100
No	2,900	22.7	20.6 - 24.9	9,900	77.3	75.1 - 79.4	12,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	33.7	28.9 - 38.9	1,800	66.3	61.1 - 71.1	2,600
No	2,300	21.2	19.0 - 23.6	8,600	78.8	76.4 - 81.0	10,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	31.2	27.9 - 34.6	3,900	68.8	65.4 - 72.1	5,700
No	1,500	18.9	16.4 - 21.6	6,600	81.1	78.4 - 83.6	8,200

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	3,300	23.3	21.4 - 25.4	10,800	76.7	74.6 - 78.6	14,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	500	31.6	27.1 - 36.5	1,100	68.4	63.5 - 72.9	1,600
Honolulu	2,300	21.7	19.4 - 24.3	8,100	78.3	75.7 - 80.6	10,400
Kauai	100	21.0	14.8 - 28.9	500	79.0	71.1 - 85.2	600
Maui	400	26.1	21.6 - 31.1	1,100	73.9	68.9 - 78.4	1,400

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	700	20.3	16.5 - 24.8	2,700	79.7	75.2 - 83.5	3,400
Native Hawaiian	1,200	32.7	30.1 - 35.4	2,400	67.3	64.6 - 69.9	3,600
Chinese	100	20.6	11.1 - 34.9	500	79.4	65.1 - 88.9	600
Filipino	700	24.9	20.3 - 30.1	2,200	75.1	69.9 - 79.7	2,900
Japanese	200	9.2	5.5 - 15.1	1,600	90.8	84.9 - 94.5	1,800
Other	400	22.7	16.5 - 30.3	1,400	77.3	69.7 - 83.5	1,800

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	40.6	34.4 - 47.0	900	59.4	53.0 - 65.6	1,600
20-24	1,000	29.9	25.9 - 34.3	2,300	70.1	65.7 - 74.1	3,300
25-34	1,300	19.2	16.5 - 22.1	5,500	80.8	77.9 - 83.5	6,800
35+	400	14.7	10.8 - 19.8	2,100	85.3	80.2 - 89.2	2,400

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	1,700	23.4	20.8 - 26.3	5,400	76.6	73.7 - 79.2	7,100
Female	1,600	23.2	20.5 - 26.2	5,300	76.8	73.8 - 79.5	6,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	18.6	16.4 - 21.0	7,900	81.4	79.0 - 83.6	9,700
No	1,500	34.0	30.3 - 37.8	2,900	66.0	62.2 - 69.7	4,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	1,800	30.3	27.2 - 33.7	4,000	69.7	66.3 - 72.8	5,800
131-185%	400	18.4	14.4 - 23.3	1,600	81.6	76.7 - 85.6	2,000
186+%	900	16.6	13.8 - 20.0	4,400	83.4	80.0 - 86.2	5,300

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,100	35.4	30.9 - 40.1	2,000	64.6	59.9 - 69.1	3,100
\$15,001-\$25,000	600	26.2	21.3 - 31.8	1,600	73.8	68.2 - 78.7	2,100
\$25,001-\$40,000	600	20.5	16.8 - 24.8	2,200	79.5	75.2 - 83.2	2,800
>=\$40,001	800	15.5	12.8 - 18.7	4,500	84.5	81.3 - 87.2	5,300

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	300	29.4	21.7 - 38.6	600	70.6	61.4 - 78.3	900
Normal birth weight	3,000	22.6	20.7 - 24.7	10,100	77.4	75.3 - 79.3	13,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	43.0	34.8 - 51.6	500	57.0	48.4 - 65.2	900
No	2,700	21.3	19.4 - 23.4	10,000	78.7	76.6 - 80.6	12,700

PRAMS Health Indicator Report - Breastfeeding (any) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	36.1	31.2 - 41.3	1,700	63.9	58.7 - 68.8	2,600
No	2,200	19.6	17.6 - 21.8	8,800	80.4	78.2 - 82.4	11,000

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	29.0	26.1 - 32.2	4,100	71.0	67.8 - 73.9	5,800
No	1,500	18.9	16.4 - 21.6	6,500	81.1	78.4 - 83.6	8,000