

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Breastfeeding (exclusive) duration 8+ weeks, among initiators	Breastfeeding (exclusive) duration 8+ weeks, among initiators

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/13/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	6,500	54.9	52.2 - 57.5	5,400	45.1	42.5 - 47.8	11,900
2001	8,200	58.5	56.0 - 60.9	5,800	41.5	39.1 - 44.0	14,000
2002	8,400	58.4	55.4 - 61.4	5,900	41.6	38.6 - 44.6	14,300
2003	8,800	60.1	57.2 - 63.0	5,800	39.9	37.0 - 42.8	14,600
2004	9,500	61.3	59.1 - 63.5	6,000	38.7	36.5 - 40.9	15,600
2005	9,200	60.5	58.0 - 62.9	6,000	39.5	37.1 - 42.0	15,200
2006	9,600	59.7	57.2 - 62.3	6,400	40.3	37.7 - 42.8	16,000
2007	9,900	60.3	57.9 - 62.7	6,500	39.7	37.3 - 42.1	16,400
2008	10,100	60.6	58.2 - 63.0	6,500	39.4	37.0 - 41.8	16,600
2009	9,900	59.7	56.5 - 62.9	6,700	40.3	37.1 - 43.5	16,500
2010	8,700	51.7	48.4 - 55.1	8,200	48.3	44.9 - 51.6	16,900
2011	9,200	53.2	49.9 - 56.5	8,100	46.8	43.5 - 50.1	17,200
2012	9,200	54.4	51.0 - 57.8	7,700	45.6	42.2 - 49.0	16,900
2013	7,900	47.5	43.9 - 51.2	8,700	52.5	48.8 - 56.1	16,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	7,900	47.5	43.9 - 51.2	8,700	52.5	48.8 - 56.1	16,700

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	47.3	41.5 - 53.3	1,100	52.7	46.7 - 58.5	2,000
Honolulu	5,700	47.4	42.5 - 52.3	6,300	52.6	47.7 - 57.5	12,000
Kauai	300	40.6	35.3 - 46.2	500	59.4	53.8 - 64.7	800
Maui	900	51.5	45.7 - 57.3	900	48.5	42.7 - 54.3	1,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	35.8	29.3 - 43.0	2,900	64.2	57.0 - 70.7	4,600
Native Hawaiian	2,500	51.5	45.1 - 57.9	2,300	48.5	42.1 - 54.9	4,800
Chinese	600	70.1	52.1 - 83.5	200	29.9	16.5 - 47.9	800
Filipino	1,600	58.7	49.3 - 67.6	1,100	41.3	32.4 - 50.7	2,700
Japanese	700	42.4	31.3 - 54.2	1,000	57.6	45.8 - 68.7	1,700
Other	900	45.2	34.9 - 56.1	1,100	54.8	43.9 - 65.1	2,100

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	53.1	37.1 - 68.4	400	46.9	31.6 - 62.9	800
20-24	1,500	39.5	32.3 - 47.2	2,300	60.5	52.8 - 67.7	3,900
25-34	4,300	47.7	42.7 - 52.7	4,800	52.3	47.3 - 57.3	9,100
35+	1,700	56.1	47.7 - 64.2	1,300	43.9	35.8 - 52.3	3,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,400	42.6	37.6 - 47.7	4,600	57.4	52.3 - 62.4	8,100
Female	4,500	52.2	47.0 - 57.4	4,100	47.8	42.6 - 53.0	8,600

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,900	45.4	40.8 - 50.2	5,900	54.6	49.8 - 59.2	10,700
No	3,000	51.3	45.4 - 57.1	2,900	48.7	42.9 - 54.6	5,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,800	47.7	41.8 - 53.7	3,100	52.3	46.3 - 58.2	5,900
131-185%	1,200	53.8	43.6 - 63.6	1,000	46.2	36.4 - 56.4	2,200
186+%	3,100	45.5	39.8 - 51.4	3,700	54.5	48.6 - 60.2	6,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Income (2012+)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	1,800	48.9	41.5 - 56.4	1,900	51.1	43.6 - 58.5	3,700
\$17,001-\$26,000	1,000	46.7	36.8 - 57.0	1,200	53.3	43.0 - 63.2	2,200
\$26,001-\$51,000	1,700	49.3	41.4 - 57.3	1,800	50.7	42.7 - 58.6	3,500
>=\$51,001	2,600	45.6	39.4 - 51.9	3,100	54.4	48.1 - 60.6	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	65.8	60.4 - 70.8	400	34.2	29.2 - 39.6	1,100
Normal birth weight	7,200	46.2	42.3 - 50.1	8,400	53.8	49.9 - 57.7	15,500

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	46.2	31.8 - 61.2	500	53.8	38.8 - 68.2	800
No	7,400	47.8	44.0 - 51.6	8,100	52.2	48.4 - 56.0	15,400

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	45.9	37.3 - 54.7	1,500	54.1	45.3 - 62.7	2,800
No	6,500	48.1	44.0 - 52.2	7,000	51.9	47.8 - 56.0	13,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,800	53.3	47.8 - 58.8	3,300	46.7	41.2 - 52.2	7,100
No	4,100	43.3	38.5 - 48.2	5,400	56.7	51.8 - 61.5	9,500

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,200	54.4	51.0 - 57.8	7,700	45.6	42.2 - 49.0	16,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	53.3	47.8 - 58.8	1,000	46.7	41.2 - 52.2	2,100
Honolulu	6,700	54.9	50.3 - 59.4	5,500	45.1	40.6 - 49.7	12,100
Kauai	300	45.4	38.9 - 52.1	400	54.6	47.9 - 61.1	800
Maui	1,000	56.3	51.1 - 61.4	800	43.7	38.6 - 48.9	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	40.0	33.5 - 46.9	2,300	60.0	53.1 - 66.5	3,900
Native Hawaiian	2,800	59.6	53.4 - 65.5	1,900	40.4	34.5 - 46.6	4,700
Chinese	500	50.5	36.3 - 64.6	500	49.5	35.4 - 63.7	1,000
Filipino	2,000	65.4	57.4 - 72.7	1,000	34.6	27.3 - 42.6	3,000
Japanese	900	59.4	47.8 - 70.1	600	40.6	29.9 - 52.2	1,500
Other	1,400	52.5	43.5 - 61.2	1,300	47.5	38.8 - 56.5	2,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	72.2	58.4 - 82.8	300	27.8	17.2 - 41.6	1,000
20-24	1,900	53.7	46.1 - 61.1	1,600	46.3	38.9 - 53.9	3,500
25-34	5,000	52.3	47.8 - 56.8	4,500	47.7	43.2 - 52.2	9,500
35+	1,700	56.3	48.3 - 64.0	1,300	43.7	36.0 - 51.7	2,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,700	55.4	50.5 - 60.1	3,800	44.6	39.9 - 49.5	8,400
Female	4,500	53.5	48.7 - 58.2	3,900	46.5	41.8 - 51.3	8,400

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,500	52.9	48.5 - 57.2	4,900	47.1	42.8 - 51.5	10,500
No	3,600	57.0	51.5 - 62.3	2,700	43.0	37.7 - 48.5	6,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,400	57.8	52.1 - 63.2	2,500	42.2	36.8 - 47.9	5,900
131-185%	1,800	62.4	54.0 - 70.2	1,100	37.6	29.8 - 46.0	2,800
186+%	3,200	48.2	42.8 - 53.6	3,400	51.8	46.4 - 57.2	6,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Income (2012+)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	2,200	59.2	51.9 - 66.2	1,500	40.8	33.8 - 48.1	3,800
\$17,001-\$26,000	1,200	55.3	46.0 - 64.3	900	44.7	35.7 - 54.0	2,100
\$26,001-\$51,000	2,300	57.1	50.0 - 63.9	1,700	42.9	36.1 - 50.0	4,000
>=\$51,001	2,800	49.8	44.0 - 55.6	2,800	50.2	44.4 - 56.0	5,600

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	65.8	60.1 - 71.1	400	34.2	28.9 - 39.9	1,100
Normal birth weight	8,400	53.5	49.9 - 57.1	7,300	46.5	42.9 - 50.1	15,700

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	67.4	51.3 - 80.3	200	32.6	19.7 - 48.7	700
No	8,500	54.2	50.6 - 57.7	7,200	45.8	42.3 - 49.4	15,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	53.3	44.8 - 61.7	1,200	46.7	38.3 - 55.2	2,600
No	7,500	54.9	51.2 - 58.7	6,200	45.1	41.3 - 48.8	13,700

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,400	59.2	54.2 - 64.1	3,100	40.8	35.9 - 45.8	7,500
No	4,700	50.5	45.9 - 55.0	4,600	49.5	45.0 - 54.1	9,300

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,200	53.2	49.9 - 56.5	8,100	46.8	43.5 - 50.1	17,200

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	54.7	49.4 - 59.9	1,000	45.3	40.1 - 50.6	2,200
Honolulu	6,700	54.2	49.7 - 58.6	5,700	45.8	41.4 - 50.3	12,300
Kauai	400	47.4	41.0 - 53.8	400	52.6	46.2 - 59.0	800
Maui	900	47.3	42.5 - 52.0	1,000	52.7	48.0 - 57.5	1,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	43.1	36.6 - 49.7	2,500	56.9	50.3 - 63.4	4,300
Native Hawaiian	2,900	59.2	53.2 - 64.9	2,000	40.8	35.1 - 46.8	4,900
Chinese	300	52.2	35.0 - 68.8	300	47.8	31.2 - 65.0	700
Filipino	1,900	63.9	56.1 - 71.1	1,100	36.1	28.9 - 43.9	3,000
Japanese	1,100	58.5	47.9 - 68.3	700	41.5	31.7 - 52.1	1,800
Other	1,100	42.8	33.9 - 52.1	1,500	57.2	47.9 - 66.1	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	63.1	49.9 - 74.7	300	36.9	25.3 - 50.1	900
20-24	2,400	61.0	53.9 - 67.7	1,600	39.0	32.3 - 46.1	4,000
25-34	4,600	49.4	44.9 - 53.9	4,700	50.6	46.1 - 55.1	9,300
35+	1,600	51.6	43.8 - 59.4	1,500	48.4	40.6 - 56.2	3,000

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,600	52.1	47.4 - 56.7	4,200	47.9	43.3 - 52.6	8,800
Female	4,600	54.3	49.6 - 59.0	3,800	45.7	41.0 - 50.4	8,400

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,400	49.4	45.1 - 53.6	5,500	50.6	46.4 - 54.9	10,900
No	3,800	59.7	54.3 - 64.8	2,600	40.3	35.2 - 45.7	6,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	58.4	52.9 - 63.7	2,500	41.6	36.3 - 47.1	5,900
131-185%	1,300	49.6	40.9 - 58.4	1,300	50.4	41.6 - 59.1	2,600
186+%	3,900	50.8	45.8 - 55.9	3,800	49.2	44.1 - 54.2	7,700

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,900	63.1	55.2 - 70.3	1,100	36.9	29.7 - 44.8	2,900
\$10,000-\$24,999	1,800	55.4	47.9 - 62.6	1,400	44.6	37.4 - 52.1	3,200
\$25,000-\$49,999	2,100	49.9	43.1 - 56.7	2,100	50.1	43.3 - 56.9	4,200
>=\$50,000	3,000	50.2	44.4 - 55.9	3,000	49.8	44.1 - 55.6	6,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	66.3	60.8 - 71.5	400	33.7	28.5 - 39.2	1,200
Normal birth weight	8,400	52.2	48.7 - 55.8	7,700	47.8	44.2 - 51.3	16,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	73.0	60.5 - 82.7	200	27.0	17.3 - 39.5	800
No	8,500	52.1	48.6 - 55.5	7,800	47.9	44.5 - 51.4	16,300

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	59.0	51.4 - 66.2	1,300	41.0	33.8 - 48.6	3,200
No	7,200	51.7	48.0 - 55.4	6,700	48.3	44.6 - 52.0	13,800

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,300	57.4	52.5 - 62.3	3,200	42.6	37.7 - 47.5	7,600
No	4,800	50.1	45.5 - 54.6	4,800	49.9	45.4 - 54.5	9,500

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,700	51.7	48.4 - 55.1	8,200	48.3	44.9 - 51.6	16,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	50.5	45.3 - 55.8	1,100	49.5	44.2 - 54.7	2,200
Honolulu	6,400	52.5	48.0 - 56.9	5,800	47.5	43.1 - 52.0	12,100
Kauai	300	45.2	39.2 - 51.4	400	54.8	48.6 - 60.8	700
Maui	900	51.0	46.1 - 56.0	900	49.0	44.0 - 53.9	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	39.2	32.8 - 46.0	2,500	60.8	54.0 - 67.2	4,100
Native Hawaiian	2,700	53.9	48.1 - 59.7	2,300	46.1	40.3 - 51.9	5,000
Chinese	300	38.4	24.1 - 55.1	500	61.6	44.9 - 75.9	700
Filipino	2,000	69.1	61.3 - 75.9	900	30.9	24.1 - 38.7	2,900
Japanese	900	55.3	44.5 - 65.7	700	44.7	34.3 - 55.5	1,700
Other	1,200	49.2	39.9 - 58.5	1,200	50.8	41.5 - 60.1	2,400

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	600	61.0	48.8 - 71.9	400	39.0	28.1 - 51.2	1,100
20-24	2,000	51.0	44.0 - 57.9	2,000	49.0	42.1 - 56.0	4,000
25-34	4,400	50.6	45.9 - 55.2	4,300	49.4	44.8 - 54.1	8,800
35+	1,600	52.9	45.1 - 60.6	1,400	47.1	39.4 - 54.9	3,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,400	52.0	47.3 - 56.6	4,100	48.0	43.4 - 52.7	8,500
Female	4,300	51.5	46.7 - 56.3	4,100	48.5	43.7 - 53.3	8,400

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,300	49.3	45.0 - 53.6	5,400	50.7	46.4 - 55.0	10,800
No	3,400	56.0	50.7 - 61.2	2,700	44.0	38.8 - 49.3	6,100

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,200	55.7	50.1 - 61.1	2,600	44.3	38.9 - 49.9	5,800
131-185%	1,500	54.0	45.6 - 62.3	1,200	46.0	37.7 - 54.4	2,700
186+%	3,500	47.9	42.8 - 53.0	3,800	52.1	47.0 - 57.2	7,400

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,800	58.2	50.7 - 65.4	1,300	41.8	34.6 - 49.3	3,100
\$10,000-\$24,999	1,500	53.3	45.3 - 61.1	1,300	46.7	38.9 - 54.7	2,900
\$25,000-\$49,999	2,200	51.7	45.0 - 58.4	2,000	48.3	41.6 - 55.0	4,200
>=\$50,000	2,800	48.1	42.4 - 53.9	3,000	51.9	46.1 - 57.6	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	61.1	55.4 - 66.5	400	38.9	33.5 - 44.6	1,100
Normal birth weight	8,100	51.1	47.5 - 54.6	7,700	48.9	45.4 - 52.5	15,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	50.9	37.9 - 63.8	600	49.1	36.2 - 62.1	1,100
No	8,100	52.2	48.7 - 55.6	7,500	47.8	44.4 - 51.3	15,600

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	58.9	51.6 - 65.8	1,400	41.1	34.2 - 48.4	3,400
No	6,700	50.4	46.6 - 54.1	6,600	49.6	45.9 - 53.4	13,300

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,900	54.3	49.3 - 59.2	3,300	45.7	40.8 - 50.7	7,200
No	4,800	50.5	46.0 - 55.0	4,700	49.5	45.0 - 54.0	9,500

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,900	59.7	56.5 - 62.9	6,700	40.3	37.1 - 43.5	16,500

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	58.1	52.8 - 63.3	900	41.9	36.7 - 47.2	2,100
Honolulu	7,100	61.0	56.5 - 65.3	4,500	39.0	34.7 - 43.5	11,700
Kauai	400	49.8	43.5 - 56.1	400	50.2	43.9 - 56.5	800
Maui	1,100	58.0	53.2 - 62.6	800	42.0	37.4 - 46.8	2,000

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	46.0	39.6 - 52.5	2,100	54.0	47.5 - 60.4	4,000
Native Hawaiian	3,300	64.8	58.9 - 70.3	1,800	35.2	29.7 - 41.1	5,000
Chinese	400	63.0	45.4 - 77.8	300	37.0	22.2 - 54.6	700
Filipino	2,000	67.1	59.3 - 74.0	1,000	32.9	26.0 - 40.7	3,100
Japanese	1,000	63.0	52.3 - 72.5	600	37.0	27.5 - 47.7	1,600
Other	1,300	59.4	50.0 - 68.1	900	40.6	31.9 - 50.0	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	64.1	52.9 - 73.9	600	35.9	26.1 - 47.1	1,700
20-24	2,400	65.2	58.4 - 71.5	1,300	34.8	28.5 - 41.6	3,700
25-34	4,600	54.9	50.3 - 59.4	3,700	45.1	40.6 - 49.7	8,300
35+	1,800	64.1	56.3 - 71.2	1,000	35.9	28.8 - 43.7	2,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,300	61.0	56.4 - 65.4	3,400	39.0	34.6 - 43.6	8,700
Female	4,500	58.3	53.6 - 62.9	3,300	41.7	37.1 - 46.4	7,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,300	53.2	48.9 - 57.4	4,700	46.8	42.6 - 51.1	10,000
No	4,500	69.8	64.9 - 74.4	2,000	30.2	25.6 - 35.1	6,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,600	63.9	58.2 - 69.2	2,000	36.1	30.8 - 41.8	5,600
131-185%	1,600	62.8	54.6 - 70.4	1,000	37.2	29.6 - 45.4	2,600
186+%	3,800	55.0	50.0 - 60.0	3,100	45.0	40.0 - 50.0	6,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,700	64.5	56.0 - 72.1	900	35.5	27.9 - 44.0	2,600
\$10,000-\$24,999	2,300	66.9	59.8 - 73.2	1,100	33.1	26.8 - 40.2	3,500
\$25,000-\$49,999	1,900	55.7	48.5 - 62.7	1,500	44.3	37.3 - 51.5	3,400
>=\$50,000	3,200	55.5	49.9 - 61.0	2,600	44.5	39.0 - 50.1	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	68.3	62.7 - 73.4	300	31.7	26.6 - 37.3	1,100
Normal birth weight	9,100	59.1	55.6 - 62.5	6,300	40.9	37.5 - 44.4	15,400

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	66.5	54.7 - 76.5	400	33.5	23.5 - 45.3	1,300
No	8,700	59.0	55.6 - 62.4	6,100	41.0	37.6 - 44.4	14,800

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	63.4	56.2 - 70.1	1,200	36.6	29.9 - 43.8	3,300
No	7,500	58.7	54.9 - 62.3	5,300	41.3	37.7 - 45.1	12,800

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,100	62.8	57.7 - 67.7	2,400	37.2	32.3 - 42.3	6,500
No	5,500	57.6	53.2 - 61.9	4,000	42.4	38.1 - 46.8	9,500

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,100	60.6	58.2 - 63.0	6,500	39.4	37.0 - 41.8	16,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	54.3	47.0 - 61.5	1,000	45.7	38.5 - 53.0	2,100
Honolulu	7,300	61.7	58.8 - 64.5	4,500	38.3	35.5 - 41.2	11,800
Kauai	500	59.6	48.3 - 70.0	300	40.4	30.0 - 51.7	800
Maui	1,200	61.1	53.6 - 68.1	700	38.9	31.9 - 46.4	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	46.9	41.9 - 51.8	2,200	53.1	48.2 - 58.1	4,100
Native Hawaiian	2,900	63.7	58.5 - 68.5	1,600	36.3	31.5 - 41.5	4,500
Chinese	400	67.0	60.3 - 73.1	200	33.0	26.9 - 39.7	600
Filipino	2,100	70.1	64.8 - 75.0	900	29.9	25.0 - 35.2	3,000
Japanese	1,300	68.5	61.6 - 74.6	600	31.5	25.4 - 38.4	2,000
Other	1,400	58.6	51.9 - 64.9	1,000	41.4	35.1 - 48.1	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	57.4	47.8 - 66.5	500	42.6	33.5 - 52.2	1,300
20-24	2,300	63.3	57.8 - 68.5	1,300	36.7	31.5 - 42.2	3,600
25-34	5,000	58.1	54.7 - 61.4	3,600	41.9	38.6 - 45.3	8,600
35+	2,100	65.7	60.5 - 70.6	1,100	34.3	29.4 - 39.5	3,200

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,300	60.6	57.2 - 63.9	3,400	39.4	36.1 - 42.8	8,700
Female	4,800	60.6	57.0 - 64.1	3,100	39.4	35.9 - 43.0	7,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,200	58.4	55.4 - 61.3	4,400	41.6	38.7 - 44.6	10,700
No	3,800	64.5	60.2 - 68.6	2,100	35.5	31.4 - 39.8	5,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,700	63.3	58.9 - 67.4	2,100	36.7	32.6 - 41.1	5,800
131-185%	1,300	56.6	50.0 - 63.1	1,000	43.4	36.9 - 50.0	2,400
186+%	4,300	58.5	54.9 - 61.9	3,100	41.5	38.1 - 45.1	7,400

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,700	65.7	59.1 - 71.7	900	34.3	28.3 - 40.9	2,600
\$10,000-\$24,999	2,100	60.2	54.5 - 65.6	1,400	39.8	34.4 - 45.5	3,500
\$25,000-\$49,999	2,400	61.7	56.6 - 66.5	1,500	38.3	33.5 - 43.4	3,800
>=\$50,000	3,300	56.8	52.8 - 60.7	2,500	43.2	39.3 - 47.2	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	73.5	64.3 - 81.0	300	26.5	19.0 - 35.7	1,100
Normal birth weight	9,200	59.6	57.1 - 62.1	6,200	40.4	37.9 - 42.9	15,500

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	74.4	65.2 - 81.8	300	25.6	18.2 - 34.8	1,300
No	9,000	59.7	57.1 - 62.2	6,100	40.3	37.8 - 42.9	15,100

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	68.7	62.7 - 74.1	900	31.3	25.9 - 37.3	2,900
No	7,900	59.1	56.4 - 61.7	5,500	40.9	38.3 - 43.6	13,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,300	63.7	59.7 - 67.5	2,400	36.3	32.5 - 40.3	6,700
No	5,700	58.9	55.8 - 62.0	4,000	41.1	38.0 - 44.2	9,700

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,900	60.3	57.9 - 62.7	6,500	39.7	37.3 - 42.1	16,400

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	48.0	40.9 - 55.2	1,100	52.0	44.8 - 59.1	2,100
Honolulu	7,300	63.5	60.6 - 66.3	4,200	36.5	33.7 - 39.4	11,500
Kauai	500	49.8	39.1 - 60.6	500	50.2	39.4 - 60.9	900
Maui	1,200	60.0	52.4 - 67.1	800	40.0	32.9 - 47.6	1,900

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,700	45.4	40.2 - 50.7	2,000	54.6	49.3 - 59.8	3,700
Native Hawaiian	2,900	63.9	58.7 - 68.7	1,600	36.1	31.3 - 41.3	4,500
Chinese	400	60.9	54.1 - 67.3	300	39.1	32.7 - 45.9	600
Filipino	2,100	70.8	65.6 - 75.5	900	29.2	24.5 - 34.4	2,900
Japanese	1,500	72.2	65.4 - 78.1	600	27.8	21.9 - 34.6	2,000
Other	1,400	54.0	47.4 - 60.4	1,200	46.0	39.6 - 52.6	2,600

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	65.0	55.1 - 73.8	500	35.0	26.2 - 44.9	1,400
20-24	2,400	60.1	54.9 - 65.1	1,600	39.9	34.9 - 45.1	4,000
25-34	4,800	58.9	55.5 - 62.3	3,300	41.1	37.7 - 44.5	8,100
35+	1,800	62.3	56.7 - 67.6	1,100	37.7	32.4 - 43.3	2,900

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,200	60.3	56.9 - 63.6	3,400	39.7	36.4 - 43.1	8,500
Female	4,700	60.4	56.8 - 63.9	3,100	39.6	36.1 - 43.2	7,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,200	59.2	56.2 - 62.1	4,300	40.8	37.9 - 43.8	10,400
No	3,700	62.4	58.0 - 66.5	2,200	37.6	33.5 - 42.0	6,000

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,700	62.2	57.8 - 66.3	2,200	37.8	33.7 - 42.2	5,900
131-185%	1,500	60.2	53.8 - 66.3	1,000	39.8	33.7 - 46.2	2,400
186+%	4,100	59.5	55.9 - 63.1	2,800	40.5	36.9 - 44.1	6,900

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,600	59.3	52.6 - 65.7	1,100	40.7	34.3 - 47.4	2,600
\$10,000-\$24,999	2,100	64.0	58.3 - 69.2	1,200	36.0	30.8 - 41.7	3,300
\$25,000-\$49,999	2,300	61.4	56.3 - 66.3	1,500	38.6	33.7 - 43.7	3,800
>=\$50,000	3,300	59.2	55.2 - 63.2	2,300	40.8	36.8 - 44.8	5,600

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	71.7	62.9 - 79.0	400	28.3	21.0 - 37.1	1,300
Normal birth weight	9,000	59.4	56.8 - 61.9	6,100	40.6	38.1 - 43.2	15,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	67.5	57.9 - 75.8	400	32.5	24.2 - 42.1	1,300
No	8,900	59.9	57.3 - 62.4	6,000	40.1	37.6 - 42.7	14,900

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	68.2	62.4 - 73.5	1,000	31.8	26.5 - 37.6	3,100
No	7,700	58.7	55.9 - 61.4	5,400	41.3	38.6 - 44.1	13,100

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,800	58.2	54.0 - 62.2	2,700	41.8	37.8 - 46.0	6,500
No	5,900	61.5	58.3 - 64.5	3,700	38.5	35.5 - 41.7	9,600

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,600	59.7	57.2 - 62.3	6,400	40.3	37.7 - 42.8	16,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	54.7	47.0 - 62.2	900	45.3	37.8 - 53.0	1,900
Honolulu	7,300	62.2	59.2 - 65.0	4,400	37.8	35.0 - 40.8	11,700
Kauai	300	44.8	33.1 - 57.1	400	55.2	42.9 - 66.9	700
Maui	900	55.3	47.0 - 63.4	700	44.7	36.6 - 53.0	1,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	46.4	41.2 - 51.7	2,100	53.6	48.3 - 58.8	3,900
Native Hawaiian	2,700	64.4	59.0 - 69.5	1,500	35.6	30.5 - 41.0	4,200
Chinese	400	67.6	60.5 - 74.1	200	32.4	25.9 - 39.5	600
Filipino	1,900	63.6	58.0 - 68.9	1,100	36.4	31.1 - 42.0	2,900
Japanese	1,300	68.8	61.5 - 75.2	600	31.2	24.8 - 38.5	1,900
Other	1,500	59.5	52.7 - 65.9	1,000	40.5	34.1 - 47.3	2,500

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	62.6	52.8 - 71.4	500	37.4	28.6 - 47.2	1,400
20-24	2,500	65.0	59.5 - 70.1	1,300	35.0	29.9 - 40.5	3,900
25-34	4,500	56.7	53.1 - 60.3	3,400	43.3	39.7 - 46.9	7,900
35+	1,700	59.7	54.0 - 65.1	1,200	40.3	34.9 - 46.0	2,900

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,200	62.8	59.3 - 66.2	3,100	37.2	33.8 - 40.7	8,300
Female	4,400	56.5	52.7 - 60.2	3,400	43.5	39.8 - 47.3	7,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,800	57.2	54.1 - 60.3	4,400	42.8	39.7 - 45.9	10,200
No	3,700	64.2	59.7 - 68.5	2,100	35.8	31.5 - 40.3	5,800

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,400	63.3	58.7 - 67.7	2,000	36.7	32.3 - 41.3	5,400
131-185%	1,200	57.8	50.3 - 64.9	900	42.2	35.1 - 49.7	2,100
186+%	4,300	57.3	53.7 - 60.8	3,200	42.7	39.2 - 46.3	7,500

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,600	62.3	55.4 - 68.6	1,000	37.7	31.4 - 44.6	2,600
\$10,000-\$24,999	2,200	64.7	58.9 - 70.1	1,200	35.3	29.9 - 41.1	3,300
\$25,000-\$49,999	2,300	60.5	55.1 - 65.6	1,500	39.5	34.4 - 44.9	3,800
>=\$50,000	2,900	54.2	49.9 - 58.4	2,400	45.8	41.6 - 50.1	5,300

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	62.1	52.6 - 70.8	500	37.9	29.2 - 47.4	1,200
Normal birth weight	8,800	59.6	56.9 - 62.2	6,000	40.4	37.8 - 43.1	14,800

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	65.7	55.8 - 74.4	400	34.3	25.6 - 44.2	1,200
No	8,600	59.1	56.4 - 61.7	5,900	40.9	38.3 - 43.6	14,500

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	65.2	59.2 - 70.8	1,100	34.8	29.2 - 40.8	3,100
No	7,300	58.2	55.3 - 61.0	5,300	41.8	39.0 - 44.7	12,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,100	62.8	58.6 - 66.8	2,400	37.2	33.2 - 41.4	6,600
No	5,200	57.0	53.7 - 60.2	3,900	43.0	39.8 - 46.3	9,200

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,200	60.5	58.0 - 62.9	6,000	39.5	37.1 - 42.0	15,200

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	56.4	49.0 - 63.6	800	43.6	36.4 - 51.0	1,900
Honolulu	7,000	63.4	60.5 - 66.1	4,100	36.6	33.9 - 39.5	11,100
Kauai	300	42.3	31.0 - 54.6	400	57.7	45.4 - 69.0	700
Maui	800	52.4	44.5 - 60.3	700	47.6	39.7 - 55.5	1,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	44.7	39.5 - 50.0	1,900	55.3	50.0 - 60.5	3,500
Native Hawaiian	2,500	61.3	56.0 - 66.4	1,600	38.7	33.6 - 44.0	4,100
Chinese	400	63.6	56.8 - 69.9	200	36.4	30.1 - 43.2	600
Filipino	2,100	72.9	67.8 - 77.4	800	27.1	22.6 - 32.2	2,800
Japanese	1,500	67.6	61.2 - 73.3	700	32.4	26.7 - 38.8	2,200
Other	1,200	60.4	53.5 - 66.8	800	39.6	33.2 - 46.5	2,000

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	64.1	54.9 - 72.3	500	35.9	27.7 - 45.1	1,300
20-24	2,400	63.5	58.3 - 68.4	1,400	36.5	31.6 - 41.7	3,700
25-34	4,500	59.2	55.7 - 62.6	3,100	40.8	37.4 - 44.3	7,500
35+	1,600	58.2	52.5 - 63.7	1,100	41.8	36.3 - 47.5	2,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,700	60.7	57.2 - 64.0	3,000	39.3	36.0 - 42.8	7,700
Female	4,500	60.3	56.8 - 63.8	3,000	39.7	36.2 - 43.2	7,500

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,700	57.8	54.7 - 60.8	4,200	42.2	39.2 - 45.3	9,900
No	3,500	65.5	61.3 - 69.6	1,800	34.5	30.4 - 38.7	5,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,400	63.9	59.6 - 68.0	1,900	36.1	32.0 - 40.4	5,300
131-185%	1,000	59.8	51.9 - 67.3	600	40.2	32.7 - 48.1	1,600
186+%	4,100	57.5	54.0 - 61.0	3,000	42.5	39.0 - 46.0	7,200

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,700	66.5	60.2 - 72.3	900	33.5	27.7 - 39.8	2,500
\$10,000-\$24,999	1,900	63.7	58.0 - 68.9	1,100	36.3	31.1 - 42.0	3,000
\$25,000-\$49,999	1,900	53.6	48.4 - 58.7	1,700	46.4	41.3 - 51.6	3,600
>=\$50,000	3,000	60.5	56.3 - 64.5	2,000	39.5	35.5 - 43.7	5,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	68.8	59.5 - 76.8	300	31.2	23.2 - 40.5	1,100
Normal birth weight	8,400	59.8	57.3 - 62.3	5,700	40.2	37.7 - 42.7	14,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	62.6	52.6 - 71.6	400	37.4	28.4 - 47.4	1,100
No	8,300	60.0	57.5 - 62.5	5,600	40.0	37.5 - 42.5	13,900

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	66.3	60.6 - 71.6	1,000	33.7	28.4 - 39.4	3,000
No	7,000	58.7	55.9 - 61.4	4,900	41.3	38.6 - 44.1	11,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,800	60.6	56.6 - 64.5	2,500	39.4	35.5 - 43.4	6,300
No	5,200	59.9	56.7 - 63.0	3,500	40.1	37.0 - 43.3	8,600

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,500	61.3	59.1 - 63.5	6,000	38.7	36.5 - 40.9	15,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	54.2	47.2 - 61.0	900	45.8	39.0 - 52.8	1,900
Honolulu	7,100	63.4	60.8 - 66.0	4,100	36.6	34.0 - 39.2	11,200
Kauai	300	44.5	34.6 - 55.0	400	55.5	45.0 - 65.4	800
Maui	1,100	62.7	55.8 - 69.1	600	37.3	30.9 - 44.2	1,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,500	43.1	38.5 - 47.9	2,000	56.9	52.1 - 61.5	3,600
Native Hawaiian	2,900	69.7	65.0 - 74.0	1,200	30.3	26.0 - 35.0	4,100
Chinese	400	64.0	59.4 - 68.3	200	36.0	31.7 - 40.6	600
Filipino	2,000	66.1	61.4 - 70.5	1,000	33.9	29.5 - 38.6	3,000
Japanese	1,500	69.7	63.5 - 75.3	600	30.3	24.7 - 36.5	2,100
Other	1,300	60.1	53.7 - 66.2	900	39.9	33.8 - 46.3	2,200

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	65.7	56.5 - 73.9	400	34.3	26.1 - 43.5	1,100
20-24	2,500	64.0	59.3 - 68.4	1,400	36.0	31.6 - 40.7	4,000
25-34	4,500	58.9	55.7 - 62.1	3,100	41.1	37.9 - 44.3	7,600
35+	1,900	62.3	57.3 - 67.1	1,100	37.7	32.9 - 42.7	3,000

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,300	62.7	59.7 - 65.7	3,100	37.3	34.3 - 40.3	8,400
Female	4,300	59.7	56.3 - 63.0	2,900	40.3	37.0 - 43.7	7,200

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,500	59.3	56.7 - 61.9	4,500	40.7	38.1 - 43.3	11,000
No	3,000	66.3	62.0 - 70.3	1,500	33.7	29.7 - 38.0	4,500

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	63.7	59.8 - 67.5	2,000	36.3	32.5 - 40.2	5,500
131-185%	1,000	55.2	48.4 - 61.7	800	44.8	38.3 - 51.6	1,800
186+%	4,300	60.6	57.3 - 63.7	2,800	39.4	36.3 - 42.7	7,100

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,600	68.5	62.5 - 73.9	700	31.5	26.1 - 37.5	2,300
\$10,000-\$24,999	2,200	59.8	55.0 - 64.5	1,500	40.2	35.5 - 45.0	3,700
\$25,000-\$49,999	2,400	57.7	53.3 - 62.0	1,700	42.3	38.0 - 46.7	4,100
>=\$50,000	2,700	61.1	57.0 - 65.0	1,700	38.9	35.0 - 43.0	4,500

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	75.2	66.4 - 82.4	200	24.8	17.6 - 33.6	900
Normal birth weight	8,800	60.4	58.1 - 62.7	5,800	39.6	37.3 - 41.9	14,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	67.6	58.5 - 75.5	400	32.4	24.5 - 41.5	1,100
No	8,700	60.9	58.6 - 63.1	5,600	39.1	36.9 - 41.4	14,300

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	65.2	59.9 - 70.2	1,100	34.8	29.8 - 40.1	3,000
No	7,500	60.4	57.9 - 62.8	4,900	39.6	37.2 - 42.1	12,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,200	61.9	58.4 - 65.3	2,600	38.1	34.7 - 41.6	6,800
No	5,200	60.9	58.0 - 63.8	3,300	39.1	36.2 - 42.0	8,500

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,800	60.1	57.2 - 63.0	5,800	39.9	37.0 - 42.8	14,600

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	48.8	43.6 - 54.0	800	51.2	46.0 - 56.4	1,500
Honolulu	6,800	63.3	59.5 - 66.9	4,000	36.7	33.1 - 40.5	10,800
Kauai	300	45.4	37.7 - 53.3	400	54.6	46.7 - 62.3	700
Maui	900	56.4	51.0 - 61.5	700	43.6	38.5 - 49.0	1,600

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,700	49.3	42.7 - 55.8	1,700	50.7	44.2 - 57.3	3,400
Native Hawaiian	2,200	59.0	55.7 - 62.2	1,600	41.0	37.8 - 44.3	3,800
Chinese	400	61.8	45.7 - 75.7	300	38.2	24.3 - 54.3	700
Filipino	2,000	65.7	58.7 - 72.0	1,100	34.3	28.0 - 41.3	3,100
Japanese	1,100	64.8	55.1 - 73.4	600	35.2	26.6 - 44.9	1,800
Other	1,300	67.7	58.2 - 75.8	600	32.3	24.2 - 41.8	1,900

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	67.4	58.7 - 75.1	400	32.6	24.9 - 41.3	1,300
20-24	2,000	60.6	54.7 - 66.3	1,300	39.4	33.7 - 45.3	3,300
25-34	4,400	59.7	55.6 - 63.8	3,000	40.3	36.2 - 44.4	7,400
35+	1,500	57.0	49.8 - 64.0	1,100	43.0	36.0 - 50.2	2,700

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,700	61.1	57.1 - 64.9	3,000	38.9	35.1 - 42.9	7,700
Female	4,100	59.1	54.8 - 63.2	2,800	40.9	36.8 - 45.2	6,900

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,800	57.8	54.1 - 61.4	4,200	42.2	38.6 - 45.9	10,000
No	3,000	65.2	60.7 - 69.4	1,600	34.8	30.6 - 39.3	4,600

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,900	63.6	59.3 - 67.7	2,200	36.4	32.3 - 40.7	6,100
131-185%	1,300	59.3	51.7 - 66.6	900	40.7	33.4 - 48.3	2,200
186+%	2,800	56.1	50.8 - 61.2	2,200	43.9	38.8 - 49.2	5,000

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,100	67.9	62.2 - 73.2	1,000	32.1	26.8 - 37.8	3,100
\$15,001-\$25,000	1,000	57.6	49.5 - 65.4	800	42.4	34.6 - 50.5	1,800
\$25,001-\$40,000	1,600	57.5	50.5 - 64.1	1,200	42.5	35.9 - 49.5	2,900
>=\$40,001	3,500	57.9	53.2 - 62.6	2,500	42.1	37.4 - 46.8	6,000

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	69.9	59.4 - 78.6	300	30.1	21.4 - 40.6	1,000
Normal birth weight	8,100	59.4	56.4 - 62.4	5,500	40.6	37.6 - 43.6	13,600

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	65.0	54.3 - 74.4	300	35.0	25.6 - 45.7	1,000
No	8,000	60.0	57.0 - 63.0	5,300	40.0	37.0 - 43.0	13,400

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	66.3	59.9 - 72.1	900	33.7	27.9 - 40.1	2,600
No	6,800	58.8	55.4 - 62.0	4,800	41.2	38.0 - 44.6	11,600

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,600	63.0	58.7 - 67.2	2,100	37.0	32.8 - 41.3	5,800
No	5,000	58.0	54.1 - 61.9	3,600	42.0	38.1 - 45.9	8,700

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,400	58.4	55.4 - 61.4	5,900	41.6	38.6 - 44.6	14,300

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	58.4	53.6 - 63.2	700	41.6	36.8 - 46.4	1,700
Honolulu	6,300	59.5	55.6 - 63.3	4,300	40.5	36.7 - 44.4	10,500
Kauai	200	43.7	35.5 - 52.3	300	56.3	47.7 - 64.5	600
Maui	900	56.3	50.9 - 61.5	700	43.7	38.5 - 49.1	1,500

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,200	40.3	33.6 - 47.3	1,800	59.7	52.7 - 66.4	3,100
Native Hawaiian	2,300	62.0	58.7 - 65.2	1,400	38.0	34.8 - 41.3	3,700
Chinese	400	77.4	59.4 - 88.9	100	22.6	11.1 - 40.6	500
Filipino	1,900	64.4	57.2 - 71.0	1,100	35.6	29.0 - 42.8	3,000
Japanese	1,600	68.4	60.1 - 75.7	700	31.6	24.3 - 39.9	2,300
Other	900	53.5	43.1 - 63.5	800	46.5	36.5 - 56.9	1,700

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	73.9	65.0 - 81.2	300	26.1	18.8 - 35.0	1,200
20-24	2,100	59.0	53.2 - 64.6	1,500	41.0	35.4 - 46.8	3,600
25-34	4,000	55.6	51.2 - 59.9	3,200	44.4	40.1 - 48.8	7,200
35+	1,300	58.1	50.4 - 65.5	1,000	41.9	34.5 - 49.6	2,300

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,100	58.5	54.3 - 62.7	2,900	41.5	37.3 - 45.7	7,000
Female	4,200	58.3	53.9 - 62.5	3,000	41.7	37.5 - 46.1	7,300

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,500	55.4	51.6 - 59.2	4,400	44.6	40.8 - 48.4	10,000
No	2,800	65.2	60.5 - 69.6	1,500	34.8	30.4 - 39.5	4,300

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,100	58.9	54.2 - 63.4	2,200	41.1	36.6 - 45.8	5,300
131-185%	1,600	54.5	47.6 - 61.3	1,400	45.5	38.7 - 52.4	3,000
186+%	2,800	57.1	51.8 - 62.3	2,100	42.9	37.7 - 48.2	4,900

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,500	60.6	54.1 - 66.7	1,000	39.4	33.3 - 45.9	2,500
\$15,001-\$25,000	1,100	52.9	45.0 - 60.6	1,000	47.1	39.4 - 55.0	2,100
\$25,001-\$40,000	1,800	59.4	52.6 - 65.8	1,200	40.6	34.2 - 47.4	3,000
>=\$40,001	3,400	57.1	52.3 - 61.9	2,500	42.9	38.1 - 47.7	5,900

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	70.1	57.0 - 80.5	300	29.9	19.5 - 43.0	900
Normal birth weight	7,700	57.6	54.5 - 60.6	5,700	42.4	39.4 - 45.5	13,400

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	66.1	55.4 - 75.4	300	33.9	24.6 - 44.6	900
No	7,700	58.3	55.1 - 61.4	5,500	41.7	38.6 - 44.9	13,200

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	62.3	55.5 - 68.6	1,000	37.7	31.4 - 44.5	2,600
No	6,600	57.7	54.3 - 61.0	4,800	42.3	39.0 - 45.7	11,400

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,500	61.6	57.0 - 66.1	2,200	38.4	33.9 - 43.0	5,600
No	4,900	56.4	52.4 - 60.3	3,800	43.6	39.7 - 47.6	8,600

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,200	58.5	56.0 - 60.9	5,800	41.5	39.1 - 44.0	14,000

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	800	52.3	47.1 - 57.5	700	47.7	42.5 - 52.9	1,500
Honolulu	6,300	60.6	57.5 - 63.6	4,100	39.4	36.4 - 42.5	10,400
Kauai	300	46.8	38.8 - 54.9	400	53.2	45.1 - 61.2	700
Maui	800	55.0	49.5 - 60.4	600	45.0	39.6 - 50.5	1,400

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,400	44.3	39.1 - 49.5	1,700	55.7	50.5 - 60.9	3,100
Native Hawaiian	2,200	61.0	58.3 - 63.6	1,400	39.0	36.4 - 41.7	3,600
Chinese	400	67.5	52.6 - 79.5	200	32.5	20.5 - 47.4	600
Filipino	1,800	61.5	55.4 - 67.2	1,100	38.5	32.8 - 44.6	3,000
Japanese	1,400	72.5	65.2 - 78.9	500	27.5	21.1 - 34.8	1,900
Other	1,000	55.2	47.3 - 62.9	800	44.8	37.1 - 52.7	1,800

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	71.8	65.4 - 77.4	400	28.2	22.6 - 34.6	1,300
20-24	2,200	61.0	56.4 - 65.4	1,400	39.0	34.6 - 43.6	3,600
25-34	3,700	54.2	50.5 - 57.7	3,100	45.8	42.3 - 49.5	6,800
35+	1,400	60.0	53.6 - 66.0	900	40.0	34.0 - 46.4	2,300

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,300	59.1	55.7 - 62.4	3,000	40.9	37.6 - 44.3	7,300
Female	3,900	57.8	54.3 - 61.3	2,800	42.2	38.7 - 45.7	6,700

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,500	56.7	53.6 - 59.7	4,200	43.3	40.3 - 46.4	9,600
No	2,700	62.5	58.6 - 66.4	1,600	37.5	33.6 - 41.4	4,400

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,000	60.5	56.6 - 64.3	2,000	39.5	35.7 - 43.4	5,000
131-185%	1,700	55.9	50.7 - 60.9	1,400	44.1	39.1 - 49.3	3,100
186+%	2,800	56.5	52.1 - 60.7	2,200	43.5	39.3 - 47.9	5,000

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,700	65.1	59.8 - 70.0	900	34.9	30.0 - 40.2	2,600
\$15,001-\$25,000	1,100	57.0	50.5 - 63.3	800	43.0	36.7 - 49.5	1,900
\$25,001-\$40,000	1,700	56.1	51.0 - 61.1	1,400	43.9	38.9 - 49.0	3,100
>=\$40,001	3,200	56.4	52.3 - 60.3	2,500	43.6	39.7 - 47.7	5,800

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	64.8	54.7 - 73.7	300	35.2	26.3 - 45.3	900
Normal birth weight	7,600	58.0	55.5 - 60.5	5,500	42.0	39.5 - 44.5	13,100

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	68.8	60.8 - 75.8	300	31.2	24.2 - 39.2	1,100
No	7,300	57.4	54.8 - 60.0	5,400	42.6	40.0 - 45.2	12,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	60.2	54.9 - 65.4	1,100	39.8	34.6 - 45.1	2,700
No	6,300	57.7	54.9 - 60.5	4,600	42.3	39.5 - 45.1	10,900

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,500	60.0	56.4 - 63.4	2,300	40.0	36.6 - 43.6	5,800
No	4,700	57.4	54.1 - 60.7	3,500	42.6	39.3 - 45.9	8,200

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	6,500	54.9	52.2 - 57.5	5,400	45.1	42.5 - 47.8	11,900

County	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	700	53.2	47.6 - 58.7	600	46.8	41.3 - 52.4	1,300
Honolulu	4,900	55.3	52.0 - 58.5	3,900	44.7	41.5 - 48.0	8,800
Kauai	200	42.5	33.6 - 52.0	300	57.5	48.0 - 66.4	500
Maui	700	58.8	52.7 - 64.7	500	41.2	35.3 - 47.3	1,200

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,300	45.0	39.6 - 50.6	1,600	55.0	49.4 - 60.4	2,800
Native Hawaiian	1,900	60.8	57.8 - 63.6	1,300	39.2	36.4 - 42.2	3,200
Chinese	300	52.1	37.0 - 66.8	200	47.9	33.2 - 63.0	500
Filipino	1,600	63.7	57.3 - 69.6	900	36.3	30.4 - 42.7	2,500
Japanese	700	50.8	41.6 - 59.9	700	49.2	40.1 - 58.4	1,300
Other	800	51.1	42.5 - 59.6	800	48.9	40.4 - 57.5	1,600

PRAMS Age Group	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	60.7	53.7 - 67.3	600	39.3	32.7 - 46.3	1,400
20-24	1,600	56.0	51.0 - 60.9	1,300	44.0	39.1 - 49.0	2,900
25-34	3,000	52.9	49.0 - 56.9	2,700	47.1	43.1 - 51.0	5,600
35+	1,000	54.5	47.7 - 61.1	900	45.5	38.9 - 52.3	1,900

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	3,300	53.6	50.0 - 57.2	2,800	46.4	42.8 - 50.0	6,100
Female	3,200	56.2	52.4 - 59.9	2,500	43.8	40.1 - 47.6	5,800

Married	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,100	51.5	48.2 - 54.8	3,900	48.5	45.2 - 51.8	8,000
No	2,400	61.8	57.5 - 65.8	1,500	38.2	34.2 - 42.5	3,900

Poverty Level	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,900	56.3	52.4 - 60.1	2,200	43.7	39.9 - 47.6	5,100
131-185%	900	56.7	49.7 - 63.3	700	43.3	36.7 - 50.3	1,600
186+%	2,200	51.0	46.5 - 55.5	2,100	49.0	44.5 - 53.5	4,400

**PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,600	58.4	53.1 - 63.5	1,100	41.6	36.5 - 46.9	2,800
\$15,001-\$25,000	1,000	54.2	47.6 - 60.8	900	45.8	39.2 - 52.4	1,900
\$25,001-\$40,000	1,300	54.8	48.9 - 60.5	1,100	45.2	39.5 - 51.1	2,300
>=\$40,001	2,300	52.4	47.9 - 56.9	2,100	47.6	43.1 - 52.1	4,300

Birthweight	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	62.6	52.1 - 72.0	300	37.4	28.0 - 47.9	800
Normal birth weight	6,000	54.3	51.6 - 57.0	5,000	45.7	43.0 - 48.4	11,000

Smoke 3rd Trimester	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	70.2	61.8 - 77.3	300	29.8	22.7 - 38.2	900
No	5,800	53.7	50.9 - 56.5	5,000	46.3	43.5 - 49.1	10,700

PRAMS Health Indicator Report - Breastfeeding (exclusive) duration 8+ weeks, among initiators- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	60.8	55.2 - 66.2	900	39.2	33.8 - 44.8	2,300
No	4,900	53.1	50.1 - 56.2	4,300	46.9	43.8 - 49.9	9,200

WIC During Pregnancy	Breastfed <8 weeks			Breastfed 8+ weeks			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	56.1	52.3 - 59.9	2,300	43.9	40.1 - 47.7	5,200
No	3,500	53.9	50.2 - 57.5	3,000	46.1	42.5 - 49.8	6,600