

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- for the State of Hawaii, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 3-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, income data does not exist for some 3-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Sleeping position	Sleeping position

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/16/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- for the State of Hawaii, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2002	10,300	64.1	62.8 - 65.4	2,700	17.1	16.0 - 18.2	2,000	12.7	11.8 - 13.6	1,000	6.2	5.6 - 6.8	16,100
2001-2003	10,700	65.9	64.5 - 67.3	2,800	17.2	16.1 - 18.4	1,900	11.9	11.0 - 12.9	800	4.9	4.3 - 5.5	16,300
2002-2004	11,300	67.7	66.3 - 69.0	2,900	17.5	16.4 - 18.6	2,000	12.2	11.3 - 13.2	400	2.7	2.3 - 3.2	16,800
2003-2005	11,500	67.8	66.5 - 69.1	2,800	16.6	15.6 - 17.7	2,000	11.9	11.1 - 12.9	600	3.6	3.1 - 4.2	17,000
2004-2006	11,600	67.3	66.0 - 68.6	2,700	15.6	14.6 - 16.6	2,100	12.0	11.2 - 13.0	900	5.1	4.5 - 5.7	17,300
2005-2007	12,000	68.7	67.4 - 70.0	2,500	14.1	13.1 - 15.1	2,000	11.6	10.8 - 12.6	1,000	5.6	5.0 - 6.3	17,500
2006-2008	12,600	70.1	68.8 - 71.4	2,600	14.4	13.4 - 15.4	2,100	11.5	10.7 - 12.5	700	3.9	3.4 - 4.6	17,900
2007-2009	13,300	73.4	72.0 - 74.7	2,300	12.8	11.8 - 13.8	1,900	10.7	9.8 - 11.7	600	3.1	2.6 - 3.7	18,100
2008-2010	13,500	74.3	72.8 - 75.8	2,300	12.6	11.5 - 13.7	1,800	10.0	9.0 - 11.1	600	3.1	2.6 - 3.7	18,200
2009-2011	13,900	76.5	74.9 - 78.1	2,200	11.9	10.8 - 13.2	1,500	8.3	7.3 - 9.4	600	3.3	2.7 - 4.0	18,200
2010-2012	14,300	78.2	76.6 - 79.7	2,200	11.8	10.6 - 13.1	1,300	7.2	6.3 - 8.2	500	2.8	2.3 - 3.5	18,200
2011-2013	14,500	79.8	78.2 - 81.3	2,100	11.4	10.2 - 12.7	1,200	6.7	5.8 - 7.8	400	2.0	1.6 - 2.7	18,100

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- for the State of Hawaii, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2011-2013

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	14,500	79.8	78.2 - 81.3	2,100	11.4	10.2 - 12.7	1,200	6.7	5.8 - 7.8	400	2.0	1.6 - 2.7	18,100

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,800	77.1	74.4 - 79.6	300	13.0	11.0 - 15.3	200	7.5	6.0 - 9.3	100	2.4	1.6 - 3.5	2,300
Honolulu	10,500	80.8	78.7 - 82.8	1,400	10.8	9.3 - 12.5	800	6.5	5.3 - 7.9	300	1.9	1.3 - 2.8	13,100
Kauai	600	77.2	74.1 - 80.0	100	13.8	11.6 - 16.4	100	7.6	5.9 - 9.7	<50	1.4	0.8 - 2.6	800
Maui	1,500	77.3	74.7 - 79.6	200	12.7	10.9 - 14.8	100	7.4	5.9 - 9.1	100	2.7	1.8 - 3.8	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2011-2013

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,800	84.6	81.6 - 87.2	400	8.5	6.6 - 10.9	200	5.5	4.0 - 7.5	100	1.4	0.7 - 2.8	4,500
Native Hawaiian	3,800	73.9	70.7 - 76.8	700	13.2	11.0 - 15.8	500	9.6	7.7 - 11.9	200	3.3	2.2 - 4.8	5,200
Chinese	800	89.7	83.0 - 93.9	100	7.1	3.7 - 13.4	<50	3.2	1.3 - 7.7	0	0.0	-	900
Filipino	2,500	81.4	77.6 - 84.7	400	12.3	9.6 - 15.6	100	4.5	2.8 - 7.0	100	1.8	1.1 - 3.2	3,100
Japanese	1,500	86.2	81.4 - 89.9	100	6.1	3.8 - 9.7	100	7.4	4.7 - 11.5	<50	0.2	0.1 - 0.9	1,800
Other	2,000	73.9	68.9 - 78.3	400	16.6	13.0 - 21.0	200	6.7	4.5 - 9.7	100	2.8	1.5 - 5.3	2,700

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	70.3	62.4 - 77.2	200	16.2	11.2 - 23.0	100	10.5	6.3 - 17.1	<50	2.9	1.3 - 6.6	1,000
20-24	3,000	75.2	71.3 - 78.6	500	12.9	10.3 - 16.1	300	8.1	6.1 - 10.7	200	3.8	2.5 - 5.7	4,000
25-34	8,000	81.4	79.3 - 83.3	1,100	10.8	9.3 - 12.6	600	6.0	4.9 - 7.4	200	1.7	1.2 - 2.5	9,900
35+	2,700	83.4	79.9 - 86.4	300	9.8	7.6 - 12.7	200	6.0	4.2 - 8.6	<50	0.7	0.3 - 1.7	3,300

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2011-2013

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,200	79.1	76.7 - 81.2	1,000	11.1	9.5 - 13.0	700	7.6	6.3 - 9.2	200	2.2	1.5 - 3.2	9,000
Female	7,300	80.5	78.3 - 82.5	1,100	11.7	10.1 - 13.6	500	5.9	4.7 - 7.3	200	1.9	1.3 - 2.7	9,100

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,300	82.1	80.1 - 83.9	1,200	10.7	9.2 - 12.3	700	5.9	4.8 - 7.1	200	1.4	0.9 - 2.1	11,300
No	5,200	75.9	73.2 - 78.5	900	12.7	10.8 - 14.8	600	8.2	6.6 - 10.1	200	3.2	2.3 - 4.4	6,800

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,000	77.1	74.3 - 79.6	800	12.9	11.0 - 15.1	500	7.2	5.7 - 9.0	200	2.8	2.0 - 4.0	6,500
131-185%	2,200	80.6	76.1 - 84.3	300	11.5	8.6 - 15.4	200	6.3	4.2 - 9.2	<50	1.6	0.8 - 3.5	2,700
186+%	6,100	83.0	80.5 - 85.1	700	9.5	7.9 - 11.5	500	6.2	4.9 - 7.9	100	1.3	0.7 - 2.2	7,400

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2011-2013

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	80.1	77.4 - 82.6	100	11.4	9.5 - 13.7	100	6.3	4.9 - 8.0	<50	2.1	1.4 - 3.3	1,200
Normal birth weight	13,500	79.8	78.1 - 81.4	1,900	11.4	10.2 - 12.8	1,100	6.8	5.8 - 7.9	300	2.0	1.5 - 2.7	16,900

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	67.5	59.3 - 74.7	200	17.9	12.4 - 25.1	100	11.0	6.7 - 17.5	<50	3.6	1.6 - 8.2	800
No	13,600	80.3	78.7 - 81.9	1,900	11.1	9.9 - 12.5	1,100	6.6	5.6 - 7.6	300	2.0	1.5 - 2.6	16,900

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	74.1	69.9 - 77.8	400	12.0	9.5 - 15.2	300	9.5	7.1 - 12.5	100	4.4	2.8 - 6.8	3,200
No	11,800	81.0	79.2 - 82.6	1,700	11.3	10.0 - 12.8	900	6.2	5.2 - 7.3	200	1.5	1.1 - 2.1	14,600

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2011-2013

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,200	76.0	73.4 - 78.3	1,100	13.6	11.8 - 15.7	600	7.5	6.1 - 9.1	200	2.9	2.1 - 4.0	8,100
No	8,200	82.9	80.8 - 84.7	1,000	9.6	8.2 - 11.3	600	6.2	5.1 - 7.6	100	1.3	0.9 - 2.1	9,900

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	14,300	78.2	76.6 - 79.7	2,200	11.8	10.6 - 13.1	1,300	7.2	6.3 - 8.2	500	2.8	2.3 - 3.5	18,200

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,700	75.1	72.4 - 77.6	300	14.3	12.3 - 16.6	200	7.4	6.0 - 9.1	100	3.1	2.2 - 4.4	2,300
Honolulu	10,400	79.3	77.2 - 81.2	1,400	11.0	9.5 - 12.7	900	7.0	5.8 - 8.3	400	2.7	2.0 - 3.7	13,100
Kauai	600	74.8	71.5 - 77.8	100	15.8	13.4 - 18.6	100	8.0	6.2 - 10.2	<50	1.4	0.8 - 2.6	800
Maui	1,500	76.2	73.8 - 78.5	200	12.3	10.6 - 14.3	200	8.1	6.7 - 9.7	100	3.4	2.5 - 4.6	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,600	82.7	79.6 - 85.3	400	9.6	7.6 - 12.0	300	6.5	4.9 - 8.6	100	1.3	0.6 - 2.5	4,300
Native Hawaiian	3,900	73.8	70.8 - 76.7	600	12.2	10.2 - 14.6	500	10.0	8.1 - 12.2	200	4.0	2.8 - 5.5	5,200
Chinese	700	85.6	78.2 - 90.7	100	8.7	5.0 - 14.9	<50	4.7	2.0 - 10.6	<50	1.0	0.1 - 6.5	900
Filipino	2,500	78.2	74.3 - 81.6	400	13.5	10.8 - 16.8	100	4.5	2.9 - 6.8	100	3.9	2.6 - 5.8	3,300
Japanese	1,500	85.3	80.6 - 89.0	100	7.1	4.6 - 10.9	100	6.8	4.4 - 10.2	<50	0.8	0.2 - 2.8	1,700
Other	2,100	73.1	68.3 - 77.4	500	16.2	12.7 - 20.5	200	7.1	4.9 - 10.3	100	3.6	2.1 - 5.9	2,800

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	73.5	66.6 - 79.4	200	16.3	11.4 - 22.6	100	6.8	4.1 - 11.0	<50	3.4	1.8 - 6.7	1,100
20-24	3,100	73.7	70.0 - 77.1	500	12.4	10.0 - 15.3	400	9.2	7.1 - 12.0	200	4.7	3.3 - 6.7	4,100
25-34	7,800	79.6	77.5 - 81.6	1,100	11.0	9.5 - 12.7	700	6.9	5.7 - 8.3	200	2.5	1.8 - 3.4	9,800
35+	2,600	81.3	77.7 - 84.5	400	12.0	9.4 - 15.2	200	5.5	3.9 - 7.8	<50	1.1	0.5 - 2.3	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,200	78.4	76.2 - 80.5	1,000	10.8	9.3 - 12.6	700	7.9	6.6 - 9.5	300	2.8	2.1 - 3.9	9,200
Female	7,100	78.0	75.8 - 80.1	1,200	12.8	11.1 - 14.6	600	6.4	5.3 - 7.8	300	2.8	2.0 - 3.7	9,100

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,200	80.8	78.8 - 82.6	1,200	10.8	9.4 - 12.4	700	6.5	5.5 - 7.8	200	1.9	1.3 - 2.7	11,300
No	5,100	74.0	71.3 - 76.5	900	13.5	11.6 - 15.6	600	8.2	6.7 - 10.0	300	4.3	3.2 - 5.7	6,900

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,800	73.8	71.0 - 76.5	900	13.9	11.9 - 16.2	500	8.4	6.8 - 10.4	300	3.9	2.9 - 5.2	6,400
131-185%	2,300	80.8	76.7 - 84.2	300	10.3	7.7 - 13.6	200	6.5	4.6 - 9.3	100	2.4	1.3 - 4.2	2,900
186+%	6,200	81.9	79.5 - 84.0	700	9.7	8.1 - 11.5	500	6.8	5.5 - 8.4	100	1.7	1.0 - 2.6	7,600

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	78.4	75.6 - 81.0	100	10.5	8.6 - 12.7	100	7.3	5.8 - 9.2	<50	3.8	2.7 - 5.3	1,200
Normal birth weight	13,300	78.2	76.5 - 79.8	2,000	11.9	10.7 - 13.2	1,200	7.2	6.2 - 8.3	500	2.7	2.2 - 3.5	17,000

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	63.8	55.9 - 71.0	200	19.7	14.1 - 26.7	100	11.9	7.5 - 18.4	<50	4.7	2.3 - 9.1	1,000
No	13,300	79.0	77.3 - 80.5	1,900	11.4	10.2 - 12.7	1,200	7.0	6.1 - 8.0	500	2.7	2.1 - 3.4	16,900

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	69.9	65.8 - 73.7	500	14.5	11.8 - 17.8	300	10.2	7.9 - 13.1	200	5.4	3.7 - 7.7	3,400
No	11,600	80.0	78.3 - 81.6	1,600	11.2	9.9 - 12.6	1,000	6.6	5.6 - 7.7	300	2.2	1.7 - 2.9	14,500

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2010-2012

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,000	73.7	71.2 - 76.1	1,100	14.1	12.2 - 16.1	600	8.0	6.6 - 9.6	300	4.3	3.3 - 5.6	8,100
No	8,100	82.0	79.9 - 83.8	1,000	9.9	8.5 - 11.5	600	6.5	5.4 - 7.9	200	1.6	1.1 - 2.4	9,900

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	13,900	76.5	74.9 - 78.1	2,200	11.9	10.8 - 13.2	1,500	8.3	7.3 - 9.4	600	3.3	2.7 - 4.0	18,200

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,700	72.7	70.0 - 75.3	400	15.2	13.2 - 17.5	200	8.1	6.6 - 9.9	100	4.0	3.0 - 5.4	2,300
Honolulu	10,200	78.1	75.9 - 80.1	1,400	10.9	9.4 - 12.6	1,000	8.0	6.7 - 9.4	400	3.1	2.3 - 4.1	13,000
Kauai	600	70.6	67.3 - 73.7	100	17.1	14.6 - 19.9	100	9.9	8.0 - 12.2	<50	2.4	1.5 - 3.7	800
Maui	1,500	73.3	70.8 - 75.7	300	12.7	11.0 - 14.6	200	10.1	8.6 - 11.9	100	3.9	2.9 - 5.1	2,000

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,600	81.3	78.3 - 84.0	400	9.8	7.9 - 12.1	300	7.5	5.8 - 9.7	100	1.3	0.7 - 2.6	4,400
Native Hawaiian	3,900	72.2	69.2 - 75.0	700	13.0	11.0 - 15.4	600	10.4	8.6 - 12.5	200	4.4	3.2 - 5.9	5,500
Chinese	600	79.6	70.7 - 86.3	100	13.2	8.0 - 21.1	<50	6.1	2.7 - 12.9	<50	1.1	0.2 - 7.3	800
Filipino	2,500	76.5	72.6 - 80.0	400	13.6	10.9 - 16.9	200	5.5	3.8 - 7.8	100	4.5	3.0 - 6.6	3,300
Japanese	1,500	85.2	80.5 - 89.0	100	5.1	3.0 - 8.3	200	8.9	6.0 - 12.9	<50	0.8	0.2 - 2.8	1,700
Other	1,900	71.1	66.2 - 75.6	400	15.0	11.6 - 19.2	200	9.0	6.4 - 12.5	100	4.9	3.1 - 7.5	2,600

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	72.7	66.3 - 78.2	200	14.6	10.3 - 20.2	100	7.9	5.3 - 11.8	100	4.9	2.7 - 8.6	1,300
20-24	3,100	72.1	68.4 - 75.5	500	12.7	10.3 - 15.6	400	10.1	8.0 - 12.8	200	5.1	3.6 - 7.1	4,300
25-34	7,400	78.1	75.9 - 80.1	1,000	11.0	9.5 - 12.7	800	8.0	6.8 - 9.5	300	2.9	2.1 - 3.9	9,400
35+	2,500	79.4	75.7 - 82.7	400	12.5	9.9 - 15.7	200	6.8	4.9 - 9.3	<50	1.3	0.6 - 2.5	3,200

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Infant Gender													
Male	7,200	76.4	74.1 - 78.5	1,100	11.6	10.0 - 13.3	800	8.8	7.4 - 10.3	300	3.3	2.5 - 4.4	9,400
Female	6,800	76.7	74.4 - 78.9	1,100	12.3	10.6 - 14.1	700	7.8	6.5 - 9.4	300	3.2	2.4 - 4.2	8,800

	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Married													
Yes	9,000	80.1	78.2 - 81.9	1,200	10.4	9.1 - 11.9	800	7.4	6.3 - 8.7	200	2.0	1.4 - 2.8	11,200
No	5,000	70.8	68.0 - 73.4	1,000	14.3	12.3 - 16.5	700	9.7	8.1 - 11.6	400	5.3	4.1 - 6.7	7,000

	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Poverty Level													
0-130%	4,500	70.3	67.4 - 73.1	900	14.8	12.7 - 17.1	700	10.3	8.5 - 12.3	300	4.6	3.5 - 6.1	6,400
131-185%	2,200	78.0	73.7 - 81.7	300	11.4	8.7 - 14.8	200	7.6	5.4 - 10.6	100	3.0	1.8 - 5.1	2,800
186+%	6,300	81.9	79.7 - 84.0	700	9.2	7.7 - 11.0	500	7.0	5.8 - 8.6	100	1.8	1.2 - 2.8	7,700

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,400	73.5	69.5 - 77.1	400	12.9	10.2 - 16.2	300	9.1	6.9 - 11.9	100	4.5	3.1 - 6.6	3,200
\$10,000-\$24,999	2,400	69.4	65.3 - 73.1	500	15.1	12.3 - 18.4	300	9.8	7.5 - 12.7	200	5.7	4.0 - 8.1	3,500
\$25,000-\$49,999	3,400	79.4	76.2 - 82.4	500	11.0	8.8 - 13.7	300	7.4	5.7 - 9.7	100	2.1	1.3 - 3.5	4,200
>=\$50,000	5,000	81.3	78.7 - 83.6	600	9.4	7.7 - 11.4	500	7.9	6.4 - 9.8	100	1.4	0.8 - 2.5	6,100

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	76.7	73.8 - 79.3	100	11.3	9.4 - 13.5	100	7.9	6.3 - 9.8	100	4.1	3.0 - 5.7	1,200
Normal birth weight	13,000	76.5	74.8 - 78.1	2,000	12.0	10.7 - 13.3	1,400	8.3	7.3 - 9.5	500	3.2	2.6 - 4.0	17,000

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	61.7	54.9 - 68.1	300	20.5	15.3 - 26.8	200	12.6	8.8 - 17.7	100	5.2	3.0 - 9.0	1,300
No	12,900	77.7	76.1 - 79.3	1,900	11.3	10.1 - 12.5	1,300	7.9	6.9 - 9.0	500	3.1	2.5 - 3.8	16,600

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	68.2	64.3 - 71.8	600	15.4	12.6 - 18.6	400	10.6	8.4 - 13.2	200	5.9	4.3 - 8.2	3,700
No	11,200	78.7	76.9 - 80.4	1,600	11.0	9.8 - 12.4	1,100	7.7	6.6 - 8.9	400	2.6	2.0 - 3.3	14,200

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,500	70.8	68.2 - 73.3	1,100	14.5	12.7 - 16.6	700	9.5	8.0 - 11.3	400	5.2	4.1 - 6.5	7,800
No	8,200	81.0	79.0 - 82.9	1,000	9.8	8.4 - 11.4	700	7.3	6.1 - 8.7	200	1.9	1.3 - 2.7	10,100

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	13,500	74.3	72.8 - 75.8	2,300	12.6	11.5 - 13.7	1,800	10.0	9.0 - 11.1	600	3.1	2.6 - 3.7	18,200

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,700	70.7	67.5 - 73.7	400	16.1	13.8 - 18.8	200	9.7	7.8 - 11.9	100	3.5	2.5 - 5.0	2,300
Honolulu	9,800	76.1	74.1 - 77.9	1,500	11.2	9.9 - 12.7	1,300	9.9	8.7 - 11.3	400	2.8	2.2 - 3.7	12,900
Kauai	600	68.7	64.2 - 72.9	100	17.7	14.4 - 21.5	100	10.2	7.8 - 13.2	<50	3.5	2.0 - 6.1	800
Maui	1,400	69.9	66.7 - 72.9	300	14.9	12.6 - 17.5	200	11.0	9.1 - 13.3	100	4.2	3.1 - 5.8	2,100

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,400	79.1	76.3 - 81.7	400	10.3	8.5 - 12.5	400	9.7	7.9 - 11.9	<50	0.8	0.4 - 1.5	4,300
Native Hawaiian	3,700	68.4	65.4 - 71.2	800	14.8	12.7 - 17.1	700	12.9	11.0 - 15.1	200	3.9	2.9 - 5.3	5,400
Chinese	500	76.2	67.9 - 82.9	100	14.4	9.1 - 22.0	100	8.0	4.4 - 14.0	<50	1.4	0.3 - 6.9	700
Filipino	2,500	75.4	71.9 - 78.6	400	13.4	11.0 - 16.3	200	6.8	5.1 - 9.2	100	4.3	2.9 - 6.3	3,300
Japanese	1,400	82.7	78.3 - 86.4	100	6.6	4.5 - 9.7	200	9.5	6.8 - 13.2	<50	1.1	0.4 - 3.2	1,700
Other	1,900	71.2	66.9 - 75.2	400	14.1	11.2 - 17.7	200	9.3	6.9 - 12.3	100	5.4	3.7 - 7.8	2,700

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	69.7	63.9 - 74.9	200	15.5	11.6 - 20.4	100	9.6	6.9 - 13.2	100	5.3	3.1 - 8.9	1,400
20-24	2,900	69.6	66.2 - 72.8	600	13.2	11.0 - 15.8	500	12.4	10.2 - 15.0	200	4.8	3.4 - 6.6	4,200
25-34	7,100	76.1	74.0 - 78.0	1,100	11.5	10.1 - 13.1	900	9.7	8.4 - 11.2	300	2.7	2.0 - 3.6	9,300
35+	2,500	77.5	74.0 - 80.6	400	13.5	11.0 - 16.5	300	8.0	6.1 - 10.4	<50	1.0	0.5 - 2.0	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,900	72.9	70.8 - 75.0	1,200	13.2	11.7 - 14.8	1,000	10.9	9.5 - 12.4	300	3.0	2.3 - 3.9	9,500
Female	6,600	75.8	73.7 - 77.9	1,000	11.9	10.5 - 13.6	800	9.1	7.8 - 10.6	300	3.1	2.4 - 4.1	8,700

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,700	77.9	76.1 - 79.5	1,200	11.1	9.9 - 12.5	1,100	9.5	8.3 - 10.8	200	1.6	1.2 - 2.2	11,200
No	4,800	68.7	66.1 - 71.2	1,000	14.9	13.0 - 17.0	800	10.9	9.3 - 12.7	400	5.5	4.3 - 6.9	7,000

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,400	68.4	65.7 - 71.0	1,000	16.1	14.1 - 18.3	700	11.4	9.7 - 13.3	300	4.1	3.1 - 5.4	6,500
131-185%	2,100	73.8	69.8 - 77.4	300	12.4	9.8 - 15.6	300	11.0	8.6 - 14.0	100	2.8	1.6 - 4.8	2,800
186+%	6,100	80.3	78.2 - 82.3	700	9.3	8.0 - 10.9	700	8.8	7.4 - 10.3	100	1.5	1.0 - 2.4	7,600

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,200	69.6	65.6 - 73.3	500	14.6	11.9 - 17.8	300	11.2	8.8 - 14.1	100	4.6	3.2 - 6.7	3,100
\$10,000-\$24,999	2,500	67.9	64.2 - 71.3	600	15.9	13.3 - 18.9	400	11.2	9.0 - 13.8	200	5.1	3.6 - 7.1	3,700
\$25,000-\$49,999	3,100	76.2	73.1 - 79.0	500	11.8	9.7 - 14.2	400	10.3	8.4 - 12.6	100	1.7	1.1 - 2.8	4,100
>=\$50,000	4,900	80.3	77.9 - 82.5	600	9.5	8.0 - 11.3	600	9.1	7.6 - 10.9	100	1.1	0.6 - 2.0	6,100

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	73.7	69.9 - 77.2	200	13.8	11.0 - 17.0	100	8.2	6.2 - 10.6	100	4.4	3.0 - 6.4	1,200
Normal birth weight	12,600	74.4	72.8 - 75.9	2,100	12.5	11.4 - 13.7	1,700	10.1	9.1 - 11.3	500	3.0	2.4 - 3.7	17,000

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	59.0	53.0 - 64.8	300	21.2	16.5 - 26.9	200	15.4	11.6 - 20.1	100	4.3	2.5 - 7.3	1,500
No	12,400	75.7	74.2 - 77.2	1,900	11.8	10.7 - 12.9	1,600	9.5	8.5 - 10.6	500	3.0	2.4 - 3.6	16,300

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	66.9	63.2 - 70.3	600	16.4	13.8 - 19.5	400	11.6	9.5 - 14.2	200	5.1	3.7 - 7.0	3,700
No	10,800	76.2	74.6 - 77.8	1,600	11.6	10.4 - 12.8	1,400	9.6	8.5 - 10.8	400	2.6	2.0 - 3.2	14,200

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,300	69.0	66.6 - 71.4	1,100	14.4	12.7 - 16.3	900	11.6	10.1 - 13.4	400	4.9	3.9 - 6.2	7,700
No	8,000	78.3	76.4 - 80.1	1,100	11.1	9.8 - 12.6	900	8.8	7.7 - 10.2	200	1.8	1.3 - 2.5	10,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	13,300	73.4	72.0 - 74.7	2,300	12.8	11.8 - 13.8	1,900	10.7	9.8 - 11.7	600	3.1	2.6 - 3.7	18,100

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,600	68.5	65.0 - 71.9	400	17.2	14.5 - 20.2	300	10.7	8.6 - 13.3	100	3.6	2.4 - 5.3	2,400
Honolulu	9,600	75.5	73.8 - 77.1	1,400	11.3	10.1 - 12.5	1,300	10.5	9.3 - 11.7	400	2.8	2.2 - 3.6	12,700
Kauai	600	68.3	63.0 - 73.3	100	15.9	12.3 - 20.3	100	9.7	7.0 - 13.2	100	6.1	3.7 - 9.9	900
Maui	1,500	68.6	64.9 - 72.0	300	15.6	12.9 - 18.6	300	12.6	10.3 - 15.3	100	3.3	2.1 - 5.0	2,100

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,300	79.3	76.7 - 81.7	400	10.4	8.7 - 12.3	400	9.1	7.5 - 11.0	100	1.2	0.7 - 2.1	4,200
Native Hawaiian	3,400	65.1	62.2 - 67.9	800	16.0	13.9 - 18.3	800	14.6	12.7 - 16.8	200	4.3	3.2 - 5.7	5,300
Chinese	600	81.3	75.3 - 86.1	100	11.9	7.8 - 17.9	<50	5.7	3.6 - 8.9	<50	1.1	0.5 - 2.3	700
Filipino	2,400	75.1	72.0 - 78.0	400	13.4	11.3 - 15.8	300	8.2	6.4 - 10.3	100	3.4	2.3 - 4.9	3,300
Japanese	1,600	84.2	80.5 - 87.3	100	6.2	4.4 - 8.6	200	9.1	6.7 - 12.4	<50	0.5	0.1 - 1.9	1,900
Other	1,900	68.8	64.9 - 72.4	400	14.3	11.7 - 17.3	300	11.0	8.7 - 13.8	200	5.9	4.2 - 8.2	2,800

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	64.9	59.4 - 70.0	300	15.9	12.2 - 20.5	200	12.7	9.6 - 16.6	100	6.5	4.2 - 10.1	1,600
20-24	2,800	67.3	64.2 - 70.2	600	14.9	12.7 - 17.3	600	13.2	11.2 - 15.5	200	4.7	3.4 - 6.3	4,200
25-34	6,900	75.7	73.8 - 77.5	1,100	11.8	10.5 - 13.2	900	10.1	8.8 - 11.4	200	2.5	1.9 - 3.2	9,100
35+	2,500	79.3	76.3 - 82.0	400	11.2	9.2 - 13.6	300	8.2	6.4 - 10.4	<50	1.3	0.7 - 2.3	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,900	72.3	70.4 - 74.2	1,300	13.4	12.0 - 14.9	1,100	11.2	9.9 - 12.6	300	3.1	2.5 - 4.0	9,500
Female	6,400	74.6	72.7 - 76.5	1,000	12.1	10.8 - 13.5	900	10.2	8.9 - 11.6	300	3.1	2.4 - 4.0	8,600

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,600	77.0	75.4 - 78.6	1,300	11.6	10.5 - 12.8	1,100	9.9	8.8 - 11.1	200	1.5	1.1 - 2.0	11,100
No	4,700	67.6	65.2 - 70.0	1,000	14.7	12.9 - 16.6	800	12.0	10.4 - 13.7	400	5.8	4.7 - 7.1	7,000

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,500	68.4	66.0 - 70.8	1,100	16.1	14.3 - 18.1	800	11.5	10.0 - 13.2	300	4.0	3.1 - 5.2	6,600
131-185%	1,900	70.5	66.7 - 74.1	400	13.9	11.3 - 16.9	400	13.5	11.0 - 16.5	100	2.1	1.2 - 3.9	2,700
186+%	6,000	80.1	78.3 - 81.9	700	8.9	7.7 - 10.2	700	9.3	8.0 - 10.7	100	1.6	1.1 - 2.4	7,500

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,100	68.9	65.2 - 72.4	500	15.5	12.9 - 18.6	400	11.8	9.6 - 14.6	100	3.8	2.5 - 5.6	3,000
\$10,000-\$24,999	2,600	67.8	64.5 - 70.8	600	15.5	13.2 - 18.0	400	11.7	9.7 - 14.0	200	5.1	3.7 - 7.0	3,800
\$25,000-\$49,999	3,000	73.8	70.9 - 76.5	500	13.4	11.4 - 15.8	500	11.3	9.5 - 13.5	100	1.4	0.9 - 2.3	4,000
>=\$50,000	4,900	80.6	78.5 - 82.5	500	8.6	7.3 - 10.0	600	9.5	8.1 - 11.2	100	1.3	0.8 - 2.1	6,100

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	73.0	68.6 - 77.0	200	13.6	10.7 - 17.3	100	9.3	6.9 - 12.3	100	4.1	2.5 - 6.7	1,300
Normal birth weight	12,300	73.4	72.0 - 74.8	2,100	12.7	11.7 - 13.8	1,800	10.8	9.9 - 11.9	500	3.0	2.5 - 3.7	16,800

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	60.9	55.5 - 66.1	300	20.2	16.0 - 25.1	200	15.1	11.7 - 19.2	100	3.8	2.2 - 6.5	1,600
No	12,100	74.7	73.3 - 76.0	1,900	11.9	11.0 - 13.0	1,700	10.3	9.3 - 11.3	500	3.1	2.6 - 3.8	16,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	67.0	63.6 - 70.2	600	15.6	13.2 - 18.4	500	12.7	10.6 - 15.1	200	4.7	3.4 - 6.6	3,600
No	10,600	75.1	73.6 - 76.6	1,700	11.9	10.8 - 13.0	1,400	10.2	9.2 - 11.3	400	2.8	2.3 - 3.4	14,100

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,100	67.7	65.4 - 69.9	1,200	15.4	13.8 - 17.2	900	11.9	10.4 - 13.6	400	5.0	4.1 - 6.2	7,500
No	7,900	77.6	75.9 - 79.3	1,100	10.8	9.6 - 12.1	1,000	9.8	8.6 - 11.0	200	1.8	1.3 - 2.5	10,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	12,600	70.1	68.8 - 71.4	2,600	14.4	13.4 - 15.4	2,100	11.5	10.7 - 12.5	700	3.9	3.4 - 4.6	17,900

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,500	67.0	63.0 - 70.7	400	18.3	15.3 - 21.7	200	10.8	8.5 - 13.6	100	4.0	2.7 - 6.0	2,300
Honolulu	9,100	71.7	70.2 - 73.2	1,700	13.1	12.0 - 14.3	1,400	11.4	10.3 - 12.5	500	3.8	3.2 - 4.5	12,700
Kauai	600	63.8	57.4 - 69.8	100	16.6	12.4 - 21.9	100	11.9	8.3 - 16.9	100	7.6	4.7 - 12.2	900
Maui	1,400	66.5	62.3 - 70.5	300	17.0	13.9 - 20.5	300	13.4	10.8 - 16.7	100	3.1	1.9 - 5.0	2,000

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,200	76.9	74.4 - 79.3	500	11.0	9.3 - 12.9	400	9.9	8.3 - 11.7	100	2.2	1.5 - 3.2	4,200
Native Hawaiian	3,100	61.4	58.5 - 64.2	800	16.8	14.7 - 19.1	800	16.5	14.5 - 18.8	300	5.3	4.1 - 6.8	5,000
Chinese	500	79.5	76.0 - 82.5	100	10.1	7.9 - 12.9	100	8.6	6.6 - 11.1	<50	1.8	1.0 - 3.3	600
Filipino	2,300	71.4	68.6 - 74.1	600	17.5	15.3 - 20.0	200	7.5	6.0 - 9.3	100	3.5	2.5 - 4.9	3,300
Japanese	1,700	81.6	78.2 - 84.6	100	7.3	5.4 - 9.9	200	9.6	7.4 - 12.4	<50	1.4	0.7 - 2.9	2,000
Other	1,800	63.5	59.9 - 67.0	500	17.5	14.9 - 20.6	300	12.0	9.8 - 14.6	200	7.0	5.3 - 9.2	2,800

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	54.3	48.9 - 59.5	300	21.8	17.8 - 26.5	300	17.5	13.8 - 22.0	100	6.4	4.2 - 9.6	1,500
20-24	2,700	64.0	61.0 - 66.8	700	16.4	14.3 - 18.8	600	14.1	12.1 - 16.3	200	5.5	4.3 - 7.1	4,300
25-34	6,500	73.2	71.4 - 74.9	1,200	13.2	11.9 - 14.6	900	10.2	9.1 - 11.5	300	3.3	2.7 - 4.1	8,900
35+	2,500	77.2	74.4 - 79.7	400	11.5	9.6 - 13.6	300	9.0	7.3 - 11.0	100	2.4	1.5 - 3.6	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,400	68.9	67.0 - 70.7	1,400	15.0	13.7 - 16.5	1,100	12.3	11.1 - 13.7	300	3.8	3.1 - 4.6	9,300
Female	6,200	71.5	69.6 - 73.3	1,200	13.7	12.3 - 15.2	900	10.7	9.5 - 12.1	400	4.1	3.4 - 5.1	8,700

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,300	73.9	72.4 - 75.4	1,400	12.7	11.6 - 13.8	1,200	10.6	9.5 - 11.7	300	2.8	2.3 - 3.5	11,300
No	4,200	63.7	61.3 - 66.0	1,200	17.3	15.5 - 19.2	900	13.2	11.6 - 14.9	400	5.8	4.8 - 7.1	6,700

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,300	64.8	62.4 - 67.1	1,100	17.5	15.7 - 19.4	800	12.7	11.1 - 14.4	300	5.0	4.0 - 6.2	6,600
131-185%	1,700	67.4	63.8 - 70.8	400	14.7	12.3 - 17.5	400	14.5	12.0 - 17.3	100	3.4	2.3 - 5.1	2,600
186+%	6,000	77.1	75.3 - 78.8	800	10.6	9.4 - 11.9	800	10.1	8.9 - 11.4	200	2.2	1.7 - 3.0	7,800

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,800	61.6	58.0 - 65.1	600	18.6	16.0 - 21.6	400	14.5	12.1 - 17.3	200	5.2	3.8 - 7.1	3,000
\$10,000-\$24,999	2,500	66.2	63.1 - 69.1	600	15.8	13.6 - 18.2	500	12.6	10.6 - 14.8	200	5.5	4.2 - 7.1	3,800
\$25,000-\$49,999	3,000	70.9	68.2 - 73.5	600	14.6	12.7 - 16.8	500	11.9	10.1 - 13.9	100	2.6	1.8 - 3.7	4,200
>=\$50,000	4,700	78.3	76.3 - 80.1	600	9.9	8.6 - 11.4	600	9.9	8.5 - 11.3	100	2.0	1.4 - 2.8	6,000

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	71.4	66.5 - 75.9	200	14.1	10.8 - 18.2	100	10.1	7.4 - 13.7	100	4.3	2.6 - 7.2	1,400
Normal birth weight	11,600	70.0	68.7 - 71.3	2,400	14.4	13.4 - 15.5	1,900	11.7	10.7 - 12.7	600	3.9	3.4 - 4.5	16,600

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	60.1	55.0 - 65.0	300	18.2	14.6 - 22.5	300	16.8	13.3 - 20.9	100	4.9	3.1 - 7.8	1,500
No	11,500	71.0	69.7 - 72.4	2,300	14.0	13.0 - 15.0	1,800	11.1	10.2 - 12.1	600	3.9	3.3 - 4.5	16,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	63.9	60.7 - 67.0	600	16.2	13.9 - 18.9	500	14.5	12.3 - 17.0	200	5.3	4.0 - 7.1	3,500
No	10,200	71.7	70.2 - 73.1	2,000	13.8	12.8 - 15.0	1,500	10.9	9.9 - 11.9	500	3.6	3.1 - 4.3	14,200

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,900	64.4	62.2 - 66.5	1,300	17.0	15.3 - 18.7	1,000	12.7	11.3 - 14.3	400	5.9	4.9 - 7.1	7,600
No	7,500	74.3	72.7 - 75.9	1,300	12.6	11.4 - 13.8	1,100	10.5	9.4 - 11.7	300	2.6	2.0 - 3.3	10,100

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	12,000	68.7	67.4 - 70.0	2,500	14.1	13.1 - 15.1	2,000	11.6	10.8 - 12.6	1,000	5.6	5.0 - 6.3	17,500

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,400	64.3	60.3 - 68.1	400	17.2	14.3 - 20.5	200	10.9	8.5 - 13.7	200	7.7	5.8 - 10.1	2,200
Honolulu	8,800	70.3	68.7 - 71.8	1,700	13.2	12.1 - 14.4	1,400	11.3	10.3 - 12.4	700	5.2	4.5 - 6.0	12,500
Kauai	500	59.6	52.9 - 66.0	100	16.6	12.2 - 22.0	100	13.6	9.6 - 19.0	100	10.2	6.7 - 15.2	800
Maui	1,300	67.4	63.1 - 71.4	300	15.2	12.3 - 18.7	300	13.9	11.1 - 17.3	100	3.5	2.2 - 5.5	1,900

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,000	76.1	73.5 - 78.6	400	10.1	8.4 - 12.0	400	10.4	8.7 - 12.3	100	3.4	2.5 - 4.7	4,000
Native Hawaiian	3,000	61.0	58.1 - 63.8	800	15.8	13.8 - 18.1	800	16.2	14.2 - 18.5	300	6.9	5.6 - 8.6	4,900
Chinese	500	79.6	76.3 - 82.6	100	9.7	7.6 - 12.3	<50	7.7	5.9 - 10.0	<50	2.9	1.9 - 4.6	600
Filipino	2,200	69.2	66.3 - 72.0	600	18.5	16.3 - 21.0	200	7.0	5.6 - 8.7	200	5.3	4.1 - 6.8	3,200
Japanese	1,700	81.6	78.3 - 84.5	100	6.4	4.7 - 8.7	200	9.6	7.4 - 12.2	100	2.4	1.4 - 3.9	2,100
Other	1,600	58.3	54.7 - 61.9	500	18.7	15.9 - 21.7	400	13.2	10.9 - 16.0	300	9.8	7.8 - 12.2	2,700

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	55.4	50.2 - 60.5	300	20.6	16.7 - 25.1	300	16.6	13.0 - 21.0	100	7.4	5.1 - 10.6	1,500
20-24	2,700	62.9	60.0 - 65.7	700	17.0	14.9 - 19.3	600	13.5	11.6 - 15.6	300	6.7	5.3 - 8.3	4,300
25-34	6,100	71.7	69.8 - 73.4	1,100	12.5	11.3 - 13.9	900	10.7	9.5 - 12.0	400	5.1	4.3 - 6.1	8,600
35+	2,300	75.2	72.3 - 77.8	300	11.1	9.3 - 13.3	300	9.3	7.6 - 11.4	100	4.4	3.2 - 5.9	3,100

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,100	68.0	66.1 - 69.8	1,300	14.1	12.8 - 15.6	1,100	12.3	11.1 - 13.7	500	5.5	4.7 - 6.5	9,000
Female	5,900	69.4	67.5 - 71.3	1,200	14.0	12.7 - 15.5	900	10.9	9.7 - 12.3	500	5.6	4.8 - 6.7	8,500

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,000	72.6	71.0 - 74.1	1,300	12.0	10.9 - 13.2	1,200	10.7	9.6 - 11.8	500	4.7	4.0 - 5.5	11,100
No	4,000	61.9	59.5 - 64.3	1,100	17.6	15.8 - 19.6	900	13.3	11.7 - 15.1	500	7.1	5.9 - 8.5	6,400

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,100	63.8	61.4 - 66.0	1,100	16.6	14.9 - 18.5	800	12.5	11.0 - 14.2	500	7.1	6.0 - 8.5	6,400
131-185%	1,500	64.8	60.9 - 68.4	400	16.2	13.5 - 19.4	300	13.6	11.1 - 16.5	100	5.4	3.9 - 7.5	2,300
186+%	5,900	75.7	73.9 - 77.5	800	10.5	9.3 - 11.8	800	10.6	9.3 - 11.9	200	3.2	2.5 - 4.0	7,700

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,900	62.7	59.2 - 66.1	500	17.6	15.1 - 20.6	400	13.1	10.8 - 15.8	200	6.6	5.0 - 8.5	3,000
\$10,000-\$24,999	2,400	65.0	62.0 - 67.9	600	15.1	13.0 - 17.5	400	12.2	10.2 - 14.4	300	7.7	6.2 - 9.6	3,700
\$25,000-\$49,999	2,800	68.8	66.0 - 71.4	600	15.4	13.4 - 17.6	500	11.7	9.9 - 13.7	200	4.2	3.2 - 5.5	4,100
>=\$50,000	4,400	76.8	74.7 - 78.8	500	9.4	8.2 - 10.9	600	10.7	9.3 - 12.3	200	3.0	2.3 - 4.0	5,700

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	68.2	63.3 - 72.8	200	13.3	10.1 - 17.2	200	11.3	8.5 - 14.9	100	7.2	4.9 - 10.5	1,400
Normal birth weight	11,100	68.7	67.4 - 70.1	2,300	14.2	13.2 - 15.2	1,900	11.7	10.7 - 12.7	900	5.4	4.8 - 6.1	16,100

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	63.0	58.0 - 67.8	200	15.1	11.8 - 19.2	200	14.3	11.1 - 18.3	100	7.5	5.2 - 10.7	1,500
No	10,900	69.3	68.0 - 70.7	2,200	13.9	12.9 - 14.9	1,800	11.4	10.4 - 12.4	900	5.4	4.8 - 6.1	15,700

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	64.1	60.9 - 67.2	500	15.2	13.0 - 17.7	500	14.1	12.0 - 16.6	200	6.6	5.1 - 8.4	3,500
No	9,600	70.1	68.6 - 71.5	1,900	13.6	12.6 - 14.8	1,500	11.0	10.0 - 12.0	700	5.3	4.7 - 6.1	13,700

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,700	63.3	61.1 - 65.4	1,200	16.7	15.0 - 18.4	900	12.6	11.1 - 14.2	600	7.5	6.4 - 8.8	7,400
No	7,100	72.7	71.1 - 74.4	1,200	12.2	11.1 - 13.5	1,100	10.8	9.7 - 12.0	400	4.2	3.5 - 5.1	9,800

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,600	67.3	66.0 - 68.6	2,700	15.6	14.6 - 16.6	2,100	12.0	11.2 - 13.0	900	5.1	4.5 - 5.7	17,300

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,400	63.8	59.8 - 67.5	400	16.7	13.9 - 19.9	300	12.5	10.1 - 15.4	200	7.1	5.3 - 9.5	2,200
Honolulu	8,700	69.0	67.5 - 70.5	1,800	14.6	13.5 - 15.8	1,500	11.6	10.6 - 12.7	600	4.8	4.1 - 5.6	12,600
Kauai	400	54.4	47.8 - 60.9	200	22.0	17.1 - 27.9	100	16.4	11.9 - 22.1	100	7.2	4.3 - 11.8	800
Maui	1,200	64.9	60.7 - 68.9	300	18.4	15.3 - 21.9	200	12.9	10.2 - 16.1	100	3.8	2.5 - 5.8	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	73.1	70.5 - 75.6	500	11.8	10.1 - 13.7	500	12.0	10.2 - 14.0	100	3.1	2.2 - 4.3	3,900
Native Hawaiian	3,000	61.1	58.3 - 63.8	800	16.4	14.4 - 18.6	800	16.2	14.3 - 18.4	300	6.3	5.1 - 7.8	4,800
Chinese	500	78.2	75.2 - 81.0	100	10.4	8.5 - 12.6	100	9.1	7.3 - 11.4	<50	2.3	1.4 - 3.7	600
Filipino	2,200	68.3	65.4 - 70.9	700	20.6	18.3 - 23.0	200	6.7	5.3 - 8.3	100	4.5	3.4 - 5.9	3,200
Japanese	1,700	79.2	75.8 - 82.2	200	7.1	5.4 - 9.4	200	11.0	8.8 - 13.7	100	2.7	1.7 - 4.2	2,200
Other	1,400	56.2	52.6 - 59.8	600	22.0	19.1 - 25.2	300	12.5	10.2 - 15.1	200	9.3	7.3 - 11.6	2,600

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	54.7	49.6 - 59.7	300	21.6	17.7 - 26.2	300	17.7	14.1 - 22.1	100	5.9	3.9 - 8.9	1,400
20-24	2,700	62.0	59.2 - 64.7	800	18.4	16.3 - 20.7	600	13.8	11.9 - 15.9	200	5.8	4.6 - 7.3	4,300
25-34	5,900	70.0	68.2 - 71.7	1,200	14.2	12.9 - 15.6	900	10.8	9.6 - 12.1	400	5.0	4.2 - 6.0	8,500
35+	2,300	72.9	70.1 - 75.5	400	12.8	10.9 - 15.0	300	10.4	8.6 - 12.4	100	3.9	2.8 - 5.3	3,100

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Infant Gender													
Male	5,900	65.0	63.1 - 66.8	1,500	16.3	15.0 - 17.8	1,200	13.6	12.3 - 14.9	500	5.1	4.3 - 6.1	9,000
Female	5,800	69.8	68.0 - 71.6	1,200	14.8	13.4 - 16.2	900	10.4	9.2 - 11.7	400	5.0	4.2 - 6.0	8,300

	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Married													
Yes	8,000	70.5	69.0 - 72.0	1,600	13.7	12.7 - 14.9	1,300	11.2	10.2 - 12.3	500	4.6	3.9 - 5.3	11,400
No	3,600	61.1	58.7 - 63.5	1,100	19.1	17.2 - 21.1	800	13.7	12.0 - 15.5	400	6.1	5.0 - 7.4	5,900

	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Poverty Level													
0-130%	3,900	61.5	59.2 - 63.8	1,200	18.8	17.1 - 20.7	800	13.1	11.5 - 14.8	400	6.6	5.5 - 7.8	6,300
131-185%	1,300	62.1	58.2 - 65.9	400	19.2	16.3 - 22.5	300	13.1	10.6 - 16.0	100	5.6	4.0 - 7.8	2,100
186+%	5,800	74.5	72.7 - 76.2	900	11.6	10.4 - 12.9	900	11.0	9.8 - 12.3	200	3.0	2.3 - 3.8	7,800

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,700	60.2	56.7 - 63.6	600	20.3	17.6 - 23.3	400	13.5	11.2 - 16.1	200	6.1	4.6 - 8.0	2,900
\$10,000-\$24,999	2,400	62.7	59.9 - 65.5	700	18.3	16.1 - 20.6	500	12.4	10.6 - 14.5	300	6.6	5.3 - 8.2	3,900
\$25,000-\$49,999	2,800	66.6	63.9 - 69.2	700	16.2	14.3 - 18.3	500	12.6	10.9 - 14.6	200	4.6	3.5 - 5.9	4,200
>=\$50,000	4,100	76.8	74.7 - 78.7	500	10.2	8.9 - 11.7	600	10.4	9.0 - 12.0	100	2.6	1.9 - 3.6	5,300

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	68.1	63.0 - 72.8	200	14.8	11.4 - 18.9	100	10.7	7.8 - 14.4	100	6.5	4.3 - 9.7	1,200
Normal birth weight	10,800	67.2	65.9 - 68.6	2,500	15.7	14.7 - 16.7	2,000	12.1	11.2 - 13.1	800	5.0	4.4 - 5.6	16,100

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	61.7	56.7 - 66.5	200	16.2	12.9 - 20.3	200	15.3	12.0 - 19.3	100	6.7	4.6 - 9.9	1,400
No	10,600	67.8	66.5 - 69.2	2,400	15.5	14.5 - 16.5	1,800	11.7	10.8 - 12.7	800	4.9	4.3 - 5.6	15,600

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	62.7	59.6 - 65.7	600	16.6	14.4 - 19.1	500	15.1	12.9 - 17.5	200	5.6	4.3 - 7.3	3,600
No	9,200	68.6	67.1 - 70.0	2,100	15.2	14.2 - 16.4	1,500	11.3	10.3 - 12.3	700	4.9	4.3 - 5.7	13,500

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,600	61.9	59.8 - 64.0	1,400	18.5	16.9 - 20.3	1,000	13.4	12.0 - 14.9	500	6.2	5.2 - 7.3	7,500
No	6,800	71.4	69.8 - 73.1	1,300	13.3	12.1 - 14.5	1,000	10.9	9.9 - 12.1	400	4.3	3.6 - 5.2	9,500

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,500	67.8	66.5 - 69.1	2,800	16.6	15.6 - 17.7	2,000	11.9	11.1 - 12.9	600	3.6	3.1 - 4.2	17,000

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,300	63.0	59.5 - 66.4	400	16.9	14.4 - 19.7	300	13.8	11.5 - 16.5	100	6.3	4.6 - 8.5	2,100
Honolulu	8,600	69.5	67.9 - 71.0	2,000	15.9	14.6 - 17.2	1,400	11.5	10.4 - 12.6	400	3.2	2.6 - 3.8	12,400
Kauai	400	57.1	51.4 - 62.6	200	24.8	20.3 - 29.9	100	13.7	10.3 - 18.1	<50	4.4	2.4 - 7.9	800
Maui	1,200	66.8	63.2 - 70.3	300	17.9	15.2 - 21.0	200	12.1	9.9 - 14.8	100	3.2	2.0 - 5.0	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	72.2	69.3 - 74.9	500	13.3	11.3 - 15.6	500	12.4	10.5 - 14.6	100	2.1	1.4 - 3.2	3,700
Native Hawaiian	2,900	61.0	58.5 - 63.4	800	17.7	15.9 - 19.6	800	16.9	15.2 - 18.9	200	4.4	3.4 - 5.6	4,700
Chinese	500	78.4	72.9 - 83.0	100	12.0	8.6 - 16.6	100	8.1	5.2 - 12.4	<50	1.6	0.9 - 2.7	700
Filipino	2,300	69.7	66.7 - 72.5	700	20.5	18.0 - 23.2	200	6.6	5.2 - 8.3	100	3.2	2.4 - 4.4	3,300
Japanese	1,800	82.8	79.6 - 85.6	100	6.6	4.8 - 8.9	200	8.6	6.6 - 11.0	<50	2.1	1.2 - 3.6	2,100
Other	1,400	55.9	51.8 - 59.8	600	24.4	21.0 - 28.1	300	12.9	10.4 - 16.0	200	6.8	5.1 - 9.0	2,400

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	60.5	55.7 - 65.2	300	18.3	14.9 - 22.4	200	17.2	13.8 - 21.1	100	4.0	2.5 - 6.2	1,400
20-24	2,600	62.5	59.7 - 65.2	800	20.1	17.9 - 22.5	600	13.5	11.7 - 15.6	200	3.9	2.9 - 5.1	4,100
25-34	5,900	70.0	68.1 - 71.8	1,300	14.9	13.5 - 16.4	1,000	11.5	10.2 - 12.8	300	3.6	2.9 - 4.5	8,400
35+	2,200	72.4	69.4 - 75.3	500	15.9	13.5 - 18.6	300	8.6	7.0 - 10.5	100	3.1	2.1 - 4.4	3,000

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,900	66.1	64.2 - 67.9	1,500	17.0	15.6 - 18.5	1,200	12.9	11.7 - 14.2	400	4.0	3.3 - 4.9	9,000
Female	5,600	69.8	67.9 - 71.6	1,300	16.2	14.7 - 17.8	900	10.8	9.6 - 12.2	300	3.1	2.5 - 3.9	8,000

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,900	70.3	68.7 - 71.8	1,800	15.5	14.3 - 16.9	1,200	11.0	10.0 - 12.2	400	3.2	2.6 - 3.8	11,300
No	3,600	63.0	60.6 - 65.3	1,100	18.8	17.0 - 20.7	800	13.7	12.2 - 15.5	300	4.5	3.6 - 5.7	5,700

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,200	63.3	61.1 - 65.4	1,400	20.2	18.4 - 22.0	800	12.0	10.7 - 13.6	300	4.5	3.7 - 5.5	6,700
131-185%	1,300	63.5	59.5 - 67.4	500	21.6	18.4 - 25.3	300	12.0	9.7 - 14.8	100	2.8	1.8 - 4.4	2,100
186+%	5,100	74.9	72.9 - 76.7	800	11.3	10.0 - 12.7	800	11.5	10.2 - 13.0	200	2.3	1.8 - 3.1	6,900

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	65.7	60.3 - 70.7	200	17.9	14.1 - 22.6	100	11.2	8.2 - 15.1	100	5.2	3.2 - 8.3	1,100
Normal birth weight	10,800	68.0	66.6 - 69.3	2,600	16.5	15.5 - 17.7	1,900	12.0	11.1 - 12.9	600	3.5	3.0 - 4.1	15,900

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	61.6	56.6 - 66.3	300	19.7	16.0 - 23.9	200	14.1	11.0 - 18.0	100	4.7	3.0 - 7.2	1,400
No	10,500	68.5	67.2 - 69.9	2,500	16.3	15.2 - 17.4	1,800	11.7	10.8 - 12.6	500	3.5	3.0 - 4.1	15,300

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	63.7	60.6 - 66.6	600	18.3	16.0 - 20.8	500	14.6	12.5 - 16.9	100	3.5	2.5 - 4.8	3,500
No	9,100	69.1	67.6 - 70.5	2,100	16.1	15.0 - 17.4	1,500	11.1	10.2 - 12.2	500	3.6	3.1 - 4.3	13,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,600	62.5	60.4 - 64.5	1,400	19.6	18.0 - 21.4	1,000	13.6	12.2 - 15.1	300	4.3	3.5 - 5.2	7,400
No	6,700	71.9	70.2 - 73.6	1,300	14.3	13.0 - 15.7	1,000	10.7	9.6 - 11.9	300	3.1	2.5 - 3.8	9,400

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,300	67.7	66.3 - 69.0	2,900	17.5	16.4 - 18.6	2,000	12.2	11.3 - 13.2	400	2.7	2.3 - 3.2	16,800

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,300	63.1	60.1 - 66.1	400	18.6	16.3 - 21.2	300	14.2	12.1 - 16.5	100	4.1	3.0 - 5.4	2,000
Honolulu	8,500	69.6	67.8 - 71.3	2,000	16.4	15.1 - 17.9	1,400	11.7	10.5 - 12.9	300	2.3	1.8 - 3.0	12,200
Kauai	400	56.7	51.7 - 61.6	200	26.1	21.9 - 30.7	100	14.2	11.1 - 18.0	<50	3.0	1.9 - 4.9	800
Maui	1,100	64.2	60.9 - 67.3	300	19.8	17.2 - 22.6	200	12.7	10.7 - 15.0	100	3.3	2.3 - 4.7	1,800

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,600	70.5	67.4 - 73.5	600	15.6	13.3 - 18.1	500	12.6	10.5 - 15.0	<50	1.3	0.8 - 2.2	3,700
Native Hawaiian	2,700	59.9	57.8 - 61.9	800	18.4	16.8 - 20.1	800	17.8	16.3 - 19.5	200	3.9	3.1 - 4.7	4,600
Chinese	500	80.5	73.9 - 85.8	100	11.3	7.6 - 16.6	<50	6.9	3.8 - 12.1	<50	1.3	0.3 - 5.7	600
Filipino	2,300	69.6	66.2 - 72.7	700	21.4	18.6 - 24.4	200	6.3	4.9 - 8.2	100	2.7	1.8 - 4.1	3,400
Japanese	1,800	81.5	77.7 - 84.7	200	7.3	5.3 - 9.9	200	9.7	7.3 - 12.7	<50	1.6	0.7 - 3.6	2,200
Other	1,400	59.4	54.7 - 63.9	600	24.1	20.4 - 28.4	300	12.7	9.9 - 16.2	100	3.8	2.3 - 6.0	2,300

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	56.8	51.9 - 61.7	300	19.8	16.1 - 24.0	300	19.8	16.2 - 24.0	100	3.6	2.1 - 5.9	1,400
20-24	2,600	63.3	60.4 - 66.1	800	20.2	18.0 - 22.7	600	14.0	12.1 - 16.1	100	2.5	1.8 - 3.5	4,100
25-34	5,900	70.2	68.2 - 72.1	1,400	16.2	14.7 - 17.8	900	11.1	9.9 - 12.5	200	2.5	1.9 - 3.2	8,300
35+	2,100	71.8	68.4 - 74.9	500	16.2	13.6 - 19.1	300	9.0	7.3 - 11.1	100	3.0	2.0 - 4.5	2,900

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,800	66.0	64.1 - 67.9	1,600	17.7	16.2 - 19.3	1,200	13.5	12.2 - 14.9	200	2.7	2.2 - 3.5	8,800
Female	5,500	69.5	67.5 - 71.5	1,400	17.2	15.6 - 18.9	800	10.7	9.4 - 12.1	200	2.6	2.0 - 3.4	7,900

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,000	70.6	68.9 - 72.3	1,800	16.1	14.7 - 17.5	1,200	10.9	9.8 - 12.1	300	2.4	1.9 - 3.1	11,400
No	3,300	61.4	59.0 - 63.8	1,100	20.5	18.5 - 22.5	800	14.9	13.3 - 16.7	200	3.2	2.4 - 4.1	5,400

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,200	62.6	60.3 - 64.8	1,400	21.0	19.2 - 22.9	900	13.1	11.6 - 14.6	200	3.4	2.6 - 4.3	6,700
131-185%	1,700	65.9	62.0 - 69.5	500	19.7	16.7 - 23.2	300	12.8	10.4 - 15.6	<50	1.6	1.0 - 2.6	2,600
186+%	4,500	74.8	72.6 - 76.9	700	12.2	10.7 - 13.9	700	10.8	9.3 - 12.4	100	2.2	1.6 - 3.2	6,000

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	68.1	62.3 - 73.4	200	20.7	16.2 - 26.2	100	8.9	6.2 - 12.7	<50	2.2	1.3 - 3.8	1,100
Normal birth weight	10,600	67.6	66.2 - 69.0	2,700	17.3	16.1 - 18.4	1,900	12.4	11.4 - 13.4	400	2.7	2.3 - 3.2	15,700

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	57.1	52.1 - 62.0	300	21.8	18.0 - 26.2	200	16.5	13.0 - 20.8	100	4.6	3.2 - 6.5	1,400
No	10,400	68.7	67.3 - 70.2	2,600	17.1	15.9 - 18.3	1,800	11.7	10.8 - 12.7	400	2.5	2.1 - 3.1	15,100

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	61.5	58.4 - 64.6	600	19.1	16.7 - 21.7	600	16.4	14.1 - 19.0	100	3.0	2.2 - 4.1	3,400
No	9,000	69.4	67.8 - 70.9	2,200	17.0	15.7 - 18.3	1,400	11.0	10.0 - 12.1	300	2.6	2.1 - 3.2	13,000

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,500	62.1	59.9 - 64.2	1,500	20.7	19.0 - 22.6	1,000	14.1	12.7 - 15.7	200	3.1	2.5 - 3.9	7,200
No	6,700	71.9	70.0 - 73.7	1,400	15.1	13.7 - 16.6	1,000	10.7	9.5 - 11.9	200	2.4	1.8 - 3.1	9,400

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,700	65.9	64.5 - 67.3	2,800	17.2	16.1 - 18.4	1,900	11.9	11.0 - 12.9	800	4.9	4.3 - 5.5	16,300

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	59.7	57.0 - 62.3	400	20.0	17.9 - 22.2	300	14.1	12.3 - 16.0	100	6.3	5.1 - 7.8	1,800
Honolulu	8,100	67.7	65.8 - 69.4	2,000	16.5	15.1 - 18.0	1,400	11.5	10.3 - 12.7	500	4.4	3.7 - 5.2	12,000
Kauai	400	55.2	50.8 - 59.6	200	24.7	21.1 - 28.9	100	13.8	11.1 - 17.0	<50	6.3	4.5 - 8.7	700
Maui	1,100	65.2	62.4 - 67.9	300	16.5	14.4 - 18.8	200	12.1	10.4 - 14.1	100	6.2	4.9 - 7.8	1,700

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,500	70.4	67.2 - 73.5	500	14.4	12.2 - 17.0	400	12.2	10.1 - 14.7	100	2.9	2.0 - 4.1	3,500
Native Hawaiian	2,500	56.9	55.3 - 58.6	800	18.3	17.0 - 19.6	800	17.9	16.7 - 19.2	300	6.9	6.1 - 7.7	4,400
Chinese	500	79.4	71.4 - 85.6	100	10.2	6.1 - 16.4	<50	7.1	3.6 - 13.6	<50	3.3	1.3 - 8.4	600
Filipino	2,300	67.7	64.2 - 71.0	700	21.2	18.3 - 24.4	200	5.7	4.3 - 7.6	200	5.4	4.1 - 7.2	3,400
Japanese	1,700	81.5	77.6 - 85.0	100	7.1	5.1 - 9.9	200	8.7	6.3 - 11.8	100	2.7	1.5 - 4.7	2,100
Other	1,200	55.3	50.4 - 60.2	600	25.2	21.1 - 29.8	300	13.6	10.5 - 17.3	100	5.9	4.0 - 8.4	2,200

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	54.9	50.4 - 59.3	300	20.6	17.2 - 24.5	300	17.1	14.0 - 20.8	100	7.4	5.6 - 9.8	1,500
20-24	2,400	59.7	56.9 - 62.5	800	20.3	18.0 - 22.8	600	15.2	13.3 - 17.3	200	4.8	3.9 - 5.8	4,000
25-34	5,600	69.2	67.1 - 71.2	1,300	15.9	14.3 - 17.5	800	10.5	9.3 - 11.9	400	4.4	3.6 - 5.4	8,100
35+	1,900	71.6	68.0 - 75.0	400	15.0	12.3 - 18.1	200	8.4	6.5 - 10.6	100	5.1	3.7 - 6.9	2,700

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,400	65.1	63.1 - 67.0	1,400	16.9	15.4 - 18.5	1,100	13.0	11.7 - 14.4	400	5.0	4.2 - 5.9	8,400
Female	5,300	66.8	64.8 - 68.8	1,400	17.6	16.0 - 19.4	900	10.8	9.6 - 12.2	400	4.8	4.0 - 5.6	7,900

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,500	69.4	67.6 - 71.2	1,700	15.6	14.3 - 17.1	1,200	10.7	9.6 - 11.9	500	4.3	3.6 - 5.0	10,900
No	3,200	59.0	56.6 - 61.3	1,100	20.5	18.6 - 22.5	800	14.4	12.9 - 16.1	300	6.1	5.2 - 7.2	5,400

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,000	60.0	57.8 - 62.2	1,400	20.6	18.8 - 22.6	900	13.4	12.0 - 14.8	400	6.0	5.1 - 7.1	6,600
131-185%	2,000	65.7	62.3 - 69.0	600	17.9	15.3 - 20.9	400	12.5	10.4 - 15.0	100	3.8	2.9 - 5.0	3,100
186+%	3,900	74.5	72.0 - 76.8	600	12.2	10.6 - 14.1	500	9.6	8.1 - 11.3	200	3.7	2.8 - 4.9	5,200

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Income (2000-2003)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,000	59.1	56.0 - 62.1	700	21.4	18.9 - 24.2	500	14.3	12.4 - 16.4	200	5.2	4.0 - 6.7	3,400
\$15,001-\$25,000	1,400	61.9	58.0 - 65.6	400	18.5	15.7 - 21.7	300	13.5	11.1 - 16.3	100	6.1	4.6 - 8.1	2,200
\$25,001-\$40,000	2,200	64.0	60.7 - 67.2	600	18.3	15.8 - 21.1	400	12.8	10.7 - 15.1	200	5.0	3.8 - 6.4	3,400
>=\$40,001	4,600	73.4	71.2 - 75.6	800	13.5	11.8 - 15.3	600	9.4	8.1 - 11.0	200	3.7	2.9 - 4.6	6,300

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	64.1	58.2 - 69.6	200	20.9	16.3 - 26.3	100	10.3	7.5 - 14.1	<50	4.6	3.0 - 7.1	1,100
Normal birth weight	10,000	66.0	64.6 - 67.5	2,600	17.0	15.9 - 18.2	1,800	12.0	11.1 - 13.0	700	4.9	4.3 - 5.5	15,200

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	53.7	48.9 - 58.4	300	19.0	15.5 - 23.1	200	17.5	14.1 - 21.5	100	9.8	7.7 - 12.5	1,400
No	9,800	67.2	65.7 - 68.7	2,500	17.0	15.8 - 18.2	1,600	11.3	10.4 - 12.3	700	4.5	3.9 - 5.1	14,600

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,900	59.6	56.4 - 62.7	600	17.6	15.3 - 20.2	500	16.1	13.9 - 18.6	200	6.6	5.4 - 8.1	3,200
No	8,500	67.7	66.1 - 69.3	2,100	17.0	15.7 - 18.4	1,400	10.8	9.8 - 11.8	600	4.5	3.9 - 5.3	12,600

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,100	60.1	57.9 - 62.2	1,300	19.4	17.7 - 21.3	1,000	14.4	13.0 - 15.9	400	6.1	5.2 - 7.1	6,900
No	6,500	70.2	68.3 - 72.0	1,500	15.7	14.2 - 17.3	900	10.1	8.9 - 11.3	400	4.1	3.4 - 4.9	9,300

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,300	64.1	62.8 - 65.4	2,700	17.1	16.0 - 18.2	2,000	12.7	11.8 - 13.6	1,000	6.2	5.6 - 6.8	16,100

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	57.2	54.5 - 59.8	400	19.1	17.1 - 21.3	300	15.8	13.9 - 17.8	100	7.9	6.6 - 9.5	1,800
Honolulu	7,900	66.2	64.5 - 67.9	1,900	16.3	15.0 - 17.7	1,400	11.9	10.8 - 13.1	700	5.6	4.8 - 6.4	11,900
Kauai	300	51.0	46.4 - 55.6	200	23.6	19.9 - 27.8	100	15.9	12.9 - 19.5	100	9.4	7.0 - 12.5	700
Maui	1,000	61.8	58.9 - 64.7	300	17.6	15.4 - 20.1	200	13.3	11.4 - 15.3	100	7.3	5.9 - 9.0	1,700

PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,500	69.2	66.3 - 72.0	500	14.5	12.4 - 16.8	500	12.9	11.0 - 15.2	100	3.4	2.5 - 4.6	3,600
Native Hawaiian	2,400	55.2	53.6 - 56.7	800	18.0	16.8 - 19.3	800	18.3	17.1 - 19.6	400	8.5	7.7 - 9.4	4,300
Chinese	500	79.9	72.5 - 85.7	100	9.1	5.4 - 14.7	<50	6.3	3.3 - 11.8	<50	4.8	2.3 - 10.0	600
Filipino	2,100	64.6	61.1 - 67.8	700	21.7	19.0 - 24.8	200	6.4	5.0 - 8.2	200	7.3	5.8 - 9.3	3,200
Japanese	1,600	77.1	73.1 - 80.7	200	8.9	6.7 - 11.8	200	10.9	8.3 - 14.2	100	3.0	1.8 - 5.1	2,100
Other	1,200	55.7	51.0 - 60.2	500	22.6	18.9 - 26.8	300	13.9	11.1 - 17.3	200	7.9	5.8 - 10.6	2,200

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	51.6	47.5 - 55.7	300	21.4	18.3 - 25.0	300	17.6	14.8 - 20.8	200	9.4	7.4 - 11.9	1,600
20-24	2,400	58.9	56.2 - 61.5	800	19.4	17.3 - 21.7	600	15.8	14.0 - 17.8	200	5.9	4.9 - 7.1	4,000
25-34	5,200	66.8	64.8 - 68.7	1,300	16.3	14.8 - 17.9	900	11.1	9.9 - 12.4	500	5.8	4.9 - 6.9	7,800
35+	1,900	71.7	68.4 - 74.9	300	13.0	10.8 - 15.7	300	9.6	7.7 - 11.9	100	5.6	4.2 - 7.5	2,600

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Infant Gender													
Male	5,100	62.9	61.0 - 64.8	1,300	16.0	14.7 - 17.5	1,200	14.7	13.4 - 16.0	500	6.4	5.5 - 7.4	8,200
Female	5,200	65.3	63.4 - 67.2	1,400	18.1	16.6 - 19.8	800	10.6	9.5 - 11.8	500	5.9	5.1 - 6.9	7,900

	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Married													
Yes	7,400	68.3	66.7 - 70.0	1,700	15.3	14.0 - 16.6	1,200	11.2	10.1 - 12.3	600	5.2	4.5 - 6.1	10,900
No	2,900	55.2	53.0 - 57.5	1,100	20.8	19.0 - 22.8	800	15.8	14.3 - 17.4	400	8.1	7.1 - 9.3	5,200

	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Poverty Level													
0-130%	3,700	57.0	54.9 - 59.1	1,300	20.7	19.0 - 22.5	900	14.5	13.2 - 15.9	500	7.8	6.8 - 9.0	6,400
131-185%	2,000	64.7	61.4 - 67.8	500	16.7	14.3 - 19.4	400	14.0	11.8 - 16.4	100	4.7	3.6 - 6.0	3,000
186+%	4,000	73.4	71.2 - 75.6	700	12.4	10.8 - 14.1	500	9.6	8.2 - 11.1	300	4.6	3.7 - 5.8	5,400

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Income (2000-2003)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,800	54.8	52.0 - 57.7	700	22.3	19.9 - 24.9	500	15.1	13.3 - 17.1	300	7.8	6.3 - 9.5	3,400
\$15,001-\$25,000	1,400	59.2	55.6 - 62.7	400	18.4	15.7 - 21.3	400	15.9	13.5 - 18.6	200	6.5	5.1 - 8.3	2,400
\$25,001-\$40,000	2,100	63.9	60.9 - 66.9	500	16.2	14.1 - 18.7	500	13.6	11.6 - 15.8	200	6.2	4.9 - 7.9	3,300
>=\$40,001	4,400	72.3	70.1 - 74.3	800	13.8	12.2 - 15.5	600	9.5	8.3 - 10.9	300	4.5	3.6 - 5.5	6,100

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	62.3	56.6 - 67.7	200	20.3	15.9 - 25.5	100	12.0	9.1 - 15.7	100	5.4	3.6 - 8.0	1,000
Normal birth weight	9,700	64.3	62.9 - 65.6	2,500	16.8	15.8 - 18.0	1,900	12.7	11.8 - 13.6	900	6.2	5.6 - 6.9	15,000

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	50.3	45.9 - 54.8	200	16.9	13.8 - 20.5	300	20.5	17.1 - 24.4	200	12.3	9.8 - 15.3	1,300
No	9,400	65.5	64.1 - 66.9	2,400	17.0	15.9 - 18.2	1,700	11.9	11.0 - 12.9	800	5.6	5.0 - 6.3	14,400

**PRAMS Health Indicator 3-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,800	56.1	53.1 - 59.1	500	16.7	14.6 - 19.1	600	18.5	16.3 - 21.0	300	8.6	7.2 - 10.3	3,200
No	8,200	66.4	64.8 - 67.9	2,100	16.9	15.7 - 18.1	1,400	11.1	10.2 - 12.1	700	5.6	5.0 - 6.4	12,400

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,000	58.0	56.0 - 60.1	1,300	19.4	17.8 - 21.2	1,000	14.7	13.4 - 16.1	500	7.8	6.9 - 8.9	6,800
No	6,200	68.7	66.9 - 70.4	1,400	15.2	13.8 - 16.6	1,000	11.2	10.1 - 12.4	400	4.9	4.2 - 5.8	9,100