

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- for the State of Hawaii, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Sleeping position	Sleeping position

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/22/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- for the State of Hawaii, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	10,800	65.8	64.7 - 66.8	2,900	17.6	16.7 - 18.4	2,000	12.4	11.8 - 13.1	700	4.2	3.8 - 4.6	16,500
2001-2005	11,100	66.6	65.5 - 67.6	2,700	16.5	15.7 - 17.4	2,000	12.0	11.3 - 12.7	800	4.9	4.5 - 5.4	16,600
2002-2006	11,400	67.4	66.4 - 68.5	2,800	16.3	15.5 - 17.1	2,000	12.0	11.3 - 12.7	700	4.3	3.8 - 4.7	17,000
2003-2007	11,800	68.5	67.4 - 69.5	2,700	15.7	15.0 - 16.6	2,000	11.8	11.1 - 12.6	700	4.0	3.5 - 4.4	17,300
2004-2008	12,200	69.1	68.1 - 70.1	2,600	14.8	14.1 - 15.6	2,100	11.8	11.1 - 12.5	800	4.4	3.9 - 4.8	17,600
2005-2009	12,600	71.0	69.9 - 72.0	2,400	13.5	12.7 - 14.3	2,000	11.1	10.4 - 11.9	800	4.5	4.0 - 5.0	17,700
2006-2010	13,000	72.4	71.3 - 73.5	2,400	13.3	12.5 - 14.2	1,900	10.7	9.9 - 11.5	700	3.6	3.2 - 4.1	18,000
2007-2011	13,500	74.6	73.4 - 75.7	2,300	12.6	11.8 - 13.5	1,700	9.5	8.8 - 10.3	600	3.3	2.8 - 3.8	18,200
2008-2012	13,900	76.6	75.5 - 77.8	2,200	12.1	11.2 - 13.0	1,500	8.4	7.7 - 9.2	500	2.8	2.4 - 3.3	18,200
2009-2013	14,200	78.2	76.9 - 79.3	2,100	11.6	10.7 - 12.5	1,400	7.8	7.0 - 8.6	500	2.5	2.1 - 3.0	18,100

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- for the State of Hawaii, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	14,200	78.2	76.9 - 79.3	2,100	11.6	10.7 - 12.5	1,400	7.8	7.0 - 8.6	500	2.5	2.1 - 3.0	18,100

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,800	75.4	73.3 - 77.4	300	13.9	12.4 - 15.7	200	7.8	6.6 - 9.1	100	2.8	2.2 - 3.8	2,300
Honolulu	10,300	79.5	77.8 - 81.0	1,400	10.7	9.5 - 12.0	1,000	7.5	6.5 - 8.6	300	2.3	1.8 - 3.0	13,000
Kauai	600	73.9	71.4 - 76.2	100	15.1	13.2 - 17.1	100	9.1	7.7 - 10.8	<50	1.9	1.3 - 2.9	800
Maui	1,500	74.7	72.7 - 76.6	300	12.9	11.5 - 14.4	200	9.1	7.9 - 10.5	100	3.3	2.6 - 4.2	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,700	82.8	80.4 - 84.8	400	9.2	7.6 - 11.0	300	7.0	5.6 - 8.6	<50	1.1	0.6 - 2.0	4,400
Native Hawaiian	3,900	72.8	70.4 - 75.0	700	13.3	11.6 - 15.2	600	10.4	8.9 - 12.1	200	3.5	2.7 - 4.6	5,400
Chinese	700	84.3	78.4 - 88.8	100	10.6	6.9 - 15.8	<50	4.5	2.3 - 8.5	<50	0.6	0.1 - 4.0	900
Filipino	2,500	78.9	75.9 - 81.6	400	12.5	10.4 - 14.9	200	5.4	4.0 - 7.3	100	3.2	2.2 - 4.5	3,200
Japanese	1,500	85.6	81.9 - 88.6	100	5.8	4.0 - 8.4	100	8.1	5.8 - 11.2	<50	0.5	0.1 - 1.7	1,700
Other	2,000	73.8	70.1 - 77.2	400	14.9	12.2 - 18.1	200	7.6	5.7 - 9.9	100	3.7	2.5 - 5.6	2,600

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	72.1	66.8 - 76.9	200	14.9	11.3 - 19.3	100	9.0	6.3 - 12.7	<50	4.1	2.4 - 6.9	1,200
20-24	3,000	74.0	71.1 - 76.6	500	12.6	10.7 - 14.9	400	9.2	7.5 - 11.2	200	4.2	3.1 - 5.7	4,100
25-34	7,700	79.7	78.0 - 81.2	1,000	10.8	9.7 - 12.1	700	7.4	6.4 - 8.5	200	2.1	1.6 - 2.7	9,600
35+	2,600	81.2	78.4 - 83.7	400	11.1	9.2 - 13.5	200	6.8	5.2 - 8.7	<50	0.9	0.5 - 1.6	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,100	77.4	75.6 - 79.1	1,000	11.4	10.2 - 12.8	800	8.7	7.6 - 10.0	200	2.5	1.9 - 3.2	9,200
Female	7,100	78.9	77.2 - 80.6	1,000	11.7	10.4 - 13.1	600	6.8	5.9 - 7.9	200	2.5	2.0 - 3.2	8,900

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	9,100	81.3	79.8 - 82.7	1,100	10.3	9.2 - 11.5	800	7.0	6.1 - 8.0	200	1.5	1.1 - 2.0	11,200
No	5,100	73.1	71.0 - 75.1	900	13.6	12.1 - 15.3	600	9.1	7.9 - 10.6	300	4.1	3.3 - 5.1	6,900

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,800	74.0	71.9 - 76.1	900	13.6	12.0 - 15.4	600	8.9	7.6 - 10.4	200	3.4	2.7 - 4.4	6,400
131-185%	2,200	78.7	75.4 - 81.7	300	11.8	9.5 - 14.5	200	7.1	5.4 - 9.4	100	2.4	1.5 - 3.8	2,800
186+%	6,100	82.3	80.5 - 84.0	700	9.3	8.0 - 10.7	500	7.0	5.9 - 8.2	100	1.4	0.9 - 2.1	7,400

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	78.3	76.2 - 80.3	100	11.3	9.8 - 13.0	100	7.1	5.9 - 8.4	<50	3.3	2.5 - 4.3	1,200
Normal birth weight	13,200	78.1	76.8 - 79.4	2,000	11.6	10.6 - 12.6	1,300	7.8	7.1 - 8.7	400	2.4	2.0 - 3.0	16,900

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	64.0	58.3 - 69.3	200	19.5	15.2 - 24.6	100	12.6	9.3 - 16.9	<50	3.9	2.3 - 6.6	1,100
No	13,200	79.1	77.8 - 80.3	1,800	11.0	10.1 - 12.0	1,200	7.5	6.7 - 8.3	400	2.4	2.0 - 2.9	16,600

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,500	71.5	68.4 - 74.4	500	13.7	11.6 - 16.2	300	10.0	8.2 - 12.1	200	4.8	3.5 - 6.4	3,400
No	11,400	79.7	78.4 - 81.0	1,600	11.1	10.1 - 12.1	1,000	7.3	6.5 - 8.2	300	1.9	1.5 - 2.4	14,300

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,800	73.7	71.7 - 75.6	1,100	13.7	12.2 - 15.3	700	8.8	7.6 - 10.2	300	3.8	3.0 - 4.7	7,900
No	8,200	81.7	80.1 - 83.2	1,000	9.8	8.7 - 11.0	700	7.0	6.0 - 8.0	200	1.5	1.1 - 2.1	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	13,900	76.6	75.5 - 77.8	2,200	12.1	11.2 - 13.0	1,500	8.4	7.7 - 9.2	500	2.8	2.4 - 3.3	18,200

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,700	72.7	70.4 - 74.9	400	15.4	13.6 - 17.3	200	8.8	7.5 - 10.4	100	3.2	2.4 - 4.2	2,300
Honolulu	10,200	78.2	76.7 - 79.6	1,400	10.9	9.8 - 12.1	1,100	8.2	7.3 - 9.2	300	2.7	2.1 - 3.3	13,000
Kauai	600	71.6	68.5 - 74.6	100	16.7	14.3 - 19.3	100	9.0	7.3 - 11.1	<50	2.7	1.6 - 4.3	800
Maui	1,500	73.2	70.9 - 75.3	300	14.0	12.4 - 15.9	200	9.3	7.9 - 10.8	100	3.5	2.7 - 4.6	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,500	81.6	79.4 - 83.6	400	9.7	8.2 - 11.4	300	7.6	6.3 - 9.2	<50	1.1	0.7 - 1.9	4,300
Native Hawaiian	3,800	70.9	68.6 - 73.1	700	13.7	12.1 - 15.5	600	11.4	9.9 - 13.1	200	3.9	3.0 - 5.0	5,300
Chinese	700	83.8	78.3 - 88.0	100	10.0	6.6 - 14.7	<50	5.5	3.3 - 9.2	<50	0.8	0.1 - 3.8	800
Filipino	2,500	77.4	74.7 - 80.0	400	13.2	11.2 - 15.5	200	5.8	4.5 - 7.5	100	3.5	2.6 - 4.9	3,300
Japanese	1,500	84.3	80.9 - 87.2	100	6.8	4.9 - 9.2	100	8.1	6.1 - 10.8	<50	0.8	0.3 - 2.0	1,700
Other	2,000	72.0	68.5 - 75.2	400	15.4	12.9 - 18.4	200	8.2	6.4 - 10.4	100	4.4	3.2 - 6.2	2,800

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	70.9	66.1 - 75.2	200	15.9	12.5 - 20.1	100	8.4	6.2 - 11.3	100	4.8	3.0 - 7.5	1,300
20-24	2,900	71.4	68.7 - 74.0	500	13.1	11.3 - 15.2	500	11.0	9.2 - 12.9	200	4.5	3.4 - 5.9	4,100
25-34	7,500	78.4	76.8 - 79.9	1,100	11.1	10.0 - 12.3	800	8.0	7.0 - 9.0	200	2.5	2.0 - 3.1	9,600
35+	2,600	80.3	77.7 - 82.7	400	12.2	10.2 - 14.4	200	6.6	5.2 - 8.3	<50	1.0	0.5 - 1.7	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,100	75.9	74.2 - 77.4	1,100	12.0	10.8 - 13.3	900	9.2	8.2 - 10.4	300	2.9	2.3 - 3.6	9,300
Female	6,900	77.4	75.8 - 79.0	1,100	12.2	11.0 - 13.5	700	7.6	6.7 - 8.7	200	2.8	2.2 - 3.5	8,900

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,900	79.5	78.0 - 80.8	1,200	11.1	10.1 - 12.2	900	7.8	7.0 - 8.8	200	1.6	1.2 - 2.1	11,200
No	5,000	72.1	70.1 - 74.0	1,000	13.7	12.2 - 15.3	700	9.4	8.2 - 10.8	300	4.8	4.0 - 5.9	7,000

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,600	71.6	69.5 - 73.6	1,000	14.9	13.4 - 16.6	600	9.6	8.4 - 11.1	200	3.8	3.1 - 4.8	6,500
131-185%	2,200	77.1	74.0 - 80.0	300	11.5	9.4 - 14.0	300	9.0	7.2 - 11.2	100	2.4	1.5 - 3.7	2,800
186+%	6,200	81.6	79.9 - 83.2	700	9.3	8.2 - 10.6	600	7.5	6.5 - 8.7	100	1.5	1.1 - 2.2	7,600

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	76.3	73.7 - 78.8	200	12.4	10.5 - 14.7	100	7.6	6.2 - 9.2	<50	3.7	2.7 - 5.0	1,200
Normal birth weight	13,000	76.6	75.4 - 77.8	2,100	12.1	11.2 - 13.0	1,400	8.5	7.7 - 9.3	500	2.8	2.3 - 3.3	17,000

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	61.2	56.0 - 66.1	300	20.4	16.4 - 25.0	200	14.0	10.8 - 17.9	100	4.5	2.8 - 7.1	1,200
No	12,900	77.7	76.5 - 78.9	1,900	11.5	10.6 - 12.4	1,300	8.1	7.3 - 8.9	500	2.7	2.3 - 3.2	16,600

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	68.8	65.9 - 71.6	500	14.9	12.8 - 17.3	400	11.0	9.3 - 13.1	200	5.2	4.0 - 6.8	3,500
No	11,300	78.4	77.2 - 79.7	1,600	11.4	10.5 - 12.4	1,100	7.9	7.1 - 8.7	300	2.3	1.9 - 2.8	14,400

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,700	71.9	70.0 - 73.7	1,100	14.2	12.8 - 15.7	800	9.6	8.5 - 10.9	300	4.3	3.5 - 5.2	7,900
No	8,100	80.4	78.9 - 81.8	1,000	10.4	9.3 - 11.5	800	7.5	6.6 - 8.5	200	1.7	1.3 - 2.3	10,000

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	13,500	74.6	73.4 - 75.7	2,300	12.6	11.8 - 13.5	1,700	9.5	8.8 - 10.3	600	3.3	2.8 - 3.8	18,200

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,700	70.4	67.9 - 72.7	400	16.2	14.3 - 18.3	200	9.5	8.0 - 11.2	100	4.0	3.0 - 5.2	2,300
Honolulu	9,800	76.2	74.7 - 77.6	1,500	11.5	10.4 - 12.6	1,200	9.3	8.4 - 10.3	400	3.0	2.5 - 3.7	12,900
Kauai	600	70.2	66.6 - 73.6	100	16.4	13.8 - 19.3	100	9.1	7.2 - 11.4	<50	4.3	2.8 - 6.7	900
Maui	1,500	71.0	68.4 - 73.4	300	14.1	12.3 - 16.1	200	11.3	9.7 - 13.2	100	3.6	2.7 - 4.8	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,400	80.0	77.8 - 81.9	400	10.3	8.9 - 12.0	400	8.4	7.1 - 9.9	100	1.4	0.8 - 2.2	4,300
Native Hawaiian	3,600	68.3	66.0 - 70.5	800	14.5	12.9 - 16.3	700	12.8	11.3 - 14.4	200	4.4	3.5 - 5.6	5,300
Chinese	600	80.2	74.6 - 84.9	100	12.1	8.4 - 17.0	<50	6.3	3.8 - 10.3	<50	1.3	0.5 - 3.9	700
Filipino	2,500	75.5	72.9 - 78.0	500	14.0	12.0 - 16.2	200	6.4	5.1 - 8.0	100	4.0	3.0 - 5.4	3,300
Japanese	1,600	84.3	81.2 - 87.0	100	6.1	4.5 - 8.3	200	8.8	6.8 - 11.4	<50	0.8	0.3 - 1.9	1,800
Other	1,900	69.2	65.8 - 72.4	400	15.4	13.0 - 18.2	300	10.1	8.1 - 12.4	100	5.3	4.0 - 7.1	2,700

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	66.9	62.4 - 71.1	200	16.1	12.9 - 19.9	200	11.5	9.0 - 14.6	100	5.6	3.8 - 8.1	1,400
20-24	3,000	69.3	66.7 - 71.8	600	13.8	12.1 - 15.8	500	11.8	10.1 - 13.7	200	5.1	4.0 - 6.4	4,300
25-34	7,100	76.6	75.0 - 78.1	1,100	11.7	10.6 - 12.9	800	8.9	8.0 - 10.0	300	2.8	2.2 - 3.4	9,300
35+	2,500	79.2	76.6 - 81.5	400	12.1	10.2 - 14.2	200	7.4	5.9 - 9.1	<50	1.4	0.8 - 2.2	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,000	74.1	72.5 - 75.6	1,200	12.6	11.5 - 13.9	900	9.9	8.9 - 11.0	300	3.4	2.8 - 4.1	9,400
Female	6,600	75.1	73.5 - 76.7	1,100	12.6	11.4 - 13.9	800	9.1	8.1 - 10.2	300	3.2	2.6 - 3.9	8,800

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,800	78.0	76.7 - 79.3	1,300	11.4	10.4 - 12.4	1,000	8.7	7.8 - 9.6	200	1.9	1.5 - 2.4	11,200
No	4,800	69.0	67.0 - 70.9	1,000	14.7	13.2 - 16.3	800	10.9	9.6 - 12.3	400	5.5	4.6 - 6.5	6,900

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,500	69.2	67.1 - 71.2	1,000	15.7	14.2 - 17.4	700	10.7	9.4 - 12.1	300	4.4	3.6 - 5.4	6,500
131-185%	2,100	74.4	71.4 - 77.2	300	12.4	10.4 - 14.8	300	10.7	8.9 - 12.9	100	2.5	1.6 - 3.8	2,800
186+%	6,200	80.4	78.8 - 81.9	700	9.4	8.3 - 10.6	600	8.4	7.4 - 9.5	100	1.8	1.4 - 2.5	7,700

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,200	70.7	67.8 - 73.5	500	14.6	12.5 - 16.9	300	10.3	8.6 - 12.4	100	4.4	3.3 - 5.8	3,100
\$10,000-\$24,999	2,500	68.4	65.6 - 71.0	600	15.5	13.5 - 17.7	400	10.8	9.1 - 12.7	200	5.3	4.1 - 6.9	3,600
\$25,000-\$49,999	3,200	76.6	74.2 - 78.8	500	12.1	10.4 - 14.0	400	9.4	8.0 - 11.0	100	1.9	1.3 - 2.8	4,200
>=\$50,000	4,900	80.3	78.5 - 82.0	600	9.3	8.1 - 10.7	500	8.9	7.8 - 10.3	100	1.4	0.9 - 2.1	6,100

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	74.4	71.4 - 77.2	200	12.6	10.6 - 15.0	100	8.8	7.1 - 10.8	100	4.2	3.0 - 5.9	1,300
Normal birth weight	12,600	74.6	73.4 - 75.8	2,100	12.6	11.7 - 13.6	1,600	9.6	8.8 - 10.4	500	3.2	2.7 - 3.7	16,900

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	61.1	56.4 - 65.6	300	20.3	16.7 - 24.5	200	14.0	11.1 - 17.5	100	4.6	3.0 - 7.0	1,400
No	12,500	75.7	74.6 - 76.9	2,000	12.0	11.1 - 12.9	1,500	9.1	8.4 - 9.9	500	3.2	2.7 - 3.7	16,500

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	66.9	64.1 - 69.6	600	15.9	13.9 - 18.2	400	11.8	10.0 - 13.7	200	5.4	4.2 - 6.9	3,600
No	10,900	76.5	75.3 - 77.7	1,700	11.8	10.9 - 12.7	1,300	9.0	8.2 - 9.8	400	2.8	2.3 - 3.3	14,300

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,300	68.9	67.0 - 70.7	1,200	15.3	13.9 - 16.8	800	10.5	9.3 - 11.8	400	5.3	4.4 - 6.2	7,800
No	8,000	78.9	77.5 - 80.3	1,100	10.5	9.5 - 11.6	900	8.7	7.8 - 9.7	200	1.8	1.4 - 2.4	10,100

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	13,000	72.4	71.3 - 73.5	2,400	13.3	12.5 - 14.2	1,900	10.7	9.9 - 11.5	700	3.6	3.2 - 4.1	18,000

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,600	69.4	66.7 - 72.0	400	17.1	15.0 - 19.4	200	9.7	8.1 - 11.6	100	3.8	2.8 - 5.1	2,300
Honolulu	9,500	74.0	72.6 - 75.4	1,500	12.1	11.1 - 13.1	1,300	10.4	9.5 - 11.4	400	3.5	2.9 - 4.1	12,800
Kauai	600	65.8	61.5 - 69.8	100	16.8	13.8 - 20.2	100	11.7	9.2 - 14.8	<50	5.8	3.8 - 8.6	900
Maui	1,400	68.3	65.5 - 71.0	300	15.4	13.4 - 17.8	300	12.7	10.9 - 14.9	100	3.5	2.6 - 4.8	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,300	78.1	76.0 - 80.1	500	10.6	9.2 - 12.2	400	9.6	8.3 - 11.2	100	1.6	1.1 - 2.2	4,300
Native Hawaiian	3,400	65.6	63.3 - 67.8	800	15.3	13.7 - 17.1	800	14.4	12.8 - 16.1	200	4.7	3.8 - 5.8	5,200
Chinese	500	77.4	72.0 - 82.0	100	13.1	9.4 - 17.9	100	7.8	5.2 - 11.5	<50	1.7	0.7 - 4.2	700
Filipino	2,400	73.0	70.4 - 75.4	500	15.7	13.8 - 17.7	200	7.2	5.9 - 8.8	100	4.2	3.1 - 5.5	3,300
Japanese	1,500	82.7	79.6 - 85.3	100	6.6	5.0 - 8.7	200	9.5	7.5 - 11.9	<50	1.2	0.6 - 2.4	1,900
Other	1,800	67.4	64.2 - 70.4	400	15.5	13.3 - 18.1	300	10.8	9.0 - 13.0	200	6.3	4.9 - 8.0	2,700

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	61.9	57.5 - 66.1	300	18.6	15.4 - 22.3	200	13.6	10.9 - 16.7	100	5.9	4.1 - 8.5	1,500
20-24	2,900	67.3	64.8 - 69.6	600	14.7	13.0 - 16.7	500	12.7	11.1 - 14.6	200	5.3	4.2 - 6.5	4,300
25-34	6,800	74.7	73.2 - 76.2	1,100	12.2	11.1 - 13.4	900	10.0	9.0 - 11.1	300	3.1	2.6 - 3.7	9,100
35+	2,500	77.4	74.9 - 79.7	400	12.1	10.4 - 14.2	300	8.5	7.1 - 10.3	100	1.9	1.3 - 2.8	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,700	71.4	69.8 - 72.9	1,300	13.7	12.6 - 15.0	1,100	11.5	10.5 - 12.7	300	3.4	2.8 - 4.1	9,300
Female	6,400	73.5	71.9 - 75.0	1,100	12.9	11.7 - 14.1	800	9.8	8.8 - 10.9	300	3.9	3.3 - 4.6	8,700

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,500	76.3	75.0 - 77.6	1,300	11.5	10.5 - 12.4	1,100	9.8	8.9 - 10.8	300	2.4	2.0 - 2.9	11,200
No	4,500	65.9	63.9 - 67.8	1,100	16.4	14.9 - 18.0	800	12.1	10.8 - 13.5	400	5.7	4.8 - 6.7	6,800

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,300	66.6	64.6 - 68.5	1,100	16.4	14.9 - 18.0	800	12.2	10.9 - 13.7	300	4.8	4.0 - 5.7	6,500
131-185%	1,900	71.2	68.2 - 74.0	400	13.6	11.5 - 15.9	300	11.8	9.9 - 14.1	100	3.4	2.4 - 4.8	2,700
186+%	6,000	78.8	77.2 - 80.2	800	9.9	8.9 - 11.1	700	9.3	8.3 - 10.4	200	2.0	1.5 - 2.6	7,700

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	65.9	62.9 - 68.7	500	16.2	14.0 - 18.5	400	13.0	11.1 - 15.2	200	5.0	3.8 - 6.5	3,100
\$10,000-\$24,999	2,500	67.5	64.8 - 70.0	600	15.4	13.5 - 17.5	400	11.7	10.0 - 13.5	200	5.5	4.4 - 7.0	3,700
\$25,000-\$49,999	3,100	73.8	71.5 - 76.0	600	13.3	11.7 - 15.1	400	10.6	9.2 - 12.3	100	2.2	1.6 - 3.0	4,200
>=\$50,000	4,800	79.1	77.4 - 80.7	600	9.7	8.5 - 11.0	600	9.5	8.3 - 10.8	100	1.8	1.3 - 2.4	6,000

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	73.0	69.7 - 76.1	200	13.0	10.7 - 15.7	100	9.4	7.5 - 11.7	100	4.6	3.2 - 6.4	1,300
Normal birth weight	12,100	72.3	71.2 - 73.5	2,200	13.3	12.5 - 14.2	1,800	10.8	10.0 - 11.6	600	3.6	3.1 - 4.1	16,700

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	60.5	56.1 - 64.7	300	19.3	15.9 - 23.1	200	15.6	12.7 - 19.0	100	4.6	3.1 - 6.8	1,500
No	11,900	73.5	72.4 - 74.6	2,100	12.7	11.9 - 13.6	1,700	10.2	9.5 - 11.0	600	3.5	3.1 - 4.0	16,200

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,400	65.8	63.1 - 68.4	600	16.1	14.1 - 18.3	500	12.9	11.2 - 14.8	200	5.3	4.1 - 6.7	3,600
No	10,400	74.1	72.9 - 75.3	1,800	12.6	11.7 - 13.5	1,400	10.1	9.3 - 11.0	500	3.2	2.8 - 3.7	14,100

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,000	66.7	64.9 - 68.5	1,200	15.7	14.3 - 17.1	900	12.0	10.8 - 13.3	400	5.6	4.8 - 6.6	7,600
No	7,800	76.6	75.2 - 78.0	1,200	11.6	10.6 - 12.7	1,000	9.6	8.6 - 10.5	200	2.2	1.8 - 2.8	10,100

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	12,600	71.0	69.9 - 72.0	2,400	13.5	12.7 - 14.3	2,000	11.1	10.4 - 11.9	800	4.5	4.0 - 5.0	17,700

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,500	66.3	63.4 - 69.1	400	17.0	14.9 - 19.4	200	10.8	9.1 - 12.9	100	5.8	4.6 - 7.4	2,300
Honolulu	9,200	72.8	71.5 - 74.0	1,500	12.2	11.3 - 13.2	1,400	10.8	10.0 - 11.8	500	4.1	3.6 - 4.7	12,600
Kauai	500	62.6	57.9 - 67.1	100	17.1	13.9 - 20.9	100	12.4	9.6 - 15.9	100	7.8	5.5 - 11.1	800
Maui	1,300	68.1	65.1 - 70.9	300	15.7	13.5 - 18.2	200	12.6	10.7 - 14.9	100	3.6	2.6 - 5.0	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,200	77.7	75.7 - 79.6	400	10.0	8.7 - 11.4	400	9.9	8.6 - 11.4	100	2.4	1.8 - 3.2	4,100
Native Hawaiian	3,200	63.4	61.2 - 65.6	800	15.9	14.3 - 17.6	800	15.1	13.5 - 16.8	300	5.6	4.7 - 6.8	5,100
Chinese	500	80.0	76.0 - 83.5	100	10.8	7.9 - 14.6	<50	7.4	5.6 - 9.6	<50	1.8	1.2 - 2.8	700
Filipino	2,300	72.1	69.8 - 74.3	500	16.2	14.4 - 18.1	200	7.3	6.1 - 8.8	100	4.4	3.5 - 5.5	3,200
Japanese	1,600	82.1	79.3 - 84.5	100	6.4	4.9 - 8.2	200	9.8	7.9 - 12.0	<50	1.8	1.1 - 2.9	2,000
Other	1,700	63.1	60.1 - 66.0	500	16.9	14.7 - 19.2	300	11.9	10.0 - 14.0	200	8.2	6.7 - 10.0	2,700

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	60.7	56.6 - 64.7	300	18.5	15.5 - 22.0	200	13.9	11.3 - 17.0	100	6.9	5.0 - 9.4	1,600
20-24	2,700	64.8	62.4 - 67.1	700	16.0	14.3 - 17.8	600	13.5	11.9 - 15.3	200	5.7	4.7 - 7.0	4,200
25-34	6,500	73.7	72.3 - 75.1	1,100	12.0	11.0 - 13.1	900	10.3	9.3 - 11.3	400	4.0	3.4 - 4.7	8,900
35+	2,400	76.6	74.3 - 78.8	400	11.7	10.1 - 13.4	300	8.8	7.4 - 10.5	100	2.9	2.1 - 3.8	3,100

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,400	69.8	68.3 - 71.2	1,300	14.0	12.9 - 15.1	1,100	11.8	10.8 - 12.9	400	4.5	3.8 - 5.2	9,200
Female	6,200	72.3	70.8 - 73.7	1,100	12.9	11.9 - 14.0	900	10.4	9.4 - 11.4	400	4.5	3.8 - 5.2	8,500

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,200	74.5	73.3 - 75.7	1,300	11.9	11.0 - 12.8	1,100	10.3	9.5 - 11.2	400	3.3	2.8 - 3.8	11,100
No	4,300	65.1	63.2 - 66.9	1,100	16.1	14.7 - 17.6	800	12.4	11.2 - 13.7	400	6.5	5.5 - 7.5	6,700

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,200	65.6	63.7 - 67.5	1,100	16.5	15.1 - 18.1	800	12.1	10.9 - 13.4	400	5.8	4.9 - 6.7	6,400
131-185%	1,700	67.7	64.7 - 70.6	400	15.0	12.9 - 17.4	300	13.3	11.3 - 15.6	100	4.0	3.0 - 5.5	2,500
186+%	6,000	77.9	76.5 - 79.3	800	9.8	8.9 - 10.9	800	9.8	8.8 - 10.9	200	2.5	2.0 - 3.0	7,700

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	65.4	62.5 - 68.1	500	16.8	14.7 - 19.1	400	12.4	10.6 - 14.5	200	5.4	4.3 - 6.9	3,000
\$10,000-\$24,999	2,500	65.8	63.4 - 68.2	600	15.6	13.8 - 17.5	500	12.1	10.5 - 13.8	200	6.5	5.3 - 7.9	3,800
\$25,000-\$49,999	2,900	71.0	68.8 - 73.1	600	14.3	12.7 - 16.1	500	11.6	10.1 - 13.2	100	3.1	2.4 - 4.0	4,100
>=\$50,000	4,600	79.0	77.3 - 80.5	500	9.1	8.1 - 10.2	600	9.8	8.7 - 11.0	100	2.1	1.6 - 2.8	5,900

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	70.1	66.5 - 73.5	200	14.1	11.6 - 17.0	100	10.0	8.0 - 12.6	100	5.8	4.2 - 8.0	1,300
Normal birth weight	11,700	71.0	69.9 - 72.1	2,200	13.4	12.6 - 14.3	1,800	11.2	10.4 - 12.0	700	4.4	3.9 - 4.9	16,400

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	61.2	57.1 - 65.2	300	17.6	14.5 - 21.1	200	15.0	12.4 - 18.1	100	6.2	4.5 - 8.4	1,500
No	11,500	71.9	70.9 - 73.0	2,100	13.0	12.2 - 13.8	1,700	10.7	10.0 - 11.5	700	4.4	3.9 - 4.9	15,900

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,300	65.4	62.9 - 67.9	500	15.3	13.5 - 17.3	500	13.4	11.7 - 15.2	200	5.9	4.8 - 7.3	3,600
No	10,000	72.5	71.3 - 73.6	1,800	12.8	12.0 - 13.7	1,500	10.5	9.7 - 11.4	600	4.1	3.7 - 4.7	13,900

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,900	65.6	63.8 - 67.3	1,200	15.8	14.5 - 17.1	900	12.5	11.3 - 13.7	500	6.2	5.4 - 7.1	7,400
No	7,500	74.9	73.6 - 76.2	1,200	11.7	10.8 - 12.7	1,000	10.0	9.2 - 11.0	300	3.3	2.8 - 3.9	10,000

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	12,200	69.1	68.1 - 70.1	2,600	14.8	14.1 - 15.6	2,100	11.8	11.1 - 12.5	800	4.4	3.9 - 4.8	17,600

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,500	65.1	62.1 - 68.0	400	17.1	14.9 - 19.6	300	12.1	10.2 - 14.2	100	5.7	4.5 - 7.4	2,300
Honolulu	8,900	70.8	69.6 - 71.9	1,700	13.7	12.9 - 14.6	1,400	11.5	10.7 - 12.3	500	4.0	3.6 - 4.6	12,600
Kauai	500	61.2	56.2 - 66.0	200	19.1	15.5 - 23.2	100	12.6	9.6 - 16.4	100	7.1	4.8 - 10.4	800
Maui	1,300	66.2	63.0 - 69.2	300	17.3	15.0 - 20.0	200	12.9	10.8 - 15.3	100	3.6	2.5 - 5.1	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,000	75.1	73.1 - 77.0	500	11.5	10.2 - 13.0	400	11.0	9.7 - 12.5	100	2.4	1.8 - 3.2	4,000
Native Hawaiian	3,000	61.4	59.2 - 63.5	800	16.6	15.1 - 18.3	800	16.4	14.8 - 18.1	300	5.6	4.6 - 6.7	4,900
Chinese	500	79.5	77.1 - 81.7	100	10.2	8.6 - 12.1	100	8.2	6.8 - 9.9	<50	2.1	1.4 - 3.1	600
Filipino	2,300	70.6	68.5 - 72.7	600	18.1	16.4 - 20.0	200	7.2	6.1 - 8.5	100	4.1	3.2 - 5.1	3,200
Japanese	1,700	80.7	78.1 - 83.0	200	7.3	5.8 - 9.0	200	10.2	8.4 - 12.2	<50	1.9	1.2 - 2.9	2,100
Other	1,600	60.6	57.9 - 63.3	500	19.5	17.3 - 21.8	300	12.1	10.3 - 14.0	200	7.8	6.4 - 9.5	2,700

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	56.7	52.7 - 60.7	300	20.1	17.1 - 23.5	200	17.0	14.2 - 20.3	100	6.1	4.5 - 8.4	1,500
20-24	2,700	63.3	61.1 - 65.4	700	17.2	15.6 - 19.0	600	14.0	12.5 - 15.6	200	5.5	4.5 - 6.6	4,300
25-34	6,300	71.7	70.4 - 73.1	1,200	13.6	12.6 - 14.7	900	10.6	9.7 - 11.6	300	4.0	3.4 - 4.7	8,700
35+	2,400	75.3	73.1 - 77.3	400	12.3	10.8 - 14.0	300	9.5	8.2 - 11.0	100	2.9	2.2 - 3.9	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,200	67.3	65.9 - 68.7	1,400	15.5	14.4 - 16.6	1,200	12.8	11.8 - 13.8	400	4.4	3.9 - 5.1	9,200
Female	6,000	71.0	69.6 - 72.4	1,200	14.1	13.1 - 15.2	900	10.7	9.7 - 11.7	400	4.2	3.6 - 4.9	8,400

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,200	72.3	71.1 - 73.4	1,500	13.3	12.5 - 14.2	1,200	10.9	10.2 - 11.8	400	3.4	3.0 - 4.0	11,400
No	4,000	63.3	61.5 - 65.1	1,100	17.5	16.1 - 18.9	800	13.2	12.0 - 14.6	400	6.0	5.1 - 7.0	6,300

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,100	64.1	62.3 - 65.8	1,200	18.1	16.7 - 19.5	800	12.3	11.2 - 13.6	400	5.5	4.7 - 6.4	6,500
131-185%	1,500	65.2	62.3 - 67.9	400	16.8	14.7 - 19.1	300	14.3	12.3 - 16.5	100	3.8	2.8 - 5.1	2,300
186+%	5,900	75.9	74.6 - 77.3	800	10.8	9.9 - 11.8	800	10.7	9.8 - 11.7	200	2.5	2.1 - 3.1	7,700

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Income (2004-2011)	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,800	62.7	60.0 - 65.3	600	19.0	16.9 - 21.3	400	13.0	11.2 - 15.0	200	5.3	4.2 - 6.7	2,900
\$10,000-\$24,999	2,500	64.5	62.2 - 66.7	700	17.4	15.7 - 19.2	500	12.3	10.8 - 13.9	200	5.9	4.9 - 7.1	3,900
\$25,000-\$49,999	2,900	68.9	66.8 - 70.8	600	15.2	13.7 - 16.8	500	12.5	11.1 - 14.0	100	3.4	2.7 - 4.3	4,200
>=\$50,000	4,400	77.7	76.2 - 79.2	600	9.8	8.7 - 10.9	600	10.4	9.4 - 11.6	100	2.1	1.6 - 2.7	5,600

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	69.4	65.6 - 73.0	200	14.7	12.0 - 17.8	100	10.4	8.1 - 13.1	100	5.6	3.9 - 7.9	1,300
Normal birth weight	11,300	69.1	68.0 - 70.1	2,400	14.8	14.1 - 15.6	1,900	11.9	11.2 - 12.6	700	4.2	3.8 - 4.7	16,400

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	61.2	57.3 - 64.9	300	17.8	15.0 - 21.0	200	15.5	12.9 - 18.6	100	5.5	3.9 - 7.7	1,500
No	11,100	69.8	68.8 - 70.9	2,300	14.5	13.7 - 15.3	1,800	11.4	10.7 - 12.2	700	4.3	3.8 - 4.8	15,900

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	63.6	61.2 - 65.9	600	16.7	14.9 - 18.7	500	14.5	12.9 - 16.4	200	5.2	4.1 - 6.4	3,500
No	9,800	70.5	69.4 - 71.6	2,000	14.2	13.4 - 15.1	1,500	11.1	10.3 - 11.9	600	4.2	3.7 - 4.7	13,800

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,800	63.6	62.0 - 65.2	1,300	17.7	16.5 - 19.0	1,000	12.8	11.7 - 14.0	400	5.9	5.1 - 6.7	7,500
No	7,200	73.2	72.0 - 74.5	1,200	12.6	11.7 - 13.6	1,100	10.9	10.0 - 11.8	300	3.3	2.8 - 3.8	9,800

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,800	68.5	67.4 - 69.5	2,700	15.7	15.0 - 16.6	2,000	11.8	11.1 - 12.6	700	4.0	3.5 - 4.4	17,300

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,400	64.8	61.9 - 67.6	400	17.6	15.5 - 20.0	300	12.1	10.3 - 14.2	100	5.4	4.2 - 7.0	2,200
Honolulu	8,700	70.0	68.8 - 71.2	1,900	14.9	13.9 - 15.8	1,400	11.4	10.6 - 12.3	500	3.7	3.2 - 4.2	12,500
Kauai	500	59.1	54.4 - 63.6	200	20.8	17.3 - 24.7	100	13.8	10.8 - 17.4	100	6.4	4.2 - 9.5	800
Maui	1,200	66.4	63.3 - 69.3	300	17.2	15.0 - 19.8	300	13.5	11.4 - 15.8	100	2.9	2.0 - 4.2	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	74.0	71.9 - 76.0	500	12.4	10.9 - 14.0	400	11.2	9.8 - 12.8	100	2.4	1.8 - 3.2	3,900
Native Hawaiian	2,900	61.0	59.0 - 63.0	800	17.1	15.6 - 18.7	800	16.9	15.4 - 18.5	200	5.1	4.2 - 6.1	4,800
Chinese	500	78.7	75.1 - 82.0	100	11.5	9.1 - 14.5	100	7.8	5.8 - 10.6	<50	1.9	1.2 - 2.9	600
Filipino	2,300	69.5	67.2 - 71.7	700	19.9	18.0 - 22.0	200	7.1	5.9 - 8.4	100	3.5	2.7 - 4.4	3,300
Japanese	1,700	82.7	80.2 - 85.0	100	6.6	5.2 - 8.4	200	8.9	7.3 - 10.8	<50	1.8	1.1 - 2.9	2,100
Other	1,500	58.5	55.5 - 61.4	600	21.4	19.0 - 24.0	300	13.0	11.0 - 15.2	200	7.1	5.8 - 8.8	2,600

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	56.6	52.6 - 60.4	300	20.3	17.3 - 23.6	300	18.0	15.2 - 21.3	100	5.2	3.7 - 7.2	1,500
20-24	2,700	63.1	60.9 - 65.3	800	18.8	17.1 - 20.6	600	13.4	11.9 - 15.0	200	4.7	3.9 - 5.8	4,200
25-34	6,100	71.1	69.6 - 72.5	1,200	14.2	13.2 - 15.4	900	11.0	10.1 - 12.1	300	3.7	3.1 - 4.3	8,500
35+	2,300	74.4	72.2 - 76.6	400	13.5	11.8 - 15.4	300	8.9	7.6 - 10.4	100	3.1	2.4 - 4.2	3,100

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,100	67.2	65.8 - 68.6	1,500	16.0	15.0 - 17.2	1,200	12.7	11.8 - 13.8	400	4.0	3.4 - 4.6	9,100
Female	5,800	69.8	68.3 - 71.3	1,300	15.4	14.3 - 16.6	900	10.8	9.9 - 11.9	300	3.9	3.4 - 4.6	8,200

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,100	71.8	70.6 - 73.0	1,600	14.1	13.2 - 15.1	1,200	10.8	10.0 - 11.6	400	3.3	2.9 - 3.8	11,300
No	3,800	62.3	60.4 - 64.1	1,100	18.7	17.3 - 20.2	800	13.8	12.6 - 15.2	300	5.2	4.4 - 6.1	6,000

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,200	63.5	61.8 - 65.3	1,200	18.9	17.5 - 20.3	800	12.6	11.5 - 13.9	300	5.0	4.2 - 5.8	6,600
131-185%	1,500	65.0	62.0 - 67.8	400	18.9	16.6 - 21.5	300	12.6	10.7 - 14.7	100	3.5	2.6 - 4.8	2,300
186+%	5,500	75.5	74.1 - 76.9	800	11.1	10.1 - 12.2	800	10.9	9.9 - 12.0	200	2.4	2.0 - 3.0	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	68.7	64.8 - 72.4	200	15.2	12.5 - 18.4	100	11.1	8.8 - 13.9	100	5.0	3.5 - 7.2	1,200
Normal birth weight	11,000	68.5	67.4 - 69.5	2,500	15.8	15.0 - 16.6	1,900	11.9	11.2 - 12.6	600	3.9	3.4 - 4.3	16,000

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	62.0	58.1 - 65.7	300	18.3	15.4 - 21.5	200	14.9	12.3 - 17.9	100	4.8	3.4 - 6.8	1,500
No	10,800	69.2	68.1 - 70.2	2,400	15.4	14.6 - 16.3	1,800	11.5	10.8 - 12.3	600	3.9	3.5 - 4.4	15,600

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,200	63.8	61.4 - 66.2	600	17.1	15.4 - 19.1	500	14.7	13.0 - 16.5	200	4.3	3.4 - 5.5	3,500
No	9,400	69.8	68.7 - 70.9	2,100	15.3	14.4 - 16.2	1,500	11.0	10.3 - 11.8	500	3.9	3.4 - 4.4	13,500

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,600	62.7	61.1 - 64.4	1,400	18.8	17.5 - 20.2	1,000	13.2	12.1 - 14.4	400	5.3	4.5 - 6.1	7,400
No	7,000	72.8	71.5 - 74.1	1,300	13.5	12.5 - 14.5	1,000	10.7	9.8 - 11.6	300	3.0	2.6 - 3.6	9,600

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,400	67.4	66.4 - 68.5	2,800	16.3	15.5 - 17.1	2,000	12.0	11.3 - 12.7	700	4.3	3.8 - 4.7	17,000

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,300	62.9	60.2 - 65.6	400	17.9	15.8 - 20.1	300	12.9	11.1 - 14.8	100	6.3	5.1 - 7.9	2,100
Honolulu	8,600	69.3	68.0 - 70.5	1,900	15.3	14.3 - 16.3	1,400	11.6	10.7 - 12.5	500	3.9	3.4 - 4.5	12,400
Kauai	400	55.0	50.4 - 59.4	200	23.5	19.9 - 27.4	100	15.4	12.3 - 19.1	<50	6.2	4.2 - 9.0	800
Maui	1,200	65.4	62.5 - 68.1	300	18.3	16.1 - 20.7	200	12.7	10.9 - 14.8	100	3.7	2.7 - 4.9	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	72.4	70.2 - 74.6	500	13.0	11.4 - 14.7	500	12.0	10.5 - 13.8	100	2.6	1.9 - 3.4	3,800
Native Hawaiian	2,800	60.2	58.3 - 62.1	800	17.3	15.9 - 18.8	800	17.0	15.6 - 18.5	300	5.5	4.7 - 6.5	4,700
Chinese	500	79.5	75.2 - 83.1	100	10.4	7.8 - 13.7	100	8.2	5.8 - 11.3	<50	2.0	1.0 - 3.9	600
Filipino	2,300	68.8	66.3 - 71.1	700	21.1	19.0 - 23.3	200	6.2	5.1 - 7.6	100	4.0	3.1 - 5.0	3,300
Japanese	1,700	80.6	77.8 - 83.1	200	7.0	5.5 - 8.9	200	10.0	8.2 - 12.2	100	2.4	1.5 - 3.7	2,200
Other	1,400	57.1	53.8 - 60.3	600	22.8	20.1 - 25.7	300	12.9	10.9 - 15.4	200	7.2	5.7 - 9.0	2,400

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	55.7	51.8 - 59.5	300	21.0	18.0 - 24.3	300	18.2	15.3 - 21.4	100	5.2	3.7 - 7.2	1,400
20-24	2,600	62.4	60.2 - 64.6	800	19.1	17.4 - 21.0	600	13.9	12.5 - 15.6	200	4.5	3.7 - 5.5	4,100
25-34	5,900	70.3	68.8 - 71.8	1,200	14.7	13.6 - 15.8	900	10.9	9.9 - 12.0	300	4.1	3.5 - 4.8	8,400
35+	2,100	72.0	69.6 - 74.4	400	14.6	12.8 - 16.7	300	9.4	8.0 - 11.0	100	4.0	3.1 - 5.2	3,000

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,800	65.8	64.4 - 67.3	1,500	16.6	15.5 - 17.8	1,200	13.3	12.3 - 14.3	400	4.3	3.7 - 4.9	8,800
Female	5,600	69.2	67.6 - 70.7	1,300	15.9	14.8 - 17.2	900	10.6	9.7 - 11.7	300	4.3	3.7 - 5.0	8,100

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,900	70.7	69.4 - 71.9	1,600	14.6	13.6 - 15.6	1,200	10.9	10.1 - 11.8	400	3.9	3.4 - 4.4	11,200
No	3,500	61.2	59.3 - 63.0	1,100	19.6	18.1 - 21.2	800	14.1	12.9 - 15.5	300	5.1	4.3 - 6.0	5,700

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,000	62.0	60.2 - 63.8	1,300	19.6	18.2 - 21.1	900	13.1	11.9 - 14.3	300	5.4	4.6 - 6.2	6,500
131-185%	1,600	64.6	61.6 - 67.6	500	18.8	16.5 - 21.5	300	12.8	10.9 - 15.0	100	3.7	2.8 - 5.0	2,400
186+%	5,100	74.7	73.2 - 76.2	800	11.7	10.7 - 12.9	700	10.7	9.6 - 11.8	200	2.9	2.3 - 3.6	6,800

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	67.0	62.8 - 71.0	200	18.0	14.8 - 21.7	100	10.0	7.7 - 12.8	100	5.0	3.5 - 7.1	1,200
Normal birth weight	10,700	67.5	66.4 - 68.6	2,600	16.2	15.3 - 17.0	1,900	12.1	11.4 - 12.9	700	4.2	3.8 - 4.7	15,800

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	59.0	55.1 - 62.8	300	18.4	15.5 - 21.6	200	15.9	13.1 - 19.0	100	6.8	5.1 - 8.9	1,400
No	10,400	68.3	67.2 - 69.4	2,500	16.0	15.2 - 16.9	1,800	11.6	10.9 - 12.4	600	4.1	3.6 - 4.5	15,300

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	62.1	59.7 - 64.6	600	17.3	15.5 - 19.3	500	15.6	13.8 - 17.5	200	5.0	4.0 - 6.1	3,400
No	9,100	69.0	67.8 - 70.1	2,100	15.9	15.0 - 16.9	1,500	11.0	10.3 - 11.8	500	4.1	3.6 - 4.6	13,200

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,500	62.2	60.5 - 63.8	1,400	18.9	17.6 - 20.3	1,000	13.8	12.6 - 15.0	400	5.1	4.4 - 5.9	7,200
No	6,800	71.4	70.0 - 72.7	1,400	14.3	13.3 - 15.4	1,000	10.6	9.7 - 11.5	400	3.7	3.2 - 4.3	9,500

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	11,100	66.6	65.5 - 67.6	2,700	16.5	15.7 - 17.4	2,000	12.0	11.3 - 12.7	800	4.9	4.5 - 5.4	16,600

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	60.7	58.2 - 63.2	400	17.9	16.0 - 19.9	300	14.1	12.4 - 16.0	100	7.3	6.0 - 8.8	2,000
Honolulu	8,300	68.3	67.0 - 69.6	1,900	15.7	14.7 - 16.8	1,400	11.5	10.7 - 12.4	500	4.4	3.9 - 5.0	12,200
Kauai	400	55.7	51.6 - 59.7	200	24.2	20.9 - 27.9	100	13.8	11.2 - 16.9	<50	6.2	4.5 - 8.6	700
Maui	1,100	65.4	62.7 - 67.9	300	17.1	15.1 - 19.2	200	12.0	10.4 - 13.9	100	5.5	4.4 - 6.9	1,700

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,600	71.1	68.8 - 73.3	500	13.6	12.0 - 15.3	500	12.5	10.9 - 14.3	100	2.8	2.2 - 3.7	3,600
Native Hawaiian	2,700	58.8	57.1 - 60.5	800	17.5	16.2 - 18.8	800	17.2	16.0 - 18.5	300	6.5	5.7 - 7.4	4,600
Chinese	500	79.5	74.6 - 83.6	100	10.3	7.5 - 13.9	<50	7.3	4.9 - 10.9	<50	3.0	1.5 - 5.7	600
Filipino	2,300	68.3	65.9 - 70.7	700	20.4	18.3 - 22.6	200	6.1	5.0 - 7.4	200	5.2	4.2 - 6.4	3,300
Japanese	1,700	80.7	77.9 - 83.2	200	7.1	5.6 - 9.0	200	9.6	7.8 - 11.8	100	2.6	1.7 - 3.9	2,200
Other	1,300	55.5	52.2 - 58.8	600	24.1	21.3 - 27.2	300	13.0	10.9 - 15.4	200	7.4	5.8 - 9.2	2,300

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	57.1	53.5 - 60.7	300	19.3	16.6 - 22.3	200	16.8	14.3 - 19.7	100	6.8	5.3 - 8.7	1,500
20-24	2,500	60.8	58.6 - 62.9	800	19.5	17.8 - 21.3	600	14.6	13.2 - 16.2	200	5.1	4.3 - 6.0	4,100
25-34	5,700	69.3	67.8 - 70.7	1,300	15.3	14.1 - 16.4	900	10.8	9.9 - 11.8	400	4.7	4.1 - 5.4	8,200
35+	2,000	71.9	69.4 - 74.3	400	14.4	12.5 - 16.5	300	9.2	7.8 - 10.8	100	4.5	3.5 - 5.7	2,800

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,600	65.0	63.5 - 66.5	1,400	16.6	15.5 - 17.8	1,100	13.2	12.2 - 14.2	500	5.2	4.6 - 5.9	8,600
Female	5,400	68.2	66.7 - 69.7	1,300	16.5	15.3 - 17.7	900	10.7	9.8 - 11.7	400	4.6	4.0 - 5.3	8,000

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,700	69.6	68.3 - 70.8	1,700	15.1	14.1 - 16.1	1,200	11.0	10.2 - 11.9	500	4.3	3.8 - 4.9	11,100
No	3,300	60.4	58.6 - 62.2	1,100	19.4	17.9 - 20.9	800	14.0	12.8 - 15.3	300	6.2	5.4 - 7.1	5,500

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,000	61.1	59.4 - 62.8	1,300	20.0	18.6 - 21.4	800	12.7	11.7 - 13.9	400	6.1	5.4 - 7.0	6,500
131-185%	1,700	64.2	61.4 - 66.9	500	18.8	16.6 - 21.2	300	12.9	11.2 - 15.0	100	4.0	3.2 - 5.0	2,600
186+%	4,600	74.4	72.7 - 75.9	700	11.7	10.6 - 12.9	700	10.6	9.5 - 11.8	200	3.4	2.8 - 4.1	6,200

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	64.8	60.5 - 68.9	200	18.7	15.5 - 22.5	100	10.5	8.2 - 13.3	100	6.0	4.3 - 8.2	1,100
Normal birth weight	10,400	66.7	65.6 - 67.7	2,500	16.4	15.5 - 17.2	1,900	12.1	11.4 - 12.8	800	4.8	4.4 - 5.3	15,500

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	57.4	53.6 - 61.0	300	18.3	15.6 - 21.4	200	15.9	13.3 - 18.8	100	8.5	6.8 - 10.6	1,400
No	10,100	67.5	66.4 - 68.6	2,400	16.3	15.4 - 17.1	1,700	11.6	10.9 - 12.3	700	4.6	4.2 - 5.1	14,900

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,100	61.1	58.7 - 63.4	600	17.5	15.8 - 19.5	500	15.5	13.8 - 17.3	200	5.9	4.9 - 7.0	3,400
No	8,800	68.1	67.0 - 69.3	2,100	16.1	15.2 - 17.1	1,400	11.0	10.3 - 11.8	600	4.7	4.2 - 5.3	12,900

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,300	61.0	59.4 - 62.7	1,400	19.2	17.9 - 20.5	1,000	13.8	12.7 - 15.0	400	6.0	5.3 - 6.7	7,100
No	6,600	70.7	69.3 - 72.0	1,300	14.5	13.4 - 15.6	1,000	10.6	9.8 - 11.6	400	4.2	3.6 - 4.8	9,300

**PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

State	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,800	65.8	64.7 - 66.8	2,900	17.6	16.7 - 18.4	2,000	12.4	11.8 - 13.1	700	4.2	3.8 - 4.6	16,500

County	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	60.8	58.5 - 63.0	400	18.8	17.0 - 20.6	300	15.1	13.6 - 16.9	100	5.3	4.4 - 6.4	1,900
Honolulu	8,200	67.6	66.3 - 68.9	2,000	16.7	15.7 - 17.8	1,400	11.8	10.9 - 12.6	500	3.9	3.4 - 4.4	12,100
Kauai	400	54.2	50.5 - 58.0	200	25.2	22.0 - 28.6	100	15.1	12.7 - 17.9	<50	5.5	4.1 - 7.3	700
Maui	1,100	63.1	60.7 - 65.5	300	18.8	16.9 - 20.9	200	13.1	11.5 - 14.8	100	5.0	4.1 - 6.2	1,700

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

DOH Race-Ethnicity	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,500	69.9	67.6 - 72.0	600	15.1	13.5 - 16.9	500	12.7	11.2 - 14.4	100	2.3	1.7 - 3.1	3,600
Native Hawaiian	2,600	57.6	56.1 - 59.1	800	18.5	17.4 - 19.7	800	18.1	17.0 - 19.3	300	5.8	5.2 - 6.5	4,500
Chinese	500	78.9	73.6 - 83.3	100	11.1	8.1 - 15.2	<50	7.0	4.4 - 10.9	<50	3.0	1.4 - 6.0	600
Filipino	2,200	66.8	64.3 - 69.2	700	21.8	19.7 - 24.1	200	6.7	5.6 - 8.1	200	4.7	3.8 - 5.9	3,300
Japanese	1,700	80.0	77.1 - 82.6	200	8.1	6.4 - 10.1	200	9.7	7.8 - 11.9	<50	2.2	1.4 - 3.5	2,100
Other	1,300	56.9	53.5 - 60.3	500	24.0	21.1 - 27.2	300	13.3	11.2 - 15.9	100	5.7	4.3 - 7.5	2,300

PRAMS Age Group	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	54.1	50.7 - 57.4	300	20.8	18.2 - 23.7	300	18.6	16.1 - 21.3	100	6.5	5.2 - 8.2	1,500
20-24	2,500	60.7	58.7 - 62.8	800	20.3	18.6 - 22.1	600	14.7	13.4 - 16.3	200	4.2	3.5 - 5.0	4,100
25-34	5,500	68.2	66.7 - 69.7	1,300	16.5	15.4 - 17.8	900	11.3	10.3 - 12.3	300	4.0	3.4 - 4.6	8,100
35+	2,000	72.4	69.9 - 74.8	400	14.8	12.9 - 16.9	300	9.1	7.7 - 10.6	100	3.7	2.8 - 4.8	2,800

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Infant Gender	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,500	64.3	62.8 - 65.7	1,500	17.2	16.1 - 18.4	1,200	14.1	13.1 - 15.1	400	4.4	3.9 - 5.0	8,600
Female	5,300	67.4	65.9 - 68.9	1,400	17.9	16.7 - 19.2	800	10.6	9.7 - 11.6	300	4.0	3.5 - 4.6	7,900

Married	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,700	69.3	68.0 - 70.5	1,800	16.1	15.1 - 17.1	1,200	11.1	10.2 - 11.9	400	3.6	3.2 - 4.1	11,200
No	3,100	58.4	56.6 - 60.2	1,100	20.8	19.3 - 22.3	800	15.3	14.1 - 16.6	300	5.5	4.8 - 6.3	5,300

Poverty Level	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,000	59.6	58.0 - 61.3	1,400	21.2	19.9 - 22.7	900	13.8	12.7 - 14.9	400	5.4	4.7 - 6.2	6,600
131-185%	1,800	64.9	62.2 - 67.5	500	18.8	16.7 - 21.1	400	13.0	11.3 - 14.9	100	3.3	2.6 - 4.3	2,700
186+%	4,300	74.2	72.5 - 75.8	700	12.2	11.1 - 13.5	600	10.5	9.4 - 11.6	200	3.1	2.5 - 3.8	5,800

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Birthweight	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	65.3	61.0 - 69.4	200	19.7	16.3 - 23.6	100	11.4	9.1 - 14.3	<50	3.5	2.4 - 5.1	1,000
Normal birth weight	10,100	65.8	64.7 - 66.9	2,700	17.4	16.6 - 18.3	1,900	12.5	11.8 - 13.2	700	4.3	3.9 - 4.7	15,400

Smoke 3rd Trimester	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	54.6	51.0 - 58.2	300	19.5	16.7 - 22.5	300	18.5	15.9 - 21.5	100	7.4	5.9 - 9.3	1,400
No	9,900	66.9	65.9 - 68.0	2,600	17.3	16.5 - 18.2	1,700	11.8	11.1 - 12.6	600	3.9	3.5 - 4.3	14,800

Smoke Before Pregnant	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,000	59.2	56.9 - 61.5	600	18.2	16.4 - 20.1	600	17.3	15.5 - 19.1	200	5.4	4.5 - 6.4	3,300
No	8,600	67.6	66.4 - 68.8	2,200	17.3	16.4 - 18.3	1,400	11.1	10.4 - 11.9	500	4.0	3.5 - 4.5	12,700

PRAMS Health Indicator 5-Year Aggregate Report - Sleeping position- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

WIC During Pregnancy	Back			Side			Stomach			More than 1 position			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,200	59.7	58.1 - 61.2	1,500	20.5	19.2 - 21.9	1,000	14.5	13.4 - 15.6	400	5.4	4.7 - 6.1	7,100
No	6,500	70.4	69.0 - 71.8	1,400	15.3	14.2 - 16.4	1,000	10.9	10.0 - 11.8	300	3.4	2.9 - 4.0	9,200