

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- for the State of Hawaii, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 3-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, income data does not exist for some 3-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Pregnancy intention	Pregnancy intention

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 9/30/15

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- for the State of Hawaii, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2002	9,500	56.0	54.6 - 57.4	1,400	8.5	7.8 - 9.3	6,000	35.5	34.2 - 36.9	16,900
2001-2003	9,500	55.8	54.4 - 57.3	1,400	8.4	7.6 - 9.2	6,100	35.8	34.3 - 37.2	17,100
2002-2004	9,800	56.3	54.9 - 57.8	1,700	9.8	9.0 - 10.6	5,900	33.9	32.5 - 35.3	17,400
2003-2005	9,700	55.4	54.0 - 56.8	2,100	12.1	11.2 - 13.0	5,700	32.5	31.2 - 33.8	17,400
2004-2006	9,600	54.7	53.3 - 56.0	2,500	14.2	13.3 - 15.2	5,500	31.1	29.9 - 32.4	17,600
2005-2007	9,600	53.6	52.3 - 55.0	2,500	14.0	13.1 - 15.0	5,800	32.3	31.0 - 33.7	17,800
2006-2008	10,000	54.6	53.2 - 55.9	2,200	12.3	11.4 - 13.3	6,000	33.1	31.8 - 34.5	18,200
2007-2009	9,900	54.1	52.5 - 55.6	2,100	11.6	10.6 - 12.6	6,300	34.3	32.9 - 35.8	18,300
2008-2010	10,000	54.5	52.8 - 56.2	1,900	10.5	9.5 - 11.6	6,400	34.9	33.3 - 36.6	18,300
2009-2011	9,900	53.9	52.1 - 55.7	2,100	11.2	10.1 - 12.4	6,400	34.9	33.1 - 36.7	18,300

PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- for the State of Hawaii, for the Year(s) 2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,900	53.9	52.1 - 55.7	2,100	11.2	10.1 - 12.4	6,400	34.9	33.1 - 36.7	18,300

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	51.4	48.5 - 54.4	300	12.9	11.0 - 15.0	800	35.7	32.9 - 38.6	2,300
Honolulu	7,100	54.5	52.0 - 56.9	1,400	10.7	9.2 - 12.3	4,600	34.9	32.5 - 37.3	13,100
Kauai	400	52.4	48.9 - 55.9	100	11.1	9.1 - 13.5	300	36.5	33.2 - 40.0	800
Maui	1,100	53.9	51.2 - 56.6	300	13.0	11.2 - 14.9	700	33.1	30.6 - 35.8	2,000

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	66.3	62.7 - 69.7	300	6.9	5.4 - 9.0	1,200	26.8	23.6 - 30.2	4,400
Native Hawaiian	2,300	41.4	38.3 - 44.7	800	14.6	12.4 - 17.1	2,400	44.0	40.7 - 47.3	5,500
Chinese	500	67.6	58.1 - 75.8	<50	6.2	2.9 - 12.9	200	26.2	18.7 - 35.4	800
Filipino	1,600	50.3	46.0 - 54.6	400	11.7	9.2 - 14.7	1,200	38.0	33.9 - 42.3	3,300
Japanese	1,200	70.4	64.5 - 75.6	100	8.0	5.2 - 12.2	400	21.6	17.0 - 27.0	1,700
Other	1,300	49.2	44.0 - 54.3	400	14.2	11.0 - 18.2	1,000	36.6	31.7 - 41.8	2,600

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	16.6	12.4 - 21.8	100	11.2	8.0 - 15.4	1,000	72.3	66.3 - 77.5	1,300
20-24	1,600	37.6	34.0 - 41.4	500	10.9	8.8 - 13.6	2,200	51.5	47.6 - 55.3	4,400
25-34	5,800	61.8	59.3 - 64.3	900	9.3	7.9 - 10.8	2,700	28.9	26.6 - 31.3	9,400
35+	2,200	68.6	64.3 - 72.5	600	17.5	14.3 - 21.2	400	14.0	11.2 - 17.3	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,300	55.7	53.1 - 58.2	900	9.7	8.3 - 11.3	3,300	34.6	32.1 - 37.1	9,400
Female	4,600	52.0	49.3 - 54.6	1,100	12.8	11.1 - 14.7	3,100	35.2	32.7 - 37.8	8,800

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,400	66.0	63.7 - 68.2	1,100	9.6	8.3 - 11.2	2,700	24.4	22.4 - 26.5	11,200
No	2,500	34.7	32.0 - 37.5	1,000	13.8	11.9 - 15.8	3,600	51.5	48.6 - 54.4	7,100

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,500	38.3	35.4 - 41.3	1,000	16.1	14.0 - 18.4	2,900	45.6	42.6 - 48.7	6,400
131-185%	1,500	51.7	46.9 - 56.5	300	9.6	7.1 - 12.9	1,100	38.7	34.1 - 43.4	2,800
186+%	5,300	69.0	66.3 - 71.6	600	7.2	5.8 - 8.9	1,800	23.8	21.4 - 26.3	7,700

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Income (2004-2011)	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,000	31.5	27.6 - 35.6	400	13.4	10.9 - 16.3	1,800	55.2	50.9 - 59.4	3,300
\$10,000-\$24,999	1,500	43.6	39.6 - 47.7	500	15.2	12.4 - 18.5	1,400	41.2	37.2 - 45.4	3,500
\$25,000-\$49,999	2,400	56.0	52.1 - 59.8	500	11.3	9.1 - 14.1	1,400	32.7	29.2 - 36.4	4,300
>=\$50,000	4,400	71.3	68.3 - 74.2	400	7.2	5.7 - 9.1	1,300	21.4	18.9 - 24.2	6,100

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	54.0	50.9 - 57.1	200	13.8	11.7 - 16.1	400	32.2	29.4 - 35.2	1,300
Normal birth weight	9,100	54.0	52.0 - 55.9	1,900	11.1	9.9 - 12.4	5,900	35.0	33.1 - 36.9	16,900

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	42.5	36.0 - 49.2	300	18.8	14.1 - 24.7	500	38.7	32.4 - 45.4	1,400
No	9,100	54.7	52.8 - 56.6	1,800	10.6	9.5 - 11.8	5,800	34.7	32.8 - 36.6	16,700

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2009-2011

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	45.6	41.7 - 49.6	600	15.2	12.5 - 18.2	1,500	39.2	35.4 - 43.2	3,800
No	8,000	56.0	53.9 - 58.0	1,400	10.2	9.0 - 11.5	4,800	33.9	31.9 - 35.9	14,300

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,200	40.6	37.9 - 43.3	1,100	14.5	12.7 - 16.6	3,500	44.9	42.2 - 47.7	7,800
No	6,500	64.2	61.7 - 66.6	900	8.7	7.4 - 10.2	2,800	27.2	25.0 - 29.5	10,200

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,000	54.5	52.8 - 56.2	1,900	10.5	9.5 - 11.6	6,400	34.9	33.3 - 36.6	18,300

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	51.0	47.7 - 54.3	300	12.6	10.5 - 15.0	900	36.4	33.2 - 39.6	2,400
Honolulu	7,200	55.4	53.2 - 57.6	1,300	9.8	8.6 - 11.3	4,500	34.8	32.7 - 36.9	13,000
Kauai	500	54.8	50.1 - 59.4	100	11.7	9.2 - 14.7	300	33.5	29.2 - 38.0	800
Maui	1,100	53.1	49.8 - 56.4	200	11.9	9.9 - 14.1	700	35.0	31.9 - 38.2	2,100

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,800	65.1	61.7 - 68.3	300	7.4	5.9 - 9.4	1,200	27.5	24.5 - 30.7	4,400
Native Hawaiian	2,400	43.5	40.5 - 46.6	700	12.9	10.9 - 15.1	2,400	43.6	40.5 - 46.7	5,500
Chinese	500	64.7	56.0 - 72.4	100	9.5	5.5 - 15.8	200	25.9	18.9 - 34.3	700
Filipino	1,600	48.9	45.0 - 52.8	400	11.0	8.8 - 13.7	1,300	40.1	36.3 - 44.0	3,300
Japanese	1,300	71.7	66.7 - 76.3	100	6.0	3.9 - 9.1	400	22.3	18.2 - 27.0	1,800
Other	1,400	52.9	48.3 - 57.5	400	13.3	10.6 - 16.6	900	33.8	29.5 - 38.3	2,700

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	19.0	14.9 - 23.9	100	9.3	6.8 - 12.6	1,100	71.7	66.3 - 76.5	1,500
20-24	1,700	41.2	37.7 - 44.8	400	9.7	7.9 - 11.9	2,100	49.1	45.5 - 52.7	4,200
25-34	5,800	61.7	59.4 - 64.0	800	8.7	7.5 - 10.1	2,800	29.5	27.4 - 31.7	9,300
35+	2,200	67.4	63.5 - 71.1	600	17.2	14.3 - 20.6	500	15.4	12.8 - 18.5	3,300

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,400	56.3	53.9 - 58.6	900	9.0	7.8 - 10.4	3,300	34.7	32.5 - 37.0	9,500
Female	4,600	52.6	50.2 - 55.1	1,100	12.1	10.6 - 13.8	3,100	35.2	32.9 - 37.6	8,800

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,400	66.2	64.1 - 68.2	1,000	8.8	7.7 - 10.1	2,800	25.0	23.1 - 26.9	11,200
No	2,500	36.0	33.5 - 38.6	900	13.2	11.5 - 15.1	3,600	50.8	48.1 - 53.5	7,100

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,700	41.8	39.0 - 44.6	1,000	14.9	13.0 - 17.0	2,800	43.3	40.5 - 46.2	6,600
131-185%	1,400	51.3	46.9 - 55.7	300	11.3	8.8 - 14.3	1,000	37.4	33.2 - 41.8	2,800
186+%	5,200	67.8	65.3 - 70.2	500	6.1	5.0 - 7.5	2,000	26.1	23.8 - 28.4	7,600

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Income (2004-2011)	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,100	35.5	31.7 - 39.5	400	13.6	11.2 - 16.5	1,600	50.9	46.8 - 55.0	3,200
\$10,000-\$24,999	1,700	47.4	43.7 - 51.2	500	13.1	10.8 - 15.8	1,500	39.5	35.9 - 43.2	3,700
\$25,000-\$49,999	2,300	54.8	51.2 - 58.4	400	10.6	8.6 - 13.1	1,400	34.5	31.2 - 38.0	4,100
>=\$50,000	4,300	69.4	66.6 - 72.0	400	6.9	5.6 - 8.5	1,500	23.7	21.3 - 26.3	6,100

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	55.6	51.7 - 59.5	200	14.1	11.5 - 17.0	400	30.3	26.9 - 34.0	1,300
Normal birth weight	9,200	54.5	52.7 - 56.3	1,700	10.3	9.2 - 11.4	6,000	35.3	33.5 - 37.0	17,000

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	43.5	37.7 - 49.4	300	16.3	12.3 - 21.3	600	40.2	34.5 - 46.2	1,600
No	9,100	55.4	53.6 - 57.2	1,600	10.0	9.0 - 11.1	5,700	34.6	32.9 - 36.4	16,500

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2008-2010

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	46.3	42.6 - 50.1	500	12.7	10.5 - 15.4	1,500	40.9	37.3 - 44.7	3,800
No	8,100	56.5	54.6 - 58.4	1,400	9.9	8.9 - 11.1	4,800	33.6	31.8 - 35.4	14,300

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,400	44.0	41.4 - 46.5	1,000	13.2	11.6 - 15.1	3,300	42.8	40.2 - 45.4	7,700
No	6,400	62.4	60.2 - 64.6	900	8.5	7.3 - 9.8	3,000	29.1	27.1 - 31.2	10,300

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,900	54.1	52.5 - 55.6	2,100	11.6	10.6 - 12.6	6,300	34.3	32.9 - 35.8	18,300

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	51.8	48.2 - 55.4	300	12.5	10.3 - 15.1	900	35.7	32.3 - 39.3	2,400
Honolulu	6,900	54.1	52.1 - 56.0	1,500	11.7	10.5 - 13.0	4,400	34.3	32.4 - 36.2	12,800
Kauai	500	56.0	50.5 - 61.3	100	10.5	7.7 - 14.2	300	33.5	28.6 - 38.9	900
Maui	1,200	55.9	52.1 - 59.5	200	10.5	8.5 - 13.0	700	33.6	30.2 - 37.3	2,100

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	64.4	61.4 - 67.3	300	7.8	6.3 - 9.6	1,200	27.8	25.1 - 30.7	4,200
Native Hawaiian	2,300	43.4	40.4 - 46.3	800	15.0	12.9 - 17.3	2,200	41.7	38.7 - 44.7	5,300
Chinese	500	66.4	59.3 - 72.8	100	10.8	6.8 - 16.5	200	22.9	17.4 - 29.4	700
Filipino	1,700	50.0	46.6 - 53.5	400	12.2	10.1 - 14.6	1,300	37.8	34.5 - 41.2	3,300
Japanese	1,300	67.8	63.3 - 72.0	100	6.7	4.7 - 9.5	500	25.5	21.6 - 29.8	1,900
Other	1,400	51.2	47.1 - 55.2	400	13.7	11.1 - 16.7	1,000	35.1	31.3 - 39.1	2,800

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	21.8	17.8 - 26.4	200	13.2	10.1 - 17.1	1,100	65.1	59.9 - 69.9	1,700
20-24	1,700	40.9	37.8 - 44.1	500	11.3	9.4 - 13.6	2,000	47.8	44.5 - 51.0	4,300
25-34	5,600	61.0	58.9 - 63.1	900	9.8	8.6 - 11.1	2,700	29.2	27.3 - 31.2	9,200
35+	2,200	68.5	65.1 - 71.8	500	16.4	13.8 - 19.4	500	15.1	12.7 - 17.8	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,400	55.8	53.7 - 57.9	900	9.7	8.5 - 11.0	3,300	34.5	32.5 - 36.6	9,600
Female	4,500	52.2	49.9 - 54.4	1,200	13.7	12.2 - 15.4	3,000	34.1	32.1 - 36.3	8,700

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,300	65.6	63.8 - 67.4	1,000	9.2	8.1 - 10.4	2,800	25.2	23.6 - 26.9	11,200
No	2,500	35.9	33.5 - 38.3	1,100	15.4	13.7 - 17.3	3,500	48.7	46.2 - 51.3	7,100

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,700	40.5	38.0 - 43.0	1,200	17.4	15.5 - 19.5	2,800	42.1	39.5 - 44.7	6,700
131-185%	1,400	51.7	47.6 - 55.7	300	12.9	10.4 - 15.8	900	35.4	31.7 - 39.4	2,700
186+%	5,100	67.9	65.7 - 70.0	400	5.9	4.8 - 7.1	2,000	26.3	24.3 - 28.4	7,500

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Income (2004-2011)	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,100	36.6	32.9 - 40.4	500	17.1	14.3 - 20.2	1,400	46.3	42.4 - 50.3	3,100
\$10,000-\$24,999	1,700	44.4	41.1 - 47.7	600	14.5	12.3 - 17.1	1,600	41.1	37.8 - 44.4	3,900
\$25,000-\$49,999	2,200	54.9	51.6 - 58.1	500	12.1	10.1 - 14.4	1,300	33.1	30.1 - 36.2	4,000
>=\$50,000	4,200	69.3	66.9 - 71.7	400	6.5	5.3 - 7.9	1,500	24.2	22.0 - 26.5	6,100

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	53.4	48.9 - 57.9	200	16.4	13.2 - 20.1	400	30.2	26.2 - 34.5	1,400
Normal birth weight	9,200	54.2	52.5 - 55.8	1,900	11.2	10.2 - 12.3	5,800	34.6	33.1 - 36.2	16,900

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	40.3	35.1 - 45.7	300	20.2	16.2 - 25.0	600	39.5	34.3 - 44.9	1,600
No	9,000	55.2	53.6 - 56.8	1,800	10.8	9.8 - 11.9	5,600	34.0	32.5 - 35.6	16,400

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2007-2009

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	45.0	41.5 - 48.5	500	14.7	12.4 - 17.4	1,500	40.3	36.9 - 43.8	3,700
No	8,000	56.1	54.4 - 57.8	1,600	10.9	9.8 - 12.0	4,700	33.0	31.4 - 34.7	14,300

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,300	43.3	41.0 - 45.7	1,200	15.4	13.7 - 17.2	3,100	41.3	38.9 - 43.7	7,600
No	6,400	61.9	59.9 - 63.8	900	9.0	7.9 - 10.2	3,000	29.2	27.3 - 31.1	10,300

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	10,000	54.6	53.2 - 55.9	2,200	12.3	11.4 - 13.3	6,000	33.1	31.8 - 34.5	18,200

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	49.7	45.7 - 53.7	300	12.0	9.5 - 14.9	900	38.3	34.5 - 42.3	2,300
Honolulu	7,100	55.0	53.4 - 56.6	1,700	12.9	11.8 - 14.0	4,200	32.1	30.6 - 33.7	12,900
Kauai	500	54.3	47.9 - 60.5	100	11.7	8.2 - 16.4	300	34.0	28.3 - 40.3	900
Maui	1,200	57.4	53.2 - 61.5	200	9.5	7.3 - 12.3	700	33.1	29.2 - 37.2	2,100

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	63.4	60.5 - 66.1	400	9.0	7.5 - 10.8	1,200	27.6	25.1 - 30.3	4,300
Native Hawaiian	2,300	46.0	43.1 - 48.8	700	14.9	13.0 - 17.1	2,000	39.1	36.4 - 42.0	5,000
Chinese	400	67.1	63.3 - 70.6	100	9.1	7.0 - 11.6	200	23.9	20.7 - 27.4	600
Filipino	1,700	51.4	48.4 - 54.4	400	12.9	11.0 - 15.0	1,200	35.7	32.9 - 38.7	3,400
Japanese	1,400	68.3	64.3 - 72.0	100	6.8	4.9 - 9.4	500	24.9	21.6 - 28.6	2,100
Other	1,400	47.6	44.0 - 51.2	500	16.7	14.1 - 19.6	1,000	35.8	32.4 - 39.3	2,900

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	27.0	22.6 - 31.8	200	14.1	10.9 - 18.2	900	58.9	53.7 - 63.9	1,600
20-24	1,800	42.5	39.6 - 45.5	500	11.8	10.0 - 13.9	2,000	45.6	42.7 - 48.6	4,300
25-34	5,300	58.9	57.0 - 60.8	1,100	12.1	10.8 - 13.5	2,600	29.0	27.3 - 30.8	9,100
35+	2,300	71.7	68.8 - 74.4	400	12.6	10.7 - 14.8	500	15.7	13.6 - 18.2	3,300

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,200	55.6	53.7 - 57.5	1,000	11.0	9.8 - 12.3	3,200	33.4	31.6 - 35.3	9,400
Female	4,700	53.4	51.4 - 55.4	1,200	13.7	12.4 - 15.2	2,900	32.8	31.0 - 34.8	8,800

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,500	65.2	63.6 - 66.8	1,100	9.7	8.7 - 10.8	2,900	25.1	23.6 - 26.6	11,400
No	2,500	36.7	34.4 - 39.0	1,100	16.6	14.9 - 18.5	3,200	46.7	44.3 - 49.1	6,800

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,800	42.8	40.4 - 45.2	1,100	16.4	14.7 - 18.3	2,700	40.7	38.4 - 43.1	6,600
131-185%	1,200	47.3	43.6 - 51.0	400	16.6	14.0 - 19.5	900	36.2	32.7 - 39.8	2,600
186+%	5,300	68.2	66.2 - 70.1	500	6.5	5.5 - 7.6	2,000	25.3	23.6 - 27.2	7,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Income (2004-2011)	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,200	39.3	35.9 - 42.9	500	17.9	15.2 - 20.8	1,300	42.8	39.3 - 46.4	3,100
\$10,000-\$24,999	1,700	44.6	41.6 - 47.7	500	14.1	12.0 - 16.4	1,600	41.3	38.3 - 44.4	3,900
\$25,000-\$49,999	2,300	54.5	51.6 - 57.3	500	12.2	10.4 - 14.2	1,400	33.4	30.7 - 36.1	4,200
>=\$50,000	4,200	69.9	67.7 - 72.0	400	7.4	6.2 - 8.7	1,400	22.7	20.8 - 24.7	6,000

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	50.9	46.0 - 55.9	200	15.3	11.9 - 19.4	500	33.8	29.2 - 38.7	1,400
Normal birth weight	9,200	54.8	53.4 - 56.2	2,000	12.1	11.1 - 13.1	5,600	33.1	31.8 - 34.5	16,800

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	39.2	34.4 - 44.2	300	21.7	17.8 - 26.2	600	39.1	34.3 - 44.1	1,600
No	9,200	56.0	54.6 - 57.5	1,900	11.5	10.6 - 12.5	5,300	32.5	31.1 - 33.9	16,400

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2006-2008

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	43.6	40.4 - 46.8	600	16.5	14.2 - 19.1	1,400	39.9	36.8 - 43.1	3,600
No	8,200	57.3	55.8 - 58.8	1,600	11.4	10.4 - 12.4	4,500	31.3	29.9 - 32.8	14,300

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,400	43.8	41.6 - 46.0	1,200	15.5	14.0 - 17.2	3,100	40.7	38.5 - 42.9	7,700
No	6,400	62.7	60.9 - 64.4	1,000	10.1	9.0 - 11.2	2,800	27.2	25.7 - 28.9	10,300

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,600	53.6	52.3 - 55.0	2,500	14.0	13.1 - 15.0	5,800	32.3	31.0 - 33.7	17,800

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	47.2	43.2 - 51.2	300	14.3	11.7 - 17.5	900	38.4	34.6 - 42.4	2,300
Honolulu	7,000	54.5	52.9 - 56.1	1,800	14.3	13.1 - 15.5	4,000	31.2	29.7 - 32.8	12,800
Kauai	400	48.7	42.3 - 55.2	100	16.8	12.4 - 22.3	300	34.5	28.6 - 41.0	900
Maui	1,100	57.8	53.5 - 62.1	200	10.7	8.3 - 13.7	600	31.5	27.5 - 35.7	1,900

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,500	61.8	58.9 - 64.6	400	10.4	8.7 - 12.3	1,100	27.8	25.3 - 30.5	4,000
Native Hawaiian	2,200	44.2	41.4 - 47.1	900	18.4	16.2 - 20.8	1,800	37.4	34.6 - 40.2	4,900
Chinese	400	69.0	65.3 - 72.4	100	8.5	6.6 - 11.0	100	22.5	19.4 - 25.9	600
Filipino	1,700	52.6	49.6 - 55.6	500	13.9	11.9 - 16.1	1,100	33.5	30.7 - 36.4	3,300
Japanese	1,400	66.0	62.1 - 69.6	200	7.1	5.3 - 9.6	600	26.9	23.5 - 30.5	2,200
Other	1,300	46.5	42.9 - 50.1	500	18.4	15.7 - 21.5	1,000	35.1	31.7 - 38.7	2,700

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	26.9	22.7 - 31.7	300	16.1	12.7 - 20.3	900	56.9	51.8 - 61.9	1,600
20-24	1,800	41.9	39.1 - 44.8	600	13.6	11.7 - 15.8	2,000	44.4	41.6 - 47.3	4,400
25-34	5,100	57.9	56.0 - 59.8	1,300	14.3	12.9 - 15.8	2,400	27.8	26.1 - 29.6	8,700
35+	2,200	71.7	68.8 - 74.4	400	12.7	10.8 - 15.0	500	15.6	13.4 - 18.0	3,100

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,900	53.8	51.8 - 55.7	1,200	12.8	11.5 - 14.2	3,100	33.4	31.6 - 35.3	9,100
Female	4,700	53.5	51.5 - 55.5	1,300	15.3	13.9 - 16.9	2,700	31.2	29.4 - 33.1	8,700

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,200	64.3	62.7 - 66.0	1,200	10.5	9.5 - 11.6	2,800	25.2	23.7 - 26.7	11,200
No	2,300	35.5	33.2 - 37.8	1,300	20.1	18.2 - 22.1	2,900	44.5	42.1 - 46.9	6,600

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,600	41.0	38.7 - 43.4	1,200	18.8	17.0 - 20.8	2,600	40.1	37.8 - 42.5	6,400
131-185%	1,100	47.3	43.5 - 51.2	400	18.1	15.3 - 21.3	800	34.6	30.9 - 38.4	2,300
186+%	5,300	67.7	65.8 - 69.6	600	7.3	6.3 - 8.5	1,900	25.0	23.2 - 26.8	7,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Income (2004-2011)	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,100	36.9	33.6 - 40.4	600	19.3	16.6 - 22.3	1,300	43.8	40.3 - 47.4	3,000
\$10,000-\$24,999	1,600	43.9	40.9 - 47.0	600	16.9	14.7 - 19.3	1,500	39.2	36.2 - 42.3	3,700
\$25,000-\$49,999	2,200	53.3	50.4 - 56.2	600	14.2	12.2 - 16.3	1,300	32.5	29.9 - 35.3	4,100
>=\$50,000	4,000	70.7	68.5 - 72.8	400	7.6	6.4 - 9.0	1,200	21.7	19.8 - 23.7	5,700

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	48.9	44.0 - 53.8	200	15.8	12.4 - 19.9	500	35.4	30.8 - 40.2	1,400
Normal birth weight	8,800	54.0	52.6 - 55.5	2,300	13.9	12.9 - 15.0	5,300	32.1	30.7 - 33.5	16,400

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	34.4	29.8 - 39.3	400	26.4	22.2 - 31.1	600	39.2	34.5 - 44.2	1,500
No	8,900	55.6	54.2 - 57.1	2,100	12.9	12.0 - 14.0	5,000	31.4	30.1 - 32.8	16,000

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2005-2007

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	40.7	37.6 - 43.8	700	20.0	17.5 - 22.7	1,400	39.3	36.3 - 42.5	3,700
No	7,900	57.2	55.7 - 58.8	1,700	12.5	11.5 - 13.6	4,200	30.3	28.8 - 31.7	13,900

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,200	42.1	40.0 - 44.3	1,400	18.0	16.3 - 19.8	3,000	39.9	37.7 - 42.1	7,500
No	6,200	62.2	60.4 - 63.9	1,100	11.2	10.1 - 12.5	2,700	26.6	25.0 - 28.2	10,000

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,600	54.7	53.3 - 56.0	2,500	14.2	13.3 - 15.2	5,500	31.1	29.9 - 32.4	17,600

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	47.9	44.0 - 51.9	400	17.1	14.3 - 20.4	800	34.9	31.2 - 38.8	2,200
Honolulu	7,200	55.7	54.1 - 57.3	1,800	13.9	12.9 - 15.1	3,900	30.3	28.9 - 31.8	12,800
Kauai	400	50.8	44.3 - 57.3	100	16.6	12.1 - 22.2	300	32.6	26.7 - 39.1	800
Maui	1,000	56.9	52.6 - 61.1	200	11.7	9.2 - 14.7	600	31.4	27.5 - 35.6	1,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,500	63.0	60.2 - 65.7	400	9.9	8.3 - 11.7	1,100	27.1	24.6 - 29.7	4,000
Native Hawaiian	2,200	45.7	42.9 - 48.5	900	17.7	15.6 - 20.0	1,800	36.6	34.0 - 39.4	4,900
Chinese	400	68.9	65.6 - 72.0	100	9.0	7.2 - 11.2	100	22.1	19.3 - 25.1	600
Filipino	1,700	53.4	50.4 - 56.3	500	14.2	12.3 - 16.4	1,100	32.4	29.8 - 35.2	3,300
Japanese	1,500	66.1	62.4 - 69.6	200	8.5	6.6 - 10.9	600	25.4	22.2 - 28.8	2,200
Other	1,200	47.2	43.7 - 50.8	500	20.5	17.7 - 23.6	800	32.3	29.1 - 35.8	2,600

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	27.9	23.6 - 32.6	200	16.0	12.7 - 20.0	800	56.1	51.1 - 61.0	1,400
20-24	1,800	42.0	39.3 - 44.8	600	14.9	12.9 - 17.1	1,900	43.1	40.3 - 45.9	4,300
25-34	5,200	60.2	58.3 - 62.1	1,200	13.6	12.3 - 15.0	2,300	26.2	24.5 - 27.9	8,600
35+	2,200	68.9	66.1 - 71.7	500	14.3	12.3 - 16.5	500	16.8	14.6 - 19.2	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,000	53.9	52.1 - 55.8	1,300	13.8	12.6 - 15.2	3,000	32.2	30.5 - 34.0	9,200
Female	4,700	55.4	53.5 - 57.4	1,200	14.7	13.3 - 16.1	2,500	29.9	28.1 - 31.7	8,400

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,400	64.4	62.8 - 65.9	1,300	10.8	9.8 - 11.9	2,900	24.8	23.4 - 26.3	11,500
No	2,200	36.1	33.8 - 38.5	1,200	20.7	18.8 - 22.8	2,600	43.2	40.8 - 45.6	6,000

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,600	41.0	38.7 - 43.3	1,200	19.0	17.2 - 20.9	2,500	40.0	37.7 - 42.3	6,300
131-185%	1,000	48.9	44.9 - 52.8	400	18.5	15.6 - 21.9	700	32.6	29.0 - 36.5	2,100
186+%	5,400	69.0	67.1 - 70.8	600	7.5	6.5 - 8.6	1,800	23.6	21.9 - 25.3	7,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Income (2004-2011)	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,100	37.0	33.7 - 40.5	500	18.4	15.8 - 21.3	1,300	44.6	41.1 - 48.1	2,900
\$10,000-\$24,999	1,700	44.7	41.8 - 47.6	700	17.9	15.8 - 20.3	1,400	37.4	34.6 - 40.3	3,900
\$25,000-\$49,999	2,300	55.1	52.3 - 57.8	600	13.1	11.3 - 15.1	1,300	31.8	29.3 - 34.5	4,200
>=\$50,000	3,900	73.1	70.9 - 75.2	400	7.9	6.7 - 9.4	1,000	19.0	17.2 - 20.9	5,400

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	53.5	48.5 - 58.5	200	13.6	10.3 - 17.6	400	32.9	28.3 - 37.8	1,300
Normal birth weight	8,900	54.7	53.3 - 56.1	2,300	14.3	13.3 - 15.3	5,000	31.0	29.7 - 32.3	16,300

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	35.6	31.1 - 40.4	400	25.0	20.9 - 29.6	600	39.4	34.7 - 44.2	1,500
No	8,900	56.6	55.2 - 58.0	2,100	13.2	12.2 - 14.2	4,800	30.3	29.0 - 31.6	15,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2004-2006

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	41.8	38.8 - 44.8	700	19.8	17.4 - 22.4	1,400	38.5	35.5 - 41.5	3,700
No	7,900	58.3	56.8 - 59.8	1,700	12.6	11.6 - 13.7	4,000	29.1	27.7 - 30.5	13,600

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,300	44.0	41.9 - 46.1	1,300	17.5	16.0 - 19.2	2,900	38.5	36.4 - 40.6	7,600
No	6,100	63.0	61.2 - 64.7	1,100	11.6	10.5 - 12.8	2,500	25.4	23.9 - 27.0	9,700

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,700	55.4	54.0 - 56.8	2,100	12.1	11.2 - 13.0	5,700	32.5	31.2 - 33.8	17,400

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	49.6	46.1 - 53.1	300	16.0	13.5 - 19.0	700	34.4	31.1 - 37.8	2,100
Honolulu	7,200	56.4	54.7 - 58.1	1,500	11.6	10.6 - 12.8	4,100	31.9	30.3 - 33.5	12,700
Kauai	400	55.5	49.8 - 61.0	100	12.4	8.9 - 17.1	300	32.1	27.0 - 37.6	800
Maui	1,000	54.7	51.0 - 58.4	200	10.6	8.4 - 13.2	600	34.7	31.2 - 38.4	1,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,500	64.9	61.9 - 67.8	300	8.1	6.6 - 9.9	1,000	27.0	24.3 - 29.8	3,800
Native Hawaiian	2,200	45.4	43.0 - 47.9	800	15.7	13.9 - 17.7	1,900	38.9	36.5 - 41.3	4,800
Chinese	500	69.2	63.4 - 74.5	<50	6.9	5.2 - 9.0	200	23.9	18.9 - 29.8	700
Filipino	1,800	53.4	50.2 - 56.6	400	12.2	10.3 - 14.4	1,200	34.4	31.4 - 37.5	3,400
Japanese	1,400	66.7	62.7 - 70.5	100	6.8	5.1 - 9.0	600	26.4	22.9 - 30.3	2,200
Other	1,200	49.2	45.2 - 53.2	400	17.2	14.4 - 20.4	800	33.6	29.9 - 37.5	2,500

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	400	25.4	21.4 - 29.8	200	15.0	11.9 - 18.9	900	59.6	54.8 - 64.2	1,500
20-24	1,800	41.6	38.9 - 44.4	600	13.1	11.3 - 15.2	1,900	45.3	42.5 - 48.1	4,300
25-34	5,400	62.4	60.5 - 64.4	900	10.5	9.4 - 11.8	2,300	27.0	25.3 - 28.8	8,600
35+	2,100	68.8	65.7 - 71.8	400	13.7	11.6 - 16.1	500	17.4	15.1 - 20.1	3,100

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,000	54.5	52.6 - 56.4	1,100	12.1	10.9 - 13.3	3,100	33.4	31.7 - 35.3	9,200
Female	4,600	56.4	54.3 - 58.4	1,000	12.2	10.9 - 13.6	2,600	31.5	29.6 - 33.4	8,200

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,500	65.0	63.3 - 66.6	1,000	8.9	8.0 - 9.9	3,000	26.1	24.6 - 27.7	11,600
No	2,100	36.3	34.1 - 38.7	1,100	18.5	16.7 - 20.5	2,600	45.2	42.8 - 47.5	5,800

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,900	42.5	40.3 - 44.8	1,100	16.2	14.6 - 17.9	2,800	41.3	39.1 - 43.6	6,800
131-185%	1,200	55.4	51.3 - 59.4	300	13.7	11.2 - 16.8	700	30.9	27.2 - 34.8	2,100
186+%	4,900	70.0	67.9 - 71.9	400	6.2	5.3 - 7.3	1,700	23.8	22.0 - 25.8	6,900

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	55.4	50.0 - 60.7	100	11.1	8.2 - 14.8	400	33.5	28.5 - 38.9	1,200
Normal birth weight	9,000	55.4	54.0 - 56.8	2,000	12.2	11.3 - 13.1	5,300	32.4	31.1 - 33.8	16,200

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	38.3	33.7 - 43.1	300	21.9	18.1 - 26.3	600	39.8	35.1 - 44.6	1,500
No	8,900	57.1	55.6 - 58.5	1,700	11.1	10.2 - 12.0	5,000	31.9	30.5 - 33.3	15,600

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	42.6	39.6 - 45.7	600	16.4	14.2 - 18.8	1,500	41.0	38.0 - 44.1	3,600
No	7,900	58.9	57.3 - 60.5	1,500	10.8	9.9 - 11.8	4,100	30.3	28.8 - 31.8	13,500

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2003-2005

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,400	45.2	43.1 - 47.3	1,100	14.5	13.1 - 16.1	3,000	40.3	38.2 - 42.4	7,500
No	6,100	63.3	61.5 - 65.1	1,000	10.1	9.0 - 11.3	2,600	26.6	24.9 - 28.3	9,700

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,800	56.3	54.9 - 57.8	1,700	9.8	9.0 - 10.6	5,900	33.9	32.5 - 35.3	17,400

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	53.0	49.9 - 56.1	300	12.1	10.1 - 14.5	700	34.9	32.0 - 37.8	2,100
Honolulu	7,200	56.9	55.1 - 58.8	1,200	9.4	8.4 - 10.5	4,300	33.6	31.9 - 35.4	12,700
Kauai	500	60.5	55.6 - 65.2	100	8.3	6.0 - 11.5	200	31.2	26.8 - 35.9	800
Maui	1,000	54.1	50.8 - 57.3	200	10.3	8.5 - 12.5	600	35.6	32.6 - 38.8	1,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,400	62.9	59.6 - 66.1	300	7.5	5.9 - 9.4	1,100	29.6	26.6 - 32.8	3,800
Native Hawaiian	2,200	46.3	44.2 - 48.4	600	12.2	10.8 - 13.7	2,000	41.5	39.5 - 43.6	4,700
Chinese	400	66.3	58.5 - 73.3	<50	5.0	3.1 - 7.9	200	28.8	22.0 - 36.7	700
Filipino	2,000	55.6	52.1 - 59.0	400	10.0	8.2 - 12.2	1,200	34.4	31.1 - 37.8	3,500
Japanese	1,500	69.4	65.1 - 73.4	200	6.8	5.0 - 9.1	500	23.9	20.1 - 28.0	2,200
Other	1,300	52.2	47.7 - 56.7	300	12.3	9.7 - 15.4	900	35.5	31.2 - 40.0	2,500

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	22.4	18.7 - 26.7	200	11.7	9.0 - 15.1	1,000	65.9	61.2 - 70.3	1,500
20-24	1,800	42.7	39.8 - 45.6	500	11.0	9.3 - 12.9	2,000	46.4	43.5 - 49.3	4,300
25-34	5,500	63.7	61.6 - 65.7	700	7.9	6.9 - 9.1	2,400	28.4	26.5 - 30.4	8,600
35+	2,100	71.1	67.8 - 74.2	400	12.5	10.5 - 15.0	500	16.4	13.9 - 19.1	3,000

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,100	55.8	53.8 - 57.7	900	10.2	9.2 - 11.4	3,100	34.0	32.1 - 35.9	9,100
Female	4,700	57.0	54.8 - 59.1	800	9.3	8.1 - 10.6	2,800	33.8	31.7 - 35.8	8,300

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,800	65.8	64.0 - 67.5	900	7.7	6.8 - 8.7	3,100	26.5	24.9 - 28.2	11,800
No	2,000	36.2	33.9 - 38.6	800	14.2	12.6 - 16.0	2,800	49.5	47.1 - 52.0	5,600

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,100	44.3	42.0 - 46.6	900	13.3	11.8 - 14.8	2,900	42.4	40.2 - 44.7	6,900
131-185%	1,500	55.7	51.7 - 59.6	300	9.9	7.9 - 12.3	900	34.4	30.7 - 38.3	2,700
186+%	4,500	72.5	70.2 - 74.6	300	5.1	4.2 - 6.3	1,400	22.4	20.4 - 24.6	6,200

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	58.6	52.8 - 64.2	100	10.3	7.5 - 14.1	400	31.0	25.9 - 36.7	1,200
Normal birth weight	9,100	56.1	54.6 - 57.6	1,600	9.7	8.9 - 10.6	5,500	34.2	32.7 - 35.6	16,200

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	41.9	37.2 - 46.9	300	18.8	15.3 - 22.9	600	39.2	34.6 - 44.1	1,500
No	9,000	57.7	56.2 - 59.3	1,400	8.7	7.9 - 9.6	5,200	33.5	32.0 - 35.0	15,600

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	43.0	39.8 - 46.2	400	12.8	10.9 - 15.1	1,500	44.2	41.0 - 47.4	3,500
No	8,100	60.0	58.3 - 61.6	1,200	8.8	7.9 - 9.7	4,200	31.2	29.7 - 32.8	13,500

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2002-2004

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,400	45.5	43.3 - 47.7	900	11.6	10.3 - 13.1	3,200	42.9	40.7 - 45.1	7,400
No	6,300	64.7	62.7 - 66.6	800	8.2	7.2 - 9.3	2,600	27.1	25.4 - 29.0	9,700

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,500	55.8	54.4 - 57.3	1,400	8.4	7.6 - 9.2	6,100	35.8	34.3 - 37.2	17,100

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	51.9	49.3 - 54.5	200	8.9	7.5 - 10.6	800	39.1	36.6 - 41.7	1,900
Honolulu	7,100	56.6	54.7 - 58.5	1,000	8.3	7.3 - 9.4	4,400	35.2	33.3 - 37.0	12,600
Kauai	400	57.2	52.9 - 61.5	100	7.0	5.1 - 9.6	300	35.8	31.7 - 40.0	800
Maui	1,000	54.4	51.5 - 57.2	200	9.3	7.7 - 11.1	600	36.3	33.6 - 39.1	1,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,300	63.9	60.5 - 67.1	300	7.2	5.7 - 9.2	1,100	28.9	25.8 - 32.2	3,600
Native Hawaiian	2,100	45.2	43.6 - 46.8	500	10.3	9.3 - 11.4	2,100	44.5	42.9 - 46.1	4,600
Chinese	400	65.7	56.9 - 73.6	<50	3.9	1.6 - 8.8	200	30.4	22.9 - 39.1	700
Filipino	1,900	54.0	50.4 - 57.5	300	8.6	6.8 - 10.9	1,400	37.4	34.0 - 40.9	3,600
Japanese	1,500	70.7	66.1 - 74.9	100	5.3	3.6 - 7.7	500	24.0	20.1 - 28.4	2,100
Other	1,200	50.9	46.1 - 55.6	200	10.2	7.7 - 13.4	900	38.9	34.3 - 43.7	2,400

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	20.3	17.0 - 23.9	100	7.8	5.8 - 10.4	1,100	72.0	67.9 - 75.7	1,600
20-24	1,800	42.8	40.0 - 45.7	400	8.8	7.3 - 10.5	2,100	48.4	45.6 - 51.3	4,200
25-34	5,400	63.4	61.3 - 65.5	600	7.1	6.1 - 8.3	2,500	29.5	27.5 - 31.5	8,500
35+	2,000	72.7	69.2 - 75.9	300	12.0	9.8 - 14.6	400	15.3	12.7 - 18.2	2,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,900	56.0	53.9 - 58.0	700	8.3	7.3 - 9.5	3,100	35.7	33.7 - 37.7	8,700
Female	4,600	55.7	53.5 - 57.8	700	8.5	7.4 - 9.7	3,000	35.8	33.8 - 37.9	8,300

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,600	66.5	64.7 - 68.3	700	6.5	5.6 - 7.4	3,100	27.0	25.4 - 28.8	11,400
No	2,000	34.4	32.2 - 36.7	700	12.3	10.8 - 14.0	3,000	53.3	50.9 - 55.6	5,700

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,000	44.0	41.7 - 46.2	800	11.1	9.8 - 12.5	3,100	45.0	42.8 - 47.2	6,900
131-185%	1,800	57.1	53.6 - 60.5	300	8.8	7.0 - 10.9	1,100	34.2	30.9 - 37.6	3,200
186+%	4,000	73.3	70.8 - 75.7	200	4.3	3.3 - 5.5	1,200	22.4	20.2 - 24.8	5,500

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Income (2000-2003)	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,400	38.1	35.2 - 41.1	500	13.7	11.8 - 15.9	1,800	48.2	45.2 - 51.2	3,700
\$15,001-\$25,000	1,100	48.6	44.7 - 52.6	200	8.4	6.5 - 10.8	1,000	43.0	39.1 - 46.9	2,300
\$25,001-\$40,000	2,000	56.4	53.0 - 59.7	200	7.2	5.7 - 9.0	1,300	36.5	33.3 - 39.8	3,500
>=\$40,001	4,700	71.5	69.2 - 73.7	400	5.6	4.6 - 6.9	1,500	22.9	20.9 - 25.1	6,500

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	55.4	49.5 - 61.1	100	9.8	7.1 - 13.3	400	34.9	29.5 - 40.6	1,200
Normal birth weight	8,900	55.8	54.3 - 57.4	1,300	8.3	7.5 - 9.2	5,700	35.9	34.4 - 37.3	15,900

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	41.6	37.1 - 46.2	200	15.7	12.6 - 19.3	600	42.8	38.3 - 47.4	1,500
No	8,700	57.4	55.8 - 58.9	1,200	7.6	6.8 - 8.4	5,300	35.1	33.6 - 36.6	15,200

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2001-2003

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	42.4	39.3 - 45.5	400	10.3	8.6 - 12.3	1,600	47.3	44.2 - 50.5	3,400
No	7,900	59.7	58.0 - 61.4	1,000	7.8	6.9 - 8.7	4,300	32.5	30.9 - 34.1	13,200

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,100	43.6	41.4 - 45.8	700	9.6	8.4 - 11.0	3,400	46.8	44.6 - 49.0	7,200
No	6,300	64.8	62.8 - 66.7	700	7.5	6.5 - 8.6	2,700	27.8	26.0 - 29.6	9,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

State	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,500	56.0	54.6 - 57.4	1,400	8.5	7.8 - 9.3	6,000	35.5	34.2 - 36.9	16,900

County	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	51.4	48.8 - 54.1	200	9.3	7.8 - 11.0	800	39.2	36.7 - 41.8	1,900
Honolulu	7,100	56.6	54.8 - 58.3	1,000	8.3	7.4 - 9.3	4,400	35.1	33.5 - 36.9	12,500
Kauai	400	57.3	52.9 - 61.7	100	8.6	6.3 - 11.5	200	34.1	30.0 - 38.4	700
Maui	1,000	56.0	53.1 - 58.9	200	9.0	7.5 - 10.9	600	35.0	32.3 - 37.8	1,700

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

DOH Race-Ethnicity	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,400	64.5	61.5 - 67.5	300	7.1	5.7 - 8.9	1,100	28.3	25.6 - 31.3	3,700
Native Hawaiian	2,100	45.6	44.0 - 47.1	500	10.1	9.2 - 11.1	2,000	44.3	42.8 - 45.8	4,600
Chinese	400	66.8	58.4 - 74.3	<50	7.3	4.0 - 13.1	200	25.8	19.1 - 33.9	700
Filipino	1,800	52.3	48.8 - 55.7	300	8.5	6.8 - 10.6	1,400	39.2	35.9 - 42.6	3,500
Japanese	1,500	69.2	65.0 - 73.2	100	5.2	3.6 - 7.6	600	25.5	21.8 - 29.6	2,200
Other	1,200	52.8	48.3 - 57.3	200	10.7	8.3 - 13.7	800	36.5	32.2 - 41.0	2,300

PRAMS Age Group	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	19.0	16.3 - 22.1	100	8.0	6.1 - 10.3	1,200	73.0	69.5 - 76.3	1,700
20-24	1,900	44.6	42.0 - 47.3	300	8.0	6.7 - 9.6	2,000	47.4	44.7 - 50.0	4,200
25-34	5,200	63.7	61.7 - 65.6	600	7.3	6.3 - 8.4	2,400	29.0	27.2 - 31.0	8,200
35+	2,000	72.5	69.3 - 75.6	400	13.0	10.9 - 15.5	400	14.4	12.1 - 17.1	2,800

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Infant Gender	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,700	55.2	53.3 - 57.1	700	8.4	7.4 - 9.5	3,100	36.4	34.5 - 38.2	8,600
Female	4,700	56.8	54.8 - 58.7	700	8.5	7.5 - 9.7	2,900	34.7	32.8 - 36.6	8,300

Married	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,700	66.8	65.1 - 68.5	800	6.8	6.0 - 7.7	3,000	26.3	24.8 - 27.9	11,500
No	1,800	33.1	31.0 - 35.2	600	11.9	10.5 - 13.5	3,000	55.0	52.7 - 57.2	5,400

Poverty Level	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	2,900	43.0	40.9 - 45.1	800	11.3	10.1 - 12.7	3,100	45.7	43.6 - 47.8	6,700
131-185%	1,800	57.0	53.7 - 60.3	300	8.1	6.5 - 10.1	1,100	34.8	31.7 - 38.1	3,100
186+%	4,200	73.3	71.1 - 75.5	300	4.9	4.0 - 6.1	1,200	21.7	19.8 - 23.9	5,700

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Income (2000-2003)	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	1,400	38.6	35.9 - 41.4	500	13.2	11.4 - 15.1	1,700	48.3	45.5 - 51.0	3,600
\$15,001-\$25,000	1,200	47.1	43.5 - 50.7	200	9.1	7.3 - 11.4	1,100	43.8	40.2 - 47.4	2,500
\$25,001-\$40,000	2,000	57.1	54.0 - 60.2	300	7.4	6.0 - 9.1	1,200	35.5	32.6 - 38.5	3,500
>=\$40,001	4,600	71.3	69.1 - 73.4	400	5.9	4.9 - 7.1	1,500	22.8	20.9 - 24.8	6,400

Birthweight	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	52.3	46.7 - 57.7	100	11.6	8.6 - 15.5	400	36.2	31.1 - 41.5	1,100
Normal birth weight	8,800	56.2	54.8 - 57.6	1,300	8.3	7.6 - 9.1	5,600	35.5	34.1 - 36.9	15,700

Smoke 3rd Trimester	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	41.5	37.3 - 45.9	200	13.8	11.3 - 16.9	600	44.6	40.3 - 49.0	1,400
No	8,700	57.7	56.2 - 59.1	1,200	7.8	7.1 - 8.6	5,200	34.5	33.1 - 36.0	15,100

**PRAMS Health Indicator 3-Year Aggregate Report - Pregnancy intention- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011**

Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years: 2000-2002

Smoke Before Pregnant	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	42.1	39.2 - 45.1	300	10.3	8.7 - 12.0	1,600	47.6	44.7 - 50.6	3,300
No	7,800	60.2	58.6 - 61.7	1,000	7.9	7.1 - 8.8	4,200	31.9	30.5 - 33.4	13,000

WIC During Pregnancy	Intended			Unwanted			Mis-timed			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,200	44.8	42.8 - 46.9	700	9.5	8.4 - 10.7	3,300	45.7	43.7 - 47.8	7,200
No	6,200	64.5	62.7 - 66.3	700	7.6	6.7 - 8.7	2,700	27.9	26.2 - 29.6	9,600