

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- for the State of Hawaii, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Discuss seat belt use with HCW during PNC	During any of your prenatal care visits, did a doctor, nurse, or other health care worker talk with you about using a seat belt during pregnancy?

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/22/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- for the State of Hawaii,
for the Year(s)**

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013


Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Years	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	8,300	49.7	48.6 - 50.8	8,400	50.3	49.2 - 51.4	16,700
2001-2005	8,500	50.4	49.3 - 51.5	8,300	49.6	48.5 - 50.7	16,800
2002-2006	8,700	50.9	49.7 - 52.0	8,400	49.1	48.0 - 50.3	17,100
2003-2007	8,800	51.0	49.9 - 52.1	8,500	49.0	47.9 - 50.1	17,300
2004-2008	8,800	49.7	48.7 - 50.8	8,800	50.3	49.2 - 51.3	17,600
2005-2009	8,800	49.6	48.4 - 50.7	9,000	50.4	49.3 - 51.6	17,700
2006-2010	8,900	49.2	48.0 - 50.5	9,100	50.8	49.5 - 52.0	18,000
2007-2011	9,000	49.5	48.2 - 50.8	9,100	50.5	49.2 - 51.8	18,100
2008-2012	9,000	49.7	48.3 - 51.1	9,100	50.3	48.9 - 51.7	18,100
2009-2013	9,000	49.5	48.1 - 51.0	9,100	50.5	49.0 - 51.9	18,100

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- for the State of Hawaii,
for the Year(s)**

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,000	49.5	48.1 - 51.0	9,100	50.5	49.0 - 51.9	18,100

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	51.8	49.5 - 54.2	1,100	48.2	45.8 - 50.5	2,300
Honolulu	6,400	49.4	47.4 - 51.4	6,600	50.6	48.6 - 52.6	13,000
Kauai	400	52.8	50.2 - 55.5	400	47.2	44.5 - 49.8	800
Maui	900	46.2	44.0 - 48.5	1,100	53.8	51.5 - 56.0	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	44.1	41.2 - 47.0	2,500	55.9	53.0 - 58.8	4,400
Native Hawaiian	3,000	54.8	52.1 - 57.3	2,400	45.2	42.7 - 47.9	5,400
Chinese	300	35.5	28.9 - 42.7	500	64.5	57.3 - 71.1	800
Filipino	1,700	55.1	51.6 - 58.5	1,400	44.9	41.5 - 48.4	3,200
Japanese	500	28.6	24.4 - 33.3	1,200	71.4	66.7 - 75.6	1,700
Other	1,500	59.2	55.1 - 63.2	1,100	40.8	36.8 - 44.9	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	66.1	60.5 - 71.2	400	33.9	28.8 - 39.5	1,100
20-24	2,300	55.3	52.1 - 58.4	1,900	44.7	41.6 - 47.9	4,100
25-34	4,600	48.0	45.9 - 50.0	5,000	52.0	50.0 - 54.1	9,600
35+	1,300	41.0	37.7 - 44.4	1,900	59.0	55.6 - 62.3	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,400	48.4	46.3 - 50.4	4,700	51.6	49.6 - 53.7	9,100
Female	4,500	50.7	48.6 - 52.8	4,400	49.3	47.2 - 51.4	8,900

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,100	45.8	43.9 - 47.7	6,100	54.2	52.3 - 56.1	11,200
No	3,800	55.6	53.3 - 57.9	3,100	44.4	42.1 - 46.7	6,900

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,600	57.0	54.6 - 59.4	2,700	43.0	40.6 - 45.4	6,400
131-185%	1,500	54.9	51.0 - 58.6	1,300	45.1	41.4 - 49.0	2,800
186+%	2,900	39.5	37.3 - 41.7	4,500	60.5	58.3 - 62.7	7,400

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	51.9	49.5 - 54.3	600	48.1	45.7 - 50.5	1,300
Normal birth weight	8,300	49.3	47.7 - 50.9	8,500	50.7	49.1 - 52.3	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	47.2	41.6 - 52.9	600	52.8	47.1 - 58.4	1,100
No	8,200	49.8	48.3 - 51.3	8,300	50.2	48.7 - 51.7	16,600

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	45.4	42.2 - 48.8	1,900	54.6	51.2 - 57.8	3,400
No	7,200	50.6	49.0 - 52.3	7,000	49.4	47.7 - 51.0	14,300

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009-2013

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,500	57.4	55.2 - 59.5	3,400	42.6	40.5 - 44.8	7,900
No	4,400	43.7	41.7 - 45.7	5,600	56.3	54.3 - 58.3	10,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,000	49.7	48.3 - 51.1	9,100	50.3	48.9 - 51.7	18,100

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	51.2	48.7 - 53.7	1,100	48.8	46.3 - 51.3	2,300
Honolulu	6,500	50.1	48.2 - 51.9	6,500	49.9	48.1 - 51.8	12,900
Kauai	400	49.6	46.3 - 53.0	400	50.4	47.0 - 53.7	800
Maui	900	45.8	43.4 - 48.3	1,100	54.2	51.7 - 56.6	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	44.7	42.0 - 47.5	2,400	55.3	52.5 - 58.0	4,300
Native Hawaiian	2,900	53.8	51.3 - 56.3	2,500	46.2	43.7 - 48.7	5,300
Chinese	300	35.1	28.9 - 41.9	500	64.9	58.1 - 71.1	800
Filipino	1,800	55.3	52.1 - 58.5	1,500	44.7	41.5 - 47.9	3,300
Japanese	500	29.6	25.7 - 33.7	1,200	70.4	66.3 - 74.3	1,700
Other	1,600	60.0	56.3 - 63.7	1,100	40.0	36.3 - 43.7	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	66.7	61.7 - 71.4	400	33.3	28.6 - 38.3	1,200
20-24	2,300	55.3	52.4 - 58.3	1,800	44.7	41.7 - 47.6	4,100
25-34	4,600	47.7	45.8 - 49.6	5,000	52.3	50.4 - 54.2	9,500
35+	1,300	41.9	38.8 - 45.1	1,900	58.1	54.9 - 61.2	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,600	49.2	47.3 - 51.1	4,700	50.8	48.9 - 52.7	9,300
Female	4,400	50.3	48.3 - 52.2	4,400	49.7	47.8 - 51.7	8,800

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,200	46.2	44.5 - 48.0	6,000	53.8	52.0 - 55.5	11,200
No	3,800	55.5	53.2 - 57.6	3,100	44.5	42.4 - 46.8	6,900

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,600	57.1	54.8 - 59.4	2,700	42.9	40.6 - 45.2	6,400
131-185%	1,500	54.2	50.6 - 57.7	1,300	45.8	42.3 - 49.4	2,800
186+%	3,000	40.3	38.2 - 42.4	4,500	59.7	57.6 - 61.8	7,600

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	50.4	47.6 - 53.2	600	49.6	46.8 - 52.4	1,300
Normal birth weight	8,300	49.6	48.2 - 51.1	8,500	50.4	48.9 - 51.8	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	48.7	43.6 - 53.9	600	51.3	46.1 - 56.4	1,200
No	8,200	49.9	48.4 - 51.3	8,300	50.1	48.7 - 51.6	16,500

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	45.8	42.7 - 48.9	1,900	54.2	51.1 - 57.3	3,500
No	7,200	50.7	49.2 - 52.3	7,000	49.3	47.7 - 50.8	14,300

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008-2012

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,500	57.6	55.6 - 59.7	3,300	42.4	40.3 - 44.4	7,900
No	4,400	43.8	42.0 - 45.7	5,600	56.2	54.3 - 58.0	10,100

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	9,000	49.5	48.2 - 50.8	9,100	50.5	49.2 - 51.8	18,100

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,200	49.6	47.0 - 52.3	1,200	50.4	47.7 - 53.0	2,400
Honolulu	6,400	50.3	48.6 - 52.0	6,400	49.7	48.0 - 51.4	12,800
Kauai	400	49.8	46.0 - 53.6	400	50.2	46.4 - 54.0	900
Maui	900	44.3	41.7 - 46.9	1,100	55.7	53.1 - 58.3	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,000	45.3	42.6 - 47.9	2,400	54.7	52.1 - 57.4	4,300
Native Hawaiian	2,800	53.2	50.8 - 55.6	2,500	46.8	44.4 - 49.2	5,300
Chinese	200	35.2	29.4 - 41.5	500	64.8	58.5 - 70.6	700
Filipino	1,800	55.6	52.6 - 58.6	1,500	44.4	41.4 - 47.4	3,300
Japanese	600	30.6	27.0 - 34.4	1,300	69.4	65.6 - 73.0	1,800
Other	1,600	58.4	54.9 - 61.9	1,100	41.6	38.1 - 45.1	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	65.5	60.9 - 69.9	500	34.5	30.1 - 39.1	1,400
20-24	2,400	55.2	52.5 - 58.0	1,900	44.8	42.0 - 47.5	4,300
25-34	4,400	47.3	45.5 - 49.1	4,900	52.7	50.9 - 54.5	9,300
35+	1,300	41.3	38.3 - 44.3	1,900	58.7	55.7 - 61.7	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,600	49.1	47.3 - 50.9	4,800	50.9	49.1 - 52.7	9,400
Female	4,400	50.0	48.1 - 51.9	4,400	50.0	48.1 - 51.9	8,700

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,200	46.1	44.5 - 47.8	6,000	53.9	52.2 - 55.5	11,200
No	3,800	55.1	53.0 - 57.2	3,100	44.9	42.8 - 47.0	6,900

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,600	56.4	54.3 - 58.6	2,800	43.6	41.4 - 45.7	6,500
131-185%	1,400	52.1	48.7 - 55.5	1,300	47.9	44.5 - 51.3	2,700
186+%	3,100	40.4	38.5 - 42.4	4,500	59.6	57.6 - 61.5	7,600

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,800	59.3	56.1 - 62.4	1,300	40.7	37.6 - 43.9	3,100
\$10,000-\$24,999	2,000	54.4	51.5 - 57.3	1,600	45.6	42.7 - 48.5	3,600
\$25,000-\$49,999	2,000	48.8	46.1 - 51.5	2,200	51.2	48.5 - 53.9	4,200
>=\$50,000	2,400	39.0	36.9 - 41.3	3,700	61.0	58.7 - 63.1	6,100

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	49.5	46.4 - 52.6	700	50.5	47.4 - 53.6	1,300
Normal birth weight	8,300	49.5	48.1 - 50.9	8,500	50.5	49.1 - 51.9	16,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	48.5	43.8 - 53.2	700	51.5	46.8 - 56.2	1,400
No	8,200	49.7	48.3 - 51.0	8,300	50.3	49.0 - 51.7	16,400

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007-2011

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	47.2	44.3 - 50.1	1,900	52.8	49.9 - 55.7	3,600
No	7,100	50.2	48.7 - 51.6	7,100	49.8	48.4 - 51.3	14,200

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,400	56.9	54.9 - 58.9	3,300	43.1	41.1 - 45.1	7,700
No	4,500	44.1	42.4 - 45.9	5,700	55.9	54.1 - 57.6	10,100

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,900	49.2	48.0 - 50.5	9,100	50.8	49.5 - 52.0	18,000

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	48.1	45.3 - 50.9	1,200	51.9	49.1 - 54.7	2,300
Honolulu	6,400	50.2	48.7 - 51.8	6,400	49.8	48.2 - 51.3	12,800
Kauai	400	50.3	46.0 - 54.5	400	49.7	45.5 - 54.0	900
Maui	900	43.8	40.9 - 46.7	1,100	56.2	53.3 - 59.1	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	45.4	42.9 - 47.9	2,300	54.6	52.1 - 57.1	4,300
Native Hawaiian	2,700	52.0	49.6 - 54.3	2,500	48.0	45.7 - 50.4	5,200
Chinese	200	35.3	30.0 - 40.9	400	64.7	59.1 - 70.0	700
Filipino	1,800	55.1	52.3 - 57.9	1,500	44.9	42.1 - 47.7	3,300
Japanese	600	33.5	30.1 - 37.1	1,300	66.5	62.9 - 69.9	1,900
Other	1,500	57.4	54.1 - 60.7	1,100	42.6	39.3 - 45.9	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	63.3	58.9 - 67.5	500	36.7	32.5 - 41.1	1,500
20-24	2,300	53.5	50.9 - 56.1	2,000	46.5	43.9 - 49.1	4,300
25-34	4,300	47.4	45.7 - 49.2	4,800	52.6	50.8 - 54.3	9,100
35+	1,300	42.0	39.2 - 44.8	1,800	58.0	55.2 - 60.8	3,200

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,500	48.4	46.7 - 50.2	4,800	51.6	49.8 - 53.3	9,300
Female	4,400	50.1	48.3 - 51.9	4,400	49.9	48.1 - 51.7	8,700

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,200	46.2	44.7 - 47.8	6,000	53.8	52.2 - 55.3	11,200
No	3,700	54.2	52.1 - 56.2	3,100	45.8	43.8 - 47.9	6,800

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	54.8	52.7 - 56.9	2,900	45.2	43.1 - 47.3	6,400
131-185%	1,400	51.4	48.1 - 54.7	1,300	48.6	45.3 - 51.9	2,700
186+%	3,100	41.3	39.5 - 43.2	4,500	58.7	56.8 - 60.5	7,600

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,700	57.3	54.3 - 60.4	1,300	42.7	39.6 - 45.7	3,000
\$10,000-\$24,999	1,900	52.9	50.2 - 55.6	1,700	47.1	44.4 - 49.8	3,700
\$25,000-\$49,999	1,900	47.2	44.6 - 49.8	2,200	52.8	50.2 - 55.4	4,100
>=\$50,000	2,500	41.0	38.9 - 43.1	3,500	59.0	56.9 - 61.1	6,000

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	49.2	45.7 - 52.6	700	50.8	47.4 - 54.3	1,400
Normal birth weight	8,200	49.2	47.9 - 50.5	8,500	50.8	49.5 - 52.1	16,600

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	48.8	44.5 - 53.2	800	51.2	46.8 - 55.5	1,500
No	8,000	49.3	48.0 - 50.6	8,200	50.7	49.4 - 52.0	16,200

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006-2010

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	47.7	45.0 - 50.5	1,900	52.3	49.5 - 55.0	3,600
No	7,000	49.6	48.2 - 51.0	7,100	50.4	49.0 - 51.8	14,100

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,300	56.0	54.1 - 57.9	3,400	44.0	42.1 - 45.9	7,600
No	4,500	44.3	42.7 - 45.9	5,700	55.7	54.1 - 57.3	10,200

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,800	49.6	48.4 - 50.7	9,000	50.4	49.3 - 51.6	17,700

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	47.3	44.3 - 50.2	1,200	52.7	49.8 - 55.7	2,300
Honolulu	6,400	50.8	49.4 - 52.2	6,200	49.2	47.8 - 50.6	12,600
Kauai	500	51.9	47.3 - 56.5	400	48.1	43.5 - 52.7	900
Maui	900	43.3	40.3 - 46.4	1,100	56.7	53.6 - 59.7	2,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	43.6	41.2 - 46.0	2,300	56.4	54.0 - 58.8	4,100
Native Hawaiian	2,700	53.0	50.7 - 55.3	2,400	47.0	44.7 - 49.3	5,100
Chinese	200	38.0	33.5 - 42.8	400	62.0	57.2 - 66.5	600
Filipino	1,900	57.0	54.4 - 59.5	1,400	43.0	40.5 - 45.6	3,200
Japanese	700	35.2	32.1 - 38.5	1,300	64.8	61.5 - 67.9	2,000
Other	1,500	56.7	53.6 - 59.7	1,100	43.3	40.3 - 46.4	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	62.4	58.2 - 66.4	600	37.6	33.6 - 41.8	1,600
20-24	2,300	53.7	51.3 - 56.1	2,000	46.3	43.9 - 48.7	4,200
25-34	4,200	47.8	46.2 - 49.4	4,600	52.2	50.6 - 53.8	8,800
35+	1,300	42.2	39.7 - 44.9	1,800	57.8	55.1 - 60.3	3,100

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,500	49.2	47.6 - 50.8	4,700	50.8	49.2 - 52.4	9,200
Female	4,300	50.0	48.3 - 51.6	4,300	50.0	48.4 - 51.7	8,500

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,200	46.8	45.4 - 48.2	5,900	53.2	51.8 - 54.6	11,100
No	3,600	54.2	52.2 - 56.1	3,100	45.8	43.9 - 47.8	6,700

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	54.5	52.5 - 56.5	2,900	45.5	43.5 - 47.5	6,400
131-185%	1,300	52.1	48.9 - 55.2	1,200	47.9	44.8 - 51.1	2,400
186+%	3,200	42.0	40.4 - 43.7	4,400	58.0	56.3 - 59.6	7,600

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,600	55.2	52.2 - 58.1	1,300	44.8	41.9 - 47.8	2,900
\$10,000-\$24,999	2,000	53.7	51.2 - 56.3	1,700	46.3	43.7 - 48.8	3,700
\$25,000-\$49,999	2,000	49.0	46.6 - 51.4	2,000	51.0	48.6 - 53.4	4,000
>=\$50,000	2,400	41.1	39.2 - 43.0	3,400	58.9	57.0 - 60.8	5,800

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	49.0	45.3 - 52.7	700	51.0	47.3 - 54.7	1,300
Normal birth weight	8,100	49.6	48.4 - 50.8	8,300	50.4	49.2 - 51.6	16,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	50.2	46.1 - 54.3	800	49.8	45.7 - 53.9	1,500
No	7,900	49.5	48.3 - 50.7	8,000	50.5	49.3 - 51.7	15,900

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005-2009

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	48.6	46.0 - 51.3	1,800	51.4	48.7 - 54.0	3,600
No	6,900	49.7	48.4 - 51.0	6,900	50.3	49.0 - 51.6	13,800

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,200	55.6	53.8 - 57.4	3,300	44.4	42.6 - 46.2	7,500
No	4,500	44.9	43.4 - 46.4	5,500	55.1	53.6 - 56.6	10,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,800	49.7	48.7 - 50.8	8,800	50.3	49.2 - 51.3	17,600

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,100	47.5	44.4 - 50.6	1,200	52.5	49.4 - 55.6	2,200
Honolulu	6,400	50.8	49.5 - 52.0	6,200	49.2	48.0 - 50.5	12,600
Kauai	500	52.2	47.3 - 57.0	400	47.8	43.0 - 52.7	900
Maui	900	44.5	41.2 - 47.7	1,100	55.5	52.3 - 58.8	1,900

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	43.2	41.0 - 45.4	2,300	56.8	54.6 - 59.0	4,100
Native Hawaiian	2,500	51.9	49.7 - 54.1	2,400	48.1	45.9 - 50.3	4,900
Chinese	200	39.1	36.3 - 41.9	400	60.9	58.1 - 63.7	600
Filipino	1,900	58.1	55.8 - 60.4	1,400	41.9	39.6 - 44.2	3,200
Japanese	800	36.2	33.3 - 39.1	1,400	63.8	60.9 - 66.7	2,100
Other	1,600	58.8	56.1 - 61.5	1,100	41.2	38.5 - 43.9	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	61.9	58.0 - 65.7	600	38.1	34.3 - 42.0	1,500
20-24	2,200	52.6	50.3 - 54.8	2,000	47.4	45.2 - 49.7	4,300
25-34	4,200	48.6	47.1 - 50.1	4,500	51.4	49.9 - 52.9	8,700
35+	1,400	43.3	41.0 - 45.7	1,800	56.7	54.3 - 59.0	3,100

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,600	49.8	48.3 - 51.2	4,600	50.2	48.8 - 51.7	9,200
Female	4,200	49.7	48.1 - 51.2	4,200	50.3	48.8 - 51.9	8,400

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,400	47.6	46.3 - 48.9	6,000	52.4	51.1 - 53.7	11,400
No	3,300	53.6	51.8 - 55.5	2,900	46.4	44.5 - 48.2	6,200

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,500	54.5	52.7 - 56.3	2,900	45.5	43.7 - 47.3	6,400
131-185%	1,200	52.4	49.4 - 55.4	1,100	47.6	44.6 - 50.6	2,300
186+%	3,300	43.2	41.6 - 44.7	4,300	56.8	55.3 - 58.4	7,600

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,600	55.0	52.2 - 57.8	1,300	45.0	42.2 - 47.8	2,900
\$10,000-\$24,999	2,100	53.7	51.4 - 56.1	1,800	46.3	43.9 - 48.6	3,800
\$25,000-\$49,999	2,000	49.3	47.1 - 51.5	2,100	50.7	48.5 - 52.9	4,100
>=\$50,000	2,400	42.1	40.4 - 43.9	3,200	57.9	56.1 - 59.6	5,600

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	47.9	44.0 - 51.9	700	52.1	48.1 - 56.0	1,300
Normal birth weight	8,100	49.9	48.8 - 51.0	8,200	50.1	49.0 - 51.2	16,300

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	50.1	46.2 - 53.9	700	49.9	46.1 - 53.8	1,500
No	7,900	49.6	48.5 - 50.7	8,000	50.4	49.3 - 51.5	15,800

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004-2008

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	49.6	47.1 - 52.1	1,800	50.4	47.9 - 52.9	3,500
No	6,800	49.6	48.5 - 50.8	6,900	50.4	49.2 - 51.5	13,800

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,200	55.1	53.4 - 56.7	3,400	44.9	43.3 - 46.6	7,600
No	4,500	45.5	44.1 - 46.9	5,400	54.5	53.1 - 55.9	9,900

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,800	51.0	49.9 - 52.1	8,500	49.0	47.9 - 50.1	17,300

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	48.7	45.7 - 51.6	1,100	51.3	48.4 - 54.3	2,100
Honolulu	6,500	51.7	50.4 - 53.1	6,000	48.3	46.9 - 49.6	12,500
Kauai	500	55.6	51.0 - 60.1	400	44.4	39.9 - 49.0	900
Maui	900	46.7	43.6 - 49.8	1,000	53.3	50.2 - 56.4	1,800

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	44.7	42.4 - 47.0	2,200	55.3	53.0 - 57.6	3,900
Native Hawaiian	2,600	53.5	51.4 - 55.5	2,200	46.5	44.5 - 48.6	4,800
Chinese	300	41.5	37.3 - 45.8	400	58.5	54.2 - 62.7	600
Filipino	2,000	59.2	56.8 - 61.6	1,400	40.8	38.4 - 43.2	3,300
Japanese	800	37.0	33.9 - 40.1	1,300	63.0	59.9 - 66.1	2,100
Other	1,500	59.3	56.4 - 62.3	1,000	40.7	37.7 - 43.6	2,500

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	59.8	55.9 - 63.6	600	40.2	36.4 - 44.1	1,500
20-24	2,300	54.0	51.8 - 56.2	2,000	46.0	43.8 - 48.2	4,300
25-34	4,300	50.3	48.8 - 51.9	4,200	49.7	48.1 - 51.2	8,500
35+	1,400	44.3	41.8 - 46.8	1,700	55.7	53.2 - 58.2	3,100

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,700	51.6	50.1 - 53.1	4,400	48.4	46.9 - 49.9	9,000
Female	4,200	50.4	48.8 - 52.0	4,100	49.6	48.0 - 51.2	8,300

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,600	49.4	48.1 - 50.8	5,700	50.6	49.2 - 51.9	11,300
No	3,300	54.0	52.1 - 55.9	2,800	46.0	44.1 - 47.9	6,000

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,600	55.6	53.8 - 57.4	2,900	44.4	42.6 - 46.2	6,600
131-185%	1,200	53.3	50.2 - 56.4	1,000	46.7	43.6 - 49.8	2,200
186+%	3,200	44.4	42.8 - 46.1	4,000	55.6	53.9 - 57.2	7,200

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	49.6	45.6 - 53.6	600	50.4	46.4 - 54.4	1,300
Normal birth weight	8,200	51.1	50.0 - 52.3	7,800	48.9	47.7 - 50.0	16,000

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	48.0	44.2 - 51.9	800	52.0	48.1 - 55.8	1,500
No	8,000	51.3	50.1 - 52.4	7,600	48.7	47.6 - 49.9	15,600

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,700	49.5	47.0 - 52.0	1,800	50.5	48.0 - 53.0	3,500
No	6,900	51.4	50.2 - 52.7	6,600	48.6	47.3 - 49.8	13,500

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003-2007

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,200	55.9	54.3 - 57.6	3,300	44.1	42.4 - 45.7	7,500
No	4,600	47.1	45.6 - 48.5	5,100	52.9	51.5 - 54.4	9,700

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,700	50.9	49.7 - 52.0	8,400	49.1	48.0 - 50.3	17,100

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	50.4	47.6 - 53.1	1,000	49.6	46.9 - 52.4	2,100
Honolulu	6,300	50.8	49.4 - 52.2	6,100	49.2	47.8 - 50.6	12,500
Kauai	400	56.4	51.9 - 60.8	300	43.6	39.2 - 48.1	800
Maui	900	49.3	46.3 - 52.2	900	50.7	47.8 - 53.7	1,800

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,700	44.5	42.1 - 47.0	2,100	55.5	53.0 - 57.9	3,800
Native Hawaiian	2,500	53.1	51.2 - 55.0	2,200	46.9	45.0 - 48.8	4,700
Chinese	300	42.0	36.8 - 47.3	400	58.0	52.7 - 63.2	600
Filipino	2,000	59.7	57.1 - 62.3	1,300	40.3	37.7 - 42.9	3,300
Japanese	800	36.5	33.3 - 39.7	1,400	63.5	60.3 - 66.7	2,200
Other	1,400	59.4	56.1 - 62.6	1,000	40.6	37.4 - 43.9	2,400

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	58.5	54.7 - 62.3	600	41.5	37.7 - 45.3	1,500
20-24	2,300	53.6	51.3 - 55.9	2,000	46.4	44.1 - 48.7	4,200
25-34	4,300	50.5	48.9 - 52.2	4,200	49.5	47.8 - 51.1	8,400
35+	1,300	44.0	41.4 - 46.7	1,700	56.0	53.3 - 58.6	3,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,600	52.0	50.4 - 53.5	4,200	48.0	46.5 - 49.6	8,800
Female	4,100	49.6	48.0 - 51.3	4,100	50.4	48.7 - 52.0	8,200

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,600	49.2	47.8 - 50.6	5,800	50.8	49.4 - 52.2	11,300
No	3,100	54.1	52.2 - 56.1	2,600	45.9	43.9 - 47.8	5,700

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,600	54.7	52.9 - 56.5	2,900	45.3	43.5 - 47.1	6,500
131-185%	1,300	53.8	50.6 - 56.9	1,100	46.2	43.1 - 49.4	2,400
186+%	3,000	44.2	42.5 - 46.0	3,800	55.8	54.0 - 57.5	6,800

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	49.3	45.0 - 53.6	600	50.7	46.4 - 55.0	1,200
Normal birth weight	8,100	51.0	49.8 - 52.2	7,800	49.0	47.8 - 50.2	15,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	45.3	41.4 - 49.2	800	54.7	50.8 - 58.6	1,400
No	7,900	51.3	50.1 - 52.5	7,500	48.7	47.5 - 49.9	15,300

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	47.4	44.9 - 49.9	1,800	52.6	50.1 - 55.1	3,400
No	6,900	51.7	50.4 - 53.0	6,400	48.3	47.0 - 49.6	13,300

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002-2006

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,100	55.4	53.7 - 57.1	3,300	44.6	42.9 - 46.3	7,400
No	4,500	47.3	45.8 - 48.8	5,000	52.7	51.2 - 54.2	9,600

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,500	50.4	49.3 - 51.5	8,300	49.6	48.5 - 50.7	16,800

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	51.2	48.7 - 53.8	1,000	48.8	46.2 - 51.3	2,000
Honolulu	6,200	50.2	48.8 - 51.5	6,100	49.8	48.5 - 51.2	12,300
Kauai	400	55.7	51.7 - 59.8	300	44.3	40.2 - 48.3	800
Maui	800	48.9	46.2 - 51.6	900	51.1	48.4 - 53.8	1,700

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	43.2	40.8 - 45.7	2,100	56.8	54.3 - 59.2	3,700
Native Hawaiian	2,400	52.9	51.2 - 54.5	2,200	47.1	45.5 - 48.8	4,600
Chinese	300	42.7	37.1 - 48.4	400	57.3	51.6 - 62.9	700
Filipino	2,000	59.8	57.2 - 62.4	1,400	40.2	37.6 - 42.8	3,400
Japanese	800	34.9	31.7 - 38.1	1,400	65.1	61.9 - 68.3	2,200
Other	1,400	59.9	56.6 - 63.2	900	40.1	36.8 - 43.4	2,300

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	57.5	54.0 - 61.0	600	42.5	39.0 - 46.0	1,500
20-24	2,200	53.5	51.3 - 55.7	1,900	46.5	44.3 - 48.7	4,200
25-34	4,100	49.8	48.2 - 51.4	4,100	50.2	48.6 - 51.8	8,300
35+	1,300	44.0	41.3 - 46.7	1,600	56.0	53.3 - 58.7	2,900

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,500	51.4	49.9 - 53.0	4,200	48.6	47.0 - 50.1	8,700
Female	4,000	49.3	47.7 - 50.9	4,100	50.7	49.1 - 52.3	8,100

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,500	48.7	47.3 - 50.1	5,800	51.3	49.9 - 52.7	11,300
No	3,000	53.8	51.9 - 55.6	2,500	46.2	44.4 - 48.1	5,500

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,600	54.9	53.1 - 56.6	2,900	45.1	43.4 - 46.9	6,500
131-185%	1,400	51.8	48.9 - 54.6	1,300	48.2	45.4 - 51.1	2,600
186+%	2,700	43.5	41.7 - 45.3	3,500	56.5	54.7 - 58.3	6,300

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	50.2	45.8 - 54.5	600	49.8	45.5 - 54.2	1,100
Normal birth weight	7,900	50.5	49.3 - 51.6	7,700	49.5	48.4 - 50.7	15,600

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	44.8	41.1 - 48.5	800	55.2	51.5 - 58.9	1,400
No	7,700	50.9	49.8 - 52.1	7,400	49.1	47.9 - 50.2	15,100

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	46.7	44.2 - 49.1	1,800	53.3	50.9 - 55.8	3,400
No	6,700	51.4	50.1 - 52.6	6,300	48.6	47.4 - 49.9	13,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001-2005

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	4,000	54.9	53.3 - 56.6	3,300	45.1	43.4 - 46.7	7,200
No	4,400	47.0	45.4 - 48.5	5,000	53.0	51.5 - 54.6	9,400

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	8,300	49.7	48.6 - 50.8	8,400	50.3	49.2 - 51.4	16,700

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	1,000	50.5	48.2 - 52.8	1,000	49.5	47.2 - 51.8	1,900
Honolulu	6,100	49.5	48.1 - 50.9	6,200	50.5	49.1 - 51.9	12,300
Kauai	400	54.4	50.7 - 58.1	300	45.6	41.9 - 49.3	700
Maui	800	48.4	46.0 - 50.9	900	51.6	49.1 - 54.0	1,700

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,600	44.0	41.6 - 46.5	2,100	56.0	53.5 - 58.4	3,700
Native Hawaiian	2,300	51.1	49.7 - 52.6	2,200	48.9	47.4 - 50.3	4,600
Chinese	300	42.4	36.5 - 48.6	400	57.6	51.4 - 63.5	700
Filipino	2,000	58.9	56.3 - 61.5	1,400	41.1	38.5 - 43.7	3,400
Japanese	700	33.8	30.6 - 37.2	1,400	66.2	62.8 - 69.4	2,100
Other	1,300	59.6	56.2 - 62.9	900	40.4	37.1 - 43.8	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	55.1	51.7 - 58.4	700	44.9	41.6 - 48.3	1,500
20-24	2,200	52.4	50.3 - 54.5	2,000	47.6	45.5 - 49.7	4,100
25-34	4,100	50.0	48.3 - 51.6	4,100	50.0	48.4 - 51.7	8,200
35+	1,200	42.3	39.7 - 45.1	1,600	57.7	54.9 - 60.3	2,900

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	4,400	50.3	48.8 - 51.9	4,300	49.7	48.1 - 51.2	8,700
Female	3,900	49.1	47.5 - 50.7	4,100	50.9	49.3 - 52.5	8,000

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,500	48.3	47.0 - 49.7	5,900	51.7	50.3 - 53.0	11,400
No	2,800	52.8	50.9 - 54.6	2,500	47.2	45.4 - 49.1	5,300

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	3,600	53.9	52.2 - 55.6	3,100	46.1	44.4 - 47.8	6,700
131-185%	1,400	51.8	49.0 - 54.6	1,300	48.2	45.4 - 51.0	2,700
186+%	2,600	42.8	41.0 - 44.7	3,400	57.2	55.3 - 59.0	6,000

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	53.1	48.7 - 57.4	500	46.9	42.6 - 51.3	1,100
Normal birth weight	7,700	49.5	48.4 - 50.6	7,900	50.5	49.4 - 51.6	15,600

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	44.9	41.4 - 48.5	800	55.1	51.5 - 58.6	1,400
No	7,500	50.3	49.1 - 51.4	7,500	49.7	48.6 - 50.9	15,000

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	46.8	44.4 - 49.1	1,800	53.2	50.9 - 55.6	3,300
No	6,500	50.5	49.3 - 51.8	6,400	49.5	48.2 - 50.7	12,900

PRAMS Health Indicator 5-Year Aggregate Report - Discuss seat belt use with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000-2004

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,900	54.6	53.0 - 56.3	3,300	45.4	43.7 - 47.0	7,200
No	4,300	46.2	44.7 - 47.7	5,000	53.8	52.3 - 55.3	9,400