

**PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.

Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.

Below are the indicators included in this report and the long description for each indicator.

PRAMS Health Indicator	Health Indicator Description
Weight - BMI status (PRAMS)	Weight - BMI status (PRAMS)

Suggested citation:

Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/13/17

For documentation on other terms and coding in this report, please go to the following links:

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact
The Hawai'i Health Data Warehouse**

profiles@hhdw.org

(855) 946-5899 x15


**PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- for the State of Hawaii, for the Year(s)
2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	1,400	8.4	7.2 - 9.8	9,700	59.1	56.9 - 61.3	3,300	20.0	18.3 - 21.8	2,100	12.6	11.2 - 14.1	16,500
2001	1,200	7.2	6.1 - 8.5	9,900	60.8	58.5 - 62.9	3,100	18.8	17.1 - 20.6	2,100	13.2	11.8 - 14.7	16,200
2002	800	5.1	4.0 - 6.4	9,800	59.6	56.8 - 62.4	3,700	22.2	20.0 - 24.7	2,100	13.0	11.3 - 15.0	16,500
2003	1,100	6.6	5.3 - 8.2	9,700	56.1	53.4 - 58.8	3,800	21.6	19.5 - 23.9	2,700	15.6	13.8 - 17.7	17,300
2004	1,200	6.7	5.7 - 7.9	9,200	54.0	51.8 - 56.1	3,900	22.8	21.0 - 24.7	2,800	16.5	14.9 - 18.3	17,100
2005	1,100	6.4	5.3 - 7.6	9,600	57.1	54.7 - 59.4	3,500	20.6	18.7 - 22.7	2,700	15.9	14.2 - 17.8	16,800
2006	1,100	5.9	4.8 - 7.1	10,500	58.3	55.8 - 60.7	3,900	21.7	19.7 - 23.8	2,600	14.2	12.5 - 16.0	18,000
2007	1,000	5.8	4.8 - 7.0	10,000	55.3	52.9 - 57.7	4,000	22.4	20.5 - 24.5	3,000	16.4	14.7 - 18.3	18,000
2008	1,000	5.7	4.7 - 6.9	10,200	55.5	53.2 - 57.8	4,100	22.1	20.2 - 24.2	3,100	16.7	14.9 - 18.6	18,400
2009	800	4.4	3.3 - 5.9	9,800	56.1	52.9 - 59.3	3,900	22.2	19.6 - 25.0	3,000	17.3	15.0 - 19.9	17,500
2010	1,000	5.3	4.0 - 7.1	10,000	55.5	52.3 - 58.7	4,400	24.2	21.5 - 27.1	2,700	14.9	12.7 - 17.3	18,000
2011	800	4.7	3.6 - 6.2	9,600	54.4	51.1 - 57.7	4,800	27.0	24.1 - 30.1	2,500	13.9	11.8 - 16.3	17,700
2012	700	4.1	3.0 - 5.6	10,600	59.3	56.1 - 62.5	3,500	19.4	17.0 - 22.1	3,100	17.2	14.8 - 19.8	17,900
2013	800	4.7	3.4 - 6.4	9,800	54.7	51.2 - 58.2	4,000	22.3	19.5 - 25.4	3,300	18.3	15.8 - 21.2	17,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	800	4.7	3.4 - 6.4	9,800	54.7	51.2 - 58.2	4,000	22.3	19.5 - 25.4	3,300	18.3	15.8 - 21.2	17,900

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.4	2.5 - 7.6	1,300	54.3	48.7 - 59.7	500	19.6	15.6 - 24.3	500	21.8	17.5 - 26.7	2,400
Honolulu	600	4.8	3.2 - 7.3	6,800	53.8	49.0 - 58.5	2,900	23.2	19.4 - 27.4	2,300	18.2	14.9 - 22.2	12,700
Kauai	<50	4.3	2.6 - 7.1	500	65.1	59.8 - 69.9	200	19.1	15.3 - 23.6	100	11.5	8.5 - 15.3	800
Maui	100	4.2	2.4 - 7.1	1,100	56.8	51.2 - 62.3	400	21.3	17.0 - 26.2	300	17.8	13.9 - 22.5	1,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	5.5	3.0 - 9.8	2,900	60.5	53.4 - 67.1	1,000	21.5	16.2 - 27.8	600	12.6	8.5 - 18.2	4,800
Native Hawaiian	200	3.7	2.0 - 6.8	2,400	43.6	37.8 - 49.6	1,300	24.5	19.6 - 30.1	1,500	28.2	23.1 - 33.9	5,500
Chinese	<50	1.2	0.5 - 3.3	700	77.0	60.8 - 87.8	100	13.6	5.8 - 28.8	100	8.1	2.5 - 23.1	900
Filipino	200	6.3	3.0 - 12.7	1,900	65.2	56.4 - 73.0	600	20.2	14.1 - 27.9	200	8.4	4.7 - 14.5	2,900
Japanese	100	6.9	2.8 - 15.7	900	56.0	43.9 - 67.4	400	22.0	13.4 - 33.9	300	15.2	8.4 - 25.8	1,700
Other	100	2.9	1.0 - 8.1	1,000	46.1	36.1 - 56.5	500	25.1	17.0 - 35.5	600	25.8	17.7 - 36.0	2,200

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	7.1	3.5 - 14.0	500	62.2	46.3 - 75.8	200	23.5	12.2 - 40.4	100	7.3	2.2 - 21.2	800
20-24	200	6.1	3.2 - 11.1	2,100	52.8	45.2 - 60.3	1,000	25.1	19.0 - 32.4	700	16.0	11.2 - 22.5	4,100
25-34	400	4.3	2.6 - 6.9	5,300	55.1	50.3 - 59.9	1,900	19.8	16.3 - 23.9	2,000	20.8	17.2 - 24.9	9,600
35+	100	3.6	1.7 - 7.5	1,800	53.9	46.1 - 61.5	900	25.6	19.3 - 33.1	600	17.0	11.8 - 23.7	3,400

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	5.0	3.2 - 7.8	4,900	55.6	50.6 - 60.5	2,000	23.2	19.2 - 27.7	1,400	16.2	12.9 - 20.2	8,800
Female	400	4.4	2.8 - 6.9	4,900	53.8	48.8 - 58.8	1,900	21.4	17.5 - 25.9	1,900	20.4	16.7 - 24.7	9,100

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	4.9	3.3 - 7.4	6,300	55.7	51.1 - 60.2	2,300	19.9	16.5 - 23.9	2,200	19.4	16.1 - 23.2	11,300
No	300	4.3	2.5 - 7.0	3,500	53.0	47.4 - 58.4	1,700	26.3	21.6 - 31.6	1,100	16.5	12.7 - 21.1	6,600

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.8	2.8 - 8.1	3,300	49.8	44.2 - 55.4	1,500	22.8	18.3 - 28.0	1,500	22.6	18.2 - 27.7	6,600
131-185%	100	6.0	2.7 - 13.0	1,300	56.3	46.5 - 65.6	500	20.9	14.0 - 30.0	400	16.8	10.7 - 25.3	2,400
186+%	300	4.6	2.7 - 7.8	4,200	57.3	51.7 - 62.8	1,700	22.8	18.5 - 27.9	1,100	15.2	11.6 - 19.7	7,300

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Income (2012+)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	200	5.4	3.0 - 9.6	2,000	50.4	43.3 - 57.5	900	21.5	16.1 - 28.1	900	22.7	17.1 - 29.5	4,100
\$17,001-\$26,000	100	5.8	2.3 - 14.0	1,400	57.3	47.6 - 66.5	500	22.3	15.3 - 31.4	300	14.5	9.2 - 22.1	2,400
\$26,001-\$51,000	200	4.4	2.1 - 8.7	2,000	53.3	45.6 - 60.9	900	22.5	16.7 - 29.8	700	19.7	14.4 - 26.4	3,800
>=\$51,001	300	4.6	2.7 - 8.0	3,500	56.5	50.4 - 62.4	1,400	23.3	18.5 - 28.8	1,000	15.6	11.6 - 20.6	6,200

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.1	4.7 - 10.6	700	55.5	50.5 - 60.4	300	20.2	16.7 - 24.4	200	17.1	13.7 - 21.2	1,300
Normal birth weight	700	4.5	3.1 - 6.4	9,000	54.7	50.9 - 58.4	3,700	22.5	19.4 - 25.8	3,000	18.4	15.6 - 21.5	16,500

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.3	3.3 - 19.6	400	41.8	29.2 - 55.6	300	26.0	15.4 - 40.4	200	23.9	14.1 - 37.4	1,000
No	700	4.4	3.1 - 6.2	9,100	55.0	51.3 - 58.6	3,700	22.4	19.4 - 25.6	3,000	18.3	15.6 - 21.3	16,500

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2013

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.3	2.6 - 10.5	1,400	43.7	35.8 - 51.9	800	23.8	17.3 - 31.8	900	27.2	20.4 - 35.3	3,200
No	600	4.4	3.1 - 6.4	8,100	56.6	52.6 - 60.5	3,200	22.3	19.1 - 25.8	2,400	16.7	13.9 - 19.8	14,300

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	4.2	2.5 - 7.0	3,700	46.9	41.7 - 52.1	1,800	23.0	18.9 - 27.7	2,000	25.9	21.5 - 30.8	7,800
No	500	5.1	3.3 - 7.6	6,000	60.5	55.8 - 65.1	2,200	21.8	18.1 - 26.1	1,300	12.6	9.8 - 16.0	10,000

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	700	4.1	3.0 - 5.6	10,600	59.3	56.1 - 62.5	3,500	19.4	17.0 - 22.1	3,100	17.2	14.8 - 19.8	17,900

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.5	2.6 - 7.4	1,200	53.0	47.6 - 58.3	500	23.0	18.8 - 27.8	400	19.6	15.6 - 24.3	2,200
Honolulu	500	4.0	2.7 - 6.1	7,900	60.6	56.2 - 64.8	2,400	18.7	15.5 - 22.4	2,200	16.7	13.6 - 20.3	13,000
Kauai	<50	4.9	2.8 - 8.6	500	58.3	51.9 - 64.5	100	17.3	12.9 - 22.6	200	19.5	14.9 - 25.1	800
Maui	100	3.4	2.0 - 5.8	1,100	58.7	53.6 - 63.6	400	21.3	17.4 - 25.7	300	16.6	13.2 - 20.7	1,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	100	3.3	1.7 - 6.4	2,500	62.1	55.3 - 68.4	900	21.3	16.3 - 27.4	500	13.3	9.3 - 18.8	4,100
Native Hawaiian	200	3.5	1.8 - 6.6	2,600	50.7	44.8 - 56.6	1,200	23.1	18.4 - 28.6	1,200	22.7	18.2 - 28.0	5,200
Chinese	100	8.5	3.2 - 20.9	800	75.7	61.8 - 85.7	100	8.0	3.2 - 18.6	100	7.7	2.8 - 19.7	1,000
Filipino	100	4.4	2.3 - 8.4	2,000	62.5	54.7 - 69.6	600	18.3	13.2 - 24.7	500	14.9	10.0 - 21.5	3,200
Japanese	100	5.8	2.3 - 13.8	1,000	67.5	56.1 - 77.2	200	12.9	7.1 - 22.4	200	13.8	7.6 - 23.8	1,500
Other	100	3.4	1.3 - 8.3	1,600	57.2	48.5 - 65.5	500	19.0	13.2 - 26.6	600	20.3	14.2 - 28.3	2,900

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.1	3.3 - 18.2	700	68.9	54.6 - 80.3	200	18.5	9.7 - 32.6	<50	4.5	1.1 - 16.3	1,000
20-24	100	3.2	1.4 - 7.2	2,100	58.1	50.7 - 65.2	800	22.1	16.6 - 28.8	600	16.6	11.8 - 22.8	3,600
25-34	400	3.9	2.5 - 5.9	5,900	58.4	54.0 - 62.7	2,000	19.3	16.0 - 23.0	1,900	18.4	15.2 - 22.1	10,100
35+	100	4.4	2.2 - 8.9	1,900	60.5	52.9 - 67.6	600	17.2	12.5 - 23.3	600	17.9	12.7 - 24.5	3,200

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.2	1.9 - 5.1	5,200	58.7	54.0 - 63.3	1,900	21.4	17.8 - 25.5	1,500	16.7	13.4 - 20.6	8,800
Female	500	4.9	3.3 - 7.4	5,500	59.9	55.3 - 64.3	1,600	17.6	14.4 - 21.2	1,600	17.6	14.4 - 21.4	9,100

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	3.8	2.5 - 5.7	6,600	59.8	55.5 - 63.9	2,100	18.8	15.7 - 22.3	1,900	17.7	14.6 - 21.2	11,000
No	300	4.5	2.8 - 7.3	4,100	58.6	53.4 - 63.6	1,400	20.5	16.7 - 25.0	1,100	16.4	12.9 - 20.5	6,900

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.3	2.5 - 7.2	3,500	55.8	50.3 - 61.1	1,300	20.8	16.8 - 25.5	1,200	19.1	15.2 - 23.7	6,300
131-185%	100	3.5	1.7 - 7.0	1,700	56.8	48.7 - 64.6	600	18.5	13.1 - 25.4	600	21.2	15.3 - 28.7	3,100
186+%	300	4.2	2.5 - 7.0	4,400	63.7	58.4 - 68.6	1,300	19.6	15.7 - 24.2	900	12.5	9.4 - 16.5	6,800

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Income (2012+)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	200	4.0	2.0 - 7.6	2,500	61.4	54.5 - 67.9	700	16.3	11.9 - 21.7	700	18.4	13.5 - 24.5	4,000
\$17,001-\$26,000	100	4.6	1.7 - 11.6	1,200	56.1	47.0 - 64.8	500	25.2	18.2 - 33.8	300	14.1	9.7 - 20.0	2,100
\$26,001-\$51,000	100	2.2	1.1 - 4.2	2,300	53.6	46.8 - 60.3	1,100	24.6	19.2 - 30.9	900	19.6	14.8 - 25.5	4,300
>=\$51,001	300	5.4	3.2 - 8.7	3,700	62.4	56.6 - 67.8	1,000	17.0	13.1 - 21.7	900	15.3	11.5 - 20.0	5,900

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.3	3.4 - 8.4	700	55.8	50.1 - 61.3	300	22.1	17.7 - 27.1	200	16.9	13.0 - 21.6	1,200
Normal birth weight	700	4.0	2.8 - 5.6	9,900	59.5	56.0 - 62.8	3,200	19.3	16.7 - 22.2	2,900	17.2	14.7 - 20.1	16,700

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	<50	1.3	0.4 - 4.4	300	35.9	23.1 - 51.0	200	25.6	15.0 - 40.2	300	37.2	23.4 - 53.5	700
No	700	4.1	3.0 - 5.7	10,000	60.5	57.1 - 63.8	3,200	19.3	16.8 - 22.2	2,700	16.0	13.7 - 18.7	16,600

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2012

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	3.6	1.7 - 7.2	1,500	52.2	44.2 - 60.2	600	22.2	16.4 - 29.2	600	22.0	16.1 - 29.4	2,800
No	600	4.1	2.9 - 5.8	8,800	60.8	57.2 - 64.3	2,800	19.1	16.4 - 22.2	2,300	16.0	13.4 - 18.8	14,500

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.6	2.2 - 5.9	4,200	52.3	47.4 - 57.2	1,700	21.1	17.4 - 25.2	1,900	23.0	19.1 - 27.4	8,100
No	400	4.3	2.9 - 6.6	6,400	65.3	60.9 - 69.4	1,700	17.9	14.7 - 21.6	1,200	12.5	9.8 - 15.7	9,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	800	4.7	3.6 - 6.2	9,600	54.4	51.1 - 57.7	4,800	27.0	24.1 - 30.1	2,500	13.9	11.8 - 16.3	17,700

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	7.0	4.8 - 10.3	1,300	57.0	51.8 - 62.1	500	21.4	17.5 - 25.9	300	14.6	11.3 - 18.6	2,300
Honolulu	500	3.9	2.6 - 6.0	6,800	53.6	49.2 - 58.0	3,700	29.3	25.4 - 33.5	1,700	13.1	10.4 - 16.4	12,700
Kauai	100	7.8	5.0 - 11.8	500	54.6	48.3 - 60.8	200	23.4	18.5 - 29.1	100	14.2	10.4 - 19.3	800
Maui	100	5.5	3.7 - 8.1	1,100	56.4	51.7 - 61.0	400	19.9	16.4 - 24.0	300	18.2	14.8 - 22.1	1,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.5	3.2 - 9.2	2,600	59.7	53.1 - 66.0	1,100	25.3	19.9 - 31.7	400	9.5	6.3 - 14.1	4,400
Native Hawaiian	200	3.2	1.8 - 5.6	2,600	52.2	46.3 - 58.1	1,200	24.3	19.6 - 29.8	1,000	20.2	16.0 - 25.3	5,100
Chinese	100	11.4	3.8 - 29.3	500	67.4	49.8 - 81.2	100	16.6	7.7 - 32.4	<50	4.6	0.8 - 21.4	700
Filipino	200	5.5	3.1 - 9.4	1,800	57.8	50.1 - 65.1	900	27.5	21.0 - 35.2	300	9.2	5.8 - 14.4	3,200
Japanese	100	5.7	2.8 - 11.3	1,000	54.4	43.9 - 64.6	500	27.0	18.6 - 37.4	200	12.9	7.2 - 22.0	1,800
Other	100	2.8	0.9 - 8.0	1,100	41.9	33.2 - 51.2	1,000	37.4	28.7 - 47.0	500	17.9	11.9 - 26.1	2,600

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	10.7	6.5 - 17.2	600	62.5	49.0 - 74.3	200	19.3	10.2 - 33.6	100	7.5	2.2 - 22.5	1,000
20-24	200	5.5	3.1 - 9.8	2,500	58.6	51.6 - 65.3	1,100	25.8	20.1 - 32.5	400	10.1	6.6 - 15.0	4,200
25-34	400	4.6	3.1 - 6.8	4,900	51.2	46.7 - 55.7	2,800	29.1	25.1 - 33.5	1,400	15.2	12.3 - 18.5	9,500
35+	100	1.8	1.0 - 3.1	1,600	56.0	48.1 - 63.6	700	24.7	18.5 - 32.2	500	17.5	12.2 - 24.5	2,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	500	5.4	3.7 - 7.7	4,900	54.2	49.5 - 58.7	2,400	26.3	22.3 - 30.6	1,300	14.2	11.3 - 17.7	9,100
Female	300	4.0	2.6 - 6.1	4,700	54.7	49.9 - 59.3	2,400	27.8	23.7 - 32.3	1,200	13.6	10.7 - 17.0	8,600

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.9	3.5 - 7.0	5,900	53.7	49.5 - 57.8	2,900	26.6	23.0 - 30.5	1,600	14.8	12.0 - 18.1	11,000
No	300	4.3	2.7 - 6.7	3,700	55.6	50.3 - 60.8	1,900	27.7	23.1 - 32.9	800	12.4	9.5 - 16.0	6,700

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	4.2	2.5 - 7.0	3,100	48.3	43.0 - 53.6	1,900	30.5	25.6 - 35.9	1,100	17.0	13.4 - 21.4	6,300
131-185%	200	6.7	3.5 - 12.2	1,500	55.3	46.6 - 63.7	700	26.7	19.6 - 35.2	300	11.3	6.9 - 18.0	2,700
186+%	300	3.3	2.0 - 5.2	4,700	59.4	54.5 - 64.2	1,900	24.5	20.4 - 29.1	1,000	12.8	9.9 - 16.5	7,800

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Income (2004-2011)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	6.2	3.4 - 10.9	1,600	50.2	42.7 - 57.8	1,000	30.6	23.8 - 38.3	400	13.0	8.9 - 18.6	3,200
\$10,000-\$24,999	200	5.2	2.7 - 9.6	1,800	53.5	46.0 - 60.8	800	25.3	19.1 - 32.6	500	16.1	11.4 - 22.3	3,300
\$25,000-\$49,999	100	3.4	1.7 - 6.7	2,300	53.2	46.4 - 59.9	1,200	27.5	21.7 - 34.1	700	15.9	11.6 - 21.3	4,300
>=\$50,000	200	3.3	2.0 - 5.5	3,600	58.3	52.6 - 63.7	1,600	25.9	21.2 - 31.3	800	12.5	9.2 - 16.7	6,100

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.3	4.7 - 11.1	800	59.0	53.4 - 64.3	300	20.8	16.8 - 25.6	200	12.9	9.7 - 17.0	1,300
Normal birth weight	700	4.5	3.3 - 6.1	8,800	54.0	50.4 - 57.5	4,500	27.5	24.4 - 30.9	2,300	14.0	11.8 - 16.6	16,400

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.3	3.7 - 17.6	400	43.6	31.5 - 56.6	300	33.8	21.5 - 48.7	100	14.2	7.4 - 25.6	900
No	800	4.5	3.3 - 6.0	9,200	55.0	51.6 - 58.3	4,500	26.8	23.8 - 30.0	2,300	13.7	11.6 - 16.2	16,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2011

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	4.5	2.5 - 7.9	1,700	49.2	41.9 - 56.6	1,000	31.0	24.4 - 38.4	500	15.3	11.1 - 20.8	3,400
No	700	4.8	3.5 - 6.5	7,900	55.6	51.9 - 59.3	3,700	26.3	23.0 - 29.7	1,900	13.3	11.0 - 16.1	14,200

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	4.5	3.0 - 6.8	3,800	48.6	43.8 - 53.5	2,300	29.3	25.0 - 34.1	1,400	17.6	14.2 - 21.5	7,900
No	500	4.9	3.3 - 7.0	5,700	58.8	54.3 - 63.2	2,500	25.3	21.5 - 29.6	1,100	11.0	8.5 - 14.1	9,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,000	5.3	4.0 - 7.1	10,000	55.5	52.3 - 58.7	4,400	24.2	21.5 - 27.1	2,700	14.9	12.7 - 17.3	18,000

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	3.8	2.3 - 6.4	1,200	54.3	49.2 - 59.4	600	25.8	21.6 - 30.6	400	16.0	12.6 - 20.1	2,300
Honolulu	700	5.5	3.8 - 7.9	7,200	55.5	51.2 - 59.8	3,200	24.7	21.2 - 28.7	1,900	14.3	11.5 - 17.6	13,100
Kauai	100	7.6	4.9 - 11.4	400	56.0	49.9 - 61.9	200	19.8	15.4 - 25.0	100	16.7	12.6 - 21.7	800
Maui	100	5.4	3.6 - 8.1	1,100	57.0	52.1 - 61.8	400	20.8	17.1 - 25.0	300	16.9	13.5 - 20.8	1,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.3	3.0 - 9.1	2,700	60.0	53.4 - 66.3	1,100	25.5	20.1 - 31.8	400	9.2	6.1 - 13.8	4,400
Native Hawaiian	200	3.9	2.2 - 6.9	2,700	50.6	44.9 - 56.3	1,400	26.2	21.4 - 31.7	1,000	19.3	15.3 - 24.0	5,400
Chinese	<50	5.2	1.1 - 21.4	600	82.3	68.8 - 90.7	100	8.9	3.8 - 19.6	<50	3.6	1.7 - 7.6	700
Filipino	100	4.2	2.1 - 8.3	1,800	55.9	48.3 - 63.3	800	26.2	20.1 - 33.4	400	13.7	9.0 - 20.1	3,200
Japanese	100	8.6	4.3 - 16.7	1,100	63.3	52.6 - 72.9	300	18.7	11.5 - 28.8	200	9.4	4.9 - 17.1	1,700
Other	200	7.9	3.9 - 15.3	1,200	45.0	36.2 - 54.1	600	23.5	16.7 - 32.0	600	23.6	16.8 - 32.2	2,600

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	12.5	6.3 - 23.4	800	66.8	54.6 - 77.1	200	17.2	9.7 - 28.5	<50	3.5	1.6 - 7.5	1,100
20-24	200	5.5	3.1 - 9.6	2,400	54.4	47.7 - 61.0	1,200	27.4	21.8 - 33.9	600	12.7	8.9 - 17.7	4,400
25-34	400	4.5	2.9 - 6.9	4,900	53.3	48.8 - 57.8	2,300	24.4	20.7 - 28.5	1,600	17.8	14.6 - 21.5	9,300
35+	200	5.1	2.5 - 10.0	2,000	59.4	51.8 - 66.6	700	21.9	16.3 - 28.8	400	13.5	9.2 - 19.5	3,300

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	600	6.1	4.2 - 8.7	5,000	54.3	49.8 - 58.8	2,100	22.7	19.0 - 26.7	1,500	17.0	13.9 - 20.6	9,100
Female	400	4.6	3.0 - 7.1	5,100	56.8	52.1 - 61.3	2,300	25.8	22.0 - 30.1	1,100	12.7	10.0 - 16.2	8,900

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	6.0	4.2 - 8.5	6,500	57.1	52.9 - 61.2	2,700	23.6	20.2 - 27.3	1,500	13.4	10.9 - 16.4	11,300
No	300	4.3	2.8 - 6.7	3,600	52.9	47.8 - 58.0	1,700	25.4	21.1 - 30.1	1,200	17.4	13.8 - 21.6	6,700

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.2	3.9 - 9.6	3,000	47.2	41.9 - 52.6	1,700	26.2	21.7 - 31.3	1,300	20.4	16.3 - 25.1	6,300
131-185%	100	3.2	1.4 - 7.3	1,700	59.4	51.3 - 67.1	700	22.9	16.7 - 30.6	400	14.5	9.8 - 20.7	2,900
186+%	400	5.8	3.8 - 8.8	4,600	59.6	54.6 - 64.4	1,800	22.9	18.9 - 27.4	900	11.7	8.9 - 15.3	7,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Income (2004-2011)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	6.3	3.5 - 11.1	1,700	49.7	42.5 - 57.0	900	26.0	20.2 - 32.8	600	18.0	12.9 - 24.6	3,400
\$10,000-\$24,999	200	6.5	3.4 - 11.9	1,500	47.8	40.3 - 55.4	900	27.2	20.6 - 35.0	600	18.6	13.5 - 25.0	3,100
\$25,000-\$49,999	200	4.6	2.5 - 8.4	2,600	58.1	51.6 - 64.3	1,000	21.7	16.9 - 27.5	700	15.6	11.5 - 20.7	4,500
>=\$50,000	300	5.2	3.2 - 8.5	3,600	59.6	53.9 - 65.0	1,400	23.2	18.8 - 28.4	700	12.0	8.8 - 16.1	6,100

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.6	5.0 - 11.2	700	55.2	49.8 - 60.5	300	22.0	17.8 - 26.8	200	15.3	11.7 - 19.6	1,300
Normal birth weight	900	5.2	3.8 - 7.1	9,300	55.5	52.0 - 58.9	4,100	24.5	21.6 - 27.6	2,500	14.9	12.6 - 17.5	16,700

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.0	3.6 - 17.2	600	40.6	29.7 - 52.6	400	27.5	18.1 - 39.5	300	23.8	15.3 - 35.1	1,400
No	800	5.2	3.8 - 7.0	9,300	56.8	53.4 - 60.1	3,900	24.0	21.2 - 27.1	2,300	14.0	11.9 - 16.5	16,400

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2010

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.2	2.9 - 9.1	1,800	47.5	40.6 - 54.5	1,000	26.4	20.6 - 33.0	800	20.9	15.7 - 27.3	3,700
No	800	5.4	3.9 - 7.5	8,100	57.6	54.0 - 61.2	3,300	23.7	20.7 - 27.0	1,900	13.2	11.0 - 15.8	14,100

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	5.1	3.4 - 7.6	3,800	48.9	44.1 - 53.7	2,000	25.9	21.8 - 30.3	1,600	20.1	16.5 - 24.3	7,800
No	500	5.3	3.6 - 7.9	6,100	60.8	56.4 - 65.0	2,300	23.1	19.6 - 27.1	1,100	10.8	8.4 - 13.7	10,000

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	800	4.4	3.3 - 5.9	9,800	56.1	52.9 - 59.3	3,900	22.2	19.6 - 25.0	3,000	17.3	15.0 - 19.9	17,500

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.0	2.4 - 6.6	1,300	53.8	48.6 - 58.9	500	22.5	18.5 - 27.1	500	19.6	15.9 - 24.0	2,300
Honolulu	500	4.3	2.9 - 6.4	6,900	55.9	51.6 - 60.2	2,800	22.2	18.8 - 26.1	2,200	17.5	14.5 - 21.1	12,400
Kauai	<50	4.2	2.4 - 7.4	500	61.4	55.3 - 67.2	200	20.5	16.0 - 25.9	100	13.9	10.1 - 18.8	800
Maui	100	5.4	3.6 - 8.1	1,100	57.8	53.0 - 62.4	400	22.4	18.6 - 26.6	300	14.4	11.4 - 18.1	2,000

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	3.6	2.1 - 6.3	2,600	63.4	57.0 - 69.3	800	20.0	15.3 - 25.8	500	13.0	9.1 - 18.2	4,200
Native Hawaiian	200	3.0	1.6 - 5.7	2,700	46.6	41.1 - 52.2	1,400	24.2	19.7 - 29.3	1,500	26.2	21.6 - 31.4	5,700
Chinese	<50	4.5	1.0 - 18.4	600	85.2	69.1 - 93.6	100	8.9	2.7 - 25.5	<50	1.5	0.4 - 5.4	700
Filipino	300	10.2	6.2 - 16.4	1,700	55.0	47.3 - 62.5	600	20.8	15.3 - 27.5	400	14.1	9.3 - 20.7	3,000
Japanese	100	4.5	2.1 - 9.4	1,000	66.4	55.4 - 75.8	300	18.7	11.3 - 29.3	200	10.4	5.5 - 19.0	1,500
Other	<50	1.6	0.8 - 3.1	1,200	52.3	43.2 - 61.2	700	29.5	21.9 - 38.5	400	16.6	10.8 - 24.8	2,400

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	9.1	4.4 - 17.9	1,100	62.2	51.5 - 71.8	400	19.7	12.5 - 29.6	200	9.0	4.6 - 16.8	1,800
20-24	100	3.4	1.8 - 6.1	2,300	55.6	48.9 - 62.2	1,000	25.1	19.6 - 31.5	600	15.9	11.6 - 21.4	4,100
25-34	300	3.9	2.6 - 6.0	4,800	54.7	50.2 - 59.1	1,900	21.5	18.1 - 25.4	1,800	19.9	16.4 - 23.8	8,800
35+	100	4.3	2.2 - 8.5	1,600	57.2	49.4 - 64.7	600	21.8	15.9 - 29.3	500	16.6	11.7 - 22.9	2,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	300	3.7	2.4 - 5.6	5,500	59.0	54.6 - 63.3	1,900	20.1	16.8 - 23.9	1,600	17.2	14.0 - 20.9	9,400
Female	400	5.2	3.5 - 7.7	4,300	52.7	48.1 - 57.3	2,000	24.6	20.8 - 28.9	1,400	17.4	14.2 - 21.2	8,200

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	3.9	2.7 - 5.7	6,000	58.0	53.9 - 62.1	2,200	21.6	18.3 - 25.2	1,700	16.4	13.5 - 19.8	10,300
No	400	5.0	3.2 - 7.9	3,900	53.4	48.3 - 58.3	1,700	23.1	19.1 - 27.7	1,300	18.5	14.9 - 22.7	7,300

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	200	3.5	2.2 - 5.7	3,200	50.8	45.4 - 56.2	1,700	26.8	22.1 - 32.0	1,200	18.9	15.1 - 23.4	6,300
131-185%	100	5.0	2.3 - 10.4	1,300	48.4	40.3 - 56.7	600	23.1	16.8 - 30.8	600	23.5	17.1 - 31.4	2,700
186+%	300	4.2	2.7 - 6.6	4,500	62.7	57.7 - 67.3	1,400	19.3	15.7 - 23.5	1,000	13.8	10.6 - 17.7	7,100

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Income (2004-2011)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	4.9	2.8 - 8.5	1,500	47.6	39.8 - 55.6	1,000	31.5	24.4 - 39.5	500	16.0	11.0 - 22.7	3,100
\$10,000-\$24,999	100	2.4	1.1 - 5.0	2,100	57.2	50.4 - 63.8	800	21.1	16.0 - 27.3	700	19.4	14.7 - 25.2	3,700
\$25,000-\$49,999	200	4.5	2.3 - 8.5	1,700	48.9	41.9 - 56.0	800	23.0	17.7 - 29.4	800	23.6	17.8 - 30.4	3,500
>=\$50,000	300	4.6	2.8 - 7.5	3,800	63.1	57.7 - 68.3	1,100	19.2	15.2 - 24.0	800	13.0	9.7 - 17.2	5,900

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.7	3.8 - 8.6	700	56.9	51.6 - 62.1	200	18.3	14.6 - 22.8	200	19.0	15.1 - 23.7	1,300
Normal birth weight	700	4.3	3.1 - 5.9	9,100	56.1	52.7 - 59.5	3,600	22.4	19.6 - 25.4	2,800	17.2	14.7 - 20.0	16,200

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.9	2.9 - 8.2	700	43.9	33.8 - 54.5	300	21.2	13.6 - 31.5	500	30.1	20.9 - 41.1	1,600
No	700	4.3	3.1 - 6.0	8,800	56.7	53.3 - 60.0	3,500	22.8	20.1 - 25.9	2,500	16.2	13.8 - 18.8	15,500

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2009

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.7	3.3 - 9.6	1,900	48.9	42.3 - 55.6	800	21.1	16.1 - 27.2	900	24.3	18.9 - 30.7	3,900
No	500	4.0	2.8 - 5.7	7,600	57.4	53.7 - 61.0	3,100	23.1	20.1 - 26.4	2,100	15.5	13.0 - 18.4	13,300

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	3.6	2.2 - 5.8	3,500	48.5	43.6 - 53.5	1,900	26.3	22.1 - 31.0	1,500	21.6	17.7 - 26.0	7,100
No	500	5.0	3.4 - 7.2	6,000	60.1	55.8 - 64.3	2,000	20.2	16.9 - 23.9	1,500	14.7	11.9 - 18.1	9,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,000	5.7	4.7 - 6.9	10,200	55.5	53.2 - 57.8	4,100	22.1	20.2 - 24.2	3,100	16.7	14.9 - 18.6	18,400

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	7.5	4.6 - 12.1	1,200	49.4	42.5 - 56.2	600	26.0	20.4 - 32.6	400	17.1	12.4 - 23.1	2,300
Honolulu	700	5.4	4.3 - 6.8	7,200	55.6	52.8 - 58.4	2,700	21.2	19.0 - 23.6	2,300	17.7	15.6 - 20.1	12,900
Kauai	<50	3.7	1.3 - 10.4	600	71.3	60.6 - 80.1	200	19.6	12.4 - 29.7	<50	5.4	2.1 - 13.0	900
Maui	100	5.9	3.4 - 10.2	1,200	54.9	47.9 - 61.7	500	24.5	19.0 - 31.0	300	14.7	10.3 - 20.5	2,200

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	6.1	4.1 - 8.8	2,900	67.9	63.2 - 72.2	700	15.8	12.6 - 19.7	400	10.2	7.7 - 13.6	4,300
Native Hawaiian	200	4.3	2.7 - 6.7	2,300	44.0	39.2 - 48.8	1,400	26.6	22.5 - 31.2	1,300	25.1	21.1 - 29.6	5,100
Chinese	100	9.1	5.8 - 13.9	400	68.8	62.3 - 74.7	100	17.7	13.2 - 23.4	<50	4.4	2.3 - 8.1	600
Filipino	200	5.9	3.9 - 8.9	2,100	61.0	55.8 - 66.0	800	22.5	18.4 - 27.1	400	10.6	7.8 - 14.3	3,400
Japanese	100	7.3	4.4 - 11.8	1,300	62.6	55.6 - 69.2	400	21.1	15.9 - 27.5	200	9.0	5.7 - 14.1	2,000
Other	200	5.5	3.4 - 8.7	1,200	42.8	36.8 - 49.0	700	25.0	20.0 - 30.8	800	26.8	21.4 - 32.9	2,800

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.9	5.8 - 16.5	800	56.6	47.5 - 65.2	300	18.7	12.6 - 26.9	200	14.8	9.3 - 22.7	1,500
20-24	300	7.9	5.5 - 11.3	2,000	50.3	45.0 - 55.6	1,000	24.9	20.5 - 29.8	700	16.9	13.2 - 21.3	3,900
25-34	500	4.8	3.6 - 6.3	5,400	56.8	53.6 - 60.0	2,000	21.3	18.8 - 24.2	1,600	17.0	14.6 - 19.7	9,600
35+	100	3.7	2.3 - 6.1	2,000	57.4	52.1 - 62.5	800	22.6	18.5 - 27.4	600	16.2	12.6 - 20.7	3,400

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	500	5.3	4.1 - 6.9	5,200	54.3	51.0 - 57.5	2,300	23.3	20.6 - 26.2	1,700	17.2	14.8 - 19.9	9,700
Female	500	6.1	4.6 - 8.0	4,900	56.9	53.4 - 60.3	1,800	20.9	18.2 - 23.9	1,400	16.1	13.7 - 18.9	8,700

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	5.3	4.2 - 6.8	6,800	59.1	56.2 - 61.9	2,300	20.3	18.1 - 22.8	1,800	15.3	13.2 - 17.6	11,600
No	400	6.2	4.6 - 8.5	3,400	49.5	45.4 - 53.5	1,700	25.2	21.8 - 29.0	1,300	19.1	16.0 - 22.5	6,800

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.2	4.5 - 8.4	3,200	47.7	43.6 - 51.7	1,700	25.2	21.8 - 28.9	1,400	21.0	17.8 - 24.6	6,800
131-185%	100	4.5	2.5 - 8.0	1,300	48.8	42.6 - 55.2	700	26.2	20.9 - 32.2	500	20.5	15.7 - 26.3	2,600
186+%	400	4.7	3.5 - 6.3	5,000	64.3	60.9 - 67.6	1,500	18.8	16.2 - 21.7	900	12.2	10.0 - 14.8	7,800

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Income (2004-2011)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	5.8	3.6 - 9.4	1,500	50.2	44.1 - 56.3	800	26.0	20.8 - 31.8	500	18.0	13.6 - 23.4	3,000
\$10,000-\$24,999	300	8.5	6.0 - 11.9	2,000	49.8	44.6 - 55.1	900	22.8	18.6 - 27.6	700	18.9	15.0 - 23.5	4,000
\$25,000-\$49,999	100	3.0	1.8 - 5.0	2,300	54.5	49.6 - 59.4	900	22.0	18.1 - 26.4	900	20.5	16.6 - 24.9	4,200
>=\$50,000	300	5.0	3.6 - 6.9	3,800	61.7	57.8 - 65.4	1,300	20.6	17.7 - 24.0	800	12.7	10.2 - 15.6	6,200

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	12.1	7.2 - 19.5	500	41.5	32.9 - 50.7	400	29.3	21.6 - 38.5	200	17.1	11.1 - 25.3	1,200
Normal birth weight	900	5.2	4.3 - 6.4	9,700	56.5	54.1 - 58.9	3,700	21.6	19.6 - 23.7	2,900	16.7	14.9 - 18.7	17,100

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.6	2.1 - 10.0	700	44.6	36.4 - 53.2	400	26.6	19.7 - 34.9	400	24.1	17.4 - 32.4	1,600
No	900	5.8	4.8 - 7.0	9,300	56.6	54.1 - 59.0	3,500	21.6	19.6 - 23.7	2,600	16.0	14.2 - 18.0	16,400

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2008

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	5.5	3.5 - 8.8	1,600	47.0	41.5 - 52.7	900	25.7	21.0 - 31.1	800	21.7	17.3 - 26.8	3,500
No	800	5.7	4.6 - 7.0	8,300	57.6	54.9 - 60.2	3,100	21.1	19.0 - 23.4	2,200	15.6	13.7 - 17.6	14,500

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	6.0	4.5 - 8.0	3,600	46.8	43.1 - 50.6	1,900	24.3	21.1 - 27.7	1,800	22.9	19.8 - 26.3	7,700
No	500	5.1	3.9 - 6.6	6,500	62.3	59.3 - 65.3	2,100	20.2	17.9 - 22.8	1,300	12.3	10.4 - 14.6	10,400

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,000	5.8	4.8 - 7.0	10,000	55.3	52.9 - 57.7	4,000	22.4	20.5 - 24.5	3,000	16.4	14.7 - 18.3	18,000

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	8.4	5.4 - 13.0	1,100	46.5	39.8 - 53.2	500	22.9	17.7 - 29.0	500	22.3	17.1 - 28.4	2,400
Honolulu	700	5.4	4.3 - 6.8	6,900	54.9	52.0 - 57.7	2,900	23.2	20.9 - 25.7	2,100	16.5	14.4 - 18.8	12,500
Kauai	100	6.0	2.5 - 13.4	700	65.2	54.8 - 74.2	200	18.8	12.0 - 28.2	100	10.0	5.4 - 17.9	1,000
Maui	100	4.9	2.6 - 9.0	1,300	63.4	56.1 - 70.1	400	18.9	13.8 - 25.5	300	12.8	8.7 - 18.4	2,100

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	300	7.0	4.8 - 10.0	2,400	58.7	53.7 - 63.5	800	20.7	16.9 - 25.1	600	13.6	10.5 - 17.5	4,100
Native Hawaiian	200	4.1	2.5 - 6.6	2,600	50.9	45.9 - 55.8	1,100	22.3	18.5 - 26.7	1,100	22.7	18.9 - 27.1	5,000
Chinese	100	13.2	9.3 - 18.4	500	69.5	63.0 - 75.3	100	12.1	8.3 - 17.2	<50	5.3	2.9 - 9.4	700
Filipino	200	5.0	3.1 - 7.8	1,900	57.7	52.6 - 62.7	800	24.8	20.6 - 29.5	400	12.6	9.5 - 16.4	3,400
Japanese	100	6.4	3.7 - 10.8	1,300	63.3	56.4 - 69.8	500	22.0	16.8 - 28.2	200	8.3	5.1 - 13.1	2,100
Other	200	5.8	3.6 - 9.2	1,300	46.5	40.4 - 52.7	700	25.2	20.1 - 31.1	600	22.4	17.6 - 28.2	2,900

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	9.1	5.2 - 15.5	1,100	69.4	60.4 - 77.2	200	15.5	9.9 - 23.5	100	5.9	2.7 - 12.3	1,600
20-24	300	7.4	5.2 - 10.4	2,400	52.8	47.9 - 57.8	1,000	23.3	19.3 - 27.7	700	16.5	13.1 - 20.6	4,500
25-34	400	4.5	3.4 - 6.0	4,700	53.2	49.8 - 56.5	2,000	22.7	20.0 - 25.7	1,700	19.6	17.0 - 22.5	8,900
35+	200	5.4	3.4 - 8.4	1,800	57.9	52.4 - 63.3	700	24.1	19.6 - 29.1	400	12.6	9.3 - 16.9	3,100

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	500	5.6	4.2 - 7.2	5,100	55.1	51.8 - 58.4	2,200	23.9	21.1 - 26.9	1,400	15.4	13.1 - 18.0	9,300
Female	500	6.0	4.6 - 7.8	4,900	55.6	52.1 - 59.0	1,800	20.9	18.2 - 23.9	1,500	17.5	15.0 - 20.3	8,800

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	5.7	4.5 - 7.2	6,200	55.3	52.3 - 58.2	2,500	22.1	19.7 - 24.6	1,900	16.9	14.8 - 19.3	11,300
No	400	5.8	4.2 - 8.0	3,800	55.5	51.3 - 59.5	1,600	23.1	19.8 - 26.8	1,100	15.6	12.9 - 18.8	6,800

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	500	6.8	5.0 - 9.1	3,400	50.4	46.3 - 54.5	1,500	23.0	19.7 - 26.7	1,300	19.8	16.7 - 23.3	6,700
131-185%	100	5.1	3.0 - 8.4	1,500	55.2	49.0 - 61.2	600	22.5	17.9 - 28.0	500	17.2	13.0 - 22.4	2,700
186+%	400	5.7	4.3 - 7.6	4,300	58.5	54.9 - 62.0	1,600	22.0	19.1 - 25.1	1,000	13.8	11.4 - 16.6	7,400

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Income (2004-2011)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	7.3	4.7 - 11.1	1,600	54.6	48.3 - 60.9	600	19.1	14.6 - 24.7	600	19.0	14.5 - 24.5	2,900
\$10,000-\$24,999	200	5.6	3.6 - 8.6	2,000	53.0	47.7 - 58.3	900	22.7	18.5 - 27.5	700	18.7	14.9 - 23.3	3,800
\$25,000-\$49,999	300	6.7	4.7 - 9.6	2,200	52.3	47.3 - 57.2	1,000	24.4	20.3 - 28.9	700	16.6	13.2 - 20.8	4,100
>=\$50,000	300	5.2	3.7 - 7.1	3,500	58.1	54.1 - 62.0	1,300	22.4	19.3 - 26.0	900	14.3	11.6 - 17.4	6,000

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.4	4.0 - 13.2	900	58.8	50.6 - 66.6	300	20.4	14.6 - 27.6	200	13.4	8.7 - 20.1	1,500
Normal birth weight	900	5.6	4.6 - 6.8	9,100	55.0	52.5 - 57.5	3,700	22.6	20.6 - 24.8	2,800	16.7	14.9 - 18.7	16,500

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	6.7	3.6 - 12.1	700	45.4	37.0 - 54.1	400	27.6	20.6 - 36.0	300	20.3	14.3 - 28.1	1,500
No	900	5.7	4.7 - 7.0	9,100	56.3	53.8 - 58.8	3,600	22.0	19.9 - 24.1	2,600	16.0	14.2 - 18.0	16,200

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2007

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	7.2	4.9 - 10.5	1,700	47.0	41.5 - 52.6	900	25.1	20.6 - 30.1	700	20.6	16.5 - 25.5	3,600
No	800	5.4	4.4 - 6.7	8,200	57.5	54.8 - 60.1	3,100	21.8	19.6 - 24.1	2,200	15.3	13.4 - 17.3	14,200

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	400	6.0	4.4 - 8.1	3,700	50.4	46.5 - 54.3	1,600	22.3	19.3 - 25.8	1,600	21.3	18.2 - 24.6	7,400
No	600	5.7	4.4 - 7.2	6,000	58.9	55.8 - 61.9	2,300	22.3	19.8 - 25.0	1,300	13.1	11.2 - 15.4	10,200

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,100	5.9	4.8 - 7.1	10,500	58.3	55.8 - 60.7	3,900	21.7	19.7 - 23.8	2,600	14.2	12.5 - 16.0	18,000

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.0	3.4 - 10.4	1,200	54.0	46.7 - 61.2	500	24.8	19.1 - 31.6	300	15.1	10.5 - 21.2	2,200
Honolulu	700	5.6	4.4 - 7.0	7,700	58.5	55.7 - 61.3	2,800	21.5	19.3 - 24.0	1,900	14.4	12.5 - 16.5	13,200
Kauai	100	9.9	4.6 - 19.9	500	59.6	47.4 - 70.8	200	25.3	16.2 - 37.3	<50	5.2	1.8 - 14.4	800
Maui	100	6.4	3.5 - 11.4	1,100	60.9	53.1 - 68.2	300	17.4	12.3 - 24.1	300	15.3	10.4 - 21.8	1,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.6	3.7 - 8.4	2,700	63.3	58.3 - 68.0	900	21.3	17.4 - 25.7	400	9.8	7.2 - 13.2	4,300
Native Hawaiian	200	4.4	2.7 - 7.1	2,500	50.5	45.4 - 55.5	1,200	24.6	20.5 - 29.2	1,000	20.5	16.7 - 24.8	5,000
Chinese	100	12.9	8.9 - 18.4	400	74.1	67.5 - 79.8	100	8.8	5.6 - 13.7	<50	4.1	2.0 - 8.0	600
Filipino	200	6.5	4.3 - 9.9	2,000	61.7	56.3 - 66.8	700	22.4	18.2 - 27.2	300	9.3	6.6 - 13.0	3,300
Japanese	100	6.0	3.4 - 10.4	1,400	65.4	58.5 - 71.7	400	19.1	14.2 - 25.1	200	9.5	6.1 - 14.6	2,100
Other	200	6.6	4.1 - 10.6	1,400	51.6	45.2 - 57.9	600	20.9	16.1 - 26.7	600	20.9	16.1 - 26.6	2,800

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	14.5	9.1 - 22.3	1,100	71.1	61.9 - 78.8	200	13.4	8.2 - 21.2	<50	1.1	0.2 - 6.7	1,500
20-24	400	7.9	5.5 - 11.1	2,500	56.6	51.4 - 61.6	900	21.0	17.1 - 25.6	600	14.6	11.2 - 18.6	4,500
25-34	400	4.5	3.3 - 6.1	5,000	56.7	53.3 - 60.1	2,000	22.9	20.1 - 26.0	1,400	15.9	13.4 - 18.6	8,800
35+	100	3.0	1.7 - 5.3	1,900	58.9	53.5 - 64.1	800	23.0	18.7 - 27.9	500	15.1	11.5 - 19.5	3,300

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	400	4.5	3.3 - 6.2	5,200	57.0	53.5 - 60.3	2,200	23.8	21.0 - 26.9	1,300	14.7	12.3 - 17.3	9,200
Female	600	7.3	5.7 - 9.3	5,300	59.6	56.1 - 63.0	1,700	19.4	16.8 - 22.4	1,200	13.7	11.4 - 16.3	8,800

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	5.2	4.1 - 6.7	6,800	59.1	56.2 - 62.0	2,600	22.6	20.2 - 25.2	1,500	13.1	11.2 - 15.2	11,500
No	500	7.0	5.1 - 9.5	3,700	56.8	52.4 - 61.0	1,300	20.1	16.8 - 23.9	1,100	16.1	13.2 - 19.6	6,500

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.2	4.5 - 8.6	3,500	56.5	52.1 - 60.7	1,300	20.2	16.9 - 23.9	1,100	17.1	14.1 - 20.7	6,300
131-185%	200	8.4	5.2 - 13.1	1,100	46.6	39.9 - 53.5	700	29.2	23.4 - 35.9	400	15.8	11.4 - 21.4	2,400
186+%	300	4.1	3.0 - 5.6	5,200	63.4	59.9 - 66.7	1,700	20.9	18.2 - 23.9	900	11.6	9.5 - 14.1	8,200

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Income (2004-2011)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	7.1	4.5 - 10.8	1,800	58.6	52.2 - 64.7	600	19.2	14.5 - 24.8	500	15.2	11.2 - 20.4	3,000
\$10,000-\$24,999	200	5.9	3.8 - 9.1	2,100	56.5	50.9 - 61.9	800	20.9	16.7 - 25.8	600	16.6	12.9 - 21.2	3,700
\$25,000-\$49,999	200	5.4	3.5 - 8.1	2,400	55.2	50.2 - 60.2	900	21.8	17.9 - 26.4	800	17.6	14.0 - 21.8	4,300
>=\$50,000	300	4.5	3.1 - 6.5	3,600	61.9	57.8 - 65.8	1,400	23.5	20.2 - 27.2	600	10.1	7.9 - 12.9	5,900

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	9.0	5.2 - 15.3	900	61.4	52.9 - 69.3	300	19.7	13.9 - 27.2	100	9.8	5.8 - 16.2	1,500
Normal birth weight	900	5.6	4.6 - 6.9	9,500	57.9	55.4 - 60.4	3,600	21.9	19.8 - 24.1	2,400	14.5	12.8 - 16.5	16,500

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.1	4.6 - 13.7	800	48.7	40.3 - 57.2	300	20.2	14.2 - 27.9	400	23.0	16.6 - 31.0	1,700
No	900	5.6	4.5 - 6.9	9,500	59.2	56.6 - 61.7	3,500	21.9	19.8 - 24.1	2,100	13.3	11.7 - 15.2	16,100

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2006

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	8.1	5.6 - 11.6	2,000	52.4	46.8 - 58.0	800	21.2	17.0 - 26.1	700	18.3	14.3 - 23.1	3,800
No	700	5.2	4.1 - 6.6	8,300	59.8	57.0 - 62.5	3,000	21.8	19.6 - 24.2	1,800	13.2	11.4 - 15.2	14,000

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	7.3	5.5 - 9.6	4,000	53.4	49.5 - 57.4	1,600	21.2	18.2 - 24.7	1,400	18.0	15.2 - 21.3	7,500
No	500	5.0	3.8 - 6.5	6,300	61.4	58.3 - 64.5	2,300	22.3	19.7 - 25.0	1,200	11.3	9.4 - 13.6	10,200

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,100	6.4	5.3 - 7.6	9,600	57.1	54.7 - 59.4	3,500	20.6	18.7 - 22.7	2,700	15.9	14.2 - 17.8	16,800

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	4.5	2.4 - 8.4	1,100	52.0	44.9 - 59.0	500	24.3	18.6 - 31.1	400	19.1	14.1 - 25.4	2,100
Honolulu	700	5.8	4.7 - 7.2	7,200	58.4	55.6 - 61.1	2,400	19.6	17.4 - 21.9	2,000	16.2	14.2 - 18.4	12,400
Kauai	100	14.2	7.6 - 25.0	300	44.8	33.7 - 56.5	200	27.3	18.1 - 39.0	100	13.7	7.1 - 24.6	700
Maui	100	9.1	5.6 - 14.5	1,000	59.2	51.3 - 66.6	300	21.1	15.4 - 28.3	200	10.7	6.5 - 16.9	1,600

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	6.2	4.1 - 9.2	2,400	65.6	60.5 - 70.4	600	16.2	12.7 - 20.4	400	12.0	9.0 - 15.8	3,700
Native Hawaiian	200	5.1	3.4 - 7.8	2,100	43.8	38.9 - 48.7	1,300	28.1	23.8 - 32.7	1,100	23.0	19.1 - 27.5	4,800
Chinese	100	12.1	8.6 - 16.9	400	69.4	63.2 - 74.9	100	12.7	9.0 - 17.6	<50	5.8	3.4 - 9.6	600
Filipino	200	6.9	4.7 - 10.0	2,100	67.3	62.3 - 72.0	500	16.4	12.9 - 20.6	300	9.4	6.8 - 12.9	3,100
Japanese	200	8.5	5.5 - 12.8	1,400	63.7	57.4 - 69.6	400	18.0	13.6 - 23.4	200	9.8	6.6 - 14.4	2,300
Other	100	4.8	2.8 - 8.3	1,100	47.8	41.5 - 54.1	500	22.7	17.7 - 28.5	600	24.7	19.5 - 30.7	2,400

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.9	5.2 - 14.8	900	64.8	56.2 - 72.5	300	18.7	12.7 - 26.5	100	7.6	4.0 - 14.0	1,500
20-24	300	6.2	4.2 - 9.0	2,500	61.0	56.0 - 65.7	900	20.7	16.9 - 25.1	500	12.1	9.1 - 15.8	4,100
25-34	500	6.3	4.9 - 8.1	4,400	53.1	49.7 - 56.4	1,700	20.8	18.1 - 23.7	1,700	19.9	17.3 - 22.8	8,300
35+	200	5.4	3.4 - 8.6	1,700	59.3	53.8 - 64.6	600	21.1	16.8 - 26.1	400	14.2	10.6 - 18.6	2,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	500	6.4	5.0 - 8.2	4,900	57.3	53.9 - 60.6	1,800	21.2	18.5 - 24.1	1,300	15.1	12.8 - 17.7	8,500
Female	500	6.3	4.8 - 8.1	4,700	56.9	53.5 - 60.2	1,700	20.1	17.4 - 23.0	1,400	16.8	14.3 - 19.6	8,300

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	6.0	4.8 - 7.6	6,300	58.1	55.2 - 61.0	2,200	20.5	18.2 - 23.1	1,700	15.3	13.3 - 17.6	10,800
No	400	6.9	5.1 - 9.3	3,300	55.3	51.1 - 59.4	1,200	20.8	17.6 - 24.4	1,000	17.0	14.0 - 20.4	6,000

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	500	8.3	6.3 - 10.8	3,100	52.1	48.0 - 56.1	1,300	21.4	18.2 - 25.0	1,100	18.2	15.2 - 21.7	6,000
131-185%	100	4.8	2.6 - 8.9	1,000	53.3	45.9 - 60.6	500	26.1	20.1 - 33.2	300	15.8	11.0 - 22.1	1,800
186+%	400	5.5	4.1 - 7.2	4,700	61.1	57.6 - 64.4	1,500	19.3	16.7 - 22.2	1,100	14.2	11.9 - 16.9	7,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Income (2004-2011)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	8.1	5.4 - 11.8	1,500	54.5	48.4 - 60.4	500	19.1	14.7 - 24.4	500	18.4	14.0 - 23.7	2,800
\$10,000-\$24,999	300	8.4	5.9 - 11.7	1,900	55.3	50.0 - 60.5	800	22.2	18.0 - 27.0	500	14.1	10.8 - 18.3	3,500
\$25,000-\$49,999	200	5.0	3.3 - 7.6	2,100	53.5	48.5 - 58.4	800	21.1	17.3 - 25.5	800	20.3	16.6 - 24.7	4,000
>=\$50,000	300	5.3	3.8 - 7.4	3,300	61.7	57.6 - 65.6	1,100	20.7	17.5 - 24.3	700	12.3	9.8 - 15.4	5,300

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	7.7	4.0 - 14.3	700	61.4	52.3 - 69.8	200	17.1	11.1 - 25.2	200	13.9	8.6 - 21.5	1,200
Normal birth weight	1,000	6.3	5.2 - 7.5	8,800	56.7	54.3 - 59.2	3,300	20.9	18.9 - 23.0	2,500	16.1	14.3 - 18.1	15,600

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	5.0	2.4 - 10.0	700	52.0	43.4 - 60.6	400	27.5	20.4 - 36.0	200	15.5	10.1 - 23.0	1,400
No	1,000	6.4	5.3 - 7.7	8,600	57.5	55.0 - 59.9	3,000	20.2	18.2 - 22.3	2,400	16.0	14.2 - 18.0	15,000

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2005

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	8.0	5.6 - 11.3	1,800	51.9	46.5 - 57.2	900	25.8	21.3 - 30.8	500	14.3	10.9 - 18.6	3,600
No	700	5.8	4.7 - 7.2	7,500	58.3	55.6 - 61.0	2,500	19.5	17.4 - 21.7	2,100	16.4	14.5 - 18.6	12,900

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	7.3	5.6 - 9.4	3,800	53.4	49.6 - 57.2	1,600	22.4	19.4 - 25.8	1,200	16.9	14.2 - 20.0	7,100
No	500	5.5	4.2 - 7.0	5,700	60.8	57.7 - 63.8	1,800	18.9	16.5 - 21.6	1,400	14.8	12.7 - 17.3	9,300

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,200	6.7	5.7 - 7.9	9,200	54.0	51.8 - 56.1	3,900	22.8	21.0 - 24.7	2,800	16.5	14.9 - 18.3	17,100

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	200	8.7	5.7 - 13.3	1,100	51.5	45.1 - 57.9	500	24.2	19.1 - 30.2	300	15.5	11.4 - 20.8	2,200
Honolulu	800	6.7	5.6 - 8.1	6,600	53.6	51.0 - 56.1	2,900	23.1	21.0 - 25.4	2,000	16.5	14.6 - 18.6	12,400
Kauai	<50	2.9	1.0 - 8.2	500	61.3	51.1 - 70.7	200	20.0	13.0 - 29.4	100	15.7	9.5 - 25.0	800
Maui	100	5.8	3.4 - 9.8	1,000	56.3	49.3 - 63.0	300	19.7	14.7 - 25.9	300	18.2	13.3 - 24.3	1,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	4.1	2.7 - 6.4	2,300	59.3	54.8 - 63.7	900	23.4	19.7 - 27.5	500	13.2	10.4 - 16.6	3,900
Native Hawaiian	300	5.5	3.7 - 8.0	2,200	46.2	41.6 - 50.8	1,100	23.1	19.4 - 27.2	1,200	25.3	21.6 - 29.5	4,700
Chinese	100	12.2	9.5 - 15.5	400	67.9	63.5 - 72.0	100	12.8	10.1 - 16.1	<50	7.2	5.1 - 10.0	700
Filipino	300	9.5	7.1 - 12.6	1,900	60.0	55.3 - 64.4	700	21.8	18.2 - 25.9	300	8.8	6.5 - 11.8	3,200
Japanese	200	7.8	5.1 - 11.8	1,300	57.9	51.6 - 64.0	500	24.5	19.4 - 30.3	200	9.8	6.7 - 14.2	2,200
Other	200	7.3	4.7 - 11.1	1,100	45.5	39.6 - 51.5	600	23.8	19.0 - 29.4	600	23.4	18.6 - 29.1	2,500

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	10.5	6.3 - 17.0	800	63.4	54.7 - 71.4	200	17.8	12.0 - 25.7	100	8.3	4.5 - 14.6	1,200
20-24	300	6.9	4.9 - 9.7	2,500	57.3	52.8 - 61.8	900	21.7	18.2 - 25.7	600	14.0	11.1 - 17.6	4,300
25-34	600	7.0	5.7 - 8.8	4,200	51.1	48.0 - 54.2	2,000	24.3	21.6 - 27.1	1,500	17.6	15.3 - 20.2	8,300
35+	100	4.3	2.8 - 6.6	1,700	53.2	48.3 - 58.0	700	22.3	18.5 - 26.7	700	20.2	16.5 - 24.6	3,300

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	600	6.1	4.9 - 7.6	4,900	53.3	50.3 - 56.2	2,100	22.2	19.7 - 24.8	1,700	18.5	16.2 - 21.0	9,300
Female	600	7.5	5.9 - 9.4	4,300	54.8	51.5 - 58.0	1,800	23.5	20.9 - 26.4	1,100	14.2	12.0 - 16.7	7,800

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	6.5	5.4 - 7.9	6,400	53.3	50.7 - 55.9	2,700	23.0	20.8 - 25.2	2,100	17.2	15.3 - 19.3	12,000
No	400	7.2	5.3 - 9.7	2,900	55.5	51.3 - 59.6	1,200	22.4	19.0 - 26.1	800	14.9	12.1 - 18.2	5,100

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	6.6	5.0 - 8.7	3,100	50.6	46.8 - 54.4	1,400	22.3	19.3 - 25.7	1,300	20.5	17.5 - 23.8	6,200
131-185%	200	7.9	5.2 - 12.0	1,100	51.1	44.8 - 57.4	500	23.4	18.4 - 29.3	400	17.5	13.2 - 23.0	2,100
186+%	500	6.1	4.8 - 7.8	4,300	57.4	54.2 - 60.6	1,700	22.8	20.2 - 25.6	1,000	13.7	11.5 - 16.1	7,600

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Income (2004-2011)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	200	6.7	4.3 - 10.3	1,300	52.4	46.4 - 58.3	600	23.2	18.5 - 28.7	500	17.8	13.6 - 22.9	2,600
\$10,000-\$24,999	300	7.8	5.7 - 10.5	2,000	49.1	44.6 - 53.7	1,000	23.1	19.4 - 27.2	800	20.0	16.5 - 24.0	4,200
\$25,000-\$49,999	200	5.4	3.8 - 7.7	2,400	55.6	51.3 - 59.9	900	21.0	17.6 - 24.7	800	18.0	14.8 - 21.6	4,400
>=\$50,000	300	6.2	4.6 - 8.3	2,700	57.0	53.0 - 60.9	1,200	24.2	20.9 - 27.9	600	12.6	10.1 - 15.6	4,800

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	6.9	3.6 - 12.8	600	54.4	45.5 - 63.0	200	16.5	11.0 - 24.1	200	22.2	15.6 - 30.5	1,100
Normal birth weight	1,100	6.7	5.7 - 7.9	8,700	53.9	51.7 - 56.2	3,700	23.2	21.3 - 25.2	2,600	16.2	14.5 - 18.0	16,000

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	4.9	2.4 - 9.9	700	52.9	44.5 - 61.2	300	26.6	19.7 - 34.8	200	15.5	10.4 - 22.6	1,300
No	1,100	7.0	5.9 - 8.2	8,400	53.9	51.6 - 56.2	3,500	22.5	20.6 - 24.5	2,600	16.6	14.9 - 18.4	15,500

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2004

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	7.2	5.0 - 10.2	1,700	50.6	45.5 - 55.7	900	26.0	21.7 - 30.8	600	16.2	12.8 - 20.4	3,400
No	900	6.7	5.6 - 8.1	7,300	54.6	52.2 - 57.1	3,000	22.0	20.0 - 24.1	2,200	16.6	14.8 - 18.6	13,400

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	6.5	5.0 - 8.3	3,800	50.4	47.0 - 53.8	1,800	23.6	20.8 - 26.7	1,500	19.5	16.9 - 22.4	7,500
No	600	6.9	5.5 - 8.5	5,300	57.3	54.4 - 60.2	2,000	22.0	19.6 - 24.6	1,300	13.8	11.9 - 16.0	9,200

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,100	6.6	5.3 - 8.2	9,700	56.1	53.4 - 58.8	3,800	21.6	19.5 - 23.9	2,700	15.6	13.8 - 17.7	17,300

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	6.8	4.7 - 9.6	1,100	54.7	50.1 - 59.2	400	22.3	18.8 - 26.2	300	16.3	13.3 - 19.8	2,000
Honolulu	800	6.5	4.9 - 8.6	7,100	55.7	52.2 - 59.1	2,800	22.2	19.4 - 25.1	2,000	15.7	13.3 - 18.4	12,700
Kauai	100	8.8	5.3 - 14.3	500	62.1	54.7 - 69.0	200	18.0	13.0 - 24.4	100	11.1	7.3 - 16.5	800
Maui	100	6.2	4.1 - 9.2	1,100	58.1	53.2 - 62.8	400	19.0	15.4 - 23.2	300	16.7	13.5 - 20.4	1,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	5.4	3.3 - 8.8	2,400	63.6	57.3 - 69.4	700	17.4	13.1 - 22.7	500	13.6	9.7 - 18.8	3,800
Native Hawaiian	200	3.5	2.5 - 4.8	2,300	48.6	45.6 - 51.6	1,300	26.6	24.0 - 29.4	1,000	21.3	19.0 - 23.9	4,700
Chinese	100	8.7	2.9 - 23.2	400	60.4	44.5 - 74.3	200	23.2	12.5 - 38.9	100	7.8	2.6 - 20.7	700
Filipino	300	8.4	5.6 - 12.6	2,200	56.8	50.5 - 63.0	800	20.9	16.2 - 26.4	500	13.9	10.0 - 18.8	3,800
Japanese	200	10.3	5.8 - 17.7	1,200	61.9	52.6 - 70.4	400	18.4	12.3 - 26.6	200	9.4	5.1 - 16.6	2,000
Other	200	7.9	4.2 - 14.2	1,300	52.2	43.6 - 60.6	500	22.0	15.7 - 29.9	400	18.0	12.3 - 25.5	2,400

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.0	4.6 - 13.7	900	59.9	51.7 - 67.5	300	18.8	13.2 - 26.2	200	13.3	8.8 - 19.7	1,600
20-24	300	6.6	4.4 - 9.8	2,300	58.1	52.8 - 63.1	800	20.9	17.1 - 25.4	600	14.4	11.3 - 18.3	4,000
25-34	600	6.7	4.8 - 9.2	4,800	54.6	50.7 - 58.4	1,900	21.5	18.5 - 24.8	1,500	17.3	14.6 - 20.4	8,700
35+	200	5.5	3.1 - 9.6	1,700	56.2	49.4 - 62.8	700	24.5	19.1 - 30.9	400	13.8	9.7 - 19.1	3,000

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	7.9	6.0 - 10.3	5,100	55.8	52.1 - 59.4	2,000	21.5	18.7 - 24.6	1,400	14.8	12.5 - 17.5	9,200
Female	400	5.1	3.6 - 7.2	4,600	56.5	52.6 - 60.4	1,800	21.8	18.7 - 25.2	1,300	16.6	13.8 - 19.7	8,100

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	5.5	4.1 - 7.5	6,500	57.2	53.7 - 60.6	2,500	21.5	18.8 - 24.5	1,800	15.8	13.4 - 18.4	11,400
No	500	8.6	6.3 - 11.6	3,200	54.2	50.0 - 58.3	1,300	21.9	18.7 - 25.4	900	15.4	12.7 - 18.5	5,900

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	500	6.4	4.7 - 8.8	3,900	50.8	46.9 - 54.7	1,800	23.9	20.7 - 27.4	1,400	18.9	16.0 - 22.1	7,700
131-185%	200	6.4	3.5 - 11.3	1,300	51.3	44.1 - 58.4	600	25.6	19.9 - 32.3	400	16.7	11.9 - 22.9	2,500
186+%	200	4.2	2.5 - 6.8	3,600	64.7	59.8 - 69.3	1,100	20.2	16.6 - 24.5	600	10.9	8.2 - 14.3	5,500

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Income (2000-2003)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	400	9.2	6.4 - 13.1	2,100	51.2	45.9 - 56.4	1,000	23.1	19.0 - 27.8	700	16.5	13.1 - 20.6	4,200
\$15,001-\$25,000	100	5.3	2.6 - 10.3	1,200	53.8	46.5 - 60.9	500	22.0	16.7 - 28.4	400	18.9	13.9 - 25.2	2,200
\$25,001-\$40,000	200	5.5	3.1 - 9.5	1,600	50.8	44.4 - 57.2	800	23.4	18.4 - 29.3	700	20.3	15.5 - 26.1	3,200
>=\$40,001	300	4.6	3.0 - 7.1	4,100	62.4	58.0 - 66.7	1,400	21.4	18.0 - 25.3	800	11.5	9.0 - 14.6	6,600

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	5.8	2.3 - 13.8	800	64.5	54.5 - 73.4	200	19.7	13.2 - 28.2	100	10.1	5.8 - 17.2	1,200
Normal birth weight	1,100	6.6	5.3 - 8.3	9,000	55.5	52.7 - 58.3	3,500	21.8	19.6 - 24.2	2,600	16.0	14.1 - 18.2	16,100

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	8.1	4.3 - 14.5	900	54.1	45.7 - 62.3	300	20.0	14.3 - 27.3	300	17.8	12.3 - 25.0	1,600
No	1,000	6.5	5.1 - 8.2	8,700	56.4	53.6 - 59.3	3,400	21.9	19.7 - 24.4	2,300	15.1	13.2 - 17.2	15,400

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2003

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	8.8	5.9 - 13.1	1,800	50.3	44.5 - 56.0	700	20.9	16.6 - 25.9	700	20.0	15.9 - 24.9	3,500
No	800	6.2	4.8 - 8.0	7,700	58.1	55.0 - 61.1	2,900	22.0	19.5 - 24.6	1,800	13.8	11.9 - 16.0	13,300

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	6.3	4.5 - 8.6	3,700	51.2	47.2 - 55.1	1,700	22.8	19.7 - 26.2	1,400	19.7	16.8 - 23.0	7,300
No	700	6.9	5.1 - 9.1	5,900	59.6	55.9 - 63.2	2,100	21.1	18.2 - 24.2	1,200	12.5	10.3 - 15.1	9,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	800	5.1	4.0 - 6.4	9,800	59.6	56.8 - 62.4	3,700	22.2	20.0 - 24.7	2,100	13.0	11.3 - 15.0	16,500

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.6	3.8 - 8.2	1,100	57.8	53.3 - 62.2	400	22.0	18.5 - 26.0	300	14.6	11.7 - 18.0	1,900
Honolulu	500	4.0	2.8 - 5.8	7,300	60.4	56.8 - 63.9	2,700	22.4	19.4 - 25.6	1,600	13.2	10.9 - 15.7	12,100
Kauai	100	9.5	5.5 - 15.7	400	58.6	50.5 - 66.2	100	18.0	12.7 - 24.9	100	14.0	9.5 - 20.1	600
Maui	200	10.1	7.3 - 13.7	1,000	56.7	51.7 - 61.6	400	23.1	19.1 - 27.6	200	10.1	7.6 - 13.3	1,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	4.8	2.9 - 7.8	2,300	65.1	58.6 - 71.1	700	20.6	15.6 - 26.7	300	9.5	6.3 - 14.1	3,500
Native Hawaiian	200	4.7	3.6 - 6.2	2,300	51.9	48.8 - 55.0	1,100	23.9	21.3 - 26.6	900	19.5	17.2 - 22.0	4,500
Chinese	100	10.0	3.4 - 26.2	500	81.0	64.4 - 90.9	<50	5.7	1.6 - 18.0	<50	3.3	0.5 - 19.4	600
Filipino	200	4.8	2.9 - 7.8	2,200	64.9	58.1 - 71.2	700	21.1	16.0 - 27.3	300	9.2	5.7 - 14.5	3,300
Japanese	100	6.0	3.0 - 11.7	1,600	69.0	60.5 - 76.4	400	18.5	12.7 - 26.1	200	6.5	3.2 - 12.7	2,300
Other	100	4.3	1.7 - 10.4	1,000	43.4	34.5 - 52.6	700	31.6	23.6 - 40.9	500	20.7	14.2 - 29.1	2,200

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	100	8.5	4.9 - 14.5	900	60.8	51.8 - 69.2	300	18.4	12.3 - 26.5	200	12.3	7.0 - 20.5	1,500
20-24	300	6.2	4.2 - 9.2	2,400	58.4	53.0 - 63.6	900	22.1	18.0 - 26.9	500	13.2	10.0 - 17.4	4,100
25-34	400	4.4	3.1 - 6.3	4,800	58.5	54.4 - 62.4	1,900	23.3	19.9 - 27.1	1,100	13.8	11.3 - 16.7	8,300
35+	100	3.4	1.6 - 7.3	1,700	64.6	57.7 - 70.9	600	21.3	16.2 - 27.5	300	10.7	7.3 - 15.3	2,600

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	500	5.6	4.1 - 7.7	4,800	58.8	54.9 - 62.6	1,600	20.1	17.1 - 23.4	1,300	15.5	12.9 - 18.5	8,200
Female	400	4.5	3.2 - 6.3	5,000	60.4	56.4 - 64.3	2,000	24.4	21.0 - 28.1	900	10.6	8.5 - 13.2	8,300

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	4.4	3.2 - 6.0	6,900	61.9	58.3 - 65.2	2,400	21.1	18.3 - 24.2	1,400	12.7	10.6 - 15.1	11,200
No	300	6.6	4.8 - 9.1	2,900	54.9	50.4 - 59.4	1,300	24.7	21.0 - 28.9	700	13.7	10.9 - 17.3	5,300

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	300	5.4	3.9 - 7.4	3,400	53.5	49.2 - 57.7	1,400	22.6	19.1 - 26.5	1,200	18.5	15.4 - 22.2	6,400
131-185%	200	7.0	4.2 - 11.5	2,000	59.1	52.6 - 65.2	700	21.2	16.5 - 26.8	400	12.7	9.1 - 17.5	3,400
186+%	200	3.4	2.0 - 5.6	3,500	67.0	61.9 - 71.7	1,200	21.9	17.8 - 26.7	400	7.7	5.5 - 10.7	5,300

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Income (2000-2003)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	200	7.2	4.7 - 10.9	1,700	52.3	46.6 - 58.0	700	21.2	16.8 - 26.4	600	19.3	15.0 - 24.4	3,300
\$15,001-\$25,000	100	4.0	2.6 - 6.3	1,400	59.7	52.4 - 66.6	500	19.8	14.6 - 26.4	400	16.4	11.8 - 22.4	2,400
\$25,001-\$40,000	200	6.4	3.8 - 10.5	2,000	54.8	48.5 - 61.0	1,000	26.7	21.5 - 32.7	400	12.0	8.6 - 16.5	3,600
>=\$40,001	200	3.8	2.4 - 5.9	4,200	65.9	61.3 - 70.2	1,300	20.1	16.5 - 24.2	600	10.2	7.8 - 13.3	6,300

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	8.8	4.1 - 17.7	700	62.5	51.3 - 72.5	200	17.7	11.0 - 27.2	100	11.1	5.8 - 20.4	1,100
Normal birth weight	700	4.8	3.8 - 6.1	9,100	59.5	56.6 - 62.3	3,400	22.5	20.1 - 25.1	2,000	13.2	11.4 - 15.2	15,200

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	5.7	2.9 - 10.8	700	50.5	41.4 - 59.4	300	26.5	19.0 - 35.5	200	17.4	11.9 - 24.8	1,300
No	800	5.1	4.0 - 6.5	9,000	60.3	57.4 - 63.2	3,300	22.0	19.6 - 24.7	1,900	12.5	10.7 - 14.6	14,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2002

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	9.0	5.7 - 13.8	1,700	53.9	47.8 - 60.0	700	22.2	17.6 - 27.6	500	14.9	10.9 - 20.0	3,200
No	500	4.2	3.3 - 5.5	7,900	61.2	58.0 - 64.3	2,800	22.1	19.4 - 24.9	1,600	12.6	10.6 - 14.8	12,800

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	6.9	5.0 - 9.4	3,800	55.2	50.9 - 59.4	1,500	22.4	19.0 - 26.2	1,100	15.6	12.8 - 18.9	6,800
No	400	3.8	2.7 - 5.3	6,000	63.0	59.3 - 66.6	2,100	21.8	18.8 - 25.1	1,100	11.3	9.2 - 13.9	9,600

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,200	7.2	6.1 - 8.5	9,900	60.8	58.5 - 62.9	3,100	18.8	17.1 - 20.6	2,100	13.2	11.8 - 14.7	16,200

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	7.3	5.2 - 10.3	1,100	61.4	56.7 - 65.8	300	17.4	14.1 - 21.2	300	13.9	11.0 - 17.5	1,800
Honolulu	900	7.1	5.7 - 8.9	7,300	60.6	57.8 - 63.3	2,300	19.0	16.9 - 21.4	1,600	13.2	11.5 - 15.2	12,000
Kauai	100	9.6	5.9 - 15.1	400	55.4	47.6 - 62.9	200	21.2	15.5 - 28.2	100	13.8	9.4 - 20.0	700
Maui	100	6.7	4.5 - 9.8	1,100	63.8	58.9 - 68.5	300	17.3	13.9 - 21.3	200	12.1	9.3 - 15.6	1,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	6.1	4.1 - 8.9	2,400	69.1	64.2 - 73.5	500	14.5	11.3 - 18.5	400	10.3	7.7 - 13.8	3,500
Native Hawaiian	200	4.7	3.8 - 5.9	2,400	53.6	51.1 - 56.0	1,000	22.7	20.7 - 24.8	800	19.0	17.1 - 21.0	4,400
Chinese	100	19.9	11.2 - 32.9	400	60.7	46.9 - 73.0	100	12.0	5.6 - 23.9	100	7.4	2.7 - 18.3	700
Filipino	200	6.2	4.1 - 9.3	2,300	69.3	63.9 - 74.2	600	17.3	13.4 - 22.0	200	7.2	4.8 - 10.7	3,400
Japanese	200	9.1	5.5 - 14.6	1,300	65.1	57.7 - 71.9	300	17.0	12.1 - 23.4	200	8.9	5.6 - 13.7	2,000
Other	200	9.8	6.2 - 15.3	1,000	45.4	38.5 - 52.6	500	23.7	17.9 - 30.6	500	21.0	15.9 - 27.4	2,300

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	200	9.9	6.9 - 14.2	1,100	67.3	61.4 - 72.6	200	15.4	11.6 - 20.2	100	7.3	4.7 - 11.2	1,600
20-24	400	8.6	6.3 - 11.7	2,400	57.7	53.3 - 61.9	800	19.3	16.1 - 23.0	600	14.4	11.7 - 17.7	4,200
25-34	500	6.0	4.5 - 8.0	4,800	61.7	58.5 - 64.9	1,400	17.9	15.5 - 20.6	1,100	14.4	12.4 - 16.6	7,800
35+	200	6.8	4.4 - 10.4	1,600	59.1	53.2 - 64.9	600	22.6	17.9 - 28.1	300	11.5	8.4 - 15.5	2,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	600	7.5	6.0 - 9.4	5,000	60.7	57.5 - 63.8	1,600	18.8	16.4 - 21.5	1,100	13.0	11.1 - 15.0	8,200
Female	600	6.9	5.3 - 8.9	4,900	60.8	57.7 - 63.9	1,500	18.8	16.5 - 21.3	1,100	13.5	11.5 - 15.8	8,000

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	6.8	5.5 - 8.5	6,700	61.9	59.1 - 64.6	1,900	17.9	15.8 - 20.2	1,500	13.4	11.7 - 15.3	10,900
No	400	8.0	6.1 - 10.4	3,100	58.5	54.9 - 62.0	1,100	20.7	17.9 - 23.8	700	12.9	10.7 - 15.4	5,400

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	400	7.1	5.4 - 9.1	3,400	55.0	51.5 - 58.4	1,200	20.3	17.7 - 23.3	1,100	17.6	15.2 - 20.4	6,100
131-185%	200	5.2	3.4 - 7.9	2,100	60.7	55.9 - 65.2	700	19.8	16.3 - 23.8	500	14.3	11.4 - 17.9	3,500
186+%	400	7.2	5.2 - 10.0	3,600	67.6	63.5 - 71.4	900	16.7	13.8 - 20.1	500	8.5	6.6 - 10.8	5,400

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Income (2000-2003)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	300	10.3	7.6 - 13.7	1,800	54.1	49.3 - 58.7	600	19.8	16.2 - 24.0	500	15.9	12.8 - 19.6	3,300
\$15,001-\$25,000	100	4.2	2.4 - 7.3	1,300	57.2	51.3 - 62.9	500	21.1	16.7 - 26.3	400	17.5	13.5 - 22.3	2,200
\$25,001-\$40,000	200	5.0	3.2 - 7.6	2,100	59.2	54.5 - 63.7	700	20.9	17.3 - 24.9	500	15.0	12.1 - 18.5	3,500
>=\$40,001	500	7.2	5.3 - 9.6	4,200	66.5	62.8 - 70.0	1,100	16.6	14.0 - 19.7	600	9.7	7.9 - 12.0	6,400

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	200	15.6	9.9 - 23.6	600	56.0	47.1 - 64.6	200	19.4	13.1 - 27.7	100	9.1	5.5 - 14.5	1,100
Normal birth weight	1,000	6.6	5.5 - 8.0	9,200	61.1	58.8 - 63.4	2,800	18.8	17.0 - 20.6	2,000	13.5	12.1 - 15.1	15,100

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	6.6	4.5 - 9.6	900	62.2	55.7 - 68.3	300	17.5	13.0 - 23.1	200	13.6	9.9 - 18.4	1,500
No	1,000	7.2	5.9 - 8.6	8,700	60.5	58.1 - 62.9	2,700	19.0	17.2 - 21.0	1,900	13.3	11.8 - 15.0	14,400

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2001

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	200	6.8	4.8 - 9.5	1,900	58.4	53.8 - 62.9	700	19.9	16.5 - 23.9	500	14.9	12.0 - 18.4	3,300
No	900	7.2	5.8 - 8.8	7,600	61.4	58.8 - 64.0	2,300	18.5	16.5 - 20.7	1,600	12.9	11.3 - 14.6	12,400

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	8.1	6.3 - 10.3	3,900	56.9	53.6 - 60.1	1,300	19.6	17.1 - 22.3	1,100	15.4	13.3 - 17.9	6,800
No	600	6.4	5.0 - 8.2	5,900	63.7	60.6 - 66.7	1,700	18.2	15.9 - 20.8	1,100	11.7	9.9 - 13.7	9,300

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

State	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Totals	1,400	8.4	7.2 - 9.8	9,700	59.1	56.9 - 61.3	3,300	20.0	18.3 - 21.8	2,100	12.6	11.2 - 14.1	16,500

County	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Hawaii	100	5.3	3.5 - 7.8	1,200	61.4	56.8 - 65.9	400	21.7	18.1 - 25.8	200	11.6	9.0 - 14.9	1,900
Honolulu	1,100	8.9	7.3 - 10.7	7,000	57.6	54.8 - 60.3	2,500	20.4	18.3 - 22.7	1,600	13.2	11.5 - 15.1	12,200
Kauai	<50	6.1	3.1 - 11.7	400	68.6	60.7 - 75.6	100	17.8	12.3 - 24.9	<50	7.5	4.5 - 12.4	700
Maui	200	9.4	6.8 - 13.1	1,100	63.6	58.6 - 68.4	300	15.9	12.5 - 20.0	200	11.0	8.3 - 14.6	1,700

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

DOH Race-Ethnicity	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	200	6.2	4.3 - 8.9	2,600	65.5	60.9 - 69.9	700	18.1	14.8 - 22.0	400	10.1	7.4 - 13.6	4,000
Native Hawaiian	200	5.4	4.4 - 6.7	2,300	51.2	48.7 - 53.7	1,100	23.5	21.4 - 25.7	900	19.9	17.9 - 22.0	4,500
Chinese	100	15.8	7.5 - 30.1	400	72.5	57.9 - 83.5	<50	2.5	0.4 - 14.6	100	9.3	3.8 - 21.0	600
Filipino	400	11.0	7.8 - 15.3	2,100	62.9	57.2 - 68.2	600	18.4	14.5 - 23.2	300	7.7	5.0 - 11.4	3,300
Japanese	300	13.1	8.8 - 19.1	1,200	59.7	52.0 - 66.8	400	20.5	14.9 - 27.5	100	6.7	3.8 - 11.7	2,000
Other	200	8.4	5.1 - 13.5	1,100	53.6	46.2 - 60.8	500	22.6	17.1 - 29.4	300	15.4	10.7 - 21.5	2,100

PRAMS Age Group	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	300	14.2	10.1 - 19.5	1,200	62.8	56.7 - 68.5	300	16.0	12.3 - 20.7	100	7.0	4.4 - 10.8	1,800
20-24	300	6.9	5.1 - 9.3	2,500	62.1	58.0 - 66.2	700	18.9	15.9 - 22.4	500	12.0	9.6 - 15.1	4,000
25-34	600	8.3	6.4 - 10.6	4,400	55.9	52.5 - 59.2	1,600	21.0	18.5 - 23.9	1,200	14.8	12.7 - 17.2	7,800
35+	200	7.0	4.5 - 10.7	1,800	61.2	55.6 - 66.5	600	21.0	16.8 - 25.9	300	10.8	7.8 - 14.6	2,900

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Infant Gender	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	700	8.7	7.0 - 10.8	4,900	58.2	55.2 - 61.2	1,700	20.6	18.2 - 23.1	1,100	12.5	10.6 - 14.6	8,500
Female	600	8.1	6.4 - 10.2	4,800	60.0	56.8 - 63.1	1,500	19.3	16.9 - 22.0	1,000	12.7	10.7 - 14.8	8,000

Married	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	7.6	6.2 - 9.4	6,800	59.5	56.8 - 62.3	2,300	20.3	18.2 - 22.6	1,400	12.5	10.8 - 14.4	11,300
No	500	10.1	7.9 - 12.8	3,000	58.0	54.4 - 61.6	1,000	19.2	16.6 - 22.1	600	12.7	10.5 - 15.2	5,100

Poverty Level	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	600	9.4	7.5 - 11.6	3,700	53.7	50.4 - 57.0	1,400	20.4	17.9 - 23.1	1,100	16.5	14.3 - 19.1	6,900
131-185%	200	6.9	4.2 - 11.2	1,300	56.9	51.1 - 62.6	500	22.5	18.0 - 27.6	300	13.7	10.3 - 18.0	2,200
186+%	500	7.6	5.6 - 10.1	4,000	65.4	61.6 - 69.0	1,100	18.8	16.0 - 22.0	500	8.2	6.3 - 10.6	6,100

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Income (2000-2003)	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	300	8.4	6.3 - 11.1	2,100	56.1	51.7 - 60.4	700	18.4	15.3 - 22.1	600	17.1	14.0 - 20.7	3,800
\$15,001-\$25,000	300	11.8	8.2 - 16.8	1,300	51.6	46.0 - 57.1	500	21.5	17.4 - 26.4	400	15.0	11.6 - 19.3	2,500
\$25,001-\$40,000	200	7.2	4.9 - 10.4	1,800	56.6	51.7 - 61.5	700	21.5	17.7 - 25.8	500	14.6	11.5 - 18.5	3,200
>=\$40,001	400	6.8	5.0 - 9.2	4,000	65.0	61.2 - 68.6	1,200	19.8	16.9 - 23.0	500	8.4	6.6 - 10.7	6,200

Birthweight	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	100	10.8	6.6 - 17.2	600	57.4	48.6 - 65.8	200	23.0	16.3 - 31.3	100	8.8	5.0 - 15.0	1,100
Normal birth weight	1,300	8.3	7.0 - 9.8	9,000	59.3	57.0 - 61.5	3,000	19.8	18.0 - 21.7	1,900	12.7	11.2 - 14.2	15,200

Smoke 3rd Trimester	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	100	7.8	4.4 - 13.3	700	49.9	42.7 - 57.1	300	18.5	13.9 - 24.3	300	23.8	17.8 - 31.0	1,400
No	1,200	8.5	7.2 - 10.1	8,700	59.8	57.5 - 62.1	2,900	20.2	18.3 - 22.1	1,700	11.5	10.1 - 13.0	14,600

PRAMS Health Indicator Report - Weight - BMI status (PRAMS)- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown

Year: 2000

Smoke Before Pregnant	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	300	9.0	6.4 - 12.6	1,700	53.9	49.1 - 58.6	600	20.0	16.6 - 23.9	500	17.1	13.7 - 21.0	3,200
No	1,000	8.2	6.8 - 9.8	7,600	60.2	57.7 - 62.7	2,500	20.1	18.1 - 22.2	1,500	11.5	10.0 - 13.2	12,600

WIC During Pregnancy	Underweight (BMI < 18.5)			Recommended range (18.5 <= BMI < 25)			Overweight (25 <= BMI < 30)			Obese (BMI>=30)			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	7.9	6.3 - 10.0	4,000	55.9	52.6 - 59.0	1,500	21.1	18.6 - 23.8	1,100	15.1	13.0 - 17.6	7,100
No	800	8.7	7.0 - 10.7	5,600	61.5	58.5 - 64.5	1,700	19.0	16.7 - 21.6	1,000	10.8	9.1 - 12.8	9,100