

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

*The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.*

*Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.*

*Below are the indicators included in this report and the long description for each indicator.*

PRAMS Health Indicator	Health Indicator Description
Took multivitamins/folic acid month prior to pregnancy	Took multivitamins/folic acid month prior to pregnancy

*Suggested citation:*

*Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/13/17*

*For documentation on other terms and coding in this report, please go to the following links:*

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact  
The Hawai'i Health Data Warehouse**

[profiles@hhdw.org](mailto:profiles@hhdw.org)

**(855) 946-5899 x15**


**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

Year	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	9,400	54.8	52.6 - 57.0	1,600	9.3	8.0 - 10.7	6,200	35.9	33.8 - 38.1	17,200
2001	10,000	59.9	57.7 - 62.1	1,300	7.6	6.5 - 9.0	5,400	32.4	30.4 - 34.6	16,700
2002	9,900	57.7	54.9 - 60.4	1,400	7.9	6.5 - 9.6	5,900	34.4	31.8 - 37.1	17,100
2003	10,400	58.8	56.1 - 61.4	1,300	7.5	6.2 - 9.1	5,900	33.7	31.1 - 36.3	17,600
2004	9,300	52.2	50.1 - 54.4	1,500	8.5	7.4 - 9.8	7,000	39.3	37.2 - 41.4	17,800
2005	9,800	56.0	53.7 - 58.3	1,300	7.2	6.1 - 8.5	6,500	36.8	34.6 - 39.0	17,500
2006	10,000	54.1	51.7 - 56.5	1,900	10.1	8.7 - 11.7	6,600	35.8	33.5 - 38.1	18,500
2007	10,900	58.3	55.9 - 60.5	1,500	8.3	7.0 - 9.7	6,300	33.5	31.4 - 35.7	18,800
2008	10,200	53.7	51.4 - 56.0	1,700	8.7	7.5 - 10.1	7,200	37.6	35.4 - 39.9	19,000
2009	10,900	59.8	56.7 - 62.9	1,200	6.5	5.1 - 8.2	6,100	33.7	30.8 - 36.7	18,300
2010	10,500	57.1	53.9 - 60.3	1,700	9.1	7.4 - 11.2	6,200	33.7	30.8 - 36.8	18,400
2011	10,700	57.7	54.5 - 60.9	1,500	7.9	6.2 - 9.9	6,400	34.4	31.5 - 37.6	18,500
2012	10,500	58.1	54.8 - 61.3	1,500	8.3	6.6 - 10.3	6,100	33.7	30.6 - 36.8	18,100
2013	10,400	57.6	54.1 - 61.0	1,300	7.0	5.4 - 9.1	6,400	35.4	32.1 - 38.8	18,000

# PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,400	57.6	54.1 - 61.0	1,300	7.0	5.4 - 9.1	6,400	35.4	32.1 - 38.8	18,000

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	67.7	62.3 - 72.6	200	6.6	4.2 - 10.0	600	25.8	21.3 - 30.9	2,300
<b>Honolulu</b>	7,200	55.5	50.7 - 60.1	900	7.3	5.2 - 10.2	4,800	37.2	32.8 - 41.9	12,900
<b>Kauai</b>	500	59.7	54.3 - 64.8	<50	4.3	2.6 - 7.2	300	36.0	31.0 - 41.3	800
<b>Maui</b>	1,100	58.9	53.3 - 64.3	100	6.8	4.5 - 10.2	700	34.3	29.2 - 39.8	1,900

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,200	46.3	39.4 - 53.2	500	9.8	6.2 - 15.1	2,100	43.9	37.2 - 50.9	4,800
Native Hawaiian	3,800	71.7	65.8 - 76.9	400	7.6	4.9 - 11.6	1,100	20.7	16.1 - 26.2	5,400
Chinese	500	51.7	35.5 - 67.5	<50	3.7	0.7 - 18.6	400	44.5	29.1 - 61.1	900
Filipino	1,600	54.1	45.4 - 62.6	100	1.9	0.6 - 5.6	1,300	44.0	35.6 - 52.8	2,900
Japanese	900	53.2	41.4 - 64.7	100	5.0	1.8 - 12.9	700	41.8	30.8 - 53.8	1,700
Other	1,300	58.5	48.2 - 68.1	200	9.1	4.4 - 17.9	700	32.4	23.7 - 42.6	2,300

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	77.8	60.7 - 88.8	100	6.3	1.7 - 21.2	100	15.9	6.8 - 32.9	900
20-24	2,800	71.1	63.5 - 77.6	300	8.2	4.7 - 13.9	800	20.8	15.1 - 27.8	4,000
25-34	5,300	53.6	48.8 - 58.4	700	7.1	5.0 - 10.1	3,800	39.3	34.7 - 44.1	9,800
35+	1,600	48.0	40.3 - 55.8	200	5.5	2.9 - 10.4	1,600	46.5	38.8 - 54.3	3,400

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,100	57.6	52.6 - 62.4	700	7.7	5.4 - 11.0	3,100	34.7	30.1 - 39.6	8,800
<b>Female</b>	5,300	57.6	52.6 - 62.5	600	6.3	4.3 - 9.3	3,300	36.1	31.4 - 41.0	9,200

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,600	49.4	44.8 - 53.9	900	7.8	5.6 - 10.8	4,900	42.8	38.4 - 47.3	11,400
<b>No</b>	4,700	71.7	66.5 - 76.4	400	5.6	3.6 - 8.8	1,500	22.6	18.3 - 27.6	6,600

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,600	70.1	64.6 - 75.1	500	7.4	4.8 - 11.2	1,500	22.5	18.1 - 27.6	6,600
<b>131-185%</b>	1,600	66.6	57.0 - 74.9	200	8.4	4.0 - 16.8	600	25.1	18.0 - 33.8	2,400
<b>186+%</b>	3,100	43.2	37.8 - 48.9	500	7.4	4.9 - 11.0	3,600	49.4	43.8 - 55.0	7,200

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

Income (2012+)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	3,200	77.3	70.8 - 82.6	200	5.3	2.8 - 9.7	700	17.4	12.8 - 23.3	4,200
\$17,001-\$26,000	1,400	58.6	48.8 - 67.8	300	12.8	7.3 - 21.5	700	28.6	20.6 - 38.1	2,400
\$26,001-\$51,000	2,400	63.3	55.7 - 70.2	200	6.0	3.2 - 11.0	1,200	30.7	24.3 - 38.0	3,800
>=\$51,001	2,500	40.7	34.8 - 46.7	500	7.7	4.9 - 11.8	3,200	51.6	45.5 - 57.7	6,100

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	55.3	50.2 - 60.3	100	4.3	2.7 - 6.9	500	40.4	35.5 - 45.5	1,300
Normal birth weight	9,600	57.8	54.0 - 61.5	1,200	7.2	5.5 - 9.5	5,800	35.0	31.4 - 38.7	16,700

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	84.1	72.4 - 91.5	<50	1.0	0.3 - 3.8	100	14.9	7.7 - 26.7	1,000
No	9,400	56.3	52.7 - 60.0	1,200	7.0	5.3 - 9.2	6,100	36.7	33.2 - 40.3	16,600

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,300	70.3	62.3 - 77.3	200	7.5	4.1 - 13.4	700	22.2	16.1 - 29.7	3,200
<b>No</b>	7,900	55.1	51.2 - 59.0	900	6.5	4.8 - 8.8	5,500	38.3	34.6 - 42.2	14,400

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,300	67.0	61.8 - 71.8	500	6.4	4.2 - 9.7	2,100	26.6	22.2 - 31.6	7,900
<b>No</b>	5,100	50.4	45.6 - 55.2	700	7.0	4.9 - 10.0	4,300	42.6	37.9 - 47.4	10,000



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,500	58.1	54.8 - 61.3	1,500	8.3	6.6 - 10.3	6,100	33.7	30.6 - 36.8	18,100

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,400	61.1	55.8 - 66.1	200	8.4	5.9 - 11.8	700	30.6	26.0 - 35.6	2,300
<b>Honolulu</b>	7,500	57.6	53.2 - 61.8	1,100	8.5	6.4 - 11.3	4,400	33.9	29.9 - 38.1	13,100
<b>Kauai</b>	500	61.6	55.3 - 67.6	100	7.5	4.7 - 11.6	300	30.9	25.4 - 37.1	800
<b>Maui</b>	1,100	56.5	51.5 - 61.5	100	6.6	4.6 - 9.5	700	36.8	32.1 - 41.8	1,900

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	43.0	36.4 - 49.8	400	10.8	7.2 - 15.8	1,900	46.3	39.7 - 53.0	4,100
Native Hawaiian	3,400	67.4	61.5 - 72.8	300	6.8	4.3 - 10.5	1,300	25.8	20.9 - 31.4	5,000
Chinese	600	60.6	46.4 - 73.2	100	4.8	1.4 - 15.8	400	34.6	22.7 - 48.8	1,100
Filipino	1,900	56.3	48.5 - 63.8	200	7.3	4.2 - 12.7	1,200	36.3	29.3 - 44.1	3,300
Japanese	800	51.2	40.0 - 62.2	100	6.7	2.9 - 14.7	700	42.2	31.6 - 53.6	1,600
Other	2,100	67.7	59.5 - 75.0	300	10.3	6.2 - 16.8	700	21.9	15.8 - 29.5	3,000

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	88.7	79.3 - 94.2	<50	3.9	0.8 - 16.2	100	7.4	3.9 - 13.3	1,000
20-24	2,500	68.4	61.0 - 75.0	300	7.6	4.3 - 13.0	900	24.0	18.2 - 31.0	3,600
25-34	5,600	54.7	50.3 - 59.0	1,000	9.6	7.3 - 12.5	3,600	35.7	31.7 - 40.0	10,200
35+	1,500	47.3	39.8 - 54.9	200	6.3	3.7 - 10.8	1,500	46.4	38.9 - 54.0	3,300

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,100	56.9	52.2 - 61.5	600	6.6	4.7 - 9.3	3,300	36.5	32.1 - 41.1	8,900
<b>Female</b>	5,400	59.2	54.6 - 63.6	900	9.9	7.4 - 13.1	2,800	30.9	26.9 - 35.2	9,100

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,600	50.4	46.2 - 54.6	1,100	9.6	7.3 - 12.5	4,400	40.0	36.0 - 44.2	11,100
<b>No</b>	4,900	70.3	65.4 - 74.8	400	6.2	4.2 - 9.0	1,600	23.5	19.4 - 28.2	7,000

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,500	70.9	65.7 - 75.7	400	5.7	3.6 - 8.9	1,500	23.4	19.0 - 28.3	6,400
<b>131-185%</b>	2,100	67.8	59.9 - 74.8	300	8.5	4.9 - 14.3	700	23.7	17.7 - 31.0	3,000
<b>186+%</b>	2,900	41.5	36.4 - 46.8	700	10.6	7.7 - 14.3	3,300	47.9	42.7 - 53.2	6,900

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

Income (2012+)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	3,000	73.3	66.6 - 79.0	300	7.7	4.6 - 12.6	800	19.0	14.1 - 25.1	4,100
\$17,001-\$26,000	1,500	66.9	57.8 - 74.9	100	4.4	2.1 - 9.0	600	28.7	21.1 - 37.7	2,200
\$26,001-\$51,000	2,700	62.9	56.3 - 69.2	400	8.2	5.3 - 12.6	1,200	28.9	23.2 - 35.3	4,300
>=\$51,001	2,400	41.2	35.6 - 47.0	600	10.5	7.4 - 14.7	2,900	48.3	42.6 - 54.0	5,900

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	53.8	48.2 - 59.3	100	6.4	4.1 - 9.8	500	39.8	34.5 - 45.4	1,200
Normal birth weight	9,800	58.4	54.9 - 61.8	1,400	8.4	6.7 - 10.6	5,600	33.2	30.0 - 36.5	16,800

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	500	65.6	49.1 - 79.0	100	13.8	5.6 - 30.2	100	20.6	10.3 - 36.9	700
No	9,800	58.4	55.1 - 61.8	1,300	7.5	5.9 - 9.5	5,700	34.0	30.9 - 37.3	16,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,000	71.1	63.1 - 78.0	300	9.1	5.1 - 15.6	600	19.8	14.1 - 27.0	2,800
<b>No</b>	8,300	56.4	52.8 - 60.0	1,100	7.5	5.9 - 9.7	5,300	36.0	32.6 - 39.6	14,700

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,700	68.7	64.0 - 73.1	600	7.7	5.4 - 10.9	1,900	23.5	19.7 - 27.9	8,200
<b>No</b>	4,800	49.0	44.5 - 53.5	900	8.8	6.6 - 11.7	4,100	42.2	37.9 - 46.7	9,700

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,700	57.7	54.5 - 60.9	1,500	7.9	6.2 - 9.9	6,400	34.4	31.5 - 37.6	18,500

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	63.4	58.4 - 68.2	100	5.9	4.0 - 8.8	700	30.6	26.2 - 35.5	2,400
<b>Honolulu</b>	7,500	56.2	51.9 - 60.4	1,100	8.5	6.3 - 11.3	4,700	35.3	31.3 - 39.5	13,300
<b>Kauai</b>	500	61.0	54.8 - 66.8	<50	5.3	3.1 - 8.9	300	33.7	28.2 - 39.8	900
<b>Maui</b>	1,200	59.3	54.7 - 63.7	100	7.1	5.1 - 9.9	700	33.6	29.4 - 38.0	2,000

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,000	44.7	38.4 - 51.1	600	12.9	9.1 - 18.1	1,900	42.4	36.2 - 48.8	4,500
Native Hawaiian	3,900	74.7	69.4 - 79.4	300	4.8	2.9 - 7.9	1,100	20.5	16.3 - 25.5	5,300
Chinese	300	44.6	29.2 - 61.1	100	17.0	7.6 - 33.8	300	38.4	23.8 - 55.4	700
Filipino	1,800	53.1	45.6 - 60.4	100	4.1	2.0 - 8.2	1,400	42.8	35.6 - 50.3	3,300
Japanese	900	47.0	36.9 - 57.3	100	4.1	1.5 - 10.5	900	48.9	38.8 - 59.2	1,900
Other	1,700	62.9	53.9 - 71.1	300	10.1	5.6 - 17.6	700	26.9	19.8 - 35.5	2,800

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	81.1	67.8 - 89.8	<50	1.4	0.4 - 5.0	200	17.5	9.0 - 31.1	1,100
20-24	3,100	71.2	64.6 - 77.0	500	10.5	6.8 - 15.9	800	18.3	13.7 - 23.9	4,400
25-34	5,300	53.7	49.2 - 58.0	800	8.0	5.9 - 10.7	3,800	38.4	34.2 - 42.7	9,900
35+	1,400	43.8	36.5 - 51.3	200	6.0	3.2 - 11.0	1,600	50.2	42.7 - 57.7	3,200

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,300	56.1	51.5 - 60.5	800	8.3	6.0 - 11.4	3,400	35.6	31.4 - 40.1	9,500
<b>Female</b>	5,400	59.4	54.9 - 63.8	700	7.4	5.3 - 10.2	3,000	33.2	29.1 - 37.6	9,100

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,700	49.7	45.6 - 53.8	1,000	8.6	6.5 - 11.3	4,800	41.7	37.8 - 45.8	11,500
<b>No</b>	5,000	71.0	66.0 - 75.5	500	6.7	4.5 - 9.7	1,600	22.4	18.3 - 27.1	7,000

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,600	69.5	64.3 - 74.3	400	6.4	4.1 - 9.7	1,600	24.1	19.8 - 29.0	6,600
<b>131-185%</b>	1,900	67.9	59.6 - 75.1	300	10.2	6.0 - 16.9	600	21.9	15.9 - 29.3	2,800
<b>186+%</b>	3,600	44.6	39.7 - 49.5	600	8.0	5.7 - 11.1	3,900	47.5	42.6 - 52.4	8,100



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

Income (2004-2011)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,500	75.0	67.6 - 81.1	200	4.7	2.3 - 9.6	700	20.3	14.7 - 27.4	3,400
\$10,000-\$24,999	2,300	66.9	59.7 - 73.4	300	7.7	4.4 - 13.0	900	25.4	19.7 - 32.1	3,400
\$25,000-\$49,999	2,900	63.8	57.3 - 69.8	400	8.0	5.1 - 12.3	1,300	28.2	22.7 - 34.4	4,600
>=\$50,000	2,400	38.9	33.6 - 44.5	600	9.1	6.3 - 13.0	3,200	52.0	46.4 - 57.5	6,200

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	57.4	51.9 - 62.6	100	5.3	3.3 - 8.2	500	37.4	32.3 - 42.7	1,300
Normal birth weight	9,900	57.6	54.2 - 61.0	1,400	8.1	6.3 - 10.2	5,900	34.3	31.1 - 37.6	17,200

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	600	68.6	55.8 - 79.1	100	9.8	4.4 - 20.5	200	21.6	13.0 - 33.7	900
No	9,900	56.9	53.6 - 60.2	1,400	7.8	6.1 - 9.9	6,200	35.3	32.2 - 38.5	17,400

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,600	74.0	67.2 - 79.9	300	7.4	4.4 - 12.2	700	18.6	13.6 - 24.9	3,500
<b>No</b>	7,900	53.6	49.9 - 57.1	1,200	8.0	6.2 - 10.3	5,700	38.4	35.0 - 42.0	14,800

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,900	71.4	66.9 - 75.5	500	6.3	4.2 - 9.2	1,900	22.3	18.6 - 26.5	8,300
<b>No</b>	4,700	46.4	42.1 - 50.8	900	9.2	6.9 - 12.2	4,500	44.3	40.0 - 48.7	10,100

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,500	57.1	53.9 - 60.3	1,700	9.1	7.4 - 11.2	6,200	33.7	30.8 - 36.8	18,400

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	63.4	58.4 - 68.2	200	7.7	5.4 - 10.9	700	28.9	24.5 - 33.7	2,300
<b>Honolulu</b>	7,400	55.7	51.5 - 59.9	1,300	9.4	7.2 - 12.2	4,600	34.9	31.0 - 39.0	13,300
<b>Kauai</b>	500	63.7	57.8 - 69.1	100	7.9	5.3 - 11.7	200	28.4	23.4 - 34.0	800
<b>Maui</b>	1,100	56.5	51.7 - 61.2	200	9.4	6.9 - 12.6	700	34.1	29.7 - 38.8	2,000

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,100	47.6	41.2 - 54.1	600	12.2	8.6 - 17.2	1,800	40.1	34.0 - 46.6	4,500
Native Hawaiian	3,600	65.9	60.3 - 71.0	500	8.7	5.9 - 12.5	1,400	25.5	20.8 - 30.8	5,500
Chinese	400	46.6	31.2 - 62.8	100	6.6	2.3 - 17.6	400	46.8	31.4 - 62.8	800
Filipino	1,900	58.7	51.2 - 65.8	300	9.9	6.3 - 15.4	1,000	31.3	24.9 - 38.5	3,200
Japanese	800	45.4	35.2 - 56.1	<50	2.9	1.0 - 8.1	900	51.7	41.2 - 62.1	1,700
Other	1,700	63.4	54.5 - 71.5	200	8.6	4.4 - 16.0	800	28.0	20.9 - 36.4	2,700

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	87.7	78.3 - 93.3	<50	3.6	1.8 - 7.0	100	8.7	3.8 - 18.7	1,100
20-24	3,300	72.8	66.5 - 78.4	200	5.4	3.1 - 9.3	1,000	21.7	16.7 - 27.8	4,500
25-34	4,900	51.9	47.5 - 56.3	1,000	10.6	8.0 - 13.8	3,600	37.5	33.4 - 41.9	9,500
35+	1,400	40.4	33.3 - 48.0	400	11.9	7.9 - 17.6	1,600	47.7	40.3 - 55.1	3,300

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,300	56.7	52.3 - 61.1	900	9.9	7.5 - 12.9	3,100	33.4	29.4 - 37.7	9,300
<b>Female</b>	5,300	57.5	52.9 - 62.0	800	8.4	6.1 - 11.3	3,100	34.1	29.9 - 38.5	9,100

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,600	48.3	44.2 - 52.5	1,100	9.8	7.6 - 12.6	4,800	41.8	37.8 - 45.9	11,500
<b>No</b>	5,000	71.7	66.9 - 76.0	600	8.0	5.6 - 11.3	1,400	20.4	16.6 - 24.7	6,900

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,400	68.7	63.6 - 73.5	500	8.2	5.6 - 11.8	1,500	23.1	18.9 - 27.8	6,400
<b>131-185%</b>	2,000	67.2	59.4 - 74.1	200	7.5	4.6 - 12.0	800	25.4	19.0 - 32.9	3,000
<b>186+%</b>	3,200	41.4	36.6 - 46.3	900	11.9	9.0 - 15.7	3,600	46.7	41.8 - 51.7	7,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

Income (2004-2011)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,500	72.5	65.4 - 78.6	200	6.2	3.4 - 10.9	700	21.3	15.9 - 28.0	3,400
\$10,000-\$24,999	2,200	67.0	59.7 - 73.6	400	11.0	6.9 - 17.1	700	22.0	16.7 - 28.3	3,200
\$25,000-\$49,999	2,900	63.3	57.0 - 69.3	300	7.3	4.8 - 10.8	1,300	29.4	23.9 - 35.6	4,600
>=\$50,000	2,300	36.7	31.5 - 42.2	800	12.5	9.1 - 16.9	3,200	50.8	45.2 - 56.4	6,200

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	55.2	50.0 - 60.4	100	8.0	5.6 - 11.3	500	36.7	31.8 - 41.9	1,300
Normal birth weight	9,800	57.4	54.0 - 60.7	1,600	9.2	7.4 - 11.5	5,700	33.4	30.2 - 36.7	17,100

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	71.9	60.4 - 81.1	100	7.5	3.1 - 17.0	300	20.6	12.9 - 31.3	1,400
No	9,400	56.0	52.6 - 59.3	1,500	9.2	7.4 - 11.3	5,800	34.8	31.7 - 38.1	16,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,700	70.1	63.5 - 76.0	300	7.7	4.8 - 12.0	800	22.2	16.9 - 28.5	3,800
<b>No</b>	7,800	53.8	50.2 - 57.4	1,400	9.4	7.5 - 11.8	5,300	36.8	33.4 - 40.3	14,400

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,400	67.8	63.3 - 72.1	700	8.5	6.1 - 11.7	1,900	23.7	20.0 - 27.8	8,000
<b>No</b>	5,000	49.1	44.7 - 53.5	1,000	9.5	7.2 - 12.4	4,200	41.4	37.2 - 45.7	10,200

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,900	59.8	56.7 - 62.9	1,200	6.5	5.1 - 8.2	6,100	33.7	30.8 - 36.7	18,300

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	63.8	58.9 - 68.5	200	7.2	5.0 - 10.2	700	29.0	24.7 - 33.7	2,400
<b>Honolulu</b>	7,500	58.7	54.5 - 62.8	800	6.3	4.5 - 8.7	4,500	35.0	31.1 - 39.1	12,900
<b>Kauai</b>	500	61.0	55.1 - 66.6	<50	5.7	3.5 - 9.1	300	33.3	28.0 - 39.1	900
<b>Maui</b>	1,300	61.5	57.0 - 65.9	200	7.2	5.2 - 10.0	700	31.2	27.2 - 35.6	2,100



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	44.2	38.1 - 50.5	400	9.2	6.1 - 13.8	2,000	46.5	40.4 - 52.8	4,200
Native Hawaiian	4,300	72.4	67.3 - 77.1	200	4.2	2.7 - 6.5	1,400	23.4	19.0 - 28.4	5,900
Chinese	400	55.0	39.0 - 70.1	<50	0.4	0.1 - 2.4	400	44.5	29.5 - 60.6	800
Filipino	1,900	59.7	52.3 - 66.7	300	8.0	4.6 - 13.4	1,000	32.3	25.9 - 39.4	3,200
Japanese	900	56.4	45.8 - 66.4	100	7.8	3.6 - 16.2	600	35.8	26.5 - 46.3	1,600
Other	1,500	60.1	51.2 - 68.4	200	6.5	3.4 - 12.2	800	33.4	25.6 - 42.2	2,500

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,500	80.1	71.0 - 86.8	100	4.1	1.7 - 9.5	300	15.9	9.8 - 24.6	1,900
20-24	3,300	77.4	71.5 - 82.4	200	3.6	2.0 - 6.6	800	19.0	14.3 - 24.6	4,200
25-34	5,000	54.5	50.1 - 58.9	700	7.6	5.5 - 10.2	3,500	37.9	33.7 - 42.3	9,200
35+	1,200	38.9	31.9 - 46.4	300	8.8	5.2 - 14.4	1,600	52.3	44.8 - 59.7	3,000

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,600	56.4	52.1 - 60.7	700	7.0	5.1 - 9.6	3,600	36.5	32.5 - 40.8	9,900
<b>Female</b>	5,400	63.8	59.4 - 68.1	500	5.9	4.1 - 8.3	2,500	30.3	26.3 - 34.6	8,400

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,200	48.8	44.7 - 52.8	800	7.3	5.4 - 9.7	4,700	44.0	40.0 - 48.1	10,800
<b>No</b>	5,700	75.7	71.4 - 79.6	400	5.4	3.7 - 7.9	1,400	18.9	15.4 - 22.9	7,500

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,800	74.3	69.5 - 78.6	400	5.7	3.6 - 8.8	1,300	20.0	16.2 - 24.4	6,500
<b>131-185%</b>	1,900	66.3	58.3 - 73.5	300	9.8	6.0 - 15.6	700	23.9	17.6 - 31.6	2,800
<b>186+%</b>	3,200	44.3	39.5 - 49.2	500	6.5	4.5 - 9.3	3,600	49.2	44.4 - 54.1	7,300

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

Income (2004-2011)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,500	79.1	72.4 - 84.4	100	2.2	0.9 - 5.5	600	18.7	13.6 - 25.1	3,100
\$10,000-\$24,999	2,600	67.8	61.2 - 73.7	400	10.3	6.8 - 15.4	800	21.9	16.9 - 27.8	3,900
\$25,000-\$49,999	2,400	65.4	58.6 - 71.6	300	7.0	4.2 - 11.5	1,000	27.5	21.8 - 34.1	3,700
>=\$50,000	2,500	41.4	36.2 - 46.9	400	6.4	4.2 - 9.6	3,100	52.2	46.7 - 57.6	6,000

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	60.3	55.1 - 65.2	100	6.6	4.5 - 9.6	400	33.1	28.5 - 38.1	1,400
Normal birth weight	10,000	59.7	56.3 - 62.9	1,100	6.5	5.0 - 8.4	5,700	33.8	30.7 - 37.1	16,800

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	78.9	69.2 - 86.1	<50	1.1	0.4 - 3.3	300	20.0	12.9 - 29.7	1,700
No	9,300	57.8	54.4 - 61.0	1,200	7.2	5.6 - 9.1	5,600	35.1	32.0 - 38.3	16,100

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,000	74.3	68.1 - 79.6	200	4.6	2.5 - 8.3	800	21.1	16.2 - 27.0	4,000
<b>No</b>	7,600	55.6	51.9 - 59.1	1,000	7.2	5.5 - 9.3	5,100	37.3	33.8 - 40.8	13,800

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,300	71.1	66.6 - 75.3	500	6.7	4.6 - 9.7	1,700	22.2	18.5 - 26.4	7,400
<b>No</b>	5,300	51.3	47.1 - 55.5	700	6.5	4.8 - 8.8	4,300	42.1	38.0 - 46.3	10,200

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,200	53.7	51.4 - 56.0	1,700	8.7	7.5 - 10.1	7,200	37.6	35.4 - 39.9	19,000

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,400	55.5	48.7 - 62.0	200	7.4	4.5 - 11.8	900	37.2	31.0 - 43.8	2,500
<b>Honolulu</b>	7,000	52.4	49.6 - 55.1	1,300	9.5	8.0 - 11.3	5,100	38.1	35.5 - 40.8	13,400
<b>Kauai</b>	500	55.7	45.0 - 65.9	100	8.7	4.2 - 17.1	300	35.6	26.2 - 46.2	900
<b>Maui</b>	1,300	58.7	51.8 - 65.3	100	5.1	2.8 - 9.1	800	36.2	29.9 - 43.0	2,300

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,000	44.7	40.0 - 49.5	500	10.3	7.7 - 13.5	2,000	45.0	40.3 - 49.8	4,500
Native Hawaiian	3,400	63.7	59.0 - 68.2	500	9.1	6.7 - 12.3	1,400	27.2	23.1 - 31.6	5,300
Chinese	200	36.3	30.2 - 43.0	100	12.0	8.4 - 17.0	300	51.6	45.0 - 58.2	600
Filipino	1,700	48.7	43.6 - 53.8	200	6.6	4.5 - 9.7	1,600	44.7	39.6 - 49.8	3,500
Japanese	1,100	53.6	46.7 - 60.3	100	5.5	3.1 - 9.5	800	41.0	34.4 - 47.8	2,100
Other	1,800	59.0	52.9 - 64.7	300	9.4	6.5 - 13.6	1,000	31.6	26.3 - 37.4	3,000

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	69.4	60.6 - 76.9	100	7.9	4.3 - 14.3	300	22.7	16.1 - 31.0	1,500
20-24	2,700	64.3	59.2 - 69.0	400	8.8	6.2 - 12.3	1,100	26.9	22.6 - 31.7	4,100
25-34	5,200	52.6	49.4 - 55.8	900	9.0	7.3 - 11.0	3,800	38.4	35.4 - 41.6	9,900
35+	1,300	37.6	32.7 - 42.7	300	8.0	5.7 - 11.3	1,900	54.4	49.2 - 59.5	3,500

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,400	53.5	50.3 - 56.7	700	7.4	5.9 - 9.3	3,900	39.1	36.0 - 42.2	10,000
<b>Female</b>	4,900	53.9	50.5 - 57.3	900	10.1	8.2 - 12.3	3,300	36.0	32.9 - 39.3	9,000

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,600	47.1	44.2 - 50.0	1,100	9.0	7.5 - 10.7	5,200	43.9	41.1 - 46.8	11,900
<b>No</b>	4,600	64.8	60.9 - 68.5	600	8.2	6.2 - 10.7	1,900	27.1	23.7 - 30.7	7,100

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,100	58.4	54.4 - 62.2	600	9.1	7.0 - 11.7	2,300	32.6	29.0 - 36.4	7,100
<b>131-185%</b>	1,800	66.3	60.2 - 71.8	200	6.8	4.4 - 10.5	700	26.9	21.8 - 32.7	2,700
<b>186+%</b>	3,500	44.4	41.0 - 47.9	700	9.3	7.5 - 11.5	3,700	46.3	42.9 - 49.7	8,000

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

Income (2004-2011)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,900	59.2	53.2 - 64.9	200	7.2	4.6 - 11.0	1,100	33.6	28.3 - 39.5	3,200
\$10,000-\$24,999	2,400	59.5	54.4 - 64.5	400	9.6	6.9 - 13.2	1,300	30.8	26.3 - 35.8	4,100
\$25,000-\$49,999	2,500	58.0	53.1 - 62.6	400	9.9	7.4 - 13.1	1,400	32.1	27.8 - 36.8	4,300
>=\$50,000	2,700	43.1	39.3 - 47.0	500	8.6	6.6 - 11.0	3,000	48.3	44.4 - 52.1	6,300

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	54.3	45.4 - 63.0	100	4.6	2.0 - 10.3	500	41.1	32.7 - 50.0	1,300
Normal birth weight	9,500	53.6	51.2 - 56.0	1,600	9.0	7.7 - 10.5	6,600	37.4	35.2 - 39.8	17,700

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	65.3	56.8 - 72.9	100	7.6	4.2 - 13.4	400	27.1	20.2 - 35.3	1,600
No	8,900	52.6	50.1 - 55.0	1,400	8.5	7.2 - 10.0	6,600	38.9	36.6 - 41.3	17,000



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,500	69.6	64.2 - 74.5	200	5.5	3.4 - 8.6	900	25.0	20.4 - 30.1	3,500
<b>No</b>	7,500	50.0	47.4 - 52.5	1,400	9.1	7.7 - 10.7	6,200	40.9	38.4 - 43.5	15,100

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,000	62.1	58.4 - 65.6	700	8.0	6.2 - 10.3	2,400	29.9	26.6 - 33.4	8,100
<b>No</b>	5,100	47.7	44.7 - 50.7	1,000	9.2	7.6 - 11.1	4,600	43.1	40.2 - 46.1	10,600

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,900	58.3	55.9 - 60.5	1,500	8.3	7.0 - 9.7	6,300	33.5	31.4 - 35.7	18,800

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	65.4	59.0 - 71.3	200	7.8	5.0 - 12.1	700	26.8	21.4 - 32.9	2,500
<b>Honolulu</b>	7,500	57.9	55.2 - 60.6	1,100	8.2	6.7 - 9.9	4,400	33.9	31.4 - 36.5	13,000
<b>Kauai</b>	600	61.3	51.2 - 70.5	100	6.4	3.0 - 13.3	300	32.3	23.7 - 42.2	1,100
<b>Maui</b>	1,100	50.5	43.4 - 57.5	200	10.2	6.6 - 15.3	900	39.4	32.7 - 46.4	2,200

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,200	51.4	46.5 - 56.3	300	8.1	5.8 - 11.3	1,700	40.5	35.8 - 45.4	4,200
Native Hawaiian	3,500	67.5	62.8 - 71.8	400	8.5	6.2 - 11.7	1,200	24.0	20.1 - 28.4	5,200
Chinese	300	46.0	39.5 - 52.7	<50	4.2	2.2 - 7.8	300	49.8	43.2 - 56.4	700
Filipino	2,000	58.3	53.3 - 63.2	200	6.1	4.1 - 9.1	1,200	35.6	30.9 - 40.5	3,500
Japanese	1,000	47.6	40.9 - 54.5	200	7.6	4.7 - 12.1	1,000	44.8	38.1 - 51.6	2,100
Other	1,900	62.2	56.2 - 67.9	400	11.8	8.3 - 16.4	800	26.0	21.1 - 31.6	3,000

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	75.2	66.8 - 82.1	100	7.7	4.1 - 14.0	300	17.1	11.4 - 24.9	1,700
20-24	3,300	70.4	65.7 - 74.6	500	9.9	7.3 - 13.3	900	19.7	16.2 - 23.8	4,700
25-34	5,100	55.3	52.0 - 58.5	700	7.7	6.1 - 9.7	3,400	37.0	33.9 - 40.2	9,100
35+	1,300	39.9	34.8 - 45.3	200	7.6	5.2 - 11.1	1,700	52.5	47.1 - 57.8	3,200

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,700	58.6	55.3 - 61.7	800	7.9	6.3 - 9.9	3,200	33.6	30.6 - 36.7	9,700
<b>Female</b>	5,300	57.9	54.6 - 61.2	800	8.7	6.9 - 10.8	3,000	33.4	30.4 - 36.6	9,100

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,000	51.4	48.6 - 54.3	1,000	8.9	7.3 - 10.7	4,600	39.7	36.9 - 42.5	11,600
<b>No</b>	5,000	69.4	65.5 - 72.9	500	7.2	5.4 - 9.7	1,700	23.4	20.2 - 27.0	7,100

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,800	68.3	64.4 - 71.9	500	7.3	5.5 - 9.8	1,700	24.4	21.1 - 28.0	7,000
<b>131-185%</b>	1,800	63.8	57.9 - 69.3	300	9.4	6.4 - 13.5	700	26.8	21.9 - 32.4	2,800
<b>186+%</b>	3,500	46.8	43.3 - 50.3	700	8.9	7.1 - 11.3	3,400	44.3	40.8 - 47.8	7,600

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

Income (2004-2011)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	64.9	58.8 - 70.5	300	8.8	5.8 - 13.1	800	26.3	21.2 - 32.0	3,200
\$10,000-\$24,999	2,800	71.0	66.0 - 75.4	200	5.6	3.6 - 8.4	900	23.5	19.4 - 28.1	3,900
\$25,000-\$49,999	2,500	59.6	54.8 - 64.3	400	8.2	5.9 - 11.3	1,400	32.1	27.8 - 36.7	4,300
>=\$50,000	2,800	46.0	42.1 - 49.9	600	9.9	7.7 - 12.7	2,700	44.1	40.3 - 48.0	6,100

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	58.7	50.6 - 66.4	100	7.0	3.8 - 12.3	500	34.3	27.1 - 42.3	1,600
Normal birth weight	10,000	58.2	55.8 - 60.6	1,400	8.4	7.1 - 9.9	5,700	33.4	31.2 - 35.7	17,200

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	79.3	71.5 - 85.4	100	6.6	3.4 - 12.4	200	14.1	9.2 - 21.1	1,500
No	9,600	56.5	54.0 - 58.9	1,400	8.4	7.1 - 10.0	5,900	35.1	32.8 - 37.4	16,900

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,600	72.0	66.8 - 76.6	300	7.7	5.2 - 11.2	700	20.4	16.4 - 25.0	3,700
<b>No</b>	8,100	55.0	52.4 - 57.5	1,300	8.5	7.1 - 10.1	5,400	36.6	34.1 - 39.1	14,800

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,200	67.2	63.5 - 70.6	400	5.5	4.0 - 7.5	2,100	27.3	24.1 - 30.8	7,800
<b>No</b>	5,400	51.6	48.6 - 54.6	1,100	10.1	8.3 - 12.1	4,000	38.3	35.4 - 41.3	10,500

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted  
total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,000	54.1	51.7 - 56.5	1,900	10.1	8.7 - 11.7	6,600	35.8	33.5 - 38.1	18,500

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,400	62.8	55.6 - 69.4	200	10.4	6.8 - 15.7	600	26.8	20.9 - 33.6	2,200
<b>Honolulu</b>	7,100	52.3	49.5 - 55.1	1,500	10.9	9.2 - 12.7	5,000	36.8	34.2 - 39.5	13,500
<b>Kauai</b>	500	56.6	45.1 - 67.4	100	7.4	3.2 - 16.3	300	35.9	25.9 - 47.4	900
<b>Maui</b>	1,100	55.7	48.2 - 63.0	100	5.6	3.0 - 10.2	800	38.7	31.7 - 46.2	2,000

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,100	47.2	42.3 - 52.2	500	11.3	8.5 - 14.8	1,800	41.5	36.7 - 46.5	4,400
Native Hawaiian	3,300	65.5	60.6 - 70.1	500	9.6	7.0 - 13.0	1,300	24.9	20.8 - 29.5	5,100
Chinese	300	41.4	34.8 - 48.4	100	11.8	8.1 - 17.1	300	46.7	39.9 - 53.6	600
Filipino	1,600	47.4	42.1 - 52.8	300	10.1	7.3 - 13.8	1,400	42.5	37.3 - 47.8	3,400
Japanese	1,000	45.8	39.1 - 52.7	200	8.8	5.6 - 13.5	1,000	45.4	38.7 - 52.3	2,200
Other	1,800	61.5	55.2 - 67.4	300	9.8	6.6 - 14.4	800	28.7	23.4 - 34.6	2,900

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	68.8	59.9 - 76.5	100	8.6	4.7 - 15.4	400	22.6	16.0 - 31.0	1,600
20-24	3,000	66.2	61.2 - 70.9	500	10.6	7.8 - 14.2	1,000	23.2	19.2 - 27.8	4,500
25-34	4,800	52.6	49.2 - 56.0	900	9.5	7.7 - 11.7	3,400	37.9	34.7 - 41.2	9,100
35+	1,200	35.1	30.2 - 40.4	400	11.7	8.7 - 15.6	1,800	53.1	47.8 - 58.4	3,400



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,000	53.1	49.7 - 56.5	1,000	10.3	8.4 - 12.5	3,500	36.6	33.4 - 39.9	9,500
<b>Female</b>	5,000	55.2	51.7 - 58.6	900	9.9	8.0 - 12.2	3,200	34.9	31.7 - 38.2	9,100

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,700	48.8	45.9 - 51.7	1,200	10.3	8.7 - 12.2	4,800	40.9	38.1 - 43.8	11,700
<b>No</b>	4,300	63.3	59.2 - 67.3	700	9.7	7.5 - 12.5	1,800	26.9	23.4 - 30.9	6,800

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,200	64.6	60.4 - 68.6	500	8.1	6.1 - 10.8	1,800	27.3	23.7 - 31.2	6,500
<b>131-185%</b>	1,300	55.0	48.2 - 61.6	200	7.7	4.8 - 12.0	900	37.3	31.0 - 44.0	2,500
<b>186+%</b>	3,700	44.7	41.3 - 48.2	1,000	12.1	10.0 - 14.5	3,600	43.2	39.8 - 46.6	8,300

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

Income (2004-2011)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	64.3	58.1 - 70.0	300	9.3	6.2 - 13.6	800	26.5	21.4 - 32.3	3,100
\$10,000-\$24,999	2,400	63.3	57.9 - 68.4	300	7.5	5.1 - 10.9	1,100	29.2	24.5 - 34.4	3,800
\$25,000-\$49,999	2,500	57.1	52.1 - 62.0	400	8.7	6.3 - 12.0	1,500	34.2	29.6 - 39.1	4,300
>=\$50,000	2,400	39.6	35.7 - 43.6	800	12.9	10.4 - 15.9	2,900	47.5	43.5 - 51.5	6,000

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	50.6	42.3 - 58.9	100	7.0	3.8 - 12.5	700	42.4	34.4 - 50.7	1,600
Normal birth weight	9,200	54.3	51.8 - 56.8	1,800	10.4	9.0 - 12.1	6,000	35.3	32.9 - 37.7	16,900

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	75.4	67.6 - 81.9	100	7.3	3.9 - 13.2	300	17.3	11.9 - 24.4	1,700
No	8,600	51.9	49.4 - 54.5	1,700	10.3	8.9 - 12.0	6,200	37.7	35.3 - 40.2	16,500

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,900	74.1	69.0 - 78.6	300	8.3	5.7 - 11.9	700	17.6	13.9 - 22.1	3,900
<b>No</b>	7,000	48.8	46.1 - 51.5	1,500	10.5	9.0 - 12.3	5,800	40.7	38.1 - 43.4	14,400

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,900	63.4	59.6 - 67.0	700	9.3	7.2 - 11.8	2,100	27.4	24.0 - 30.9	7,800
<b>No</b>	5,000	47.4	44.2 - 50.5	1,100	10.7	8.9 - 12.8	4,400	41.9	38.9 - 45.0	10,500

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	9,800	56.0	53.7 - 58.3	1,300	7.2	6.1 - 8.5	6,500	36.8	34.6 - 39.0	17,500

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,400	62.5	55.8 - 68.8	200	9.5	6.2 - 14.1	600	28.1	22.4 - 34.5	2,300
<b>Honolulu</b>	7,100	55.1	52.4 - 57.7	900	7.1	5.9 - 8.6	4,800	37.8	35.2 - 40.4	12,800
<b>Kauai</b>	400	53.1	41.5 - 64.4	100	9.9	4.8 - 19.4	300	37.0	26.6 - 48.8	700
<b>Maui</b>	1,000	55.6	48.0 - 63.0	100	3.7	1.7 - 8.1	700	40.7	33.5 - 48.2	1,700

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	49.2	44.1 - 54.4	300	8.6	6.2 - 12.0	1,600	42.1	37.2 - 47.2	3,800
Native Hawaiian	3,300	66.4	61.7 - 70.8	300	5.5	3.7 - 8.2	1,400	28.1	23.9 - 32.6	5,000
Chinese	300	40.3	34.3 - 46.7	100	10.8	7.5 - 15.4	300	48.9	42.6 - 55.2	700
Filipino	1,700	53.1	48.1 - 58.1	300	7.8	5.5 - 11.0	1,300	39.0	34.3 - 44.1	3,300
Japanese	1,100	48.9	42.6 - 55.2	200	6.7	4.2 - 10.6	1,000	44.4	38.3 - 50.7	2,300
Other	1,500	59.8	53.7 - 65.7	200	7.2	4.7 - 10.9	800	33.0	27.5 - 39.0	2,600

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	68.5	60.2 - 75.8	100	8.5	4.8 - 14.5	300	23.0	16.6 - 30.8	1,500
20-24	3,000	68.4	63.7 - 72.8	300	6.3	4.3 - 9.2	1,100	25.3	21.3 - 29.7	4,300
25-34	4,700	54.1	50.8 - 57.4	700	7.7	6.2 - 9.6	3,300	38.2	35.0 - 41.4	8,700
35+	1,100	37.0	31.9 - 42.4	200	6.5	4.3 - 9.7	1,700	56.5	51.0 - 61.8	3,000

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	4,900	54.6	51.3 - 57.8	700	8.3	6.7 - 10.3	3,300	37.1	34.0 - 40.3	8,900
<b>Female</b>	5,000	57.4	54.1 - 60.7	500	6.1	4.7 - 7.9	3,100	36.4	33.3 - 39.7	8,600

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,400	48.3	45.4 - 51.2	1,000	8.7	7.2 - 10.5	4,800	43.0	40.2 - 45.9	11,200
<b>No</b>	4,400	69.5	65.6 - 73.0	300	4.6	3.2 - 6.6	1,700	26.0	22.6 - 29.6	6,400

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,100	65.3	61.5 - 69.0	400	6.2	4.6 - 8.4	1,800	28.4	25.0 - 32.1	6,300
<b>131-185%</b>	1,200	62.6	55.3 - 69.3	100	6.7	3.9 - 11.2	600	30.7	24.4 - 37.8	1,900
<b>186+%</b>	3,600	46.3	42.9 - 49.8	700	8.5	6.7 - 10.6	3,500	45.2	41.9 - 48.7	7,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

Income (2004-2011)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,000	65.5	59.8 - 70.8	200	5.2	3.2 - 8.4	900	29.3	24.3 - 34.8	3,000
\$10,000-\$24,999	2,400	65.0	60.0 - 69.7	300	7.7	5.4 - 10.9	1,000	27.2	22.9 - 32.0	3,700
\$25,000-\$49,999	2,300	57.9	53.0 - 62.7	300	7.1	5.0 - 9.9	1,400	35.1	30.5 - 39.9	4,000
>=\$50,000	2,400	43.4	39.4 - 47.5	500	8.4	6.4 - 10.9	2,600	48.2	44.2 - 52.3	5,400

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	57.7	49.1 - 65.9	100	5.7	2.8 - 11.3	500	36.6	28.8 - 45.1	1,300
Normal birth weight	9,100	55.9	53.4 - 58.2	1,200	7.4	6.2 - 8.7	6,000	36.8	34.5 - 39.2	16,200

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	75.5	67.6 - 82.0	100	6.0	3.1 - 11.5	300	18.5	12.8 - 25.9	1,400
No	8,500	54.0	51.5 - 56.4	1,200	7.4	6.2 - 8.8	6,100	38.6	36.2 - 41.0	15,700

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,600	71.8	66.8 - 76.3	200	5.1	3.2 - 8.1	800	23.1	19.0 - 27.9	3,600
<b>No</b>	6,900	51.4	48.7 - 54.0	1,100	7.9	6.6 - 9.4	5,500	40.7	38.2 - 43.3	13,500

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,900	65.4	61.8 - 68.8	500	6.8	5.2 - 8.8	2,100	27.8	24.7 - 31.2	7,500
<b>No</b>	4,700	48.8	45.7 - 51.9	700	7.6	6.2 - 9.4	4,200	43.6	40.5 - 46.7	9,700



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted  
total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	9,300	52.2	50.1 - 54.4	1,500	8.5	7.4 - 9.8	7,000	39.3	37.2 - 41.4	17,800

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,400	60.9	54.6 - 66.9	100	6.5	4.0 - 10.2	700	32.6	27.0 - 38.7	2,300
<b>Honolulu</b>	6,600	50.9	48.4 - 53.4	1,200	9.0	7.6 - 10.5	5,200	40.2	37.7 - 42.6	13,000
<b>Kauai</b>	500	57.9	47.8 - 67.3	100	6.4	3.0 - 13.1	300	35.7	26.8 - 45.7	800
<b>Maui</b>	900	48.5	41.8 - 55.3	200	8.8	5.6 - 13.5	800	42.7	36.2 - 49.5	1,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	45.2	40.7 - 49.7	400	9.8	7.4 - 12.9	1,800	45.0	40.6 - 49.6	3,900
Native Hawaiian	3,100	64.1	59.6 - 68.3	300	6.8	4.9 - 9.4	1,400	29.1	25.2 - 33.4	4,900
Chinese	300	40.2	35.9 - 44.7	100	7.6	5.6 - 10.4	400	52.1	47.6 - 56.6	700
Filipino	1,600	47.3	42.8 - 51.9	300	9.1	6.8 - 12.1	1,500	43.6	39.1 - 48.1	3,300
Japanese	1,000	43.3	37.3 - 49.5	200	7.8	5.1 - 11.8	1,100	48.9	42.8 - 55.1	2,300
Other	1,500	57.9	52.0 - 63.5	300	9.8	6.9 - 13.7	900	32.4	27.1 - 38.1	2,700

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	74.0	66.0 - 80.7	100	8.9	5.1 - 15.1	200	17.1	11.7 - 24.3	1,300
20-24	2,900	65.3	61.0 - 69.4	400	9.0	6.8 - 11.8	1,200	25.7	22.0 - 29.8	4,500
25-34	4,200	48.3	45.3 - 51.4	700	8.4	6.8 - 10.2	3,800	43.3	40.3 - 46.4	8,700
35+	1,300	36.7	32.2 - 41.5	300	8.1	5.9 - 11.1	1,900	55.1	50.3 - 59.9	3,400

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,000	51.1	48.2 - 54.0	900	8.8	7.3 - 10.6	3,900	40.1	37.2 - 42.9	9,800
<b>Female</b>	4,300	53.5	50.3 - 56.7	700	8.1	6.6 - 10.0	3,100	38.3	35.3 - 41.4	8,100

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,700	46.0	43.5 - 48.6	1,100	9.0	7.7 - 10.6	5,600	44.9	42.4 - 47.5	12,500
<b>No</b>	3,600	66.6	62.6 - 70.3	400	7.3	5.5 - 9.7	1,400	26.1	22.7 - 29.9	5,400

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,100	62.8	59.2 - 66.3	500	6.9	5.3 - 9.0	2,000	30.3	27.0 - 33.7	6,600
<b>131-185%</b>	1,200	55.6	49.3 - 61.7	200	10.2	7.0 - 14.6	700	34.2	28.5 - 40.4	2,100
<b>186+%</b>	3,200	41.8	38.7 - 44.9	700	9.5	7.8 - 11.5	3,800	48.7	45.6 - 51.9	7,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

Income (2004-2011)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	1,800	66.2	60.6 - 71.4	100	4.6	2.7 - 7.6	800	29.2	24.2 - 34.7	2,700
\$10,000-\$24,999	2,600	60.3	55.9 - 64.5	400	9.4	7.1 - 12.4	1,300	30.3	26.4 - 34.5	4,400
\$25,000-\$49,999	2,300	51.9	47.6 - 56.2	400	9.3	7.1 - 12.0	1,700	38.8	34.7 - 43.1	4,500
>=\$50,000	1,800	36.7	32.9 - 40.5	500	9.3	7.2 - 11.8	2,700	54.1	50.1 - 58.0	4,900

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	43.6	35.2 - 52.3	100	6.6	3.4 - 12.5	600	49.8	41.3 - 58.4	1,100
Normal birth weight	8,800	52.8	50.6 - 55.0	1,400	8.6	7.4 - 9.9	6,400	38.6	36.4 - 40.7	16,700

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	65.5	57.5 - 72.8	100	8.8	5.2 - 14.4	400	25.7	19.2 - 33.4	1,400
No	8,200	50.9	48.6 - 53.1	1,400	8.4	7.2 - 9.7	6,600	40.7	38.6 - 43.0	16,100

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,400	67.3	62.5 - 71.9	300	7.8	5.5 - 10.8	900	24.9	20.8 - 29.5	3,600
<b>No</b>	6,700	48.1	45.7 - 50.5	1,200	8.6	7.3 - 10.0	6,000	43.3	41.0 - 45.7	13,900

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,700	59.8	56.5 - 63.0	600	7.7	6.1 - 9.7	2,600	32.5	29.4 - 35.7	7,900
<b>No</b>	4,400	46.0	43.1 - 48.9	800	8.9	7.4 - 10.7	4,300	45.1	42.3 - 48.0	9,500

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,400	58.8	56.1 - 61.4	1,300	7.5	6.2 - 9.1	5,900	33.7	31.1 - 36.3	17,600

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,200	62.1	57.6 - 66.4	200	9.1	6.7 - 12.2	600	28.8	24.8 - 33.1	2,000
<b>Honolulu</b>	7,500	57.9	54.4 - 61.4	900	7.3	5.7 - 9.4	4,500	34.8	31.5 - 38.2	12,900
<b>Kauai</b>	500	58.9	51.6 - 65.8	100	8.3	5.1 - 13.3	300	32.8	26.3 - 40.0	900
<b>Maui</b>	1,200	61.3	56.5 - 65.9	100	7.0	4.9 - 10.0	600	31.7	27.3 - 36.4	1,900

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,800	46.5	40.3 - 52.7	400	11.1	7.8 - 15.7	1,600	42.4	36.4 - 48.7	3,800
Native Hawaiian	3,400	70.8	68.1 - 73.5	300	6.7	5.3 - 8.3	1,100	22.5	20.1 - 25.1	4,800
Chinese	300	48.6	33.6 - 63.8	100	20.2	10.4 - 35.5	200	31.2	18.7 - 47.2	700
Filipino	2,400	62.0	55.8 - 67.8	100	3.1	1.8 - 5.5	1,300	34.9	29.1 - 41.1	3,900
Japanese	900	45.0	36.2 - 54.1	100	5.3	2.5 - 10.9	1,000	49.7	40.7 - 58.8	2,000
Other	1,600	63.8	55.4 - 71.4	200	8.7	5.0 - 14.8	700	27.4	20.6 - 35.5	2,500

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	75.7	67.9 - 82.1	100	5.9	2.9 - 11.7	300	18.4	12.9 - 25.6	1,600
20-24	3,300	78.9	74.3 - 82.9	200	5.6	3.8 - 8.2	600	15.5	12.0 - 19.9	4,200
25-34	4,700	53.7	49.8 - 57.5	700	7.4	5.6 - 9.7	3,400	38.9	35.2 - 42.8	8,800
35+	1,100	37.4	31.1 - 44.1	300	11.4	7.7 - 16.5	1,600	51.2	44.4 - 57.9	3,000

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,600	59.3	55.6 - 62.9	700	7.1	5.4 - 9.1	3,200	33.7	30.2 - 37.3	9,400
<b>Female</b>	4,800	58.3	54.3 - 62.1	700	8.1	6.1 - 10.5	2,800	33.7	30.0 - 37.5	8,300

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,900	50.5	47.0 - 54.0	1,000	8.6	6.9 - 10.8	4,700	40.8	37.5 - 44.3	11,600
<b>No</b>	4,500	74.7	70.9 - 78.2	300	5.4	3.9 - 7.4	1,200	19.9	16.7 - 23.5	6,000

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	5,400	68.7	64.9 - 72.2	500	5.9	4.4 - 7.9	2,000	25.4	22.1 - 29.0	7,800
<b>131-185%</b>	1,400	54.6	47.4 - 61.7	200	8.3	5.2 - 13.1	900	37.0	30.3 - 44.3	2,500
<b>186+%</b>	2,500	44.5	39.6 - 49.6	500	9.3	6.8 - 12.7	2,500	46.1	41.1 - 51.2	5,500



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

Income (2000-2003)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	3,000	70.8	65.8 - 75.3	200	5.0	3.3 - 7.5	1,000	24.2	19.9 - 29.0	4,300
\$15,001-\$25,000	1,600	70.6	63.7 - 76.7	200	7.1	4.1 - 12.0	500	22.3	17.0 - 28.7	2,300
\$25,001-\$40,000	2,000	60.9	54.4 - 67.0	200	7.1	4.6 - 10.9	1,000	32.0	26.2 - 38.4	3,200
>=\$40,001	2,900	43.7	39.3 - 48.2	600	9.4	7.1 - 12.4	3,100	46.9	42.4 - 51.5	6,700

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	700	53.3	43.1 - 63.2	200	12.6	7.1 - 21.5	400	34.1	25.2 - 44.3	1,200
Normal birth weight	9,700	59.2	56.4 - 61.9	1,200	7.1	5.9 - 8.7	5,500	33.6	31.0 - 36.3	16,400

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	75.3	67.2 - 82.0	100	4.5	2.1 - 9.1	300	20.2	14.1 - 28.1	1,600
No	9,000	57.5	54.6 - 60.3	1,200	7.6	6.2 - 9.3	5,500	34.9	32.2 - 37.7	15,700

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,600	74.0	68.5 - 78.8	200	5.7	3.5 - 9.2	700	20.3	16.0 - 25.5	3,600
<b>No</b>	7,500	55.0	51.9 - 58.1	1,100	7.9	6.4 - 9.7	5,000	37.1	34.1 - 40.2	13,600

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,200	70.4	66.7 - 73.8	400	5.7	4.2 - 7.7	1,800	23.9	20.7 - 27.5	7,400
<b>No</b>	5,100	50.8	47.2 - 54.5	900	8.7	6.8 - 11.0	4,100	40.5	36.9 - 44.2	10,100

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	9,900	57.7	54.9 - 60.4	1,400	7.9	6.5 - 9.6	5,900	34.4	31.8 - 37.1	17,100

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,200	61.9	57.5 - 66.2	100	6.4	4.4 - 9.0	600	31.7	27.7 - 36.0	2,000
<b>Honolulu</b>	7,200	56.7	53.1 - 60.2	1,100	8.3	6.6 - 10.5	4,400	35.0	31.6 - 38.5	12,600
<b>Kauai</b>	400	58.6	50.7 - 66.1	100	7.7	4.5 - 13.0	200	33.7	26.7 - 41.5	700
<b>Maui</b>	1,100	59.8	54.7 - 64.6	100	6.9	4.7 - 9.9	600	33.4	28.8 - 38.3	1,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	52.7	46.3 - 59.1	400	10.0	6.9 - 14.4	1,300	37.2	31.3 - 43.6	3,600
Native Hawaiian	3,200	68.3	65.5 - 71.1	300	6.6	5.3 - 8.3	1,200	25.1	22.5 - 27.8	4,600
Chinese	300	54.7	38.0 - 70.5	100	14.7	5.7 - 32.8	200	30.6	17.7 - 47.4	600
Filipino	1,800	52.1	45.4 - 58.8	200	5.9	3.3 - 10.2	1,500	42.0	35.4 - 48.8	3,500
Japanese	1,000	40.8	32.9 - 49.2	200	8.4	4.8 - 14.3	1,200	50.8	42.5 - 59.1	2,400
Other	1,700	70.4	61.8 - 77.7	200	8.1	4.5 - 14.2	500	21.5	15.2 - 29.5	2,300

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	77.3	68.8 - 84.1	100	3.6	1.6 - 7.8	300	19.0	12.7 - 27.5	1,500
20-24	3,100	71.9	66.9 - 76.5	300	6.8	4.7 - 9.9	900	21.2	17.1 - 26.0	4,200
25-34	4,500	52.9	48.9 - 56.9	800	9.2	7.1 - 11.8	3,200	37.9	34.1 - 41.9	8,500
35+	1,100	39.5	33.0 - 46.5	200	8.2	4.9 - 13.4	1,400	52.3	45.2 - 59.3	2,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,100	59.5	55.6 - 63.3	700	8.3	6.4 - 10.8	2,700	32.2	28.6 - 36.0	8,500
<b>Female</b>	4,800	55.9	51.9 - 59.8	600	7.5	5.7 - 9.9	3,100	36.6	32.8 - 40.5	8,600

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,000	51.9	48.3 - 55.4	1,000	9.0	7.1 - 11.2	4,600	39.2	35.8 - 42.7	11,600
<b>No</b>	3,800	70.1	65.7 - 74.2	300	5.7	4.1 - 7.9	1,300	24.2	20.3 - 28.5	5,500

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,500	67.4	63.2 - 71.4	500	7.6	5.6 - 10.2	1,700	25.0	21.4 - 29.0	6,700
<b>131-185%</b>	2,100	59.8	53.3 - 65.9	300	7.7	5.0 - 11.7	1,100	32.5	26.8 - 38.9	3,500
<b>186+%</b>	2,300	42.8	37.8 - 47.9	500	8.6	6.1 - 12.1	2,600	48.6	43.5 - 53.7	5,400

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

Income (2000-2003)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,400	70.3	64.9 - 75.2	200	7.0	4.7 - 10.4	800	22.7	18.2 - 27.9	3,400
\$15,001-\$25,000	1,700	66.1	58.8 - 72.8	200	9.7	5.9 - 15.5	600	24.2	18.5 - 31.0	2,500
\$25,001-\$40,000	2,300	63.3	57.0 - 69.1	300	6.9	4.4 - 10.6	1,100	29.8	24.3 - 36.0	3,700
>=\$40,001	2,800	42.9	38.4 - 47.6	500	8.1	5.9 - 11.0	3,200	49.0	44.3 - 53.6	6,500

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	500	46.0	35.5 - 56.9	100	9.5	5.3 - 16.6	500	44.5	33.8 - 55.7	1,200
Normal birth weight	9,300	58.5	55.6 - 61.3	1,200	7.7	6.3 - 9.4	5,400	33.8	31.1 - 36.6	15,800

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	77.7	70.0 - 83.9	100	6.0	2.9 - 12.0	200	16.3	11.2 - 23.1	1,400
No	8,600	56.1	53.1 - 59.0	1,300	8.2	6.7 - 10.0	5,500	35.7	32.9 - 38.6	15,400

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,500	75.5	70.1 - 80.3	100	4.5	2.9 - 7.1	700	19.9	15.5 - 25.2	3,300
<b>No</b>	7,100	53.2	50.0 - 56.4	1,200	8.9	7.2 - 10.9	5,000	37.9	34.8 - 41.1	13,300

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,600	63.9	59.7 - 67.9	600	8.1	6.0 - 10.8	2,000	28.0	24.2 - 32.0	7,100
<b>No</b>	5,200	53.3	49.5 - 57.0	800	7.8	6.1 - 10.1	3,800	38.9	35.3 - 42.6	9,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	10,000	59.9	57.7 - 62.1	1,300	7.6	6.5 - 9.0	5,400	32.4	30.4 - 34.6	16,700

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,200	64.4	59.8 - 68.7	200	9.1	6.7 - 12.2	500	26.6	22.6 - 30.9	1,900
<b>Honolulu</b>	7,300	59.4	56.6 - 62.2	900	7.6	6.2 - 9.3	4,100	33.0	30.4 - 35.8	12,300
<b>Kauai</b>	500	61.3	53.7 - 68.4	100	7.8	4.5 - 13.0	200	30.9	24.4 - 38.4	800
<b>Maui</b>	1,000	58.4	53.4 - 63.3	100	6.7	4.5 - 9.9	600	34.9	30.2 - 39.8	1,700



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	54.1	49.2 - 59.0	300	8.6	6.5 - 11.3	1,300	37.2	32.6 - 42.1	3,600
Native Hawaiian	3,200	70.5	68.2 - 72.7	300	5.8	4.8 - 7.1	1,100	23.7	21.7 - 25.8	4,500
Chinese	300	38.5	26.5 - 52.0	100	13.8	6.7 - 26.2	300	47.7	34.8 - 60.9	700
Filipino	2,100	61.0	55.5 - 66.2	200	6.5	4.1 - 10.3	1,100	32.5	27.6 - 37.8	3,500
Japanese	1,000	46.5	39.2 - 54.0	100	7.2	4.2 - 12.0	1,000	46.3	39.0 - 53.8	2,100
Other	1,500	65.3	58.4 - 71.6	200	9.8	6.3 - 15.0	600	24.9	19.5 - 31.3	2,300

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	78.1	72.9 - 82.6	100	4.7	2.9 - 7.4	300	17.2	13.1 - 22.2	1,600
20-24	3,100	72.8	68.9 - 76.3	200	5.5	3.9 - 7.8	900	21.7	18.5 - 25.3	4,300
25-34	4,300	53.9	50.5 - 57.1	800	9.9	8.1 - 12.2	2,900	36.2	33.1 - 39.5	8,100
35+	1,300	46.9	41.0 - 52.9	200	5.9	3.6 - 9.6	1,300	47.2	41.3 - 53.1	2,700

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	5,000	58.8	55.6 - 61.9	900	10.1	8.3 - 12.3	2,600	31.1	28.2 - 34.1	8,500
<b>Female</b>	5,000	61.1	57.9 - 64.2	400	5.1	3.9 - 6.6	2,800	33.8	30.9 - 36.9	8,200

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,900	52.9	50.1 - 55.8	900	8.5	7.0 - 10.3	4,300	38.6	35.8 - 41.4	11,200
<b>No</b>	4,100	74.0	70.7 - 77.0	300	5.9	4.4 - 7.9	1,100	20.1	17.4 - 23.1	5,500

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,500	70.9	67.8 - 73.9	400	6.9	5.3 - 9.0	1,400	22.1	19.5 - 25.0	6,300
<b>131-185%</b>	2,200	61.4	56.6 - 66.0	300	9.0	6.6 - 12.1	1,100	29.6	25.3 - 34.2	3,600
<b>186+%</b>	2,500	45.0	40.9 - 49.2	400	7.5	5.5 - 10.0	2,600	47.5	43.4 - 51.7	5,500

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

Income (2000-2003)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,400	71.3	67.0 - 75.3	200	6.5	4.5 - 9.3	800	22.2	18.6 - 26.1	3,400
\$15,001-\$25,000	1,600	69.1	63.5 - 74.2	200	8.3	5.5 - 12.3	500	22.6	18.2 - 27.7	2,300
\$25,001-\$40,000	2,100	59.7	55.0 - 64.2	300	8.2	6.0 - 11.2	1,100	32.1	27.8 - 36.7	3,600
>=\$40,001	3,200	49.0	45.2 - 52.8	500	7.9	6.0 - 10.3	2,800	43.1	39.4 - 47.0	6,500

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	58.3	49.3 - 66.8	100	5.0	2.2 - 10.9	400	36.7	28.4 - 45.8	1,100
Normal birth weight	9,300	60.0	57.7 - 62.3	1,200	7.8	6.7 - 9.2	5,000	32.2	30.0 - 34.4	15,600

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	75.2	69.2 - 80.3	100	5.0	2.9 - 8.5	300	19.9	15.2 - 25.5	1,600
No	8,600	58.0	55.6 - 60.4	1,200	8.1	6.8 - 9.5	5,000	33.9	31.7 - 36.3	14,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,500	73.8	69.5 - 77.6	200	6.2	4.3 - 9.0	700	20.0	16.6 - 23.9	3,400
<b>No</b>	7,100	55.4	52.8 - 58.0	1,100	8.3	6.9 - 9.9	4,600	36.3	33.8 - 38.9	12,800

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,900	69.2	66.2 - 72.2	500	6.6	5.1 - 8.4	1,700	24.2	21.5 - 27.1	7,000
<b>No</b>	5,000	52.9	49.8 - 56.0	800	8.5	6.9 - 10.5	3,700	38.6	35.6 - 41.7	9,500

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

State	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	9,400	54.8	52.6 - 57.0	1,600	9.3	8.0 - 10.7	6,200	35.9	33.8 - 38.1	17,200

County	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,200	59.2	54.6 - 63.7	200	8.8	6.5 - 11.9	600	32.0	27.8 - 36.4	2,000
<b>Honolulu</b>	6,900	54.1	51.3 - 56.8	1,300	9.8	8.2 - 11.7	4,600	36.1	33.5 - 38.8	12,800
<b>Kauai</b>	400	59.9	51.9 - 67.3	100	8.4	5.0 - 13.7	200	31.7	24.8 - 39.5	700
<b>Maui</b>	900	52.9	47.8 - 57.9	100	6.3	4.3 - 9.2	700	40.8	35.9 - 45.9	1,800

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

DOH Race-Ethnicity	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	1,900	46.5	42.0 - 51.1	400	10.4	7.8 - 13.7	1,800	43.1	38.6 - 47.8	4,100
Native Hawaiian	3,100	66.5	64.1 - 68.8	400	7.9	6.6 - 9.3	1,200	25.7	23.6 - 27.9	4,600
Chinese	300	49.9	36.7 - 63.0	100	17.4	9.4 - 30.0	200	32.8	21.5 - 46.5	700
Filipino	2,000	58.2	52.7 - 63.5	300	8.5	5.7 - 12.4	1,200	33.3	28.4 - 38.7	3,500
Japanese	800	40.4	33.4 - 47.7	200	9.6	6.0 - 15.0	1,000	50.0	42.7 - 57.3	2,000
Other	1,300	55.0	47.8 - 61.9	200	8.8	5.4 - 14.0	800	36.2	29.7 - 43.3	2,300

PRAMS Age Group	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,500	78.4	73.0 - 83.0	100	4.7	2.6 - 8.3	300	16.9	12.9 - 21.8	1,900
20-24	2,800	66.6	62.4 - 70.6	400	10.5	7.9 - 13.8	900	22.9	19.5 - 26.6	4,100
25-34	4,100	49.8	46.5 - 53.0	800	9.7	7.9 - 11.8	3,300	40.6	37.4 - 43.9	8,200
35+	1,100	37.3	32.3 - 42.7	300	9.4	6.5 - 13.3	1,600	53.3	47.8 - 58.7	3,000

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

Infant Gender	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	4,800	54.6	51.6 - 57.6	700	8.3	6.8 - 10.1	3,300	37.1	34.2 - 40.1	8,900
<b>Female</b>	4,600	55.0	51.8 - 58.2	900	10.3	8.4 - 12.7	2,900	34.7	31.7 - 37.8	8,300

Married	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,800	49.4	46.7 - 52.1	1,100	9.0	7.5 - 10.7	4,900	41.6	38.9 - 44.4	11,800
<b>No</b>	3,600	66.6	63.1 - 69.9	500	9.9	7.7 - 12.7	1,300	23.5	20.6 - 26.6	5,400

Poverty Level	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	4,600	62.8	59.6 - 66.0	600	8.8	7.0 - 10.9	2,100	28.4	25.5 - 31.5	7,300
<b>131-185%</b>	1,500	63.8	58.0 - 69.2	200	6.9	4.5 - 10.4	700	29.3	24.3 - 34.9	2,300
<b>186+%</b>	2,600	41.2	37.5 - 45.0	700	11.7	9.3 - 14.5	2,900	47.1	43.3 - 51.0	6,200

**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

Income (2000-2003)	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,500	62.3	57.9 - 66.5	400	9.4	7.0 - 12.5	1,100	28.3	24.4 - 32.5	4,000
\$15,001-\$25,000	1,700	64.6	59.1 - 69.6	200	8.6	6.1 - 12.1	700	26.8	22.2 - 32.0	2,700
\$25,001-\$40,000	1,900	59.5	54.5 - 64.3	300	7.7	5.3 - 11.0	1,100	32.8	28.3 - 37.7	3,300
>=\$40,001	2,700	42.4	38.8 - 46.2	700	10.9	8.6 - 13.6	2,900	46.7	42.9 - 50.5	6,300

Birthweight	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	600	56.8	48.1 - 65.1	100	8.5	4.3 - 15.9	400	34.8	27.0 - 43.4	1,100
Normal birth weight	8,700	54.5	52.2 - 56.8	1,500	9.3	8.0 - 10.8	5,800	36.2	34.0 - 38.4	15,900

Smoke 3rd Trimester	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	64.9	57.7 - 71.4	200	11.0	7.0 - 16.6	300	24.2	18.6 - 30.9	1,400
No	8,200	53.8	51.4 - 56.1	1,400	9.0	7.7 - 10.5	5,700	37.2	35.0 - 39.5	15,200



**PRAMS Health Indicator Report - Took multivitamins/folic acid month prior to pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the  
Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

Smoke Before Pregnant	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,300	70.1	65.5 - 74.4	300	9.1	6.5 - 12.6	700	20.8	17.2 - 24.9	3,300
<b>No</b>	6,700	50.8	48.3 - 53.4	1,200	9.2	7.8 - 10.8	5,300	40.0	37.5 - 42.5	13,200

WIC During Pregnancy	None			1-3 times per week			4 or more times per week			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	4,700	63.7	60.5 - 66.7	500	7.0	5.5 - 8.9	2,200	29.3	26.4 - 32.3	7,400
<b>No</b>	4,500	47.8	44.8 - 50.9	1,000	10.9	9.1 - 13.1	3,900	41.2	38.2 - 44.3	9,500