

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- for the State of Hawaii, for the Year(s)**

**2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

*The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 3-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.*

*Note: Data collection for household income changed in 2004 and again in 2012. Because of this, income data does not exist for some 3-year aggregate periods.*

*Below are the indicators included in this report and the long description for each indicator.*

PRAMS Health Indicator	Health Indicator Description
Discuss effects of smoking with HCW during PNC	Discuss effects of smoking with HCW during PNC

*Suggested citation:*

*Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/21/17*

*For documentation on other terms and coding in this report, please go to the following links:*

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact  
The Hawai'i Health Data Warehouse**

[profiles@hhdw.org](mailto:profiles@hhdw.org)

**(855) 946-5899 x15**

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- for the State of Hawaii,  
for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**


*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

Years	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2002	12,100	73.4	72.1 - 74.7	4,400	26.6	25.3 - 27.9	16,500
2001-2003	12,400	74.5	73.2 - 75.8	4,300	25.5	24.2 - 26.8	16,700
2002-2004	12,600	74.2	72.9 - 75.5	4,400	25.8	24.5 - 27.1	17,000
2003-2005	12,600	73.8	72.6 - 75.1	4,500	26.2	24.9 - 27.4	17,100
2004-2006	12,800	73.7	72.5 - 74.9	4,600	26.3	25.1 - 27.5	17,300
2005-2007	12,900	73.4	72.2 - 74.6	4,700	26.6	25.4 - 27.8	17,600
2006-2008	13,100	72.6	71.4 - 73.8	4,900	27.4	26.2 - 28.6	18,000
2007-2009	12,900	71.4	70.0 - 72.7	5,200	28.6	27.3 - 30.0	18,000
2008-2010	13,000	71.5	70.0 - 73.0	5,200	28.5	27.0 - 30.0	18,200
2009-2011	13,100	72.5	70.8 - 74.1	5,000	27.5	25.9 - 29.2	18,100
2010-2012	13,100	72.2	70.5 - 73.8	5,100	27.8	26.2 - 29.5	18,200
2011-2013	12,900	71.1	69.3 - 72.8	5,200	28.9	27.2 - 30.7	18,100

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- for the State of Hawaii, for the Year(s)**

**2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*



**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2011-2013**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,900	71.1	69.3 - 72.8	5,200	28.9	27.2 - 30.7	18,100

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,700	73.9	71.1 - 76.5	600	26.1	23.5 - 28.9	2,300
<b>Honolulu</b>	9,200	70.5	68.1 - 72.8	3,800	29.5	27.2 - 31.9	13,000
<b>Kauai</b>	600	73.7	70.6 - 76.7	200	26.3	23.3 - 29.4	800
<b>Maui</b>	1,400	71.1	68.3 - 73.7	600	28.9	26.3 - 31.7	1,900

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2011-2013**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,800	63.8	59.9 - 67.4	1,600	36.2	32.6 - 40.1	4,500
Native Hawaiian	4,200	79.9	77.1 - 82.5	1,100	20.1	17.5 - 22.9	5,300
Chinese	600	66.3	57.2 - 74.4	300	33.7	25.6 - 42.8	900
Filipino	2,300	72.6	68.3 - 76.5	900	27.4	23.5 - 31.7	3,200
Japanese	900	54.1	47.5 - 60.5	800	45.9	39.5 - 52.5	1,700
Other	2,000	76.9	72.1 - 81.1	600	23.1	18.9 - 27.9	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	85.6	78.7 - 90.6	100	14.4	9.4 - 21.3	900
20-24	3,300	80.7	77.1 - 83.8	800	19.3	16.2 - 22.9	4,000
25-34	6,900	70.0	67.5 - 72.3	3,000	30.0	27.7 - 32.5	9,900
35+	1,900	58.6	54.1 - 62.9	1,300	41.4	37.1 - 45.9	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2011-2013**

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Infant Gender</b>							
<b>Male</b>	6,300	69.4	66.8 - 71.8	2,800	30.6	28.2 - 33.2	9,000
<b>Female</b>	6,600	72.9	70.4 - 75.2	2,500	27.1	24.8 - 29.6	9,100

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Married</b>							
<b>Yes</b>	7,500	66.0	63.6 - 68.3	3,900	34.0	31.7 - 36.4	11,300
<b>No</b>	5,400	79.7	77.1 - 82.0	1,400	20.3	18.0 - 22.9	6,800

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Poverty Level</b>							
<b>0-130%</b>	5,200	79.8	77.2 - 82.2	1,300	20.2	17.8 - 22.8	6,500
<b>131-185%</b>	2,100	75.4	70.9 - 79.4	700	24.6	20.6 - 29.1	2,800
<b>186+%</b>	4,600	61.8	58.8 - 64.8	2,800	38.2	35.2 - 41.2	7,400

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2011-2013**

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	74.4	71.7 - 77.0	300	25.6	23.0 - 28.3	1,300
Normal birth weight	11,900	70.9	69.0 - 72.7	4,900	29.1	27.3 - 31.0	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	88.8	82.1 - 93.2	100	11.2	6.8 - 17.9	900
No	11,800	70.1	68.3 - 71.9	5,000	29.9	28.1 - 31.7	16,800

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,700	83.5	79.8 - 86.6	500	16.5	13.4 - 20.2	3,200
No	9,900	68.4	66.3 - 70.3	4,600	31.6	29.7 - 33.7	14,500

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2011-2013**

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,400	78.8	76.4 - 81.0	1,700	21.2	19.0 - 23.6	8,100
<b>No</b>	6,500	65.0	62.4 - 67.4	3,500	35.0	32.6 - 37.6	10,000



**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2010-2012**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,100	72.2	70.5 - 73.8	5,100	27.8	26.2 - 29.5	18,200

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,800	76.9	74.4 - 79.3	500	23.1	20.7 - 25.6	2,300
<b>Honolulu</b>	9,300	71.1	68.8 - 73.3	3,800	28.9	26.7 - 31.2	13,100
<b>Kauai</b>	600	72.9	69.6 - 75.9	200	27.1	24.1 - 30.4	800
<b>Maui</b>	1,400	73.1	70.5 - 75.5	500	26.9	24.5 - 29.5	1,900

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2010-2012**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	66.7	63.1 - 70.1	1,500	33.3	29.9 - 36.9	4,400
Native Hawaiian	4,200	79.4	76.6 - 82.0	1,100	20.6	18.0 - 23.4	5,300
Chinese	500	65.8	56.8 - 73.7	300	34.2	26.3 - 43.2	800
Filipino	2,300	71.6	67.4 - 75.5	900	28.4	24.5 - 32.6	3,300
Japanese	900	52.6	46.3 - 58.9	800	47.4	41.1 - 53.7	1,700
Other	2,300	81.3	77.1 - 84.9	500	18.7	15.1 - 22.9	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	87.3	81.0 - 91.8	100	12.7	8.2 - 19.0	1,000
20-24	3,500	83.4	80.3 - 86.2	700	16.6	13.8 - 19.7	4,200
25-34	6,900	70.2	67.8 - 72.5	2,900	29.8	27.5 - 32.2	9,800
35+	1,900	58.9	54.5 - 63.2	1,300	41.1	36.8 - 45.5	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2010-2012**

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,600	71.9	69.4 - 74.2	2,600	28.1	25.8 - 30.6	9,100
Female	6,600	72.4	70.0 - 74.8	2,500	27.6	25.2 - 30.0	9,000

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,700	67.7	65.5 - 69.9	3,700	32.3	30.1 - 34.5	11,400
No	5,400	79.5	77.0 - 81.9	1,400	20.5	18.1 - 23.0	6,800

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,200	81.5	79.0 - 83.8	1,200	18.5	16.2 - 21.0	6,400
131-185%	2,200	74.3	69.9 - 78.3	800	25.7	21.7 - 30.1	2,900
186+%	4,800	63.2	60.3 - 65.9	2,800	36.8	34.1 - 39.7	7,600

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2010-2012**

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	73.2	70.3 - 75.9	300	26.8	24.1 - 29.7	1,300
Normal birth weight	12,200	72.1	70.2 - 73.8	4,700	27.9	26.2 - 29.8	16,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	900	91.7	86.1 - 95.2	100	8.3	4.8 - 13.9	1,000
No	11,900	71.0	69.2 - 72.7	4,900	29.0	27.3 - 30.8	16,800

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	83.8	80.3 - 86.7	500	16.2	13.3 - 19.7	3,400
No	10,100	69.5	67.5 - 71.4	4,400	30.5	28.6 - 32.5	14,500

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2010-2012**

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,500	79.8	77.5 - 81.9	1,600	20.2	18.1 - 22.5	8,100
<b>No</b>	6,600	65.9	63.4 - 68.3	3,400	34.1	31.7 - 36.6	10,000

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2009-2011**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,100	72.5	70.8 - 74.1	5,000	27.5	25.9 - 29.2	18,100

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,800	77.3	74.8 - 79.7	500	22.7	20.3 - 25.2	2,300
<b>Honolulu</b>	9,300	71.4	69.1 - 73.6	3,700	28.6	26.4 - 30.9	13,000
<b>Kauai</b>	600	74.0	70.9 - 76.9	200	26.0	23.1 - 29.1	800
<b>Maui</b>	1,500	73.3	70.9 - 75.7	500	26.7	24.3 - 29.1	2,000

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2009-2011**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,000	68.0	64.5 - 71.3	1,400	32.0	28.7 - 35.5	4,400
Native Hawaiian	4,400	79.4	76.7 - 82.0	1,100	20.6	18.0 - 23.3	5,500
Chinese	500	62.3	52.8 - 71.0	300	37.7	29.0 - 47.2	700
Filipino	2,400	74.2	70.2 - 77.9	800	25.8	22.1 - 29.8	3,200
Japanese	900	54.9	48.7 - 60.9	800	45.1	39.1 - 51.3	1,700
Other	2,000	77.8	73.1 - 81.8	600	22.2	18.2 - 26.9	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	91.9	87.8 - 94.7	100	8.1	5.3 - 12.2	1,300
20-24	3,600	84.7	81.8 - 87.3	700	15.3	12.7 - 18.2	4,300
25-34	6,500	69.0	66.6 - 71.3	2,900	31.0	28.7 - 33.4	9,400
35+	1,800	58.4	54.0 - 62.7	1,300	41.6	37.3 - 46.0	3,100

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2009-2011**

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Infant Gender</b>							
<b>Male</b>	6,700	72.0	69.7 - 74.3	2,600	28.0	25.7 - 30.3	9,300
<b>Female</b>	6,400	73.0	70.6 - 75.3	2,400	27.0	24.7 - 29.4	8,800

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Married</b>							
<b>Yes</b>	7,400	66.6	64.3 - 68.8	3,700	33.4	31.2 - 35.7	11,200
<b>No</b>	5,700	82.0	79.6 - 84.2	1,300	18.0	15.8 - 20.4	7,000

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Poverty Level</b>							
<b>0-130%</b>	5,200	82.7	80.3 - 85.0	1,100	17.3	15.0 - 19.7	6,300
<b>131-185%</b>	2,100	74.8	70.4 - 78.7	700	25.2	21.3 - 29.6	2,800
<b>186+%</b>	4,700	61.3	58.5 - 64.1	3,000	38.7	35.9 - 41.5	7,700



**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2009-2011**

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,700	85.5	82.1 - 88.3	500	14.5	11.7 - 17.9	3,200
\$10,000-\$24,999	2,800	82.4	79.1 - 85.3	600	17.6	14.7 - 20.9	3,400
\$25,000-\$49,999	3,100	72.6	69.0 - 75.9	1,200	27.4	24.1 - 31.0	4,300
>=\$50,000	3,500	57.8	54.6 - 61.0	2,600	42.2	39.0 - 45.4	6,100

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	74.9	72.2 - 77.5	300	25.1	22.5 - 27.8	1,300
Normal birth weight	12,100	72.3	70.4 - 74.0	4,700	27.7	26.0 - 29.6	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	92.7	88.4 - 95.5	100	7.3	4.5 - 11.6	1,300
No	11,700	70.9	69.1 - 72.6	4,800	29.1	27.4 - 30.9	16,500

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2009-2011**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,200	86.5	83.5 - 89.0	500	13.5	11.0 - 16.5	3,700
No	9,700	68.8	66.8 - 70.8	4,400	31.2	29.2 - 33.2	14,100

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,300	80.9	78.6 - 82.9	1,500	19.1	17.1 - 21.4	7,800
No	6,600	66.0	63.6 - 68.4	3,400	34.0	31.6 - 36.4	10,100

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2008-2010**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,000	71.5	70.0 - 73.0	5,200	28.5	27.0 - 30.0	18,200

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,800	76.7	73.8 - 79.4	600	23.3	20.6 - 26.2	2,400
<b>Honolulu</b>	9,100	70.6	68.5 - 72.6	3,800	29.4	27.4 - 31.5	12,900
<b>Kauai</b>	600	71.7	67.2 - 75.8	200	28.3	24.2 - 32.8	800
<b>Maui</b>	1,500	71.5	68.5 - 74.4	600	28.5	25.6 - 31.5	2,100

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2008-2010**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,000	68.4	65.2 - 71.5	1,400	31.6	28.5 - 34.8	4,300
Native Hawaiian	4,200	76.9	74.2 - 79.4	1,300	23.1	20.6 - 25.8	5,500
Chinese	400	55.3	46.7 - 63.6	300	44.7	36.4 - 53.3	700
Filipino	2,400	72.9	69.1 - 76.3	900	27.1	23.7 - 30.9	3,300
Japanese	1,000	57.0	51.6 - 62.3	800	43.0	37.7 - 48.4	1,800
Other	2,000	78.0	74.0 - 81.6	600	22.0	18.4 - 26.0	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	90.3	86.3 - 93.2	100	9.7	6.8 - 13.7	1,400
20-24	3,500	82.5	79.6 - 85.1	700	17.5	14.9 - 20.4	4,200
25-34	6,300	67.7	65.5 - 69.9	3,000	32.3	30.1 - 34.5	9,300
35+	1,900	60.0	56.0 - 63.8	1,300	40.0	36.2 - 44.0	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2008-2010**

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Infant Gender</b>							
<b>Male</b>	6,600	69.8	67.6 - 71.9	2,900	30.2	28.1 - 32.4	9,400
<b>Female</b>	6,400	73.4	71.2 - 75.5	2,300	26.6	24.5 - 28.8	8,700

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Married</b>							
<b>Yes</b>	7,400	66.3	64.2 - 68.3	3,800	33.7	31.7 - 35.8	11,200
<b>No</b>	5,600	80.0	77.7 - 82.2	1,400	20.0	17.8 - 22.3	6,900

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Poverty Level</b>							
<b>0-130%</b>	5,100	80.0	77.7 - 82.2	1,300	20.0	17.8 - 22.3	6,400
<b>131-185%</b>	2,100	74.6	70.5 - 78.2	700	25.4	21.8 - 29.5	2,800
<b>186+%</b>	4,700	61.2	58.7 - 63.7	2,900	38.8	36.3 - 41.3	7,600

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2008-2010**

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,500	82.8	79.5 - 85.8	500	17.2	14.2 - 20.5	3,100
\$10,000-\$24,999	2,900	80.6	77.6 - 83.3	700	19.4	16.7 - 22.4	3,700
\$25,000-\$49,999	2,900	70.5	67.2 - 73.7	1,200	29.5	26.3 - 32.8	4,100
>=\$50,000	3,600	58.8	55.9 - 61.7	2,500	41.2	38.3 - 44.1	6,100

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	72.7	69.0 - 76.0	300	27.3	24.0 - 31.0	1,300
Normal birth weight	12,000	71.4	69.8 - 73.0	4,800	28.6	27.0 - 30.2	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	91.7	88.2 - 94.3	100	8.3	5.7 - 11.8	1,500
No	11,300	69.6	67.9 - 71.2	5,000	30.4	28.8 - 32.1	16,300

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2008-2010**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,100	85.2	82.4 - 87.7	500	14.8	12.3 - 17.6	3,700
<b>No</b>	9,600	67.9	66.0 - 69.6	4,500	32.1	30.4 - 34.0	14,100

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,100	79.6	77.4 - 81.6	1,600	20.4	18.4 - 22.6	7,700
<b>No</b>	6,700	65.5	63.3 - 67.6	3,500	34.5	32.4 - 36.7	10,200

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2007-2009**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,900	71.4	70.0 - 72.7	5,200	28.6	27.3 - 30.0	18,000

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,700	71.0	67.6 - 74.2	700	29.0	25.8 - 32.4	2,400
<b>Honolulu</b>	9,000	71.6	69.9 - 73.4	3,600	28.4	26.6 - 30.1	12,600
<b>Kauai</b>	700	72.5	67.4 - 77.2	300	27.5	22.8 - 32.6	900
<b>Maui</b>	1,500	69.7	66.2 - 73.0	600	30.3	27.0 - 33.8	2,100



**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2007-2009**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	68.2	65.3 - 71.0	1,300	31.8	29.0 - 34.7	4,200
Native Hawaiian	4,100	77.0	74.5 - 79.4	1,200	23.0	20.6 - 25.5	5,300
Chinese	400	56.1	49.1 - 63.0	300	43.9	37.0 - 50.9	700
Filipino	2,400	73.1	69.9 - 76.1	900	26.9	23.9 - 30.1	3,300
Japanese	1,100	59.9	55.3 - 64.3	800	40.1	35.7 - 44.7	1,900
Other	2,000	75.0	71.3 - 78.5	700	25.0	21.5 - 28.7	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,500	89.6	86.2 - 92.3	200	10.4	7.7 - 13.8	1,600
20-24	3,300	79.5	76.8 - 82.0	900	20.5	18.0 - 23.2	4,200
25-34	6,200	67.8	65.7 - 69.7	2,900	32.2	30.3 - 34.3	9,100
35+	1,900	61.5	57.9 - 64.9	1,200	38.5	35.1 - 42.1	3,100

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2007-2009**

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Infant Gender</b>							
<b>Male</b>	6,600	69.6	67.6 - 71.5	2,900	30.4	28.5 - 32.4	9,500
<b>Female</b>	6,300	73.3	71.3 - 75.2	2,300	26.7	24.8 - 28.7	8,600

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Married</b>							
<b>Yes</b>	7,300	65.7	63.8 - 67.5	3,800	34.3	32.5 - 36.2	11,100
<b>No</b>	5,600	80.6	78.5 - 82.5	1,300	19.4	17.5 - 21.5	6,900

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Poverty Level</b>							
<b>0-130%</b>	5,100	78.3	76.1 - 80.3	1,400	21.7	19.7 - 23.9	6,500
<b>131-185%</b>	2,000	73.1	69.4 - 76.5	700	26.9	23.5 - 30.6	2,700
<b>186+%</b>	4,700	62.9	60.6 - 65.1	2,800	37.1	34.9 - 39.4	7,500

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2007-2009**

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,400	81.6	78.3 - 84.4	500	18.4	15.6 - 21.7	3,000
\$10,000-\$24,999	3,000	78.9	76.2 - 81.5	800	21.1	18.5 - 23.8	3,900
\$25,000-\$49,999	2,800	70.4	67.4 - 73.3	1,200	29.6	26.7 - 32.6	4,000
>=\$50,000	3,600	60.3	57.7 - 62.8	2,400	39.7	37.2 - 42.3	6,000

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	72.7	68.6 - 76.5	400	27.3	23.5 - 31.4	1,400
Normal birth weight	11,900	71.2	69.8 - 72.7	4,800	28.8	27.3 - 30.2	16,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	90.8	87.6 - 93.3	100	9.2	6.7 - 12.4	1,500
No	11,200	69.5	68.0 - 71.0	4,900	30.5	29.0 - 32.0	16,100

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2007-2009**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,100	85.8	83.3 - 87.9	500	14.2	12.1 - 16.7	3,600
<b>No</b>	9,500	67.7	66.0 - 69.3	4,600	32.3	30.7 - 34.0	14,100

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,000	79.0	77.0 - 80.8	1,600	21.0	19.2 - 23.0	7,600
<b>No</b>	6,700	65.8	63.9 - 67.7	3,500	34.2	32.3 - 36.1	10,200

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2006-2008**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,100	72.6	71.4 - 73.8	4,900	27.4	26.2 - 28.6	18,000

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,700	71.8	68.1 - 75.2	700	28.2	24.8 - 31.9	2,300
<b>Honolulu</b>	9,300	73.0	71.5 - 74.4	3,400	27.0	25.6 - 28.5	12,700
<b>Kauai</b>	700	73.3	67.5 - 78.4	300	26.7	21.6 - 32.5	900
<b>Maui</b>	1,400	70.8	66.8 - 74.5	600	29.2	25.5 - 33.2	2,000

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2006-2008**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	69.1	66.4 - 71.7	1,300	30.9	28.3 - 33.6	4,200
Native Hawaiian	3,800	77.1	74.6 - 79.4	1,100	22.9	20.6 - 25.4	5,000
Chinese	400	60.6	56.6 - 64.4	200	39.4	35.6 - 43.4	600
Filipino	2,500	75.8	73.1 - 78.3	800	24.2	21.7 - 26.9	3,300
Japanese	1,200	59.6	55.6 - 63.4	800	40.4	36.6 - 44.4	2,100
Other	2,200	78.4	75.4 - 81.2	600	21.6	18.8 - 24.6	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	87.4	83.6 - 90.5	200	12.6	9.5 - 16.4	1,500
20-24	3,300	78.0	75.4 - 80.3	900	22.0	19.7 - 24.6	4,300
25-34	6,400	70.9	69.1 - 72.6	2,600	29.1	27.4 - 30.9	9,000
35+	2,000	63.1	60.1 - 66.1	1,200	36.9	33.9 - 39.9	3,200

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2006-2008**

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,600	71.5	69.7 - 73.2	2,700	28.5	26.8 - 30.3	9,300
Female	6,400	73.8	72.0 - 75.5	2,300	26.2	24.5 - 28.0	8,700

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,800	68.2	66.6 - 69.7	3,600	31.8	30.3 - 33.4	11,400
No	5,300	80.2	78.2 - 82.0	1,300	19.8	18.0 - 21.8	6,600

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,100	78.6	76.6 - 80.5	1,400	21.4	19.5 - 23.4	6,500
131-185%	1,900	73.5	70.1 - 76.7	700	26.5	23.3 - 29.9	2,500
186+%	5,100	66.2	64.3 - 68.1	2,600	33.8	31.9 - 35.7	7,700

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2006-2008**

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,400	80.3	77.3 - 83.0	600	19.7	17.0 - 22.7	2,900
\$10,000-\$24,999	3,000	78.8	76.2 - 81.2	800	21.2	18.8 - 23.8	3,800
\$25,000-\$49,999	2,900	71.4	68.7 - 73.9	1,200	28.6	26.1 - 31.3	4,100
>=\$50,000	3,800	64.5	62.3 - 66.7	2,100	35.5	33.3 - 37.7	6,000

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	74.3	69.8 - 78.4	400	25.7	21.6 - 30.2	1,400
Normal birth weight	12,000	72.5	71.2 - 73.7	4,600	27.5	26.3 - 28.8	16,600

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	90.6	87.3 - 93.1	100	9.4	6.9 - 12.7	1,500
No	11,500	71.0	69.6 - 72.3	4,700	29.0	27.7 - 30.4	16,200



**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2006-2008**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,000	85.1	82.7 - 87.3	500	14.9	12.7 - 17.3	3,500
No	9,900	69.6	68.1 - 70.9	4,300	30.4	29.1 - 31.9	14,200

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,000	79.0	77.1 - 80.7	1,600	21.0	19.3 - 22.9	7,700
No	7,000	67.8	66.1 - 69.5	3,300	32.2	30.5 - 33.9	10,300

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2005-2007**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,900	73.4	72.2 - 74.6	4,700	26.6	25.4 - 27.8	17,600

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	72.5	68.8 - 76.0	600	27.5	24.0 - 31.2	2,200
<b>Honolulu</b>	9,200	73.5	72.0 - 74.9	3,300	26.5	25.1 - 28.0	12,600
<b>Kauai</b>	700	77.7	72.1 - 82.5	200	22.3	17.5 - 27.9	900
<b>Maui</b>	1,400	72.0	68.0 - 75.8	500	28.0	24.2 - 32.0	1,900

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2005-2007**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,800	69.1	66.3 - 71.7	1,200	30.9	28.3 - 33.7	4,000
Native Hawaiian	3,900	80.8	78.4 - 82.9	900	19.2	17.1 - 21.6	4,900
Chinese	400	63.3	59.4 - 67.0	200	36.7	33.0 - 40.6	600
Filipino	2,400	75.3	72.6 - 77.8	800	24.7	22.2 - 27.4	3,200
Japanese	1,300	59.1	55.2 - 62.9	900	40.9	37.1 - 44.8	2,100
Other	2,100	78.0	74.9 - 80.8	600	22.0	19.2 - 25.1	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,400	87.4	83.7 - 90.4	200	12.6	9.6 - 16.3	1,600
20-24	3,400	78.8	76.3 - 81.0	900	21.2	19.0 - 23.7	4,300
25-34	6,200	72.1	70.3 - 73.8	2,400	27.9	26.2 - 29.7	8,600
35+	1,900	62.3	59.2 - 65.4	1,100	37.7	34.6 - 40.8	3,100

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2005-2007**

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Infant Gender</b>							
<b>Male</b>	6,600	73.6	71.9 - 75.3	2,400	26.4	24.7 - 28.1	9,000
<b>Female</b>	6,300	73.2	71.4 - 74.9	2,300	26.8	25.1 - 28.6	8,600

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Married</b>							
<b>Yes</b>	7,600	68.6	67.1 - 70.2	3,500	31.4	29.8 - 32.9	11,100
<b>No</b>	5,200	81.7	79.7 - 83.4	1,200	18.3	16.6 - 20.3	6,400

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Poverty Level</b>							
<b>0-130%</b>	5,000	79.5	77.5 - 81.3	1,300	20.5	18.7 - 22.5	6,300
<b>131-185%</b>	1,700	74.0	70.4 - 77.3	600	26.0	22.7 - 29.6	2,300
<b>186+%</b>	5,200	67.2	65.3 - 69.1	2,500	32.8	30.9 - 34.7	7,700

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2005-2007**

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,400	81.6	78.7 - 84.2	500	18.4	15.8 - 21.3	2,900
\$10,000-\$24,999	2,900	79.2	76.6 - 81.6	800	20.8	18.4 - 23.4	3,700
\$25,000-\$49,999	2,900	72.4	69.7 - 74.9	1,100	27.6	25.1 - 30.3	4,100
>=\$50,000	3,700	65.0	62.7 - 67.2	2,000	35.0	32.8 - 37.3	5,700

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,100	76.5	72.1 - 80.3	300	23.5	19.7 - 27.9	1,400
Normal birth weight	11,800	73.1	71.9 - 74.4	4,300	26.9	25.6 - 28.1	16,200

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	93.0	90.0 - 95.1	100	7.0	4.9 - 10.0	1,500
No	11,300	71.7	70.4 - 73.0	4,500	28.3	27.0 - 29.6	15,800

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2005-2007**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,000	85.5	83.2 - 87.6	500	14.5	12.4 - 16.8	3,600
<b>No</b>	9,600	70.4	68.9 - 71.8	4,100	29.6	28.2 - 31.1	13,700

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,900	79.4	77.5 - 81.1	1,500	20.6	18.9 - 22.5	7,500
<b>No</b>	6,800	68.8	67.2 - 70.5	3,100	31.2	29.5 - 32.8	9,900

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2004-2006**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,800	73.7	72.5 - 74.9	4,600	26.3	25.1 - 27.5	17,300

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	75.3	71.8 - 78.6	500	24.7	21.4 - 28.2	2,100
<b>Honolulu</b>	9,200	73.2	71.8 - 74.6	3,400	26.8	25.4 - 28.2	12,600
<b>Kauai</b>	600	77.6	71.9 - 82.5	200	22.4	17.5 - 28.1	800
<b>Maui</b>	1,300	73.6	69.8 - 77.2	500	26.4	22.8 - 30.2	1,800

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2004-2006**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	68.5	65.8 - 71.1	1,200	31.5	28.9 - 34.2	3,900
Native Hawaiian	3,900	80.7	78.4 - 82.8	900	19.3	17.2 - 21.6	4,800
Chinese	400	63.8	60.3 - 67.1	200	36.2	32.9 - 39.7	600
Filipino	2,500	77.8	75.3 - 80.1	700	22.2	19.9 - 24.7	3,200
Japanese	1,300	58.6	54.8 - 62.3	900	41.4	37.7 - 45.2	2,200
Other	2,000	78.9	75.9 - 81.6	500	21.1	18.4 - 24.1	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	86.9	83.2 - 89.9	200	13.1	10.1 - 16.8	1,400
20-24	3,400	79.7	77.4 - 81.9	900	20.3	18.1 - 22.6	4,300
25-34	6,100	72.5	70.7 - 74.1	2,300	27.5	25.9 - 29.3	8,500
35+	2,000	62.8	59.9 - 65.7	1,200	37.2	34.3 - 40.1	3,100



**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2004-2006**

Infant Gender	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,700	74.4	72.8 - 76.0	2,300	25.6	24.0 - 27.2	9,100
Female	6,000	73.0	71.2 - 74.7	2,200	27.0	25.3 - 28.8	8,300

Married	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,000	69.6	68.1 - 71.1	3,500	30.4	28.9 - 31.9	11,400
No	4,800	81.6	79.7 - 83.4	1,100	18.4	16.6 - 20.3	5,900

Poverty Level	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,000	80.4	78.5 - 82.2	1,200	19.6	17.8 - 21.5	6,200
131-185%	1,600	75.9	72.3 - 79.1	500	24.1	20.9 - 27.7	2,100
186+%	5,200	66.8	65.0 - 68.7	2,600	33.2	31.3 - 35.0	7,700

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2004-2006**

Income (2004-2011)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,300	82.1	79.3 - 84.6	500	17.9	15.4 - 20.7	2,800
\$10,000-\$24,999	3,000	79.4	77.0 - 81.7	800	20.6	18.3 - 23.0	3,800
\$25,000-\$49,999	3,000	73.1	70.6 - 75.5	1,100	26.9	24.5 - 29.4	4,100
>=\$50,000	3,400	64.3	62.1 - 66.5	1,900	35.7	33.5 - 37.9	5,300

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	77.2	72.7 - 81.1	300	22.8	18.9 - 27.3	1,200
Normal birth weight	11,800	73.5	72.2 - 74.7	4,300	26.5	25.3 - 27.8	16,100

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	92.9	90.0 - 95.0	100	7.1	5.0 - 10.0	1,400
No	11,200	71.9	70.6 - 73.2	4,400	28.1	26.8 - 29.4	15,600

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2004-2006**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,100	85.6	83.4 - 87.7	500	14.4	12.3 - 16.6	3,600
No	9,500	70.5	69.1 - 71.9	4,000	29.5	28.1 - 30.9	13,500

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,000	79.9	78.2 - 81.5	1,500	20.1	18.5 - 21.8	7,600
No	6,600	69.0	67.4 - 70.6	3,000	31.0	29.4 - 32.6	9,600

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2003-2005**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,600	73.8	72.6 - 75.1	4,500	26.2	24.9 - 27.4	17,100

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	75.8	72.7 - 78.7	500	24.2	21.3 - 27.3	2,100
<b>Honolulu</b>	9,100	73.1	71.6 - 74.6	3,400	26.9	25.4 - 28.4	12,500
<b>Kauai</b>	600	79.5	74.6 - 83.7	200	20.5	16.3 - 25.4	800
<b>Maui</b>	1,300	74.1	70.7 - 77.2	500	25.9	22.8 - 29.3	1,800

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2003-2005**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,600	69.6	66.8 - 72.4	1,100	30.4	27.6 - 33.2	3,800
Native Hawaiian	3,800	80.8	78.8 - 82.7	900	19.2	17.3 - 21.2	4,800
Chinese	400	66.8	60.8 - 72.3	200	33.2	27.7 - 39.2	600
Filipino	2,600	76.1	73.2 - 78.7	800	23.9	21.3 - 26.8	3,400
Japanese	1,300	58.1	53.9 - 62.2	900	41.9	37.8 - 46.1	2,200
Other	1,900	79.5	76.2 - 82.5	500	20.5	17.5 - 23.8	2,400

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	87.7	84.3 - 90.4	200	12.3	9.6 - 15.7	1,400
20-24	3,500	82.2	79.9 - 84.2	800	17.8	15.8 - 20.1	4,200
25-34	6,000	71.0	69.1 - 72.8	2,400	29.0	27.2 - 30.9	8,400
35+	1,900	63.6	60.4 - 66.8	1,100	36.4	33.2 - 39.6	3,000

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2003-2005**

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Infant Gender</b>							
<b>Male</b>	6,800	74.9	73.2 - 76.5	2,300	25.1	23.5 - 26.8	9,100
<b>Female</b>	5,800	72.7	70.8 - 74.5	2,200	27.3	25.5 - 29.2	8,000

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Married</b>							
<b>Yes</b>	8,000	70.3	68.7 - 71.8	3,400	29.7	28.2 - 31.3	11,400
<b>No</b>	4,600	81.0	79.0 - 82.8	1,100	19.0	17.2 - 21.0	5,700

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Poverty Level</b>							
<b>0-130%</b>	5,300	79.4	77.5 - 81.2	1,400	20.6	18.8 - 22.5	6,700
<b>131-185%</b>	1,600	74.2	70.4 - 77.6	500	25.8	22.4 - 29.6	2,100
<b>186+%</b>	4,600	66.9	64.8 - 68.9	2,300	33.1	31.1 - 35.2	6,900

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2003-2005**

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	76.8	71.8 - 81.0	300	23.2	19.0 - 28.2	1,100
Normal birth weight	11,700	73.6	72.3 - 74.9	4,200	26.4	25.1 - 27.7	15,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	90.8	87.5 - 93.4	100	9.2	6.6 - 12.5	1,400
No	11,100	72.2	70.8 - 73.5	4,300	27.8	26.5 - 29.2	15,400

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,000	85.4	83.1 - 87.5	500	14.6	12.5 - 16.9	3,500
No	9,400	70.7	69.3 - 72.2	3,900	29.3	27.8 - 30.7	13,200

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2003-2005**

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,000	80.1	78.4 - 81.8	1,500	19.9	18.2 - 21.6	7,500
<b>No</b>	6,500	69.0	67.2 - 70.7	2,900	31.0	29.3 - 32.8	9,400



**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2002-2004**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,600	74.2	72.9 - 75.5	4,400	25.8	24.5 - 27.1	17,000

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	75.6	72.8 - 78.2	500	24.4	21.8 - 27.2	2,000
<b>Honolulu</b>	9,200	73.8	72.1 - 75.4	3,300	26.2	24.6 - 27.9	12,400
<b>Kauai</b>	600	77.7	73.2 - 81.6	200	22.3	18.4 - 26.8	800
<b>Maui</b>	1,300	74.3	71.3 - 77.0	500	25.7	23.0 - 28.7	1,800

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2002-2004**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,600	70.3	67.2 - 73.3	1,100	29.7	26.7 - 32.8	3,700
Native Hawaiian	3,700	79.5	77.7 - 81.1	1,000	20.5	18.9 - 22.3	4,700
Chinese	400	65.1	57.3 - 72.2	200	34.9	27.8 - 42.7	600
Filipino	2,700	77.7	74.7 - 80.5	800	22.3	19.5 - 25.3	3,400
Japanese	1,300	57.5	52.9 - 62.0	900	42.5	38.0 - 47.1	2,200
Other	1,900	82.8	79.3 - 85.8	400	17.2	14.2 - 20.7	2,300

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	86.7	82.9 - 89.7	200	13.3	10.3 - 17.1	1,400
20-24	3,400	82.7	80.4 - 84.8	700	17.3	15.2 - 19.6	4,200
25-34	6,000	71.6	69.6 - 73.5	2,400	28.4	26.5 - 30.4	8,400
35+	1,900	63.6	60.0 - 67.0	1,100	36.4	33.0 - 40.0	2,900

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2002-2004**

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Infant Gender</b>							
<b>Male</b>	6,700	75.3	73.6 - 77.0	2,200	24.7	23.0 - 26.4	8,900
<b>Female</b>	5,900	73.0	70.9 - 74.9	2,200	27.0	25.1 - 29.1	8,100

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Married</b>							
<b>Yes</b>	8,200	70.7	69.0 - 72.3	3,400	29.3	27.7 - 31.0	11,600
<b>No</b>	4,400	81.8	79.8 - 83.6	1,000	18.2	16.4 - 20.2	5,400

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Poverty Level</b>							
<b>0-130%</b>	5,500	80.2	78.3 - 82.0	1,300	19.8	18.0 - 21.7	6,800
<b>131-185%</b>	2,000	74.1	70.5 - 77.4	700	25.9	22.6 - 29.5	2,600
<b>186+%</b>	4,000	66.2	63.8 - 68.5	2,100	33.8	31.5 - 36.2	6,100

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2002-2004**

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	75.7	70.2 - 80.5	300	24.3	19.5 - 29.8	1,100
Normal birth weight	11,700	74.1	72.7 - 75.5	4,100	25.9	24.5 - 27.3	15,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	88.9	85.4 - 91.7	200	11.1	8.3 - 14.6	1,400
No	11,100	72.8	71.3 - 74.1	4,200	27.2	25.9 - 28.7	15,300

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	85.6	83.1 - 87.7	500	14.4	12.3 - 16.9	3,400
No	9,400	71.2	69.6 - 72.7	3,800	28.8	27.3 - 30.4	13,200

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2002-2004**

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,000	81.1	79.3 - 82.8	1,400	18.9	17.2 - 20.7	7,400
<b>No</b>	6,500	68.9	67.0 - 70.8	2,900	31.1	29.2 - 33.0	9,500

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2001-2003**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,400	74.5	73.2 - 75.8	4,300	25.5	24.2 - 26.8	16,700

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	77.2	74.9 - 79.4	400	22.8	20.6 - 25.1	1,900
<b>Honolulu</b>	9,100	73.8	72.1 - 75.5	3,200	26.2	24.5 - 27.9	12,300
<b>Kauai</b>	600	78.2	74.4 - 81.6	200	21.8	18.4 - 25.6	800
<b>Maui</b>	1,300	74.8	72.2 - 77.3	400	25.2	22.7 - 27.8	1,700

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2001-2003**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,500	70.0	66.8 - 73.1	1,100	30.0	26.9 - 33.2	3,600
Native Hawaiian	3,700	80.1	78.8 - 81.4	900	19.9	18.6 - 21.2	4,600
Chinese	500	66.8	58.1 - 74.4	200	33.2	25.6 - 41.9	700
Filipino	2,800	78.3	75.1 - 81.2	800	21.7	18.8 - 24.9	3,500
Japanese	1,200	57.2	52.3 - 61.9	900	42.8	38.1 - 47.7	2,100
Other	1,800	83.2	79.3 - 86.4	400	16.8	13.6 - 20.7	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	87.4	84.1 - 90.1	200	12.6	9.9 - 15.9	1,500
20-24	3,400	82.7	80.5 - 84.7	700	17.3	15.3 - 19.5	4,100
25-34	6,000	72.5	70.4 - 74.4	2,300	27.5	25.6 - 29.6	8,300
35+	1,700	61.2	57.4 - 65.0	1,100	38.8	35.0 - 42.6	2,700

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2001-2003**

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Infant Gender</b>							
<b>Male</b>	6,400	75.6	73.7 - 77.3	2,100	24.4	22.7 - 26.3	8,500
<b>Female</b>	6,000	73.4	71.4 - 75.3	2,200	26.6	24.7 - 28.6	8,200

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Married</b>							
<b>Yes</b>	7,900	70.5	68.7 - 72.3	3,300	29.5	27.7 - 31.3	11,200
<b>No</b>	4,500	82.7	80.8 - 84.4	900	17.3	15.6 - 19.2	5,500

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Poverty Level</b>							
<b>0-130%</b>	5,400	80.8	78.9 - 82.6	1,300	19.2	17.4 - 21.1	6,700
<b>131-185%</b>	2,300	74.0	70.8 - 77.0	800	26.0	23.0 - 29.2	3,100
<b>186+%</b>	3,500	65.2	62.5 - 67.9	1,900	34.8	32.1 - 37.5	5,400



**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2001-2003**

Income (2000-2003)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>&lt;=\$15,000</b>	2,900	82.8	80.3 - 85.0	600	17.2	15.0 - 19.7	3,500
<b>\$15,001-\$25,000</b>	1,800	80.7	77.5 - 83.6	400	19.3	16.4 - 22.5	2,300
<b>\$25,001-\$40,000</b>	2,600	74.8	71.8 - 77.7	900	25.2	22.3 - 28.2	3,400
<b>&gt;=\$40,001</b>	4,300	66.1	63.7 - 68.5	2,200	33.9	31.5 - 36.3	6,400

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Low birth weight</b>	900	77.3	71.8 - 82.0	300	22.7	18.0 - 28.2	1,100
<b>Normal birth weight</b>	11,600	74.4	73.0 - 75.7	4,000	25.6	24.3 - 27.0	15,500

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,300	89.2	85.7 - 91.9	200	10.8	8.1 - 14.3	1,400
<b>No</b>	10,900	73.1	71.6 - 74.5	4,000	26.9	25.5 - 28.4	14,900

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2001-2003**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	85.0	82.5 - 87.1	500	15.0	12.9 - 17.5	3,300
No	9,200	71.7	70.1 - 73.3	3,600	28.3	26.7 - 29.9	12,900

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,800	81.7	79.9 - 83.4	1,300	18.3	16.6 - 20.1	7,100
No	6,600	69.2	67.2 - 71.1	2,900	30.8	28.9 - 32.8	9,500

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2000-2002**

State	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,100	73.4	72.1 - 74.7	4,400	26.6	25.3 - 27.9	16,500

County	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,400	75.5	73.1 - 77.7	500	24.5	22.3 - 26.9	1,900
<b>Honolulu</b>	8,900	72.7	71.1 - 74.3	3,300	27.3	25.7 - 28.9	12,300
<b>Kauai</b>	500	77.4	73.4 - 80.9	200	22.6	19.1 - 26.6	700
<b>Maui</b>	1,300	74.6	72.0 - 77.1	400	25.4	22.9 - 28.0	1,700

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2000-2002**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,500	67.4	64.4 - 70.3	1,200	32.6	29.7 - 35.6	3,700
Native Hawaiian	3,600	79.7	78.4 - 80.9	900	20.3	19.1 - 21.6	4,500
Chinese	400	66.6	58.4 - 74.0	200	33.4	26.0 - 41.6	700
Filipino	2,600	77.9	74.8 - 80.6	800	22.1	19.4 - 25.2	3,400
Japanese	1,200	57.0	52.4 - 61.4	900	43.0	38.6 - 47.6	2,100
Other	1,800	81.7	78.1 - 84.9	400	18.3	15.1 - 21.9	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,400	85.9	82.6 - 88.7	200	14.1	11.3 - 17.4	1,600
20-24	3,300	80.4	78.2 - 82.4	800	19.6	17.6 - 21.8	4,100
25-34	5,800	71.7	69.8 - 73.5	2,300	28.3	26.5 - 30.2	8,100
35+	1,700	60.6	57.0 - 64.2	1,100	39.4	35.8 - 43.0	2,700

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2000-2002**

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Infant Gender</b>							
<b>Male</b>	6,200	74.4	72.6 - 76.1	2,200	25.6	23.9 - 27.4	8,400
<b>Female</b>	5,900	72.4	70.5 - 74.2	2,200	27.6	25.8 - 29.5	8,100

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Married</b>							
<b>Yes</b>	7,800	69.2	67.5 - 70.9	3,500	30.8	29.1 - 32.5	11,300
<b>No</b>	4,300	82.4	80.7 - 84.1	900	17.6	15.9 - 19.3	5,200

	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Poverty Level</b>							
<b>0-130%</b>	5,200	80.0	78.2 - 81.7	1,300	20.0	18.3 - 21.8	6,500
<b>131-185%</b>	2,300	75.0	72.0 - 77.8	800	25.0	22.2 - 28.0	3,100
<b>186+%</b>	3,600	64.0	61.5 - 66.4	2,000	36.0	33.6 - 38.5	5,600

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2000-2002**

Income (2000-2003)	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>&lt;=\$15,000</b>	2,800	81.9	79.6 - 84.0	600	18.1	16.0 - 20.4	3,400
<b>\$15,001-\$25,000</b>	1,900	78.6	75.5 - 81.4	500	21.4	18.6 - 24.5	2,400
<b>\$25,001-\$40,000</b>	2,600	75.2	72.3 - 77.8	900	24.8	22.2 - 27.7	3,500
<b>&gt;=\$40,001</b>	4,100	64.9	62.5 - 67.1	2,200	35.1	32.9 - 37.5	6,300

Birthweight	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Low birth weight</b>	800	77.7	72.5 - 82.2	200	22.3	17.8 - 27.5	1,100
<b>Normal birth weight</b>	11,300	73.1	71.8 - 74.4	4,100	26.9	25.6 - 28.2	15,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	1,200	88.4	85.2 - 91.0	200	11.6	9.0 - 14.8	1,400
<b>No</b>	10,700	72.0	70.6 - 73.4	4,100	28.0	26.6 - 29.4	14,800

**PRAMS Health Indicator 3-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2002;2001-2003;2002-2004;2003-2005;2004-2006;2005-2007;2006-2008;2007-2009;2008-2010;2009-2011;2010-2012;2011-2013**

*Note: Weighted counts (#) represent 3-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Years: 2000-2002**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,700	84.3	81.9 - 86.4	500	15.7	13.6 - 18.1	3,200
<b>No</b>	9,000	70.4	68.9 - 71.9	3,800	29.6	28.1 - 31.1	12,800

WIC During Pregnancy	Yes			No			Weighted Total Responses (3-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,700	80.5	78.8 - 82.1	1,400	19.5	17.9 - 21.2	7,000
<b>No</b>	6,400	68.2	66.3 - 70.0	3,000	31.8	30.0 - 33.7	9,300