

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

*The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.*

*Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year aggregate periods.*

*Below are the indicators included in this report and the long description for each indicator.*

PRAMS Health Indicator	Health Indicator Description
Discuss effects of smoking with HCW during PNC	During any of your prenatal care visits, did a doctor, nurse, or other health care worker talk with you about how smoking during pregnancy could affect your baby?

*Suggested citation:*

*Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/23/17*

*For documentation on other terms and coding in this report, please go to the following links:*

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact  
The Hawai'i Health Data Warehouse**

[profiles@hhdw.org](mailto:profiles@hhdw.org)

**(855) 946-5899 x15**

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**


*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

Years	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000-2004	12,400	73.8	72.8 - 74.7	4,400	26.2	25.3 - 27.2	16,800
2001-2005	12,400	73.8	72.8 - 74.8	4,400	26.2	25.2 - 27.2	16,800
2002-2006	12,700	74.3	73.3 - 75.2	4,400	25.7	24.8 - 26.7	17,100
2003-2007	12,800	73.7	72.8 - 74.7	4,600	26.3	25.3 - 27.2	17,400
2004-2008	12,800	72.7	71.7 - 73.6	4,800	27.3	26.4 - 28.3	17,600
2005-2009	12,900	72.5	71.5 - 73.5	4,900	27.5	26.5 - 28.5	17,800
2006-2010	13,100	72.4	71.2 - 73.4	5,000	27.6	26.6 - 28.8	18,000
2007-2011	13,000	72.0	70.8 - 73.1	5,100	28.0	26.9 - 29.2	18,100
2008-2012	13,000	71.8	70.6 - 73.0	5,100	28.2	27.0 - 29.4	18,100
2009-2013	12,900	71.5	70.1 - 72.8	5,200	28.5	27.2 - 29.9	18,100

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*



**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,900	71.5	70.1 - 72.8	5,200	28.5	27.2 - 29.9	18,100

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,700	75.2	73.1 - 77.1	600	24.8	22.9 - 26.9	2,300
<b>Honolulu</b>	9,200	70.6	68.8 - 72.3	3,800	29.4	27.7 - 31.2	13,000
<b>Kauai</b>	600	74.3	71.9 - 76.6	200	25.7	23.4 - 28.1	800
<b>Maui</b>	1,400	71.7	69.7 - 73.7	600	28.3	26.3 - 30.3	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	66.1	63.2 - 68.8	1,500	33.9	31.2 - 36.8	4,400
Native Hawaiian	4,300	79.5	77.4 - 81.5	1,100	20.5	18.5 - 22.6	5,400
Chinese	500	61.6	54.4 - 68.3	300	38.4	31.7 - 45.6	800
Filipino	2,300	71.8	68.5 - 74.8	900	28.2	25.2 - 31.5	3,200
Japanese	900	55.0	50.0 - 59.9	800	45.0	40.1 - 50.0	1,700
Other	2,000	77.3	73.7 - 80.5	600	22.7	19.5 - 26.3	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	88.8	84.7 - 91.9	100	11.2	8.1 - 15.3	1,100
20-24	3,400	82.1	79.5 - 84.4	700	17.9	15.6 - 20.5	4,100
25-34	6,700	69.2	67.3 - 71.1	3,000	30.8	28.9 - 32.7	9,600
35+	1,900	58.4	54.9 - 61.7	1,300	41.6	38.3 - 45.1	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,400	69.9	68.0 - 71.8	2,800	30.1	28.2 - 32.0	9,200
Female	6,500	73.1	71.2 - 74.9	2,400	26.9	25.1 - 28.8	8,900

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,400	66.1	64.3 - 67.8	3,800	33.9	32.2 - 35.7	11,200
No	5,500	80.3	78.3 - 82.1	1,400	19.7	17.9 - 21.7	6,900

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,200	80.7	78.7 - 82.5	1,200	19.3	17.5 - 21.3	6,400
131-185%	2,100	75.4	72.0 - 78.5	700	24.6	21.5 - 28.0	2,800
186+%	4,600	61.1	58.9 - 63.4	2,900	38.9	36.6 - 41.1	7,400

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	74.4	72.3 - 76.4	300	25.6	23.6 - 27.7	1,300
Normal birth weight	12,000	71.2	69.8 - 72.6	4,800	28.8	27.4 - 30.2	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,000	91.3	87.5 - 94.0	100	8.7	6.0 - 12.5	1,100
No	11,600	70.1	68.6 - 71.5	5,000	29.9	28.5 - 31.4	16,600

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	84.4	81.9 - 86.7	500	15.6	13.3 - 18.1	3,400
No	9,800	68.3	66.7 - 69.8	4,500	31.7	30.2 - 33.3	14,300

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,300	79.2	77.4 - 80.9	1,600	20.8	19.1 - 22.6	7,900
No	6,600	65.4	63.4 - 67.2	3,500	34.6	32.8 - 36.6	10,000



**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,000	71.8	70.6 - 73.0	5,100	28.2	27.0 - 29.4	18,100

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,800	75.9	73.7 - 77.9	600	24.1	22.1 - 26.3	2,400
<b>Honolulu</b>	9,200	71.0	69.3 - 72.6	3,800	29.0	27.4 - 30.7	12,900
<b>Kauai</b>	600	72.0	68.9 - 75.0	200	28.0	25.0 - 31.1	800
<b>Maui</b>	1,400	72.1	69.9 - 74.2	600	27.9	25.8 - 30.1	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	67.0	64.3 - 69.5	1,400	33.0	30.5 - 35.7	4,300
Native Hawaiian	4,200	78.1	76.0 - 80.1	1,200	21.9	19.9 - 24.0	5,400
Chinese	500	61.8	55.1 - 68.1	300	38.2	31.9 - 44.9	800
Filipino	2,400	73.0	70.0 - 75.8	900	27.0	24.2 - 30.0	3,300
Japanese	1,000	56.2	51.7 - 60.6	800	43.8	39.4 - 48.3	1,700
Other	2,100	78.5	75.3 - 81.4	600	21.5	18.6 - 24.7	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	89.1	85.5 - 91.8	100	10.9	8.2 - 14.5	1,200
20-24	3,400	82.5	80.1 - 84.6	700	17.5	15.4 - 19.9	4,100
25-34	6,600	69.0	67.2 - 70.7	3,000	31.0	29.3 - 32.8	9,600
35+	1,900	59.9	56.7 - 63.0	1,300	40.1	37.0 - 43.3	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,600	70.8	69.0 - 72.5	2,700	29.2	27.5 - 31.0	9,300
Female	6,400	72.9	71.1 - 74.6	2,400	27.1	25.4 - 28.9	8,800

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,500	66.7	65.0 - 68.3	3,800	33.3	31.7 - 35.0	11,200
No	5,500	80.2	78.4 - 81.9	1,400	19.8	18.1 - 21.6	6,900

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,200	80.5	78.7 - 82.3	1,200	19.5	17.7 - 21.3	6,400
131-185%	2,100	74.5	71.3 - 77.5	700	25.5	22.5 - 28.7	2,800
186+%	4,700	62.3	60.2 - 64.3	2,900	37.7	35.7 - 39.8	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	72.8	70.2 - 75.2	300	27.2	24.8 - 29.8	1,300
Normal birth weight	12,100	71.7	70.4 - 73.0	4,800	28.3	27.0 - 29.6	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	91.6	88.5 - 94.0	100	8.4	6.0 - 11.5	1,200
No	11,600	70.3	68.9 - 71.6	4,900	29.7	28.4 - 31.1	16,500

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	85.1	82.7 - 87.1	500	14.9	12.9 - 17.3	3,500
No	9,800	68.5	67.1 - 70.0	4,500	31.5	30.0 - 32.9	14,300

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,300	79.8	78.1 - 81.4	1,600	20.2	18.6 - 21.9	7,900
<b>No</b>	6,600	65.6	63.8 - 67.3	3,500	34.4	32.7 - 36.2	10,100

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,000	72.0	70.8 - 73.1	5,100	28.0	26.9 - 29.2	18,100

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,800	74.4	72.0 - 76.6	600	25.6	23.4 - 28.0	2,400
<b>Honolulu</b>	9,200	71.6	70.0 - 73.1	3,600	28.4	26.9 - 30.0	12,800
<b>Kauai</b>	700	72.6	69.1 - 75.8	200	27.4	24.2 - 30.9	900
<b>Maui</b>	1,500	71.4	68.9 - 73.7	600	28.6	26.3 - 31.1	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	68.0	65.5 - 70.4	1,400	32.0	29.6 - 34.5	4,300
Native Hawaiian	4,100	78.0	75.9 - 79.9	1,200	22.0	20.1 - 24.1	5,300
Chinese	400	61.4	55.1 - 67.4	300	38.6	32.6 - 44.9	700
Filipino	2,400	73.8	71.0 - 76.3	900	26.2	23.7 - 29.0	3,300
Japanese	1,000	56.2	52.1 - 60.2	800	43.8	39.8 - 47.9	1,800
Other	2,100	77.7	74.6 - 80.5	600	22.3	19.5 - 25.4	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	89.9	86.9 - 92.2	100	10.1	7.8 - 13.1	1,400
20-24	3,500	82.1	79.9 - 84.0	800	17.9	16.0 - 20.1	4,300
25-34	6,400	68.7	67.0 - 70.3	2,900	31.3	29.7 - 33.0	9,300
35+	1,900	60.1	57.1 - 63.1	1,300	39.9	36.9 - 42.9	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,600	70.9	69.3 - 72.5	2,700	29.1	27.5 - 30.7	9,400
Female	6,400	73.0	71.4 - 74.7	2,400	27.0	25.3 - 28.6	8,800

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,500	66.6	65.1 - 68.2	3,800	33.4	31.8 - 34.9	11,200
No	5,500	80.7	78.9 - 82.3	1,300	19.3	17.7 - 21.1	6,900

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,200	80.2	78.5 - 81.9	1,300	19.8	18.1 - 21.5	6,500
131-185%	2,000	73.4	70.2 - 76.2	700	26.6	23.8 - 29.8	2,700
186+%	4,800	62.8	60.9 - 64.7	2,800	37.2	35.3 - 39.1	7,600



**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,600	83.2	80.8 - 85.4	500	16.8	14.6 - 19.2	3,100
\$10,000-\$24,999	2,900	80.0	77.6 - 82.1	700	20.0	17.9 - 22.4	3,600
\$25,000-\$49,999	3,000	72.0	69.5 - 74.3	1,200	28.0	25.7 - 30.5	4,200
>=\$50,000	3,600	59.6	57.4 - 61.8	2,500	40.4	38.2 - 42.6	6,100

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	73.6	70.7 - 76.2	400	26.4	23.8 - 29.3	1,300
Normal birth weight	12,000	71.8	70.5 - 73.0	4,700	28.2	27.0 - 29.5	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	91.1	88.2 - 93.3	100	8.9	6.7 - 11.8	1,400
No	11,600	70.4	69.1 - 71.6	4,900	29.6	28.4 - 30.9	16,400

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,100	85.5	83.4 - 87.4	500	14.5	12.6 - 16.6	3,600
<b>No</b>	9,700	68.5	67.1 - 69.9	4,500	31.5	30.1 - 32.9	14,200

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,200	79.7	78.1 - 81.3	1,600	20.3	18.7 - 21.9	7,700
<b>No</b>	6,700	66.0	64.3 - 67.6	3,500	34.0	32.4 - 35.7	10,200

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,100	72.4	71.2 - 73.4	5,000	27.6	26.6 - 28.8	18,000

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,700	73.9	71.4 - 76.3	600	26.1	23.7 - 28.6	2,300
<b>Honolulu</b>	9,200	72.1	70.7 - 73.5	3,600	27.9	26.5 - 29.3	12,800
<b>Kauai</b>	700	74.0	70.2 - 77.5	200	26.0	22.5 - 29.8	900
<b>Maui</b>	1,500	71.5	68.9 - 74.0	600	28.5	26.0 - 31.1	2,000

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,000	69.3	67.0 - 71.5	1,300	30.7	28.5 - 33.0	4,300
Native Hawaiian	4,100	77.9	75.9 - 79.8	1,200	22.1	20.2 - 24.1	5,200
Chinese	400	57.3	51.6 - 62.9	300	42.7	37.1 - 48.4	700
Filipino	2,400	73.7	71.1 - 76.2	900	26.3	23.8 - 28.9	3,300
Japanese	1,100	58.5	54.8 - 62.1	800	41.5	37.9 - 45.2	1,900
Other	2,100	78.2	75.4 - 80.8	600	21.8	19.2 - 24.6	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	89.1	86.2 - 91.5	200	10.9	8.5 - 13.8	1,500
20-24	3,400	80.4	78.3 - 82.4	800	19.6	17.6 - 21.7	4,300
25-34	6,300	69.7	68.1 - 71.3	2,700	30.3	28.7 - 31.9	9,100
35+	1,900	61.1	58.3 - 63.8	1,200	38.9	36.2 - 41.7	3,200

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,600	71.2	69.6 - 72.7	2,700	28.8	27.3 - 30.4	9,300
Female	6,400	73.6	72.0 - 75.2	2,300	26.4	24.8 - 28.0	8,700

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,600	67.4	65.9 - 68.8	3,700	32.6	31.2 - 34.1	11,300
No	5,500	80.6	78.9 - 82.1	1,300	19.4	17.9 - 21.1	6,800

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,100	79.9	78.2 - 81.5	1,300	20.1	18.5 - 21.8	6,400
131-185%	2,000	74.3	71.4 - 77.1	700	25.7	22.9 - 28.6	2,700
186+%	4,900	63.8	62.0 - 65.6	2,800	36.2	34.4 - 38.0	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,500	82.1	79.7 - 84.4	500	17.9	15.6 - 20.3	3,000
\$10,000-\$24,999	3,000	80.5	78.3 - 82.5	700	19.5	17.5 - 21.7	3,700
\$25,000-\$49,999	2,900	70.8	68.4 - 73.1	1,200	29.2	26.9 - 31.6	4,100
>=\$50,000	3,700	61.8	59.7 - 63.8	2,300	38.2	36.2 - 40.3	6,000

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	74.3	71.3 - 77.1	300	25.7	22.9 - 28.7	1,400
Normal birth weight	12,000	72.2	71.0 - 73.3	4,600	27.8	26.7 - 29.0	16,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	91.7	89.2 - 93.7	100	8.3	6.3 - 10.8	1,500
No	11,400	70.6	69.4 - 71.7	4,800	29.4	28.3 - 30.6	16,200

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,100	85.3	83.3 - 87.2	500	14.7	12.8 - 16.7	3,600
<b>No</b>	9,700	69.0	67.7 - 70.3	4,400	31.0	29.7 - 32.3	14,100

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,100	79.3	77.8 - 80.8	1,600	20.7	19.2 - 22.2	7,600
<b>No</b>	6,800	67.1	65.5 - 68.6	3,400	32.9	31.4 - 34.5	10,200

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,900	72.5	71.5 - 73.5	4,900	27.5	26.5 - 28.5	17,800

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,700	73.3	70.7 - 75.8	600	26.7	24.2 - 29.3	2,300
<b>Honolulu</b>	9,100	72.4	71.1 - 73.6	3,500	27.6	26.4 - 28.9	12,600
<b>Kauai</b>	700	75.0	70.9 - 78.7	200	25.0	21.3 - 29.1	900
<b>Maui</b>	1,400	71.5	68.6 - 74.2	600	28.5	25.8 - 31.4	2,000



**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,800	68.4	66.2 - 70.5	1,300	31.6	29.5 - 33.8	4,100
Native Hawaiian	4,000	78.8	76.9 - 80.6	1,100	21.2	19.4 - 23.1	5,100
Chinese	400	59.6	54.7 - 64.2	300	40.4	35.8 - 45.3	600
Filipino	2,400	75.3	73.0 - 77.4	800	24.7	22.6 - 27.0	3,300
Japanese	1,200	59.8	56.5 - 63.0	800	40.2	37.0 - 43.5	2,000
Other	2,000	76.4	73.7 - 78.9	600	23.6	21.1 - 26.3	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,400	88.7	86.1 - 91.0	200	11.3	9.0 - 13.9	1,600
20-24	3,400	79.6	77.6 - 81.5	900	20.4	18.5 - 22.4	4,200
25-34	6,200	70.0	68.5 - 71.4	2,700	30.0	28.6 - 31.5	8,900
35+	1,900	61.9	59.3 - 64.5	1,200	38.1	35.5 - 40.7	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,600	71.8	70.3 - 73.2	2,600	28.2	26.8 - 29.7	9,200
Female	6,300	73.4	71.9 - 74.8	2,300	26.6	25.2 - 28.1	8,600

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,500	67.2	65.8 - 68.5	3,600	32.8	31.5 - 34.2	11,100
No	5,400	81.5	79.9 - 82.9	1,200	18.5	17.1 - 20.1	6,700

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,100	79.3	77.7 - 80.8	1,300	20.7	19.2 - 22.3	6,400
131-185%	1,800	74.4	71.6 - 77.0	600	25.6	23.0 - 28.4	2,400
186+%	4,900	64.7	63.1 - 66.3	2,700	35.3	33.7 - 36.9	7,600

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,400	81.8	79.4 - 84.0	500	18.2	16.0 - 20.6	2,900
\$10,000-\$24,999	3,000	79.4	77.3 - 81.3	800	20.6	18.7 - 22.7	3,800
\$25,000-\$49,999	2,900	71.6	69.4 - 73.8	1,100	28.4	26.2 - 30.6	4,000
>=\$50,000	3,600	62.3	60.4 - 64.2	2,200	37.7	35.8 - 39.6	5,800

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	74.8	71.6 - 77.8	300	25.2	22.2 - 28.4	1,300
Normal birth weight	11,900	72.3	71.3 - 73.4	4,500	27.7	26.6 - 28.7	16,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	92.4	90.1 - 94.1	100	7.6	5.9 - 9.9	1,500
No	11,300	70.7	69.5 - 71.7	4,700	29.3	28.3 - 30.5	15,900

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,100	86.1	84.2 - 87.7	500	13.9	12.3 - 15.8	3,600
<b>No</b>	9,600	69.0	67.8 - 70.2	4,300	31.0	29.8 - 32.2	13,800

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,000	79.5	78.1 - 80.9	1,500	20.5	19.1 - 21.9	7,500
<b>No</b>	6,800	67.3	65.9 - 68.7	3,300	32.7	31.3 - 34.1	10,100

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,800	72.7	71.7 - 73.6	4,800	27.3	26.4 - 28.3	17,600

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	73.1	70.3 - 75.7	600	26.9	24.3 - 29.7	2,200
<b>Honolulu</b>	9,200	72.6	71.5 - 73.7	3,400	27.4	26.3 - 28.5	12,600
<b>Kauai</b>	600	74.5	70.1 - 78.5	200	25.5	21.5 - 29.9	900
<b>Maui</b>	1,400	71.4	68.4 - 74.3	500	28.6	25.7 - 31.6	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,800	68.3	66.2 - 70.3	1,300	31.7	29.7 - 33.8	4,100
Native Hawaiian	3,900	78.6	76.8 - 80.3	1,100	21.4	19.7 - 23.2	4,900
Chinese	400	62.2	59.3 - 64.9	200	37.8	35.1 - 40.7	600
Filipino	2,500	75.9	73.8 - 77.8	800	24.1	22.2 - 26.2	3,300
Japanese	1,200	58.3	55.3 - 61.2	900	41.7	38.8 - 44.7	2,100
Other	2,100	78.4	76.1 - 80.5	600	21.6	19.5 - 23.9	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	87.1	84.2 - 89.5	200	12.9	10.5 - 15.8	1,500
20-24	3,400	79.0	77.2 - 80.8	900	21.0	19.2 - 22.8	4,300
25-34	6,200	70.7	69.3 - 72.0	2,600	29.3	28.0 - 30.7	8,700
35+	2,000	62.8	60.5 - 65.0	1,200	37.2	35.0 - 39.5	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,700	72.3	71.0 - 73.6	2,500	27.7	26.4 - 29.0	9,200
Female	6,200	73.0	71.7 - 74.4	2,300	27.0	25.6 - 28.3	8,400

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,800	68.5	67.3 - 69.6	3,600	31.5	30.4 - 32.7	11,400
No	5,000	80.3	78.8 - 81.8	1,200	19.7	18.2 - 21.2	6,200

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,000	78.8	77.3 - 80.3	1,400	21.2	19.7 - 22.7	6,400
131-185%	1,700	73.7	71.0 - 76.2	600	26.3	23.8 - 29.0	2,300
186+%	5,100	66.2	64.7 - 67.6	2,600	33.8	32.4 - 35.3	7,700

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Income (2004-2011)	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,300	81.1	78.9 - 83.2	500	18.9	16.8 - 21.1	2,900
\$10,000-\$24,999	3,000	78.3	76.4 - 80.2	800	21.7	19.8 - 23.6	3,800
\$25,000-\$49,999	3,000	72.3	70.3 - 74.2	1,100	27.7	25.8 - 29.7	4,100
>=\$50,000	3,600	63.5	61.7 - 65.2	2,000	36.5	34.8 - 38.3	5,600

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	74.8	71.3 - 78.0	300	25.2	22.0 - 28.7	1,300
Normal birth weight	11,800	72.5	71.5 - 73.5	4,500	27.5	26.5 - 28.5	16,300

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	91.2	88.9 - 93.2	100	8.8	6.8 - 11.1	1,500
No	11,300	70.9	69.9 - 71.9	4,600	29.1	28.1 - 30.1	15,900



**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,000	85.0	83.2 - 86.6	500	15.0	13.4 - 16.8	3,500
<b>No</b>	9,600	69.5	68.4 - 70.6	4,200	30.5	29.4 - 31.6	13,800

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,000	79.1	77.8 - 80.5	1,600	20.9	19.5 - 22.2	7,600
<b>No</b>	6,700	67.7	66.4 - 69.0	3,200	32.3	31.0 - 33.6	9,900

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,800	73.7	72.8 - 74.7	4,600	26.3	25.3 - 27.2	17,400

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	73.2	70.6 - 75.7	600	26.8	24.3 - 29.4	2,100
<b>Honolulu</b>	9,200	73.6	72.5 - 74.8	3,300	26.4	25.2 - 27.5	12,600
<b>Kauai</b>	700	78.4	74.4 - 81.9	200	21.6	18.1 - 25.6	900
<b>Maui</b>	1,400	73.0	70.1 - 75.6	500	27.0	24.4 - 29.9	1,900

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,800	70.1	67.9 - 72.1	1,200	29.9	27.9 - 32.1	3,900
Native Hawaiian	3,900	80.3	78.6 - 81.9	900	19.7	18.1 - 21.4	4,800
Chinese	400	64.5	60.4 - 68.4	200	35.5	31.6 - 39.6	600
Filipino	2,500	75.7	73.5 - 77.7	800	24.3	22.3 - 26.5	3,300
Japanese	1,300	59.0	55.9 - 62.1	900	41.0	37.9 - 44.1	2,100
Other	2,000	79.1	76.6 - 81.4	500	20.9	18.6 - 23.4	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	87.6	84.9 - 89.9	200	12.4	10.1 - 15.1	1,500
20-24	3,400	80.2	78.4 - 81.9	800	19.8	18.1 - 21.6	4,300
25-34	6,200	71.8	70.4 - 73.2	2,400	28.2	26.8 - 29.6	8,600
35+	1,900	63.3	60.8 - 65.7	1,100	36.7	34.3 - 39.2	3,100

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,700	74.2	72.9 - 75.5	2,300	25.8	24.5 - 27.1	9,100
Female	6,100	73.2	71.8 - 74.6	2,200	26.8	25.4 - 28.2	8,300

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,900	69.8	68.6 - 71.0	3,400	30.2	29.0 - 31.4	11,400
No	4,900	81.2	79.7 - 82.6	1,100	18.8	17.4 - 20.3	6,100

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,200	79.6	78.1 - 81.0	1,300	20.4	19.0 - 21.9	6,600
131-185%	1,700	74.1	71.3 - 76.7	600	25.9	23.3 - 28.7	2,300
186+%	4,800	67.2	65.6 - 68.7	2,400	32.8	31.3 - 34.4	7,200

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	76.6	73.1 - 79.8	300	23.4	20.2 - 26.9	1,300
Normal birth weight	11,800	73.5	72.5 - 74.5	4,300	26.5	25.5 - 27.5	16,100

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	91.2	88.7 - 93.1	100	8.8	6.9 - 11.3	1,500
No	11,300	72.1	71.1 - 73.1	4,400	27.9	26.9 - 28.9	15,700

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,000	85.5	83.6 - 87.1	500	14.5	12.9 - 16.4	3,500
No	9,600	70.8	69.6 - 71.9	4,000	29.2	28.1 - 30.4	13,500

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,000	79.7	78.3 - 81.0	1,500	20.3	19.0 - 21.7	7,500
No	6,700	69.2	67.9 - 70.5	3,000	30.8	29.5 - 32.1	9,700

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,700	74.3	73.3 - 75.2	4,400	25.7	24.8 - 26.7	17,100

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	76.3	73.9 - 78.6	500	23.7	21.4 - 26.1	2,100
<b>Honolulu</b>	9,200	73.7	72.4 - 74.9	3,300	26.3	25.1 - 27.6	12,500
<b>Kauai</b>	600	78.3	74.5 - 81.8	200	21.7	18.2 - 25.5	800
<b>Maui</b>	1,300	74.4	71.8 - 76.9	500	25.6	23.1 - 28.2	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	69.6	67.3 - 71.8	1,200	30.4	28.2 - 32.7	3,800
Native Hawaiian	3,800	80.4	78.8 - 81.8	900	19.6	18.2 - 21.2	4,700
Chinese	400	65.2	60.1 - 70.0	200	34.8	30.0 - 39.9	600
Filipino	2,600	78.1	75.9 - 80.2	700	21.9	19.8 - 24.1	3,400
Japanese	1,300	58.4	55.1 - 61.7	900	41.6	38.3 - 44.9	2,200
Other	2,000	81.0	78.4 - 83.3	500	19.0	16.7 - 21.6	2,400

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	87.0	84.1 - 89.4	200	13.0	10.6 - 15.9	1,500
20-24	3,400	81.5	79.7 - 83.1	800	18.5	16.9 - 20.3	4,200
25-34	6,100	72.4	70.9 - 73.8	2,300	27.6	26.2 - 29.1	8,500
35+	1,900	63.2	60.6 - 65.7	1,100	36.8	34.3 - 39.4	3,000



**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,700	75.3	74.0 - 76.6	2,200	24.7	23.4 - 26.0	8,900
Female	6,000	73.1	71.7 - 74.6	2,200	26.9	25.4 - 28.3	8,300

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,000	70.2	68.9 - 71.5	3,400	29.8	28.5 - 31.1	11,400
No	4,700	82.3	80.8 - 83.7	1,000	17.7	16.3 - 19.2	5,800

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,300	80.5	79.0 - 81.9	1,300	19.5	18.1 - 21.0	6,500
131-185%	1,800	75.1	72.3 - 77.7	600	24.9	22.3 - 27.7	2,400
186+%	4,500	66.7	65.1 - 68.4	2,300	33.3	31.6 - 34.9	6,800

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	76.8	72.9 - 80.2	300	23.2	19.8 - 27.1	1,200
Normal birth weight	11,800	74.1	73.1 - 75.1	4,100	25.9	24.9 - 26.9	15,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	91.4	89.0 - 93.3	100	8.6	6.7 - 11.0	1,400
No	11,200	72.6	71.6 - 73.7	4,200	27.4	26.3 - 28.4	15,400

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	3,000	86.0	84.1 - 87.6	500	14.0	12.4 - 15.9	3,500
No	9,500	71.2	70.0 - 72.3	3,800	28.8	27.7 - 30.0	13,300

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,000	80.9	79.5 - 82.2	1,400	19.1	17.8 - 20.5	7,400
No	6,600	69.2	67.8 - 70.6	2,900	30.8	29.4 - 32.2	9,600

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,400	73.8	72.8 - 74.8	4,400	26.2	25.2 - 27.2	16,800

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	76.3	74.1 - 78.4	500	23.7	21.6 - 25.9	2,000
<b>Honolulu</b>	9,000	73.1	71.9 - 74.4	3,300	26.9	25.6 - 28.1	12,400
<b>Kauai</b>	600	77.7	74.1 - 80.8	200	22.3	19.2 - 25.9	800
<b>Maui</b>	1,300	73.9	71.5 - 76.2	500	26.1	23.8 - 28.5	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,500	68.8	66.4 - 71.0	1,100	31.2	29.0 - 33.6	3,700
Native Hawaiian	3,700	80.5	79.1 - 81.8	900	19.5	18.2 - 20.9	4,600
Chinese	400	65.9	60.4 - 71.0	200	34.1	29.0 - 39.6	700
Filipino	2,600	77.4	75.2 - 79.5	800	22.6	20.5 - 24.8	3,400
Japanese	1,200	56.9	53.5 - 60.2	900	43.1	39.8 - 46.5	2,200
Other	1,900	81.1	78.4 - 83.5	400	18.9	16.5 - 21.6	2,300

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	87.0	84.5 - 89.3	200	13.0	10.7 - 15.5	1,500
20-24	3,400	81.8	80.1 - 83.4	800	18.2	16.6 - 19.9	4,200
25-34	5,900	71.6	70.1 - 73.0	2,400	28.4	27.0 - 29.9	8,300
35+	1,800	61.7	59.0 - 64.3	1,100	38.3	35.7 - 41.0	2,900

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,500	74.7	73.4 - 76.0	2,200	25.3	24.0 - 26.6	8,700
Female	5,900	72.8	71.3 - 74.2	2,200	27.2	25.8 - 28.7	8,100

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,900	69.9	68.6 - 71.1	3,400	30.1	28.9 - 31.4	11,300
No	4,500	81.8	80.3 - 83.2	1,000	18.2	16.8 - 19.7	5,500

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,200	80.2	78.7 - 81.5	1,300	19.8	18.5 - 21.3	6,500
131-185%	2,000	74.0	71.4 - 76.4	700	26.0	23.6 - 28.6	2,700
186+%	4,100	65.6	63.9 - 67.4	2,200	34.4	32.6 - 36.1	6,300

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	76.7	72.7 - 80.2	300	23.3	19.8 - 27.3	1,100
Normal birth weight	11,600	73.6	72.6 - 74.6	4,100	26.4	25.4 - 27.4	15,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	90.4	87.9 - 92.4	100	9.6	7.6 - 12.1	1,400
No	10,900	72.2	71.1 - 73.2	4,200	27.8	26.8 - 28.9	15,100

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,900	84.9	83.1 - 86.5	500	15.1	13.5 - 16.9	3,400
No	9,200	70.8	69.6 - 71.9	3,800	29.2	28.1 - 30.4	13,000

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	5,900	80.8	79.4 - 82.0	1,400	19.2	18.0 - 20.6	7,300
No	6,500	68.5	67.1 - 69.9	3,000	31.5	30.1 - 32.9	9,400



**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

State	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,400	73.8	72.8 - 74.7	4,400	26.2	25.3 - 27.2	16,800

County	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	75.0	73.0 - 77.0	500	25.0	23.0 - 27.0	1,900
<b>Honolulu</b>	9,000	73.2	71.9 - 74.4	3,300	26.8	25.6 - 28.1	12,400
<b>Kauai</b>	600	78.3	75.0 - 81.2	200	21.7	18.8 - 25.0	700
<b>Maui</b>	1,300	74.5	72.3 - 76.6	400	25.5	23.4 - 27.7	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,600	69.1	66.8 - 71.3	1,200	30.9	28.7 - 33.2	3,700
Native Hawaiian	3,700	79.7	78.5 - 80.8	900	20.3	19.2 - 21.5	4,600
Chinese	400	66.5	60.5 - 71.9	200	33.5	28.1 - 39.5	700
Filipino	2,600	77.2	74.9 - 79.3	800	22.8	20.7 - 25.1	3,400
Japanese	1,200	57.7	54.2 - 61.2	900	42.3	38.8 - 45.8	2,100
Other	1,800	81.4	78.6 - 83.8	400	18.6	16.2 - 21.4	2,300

PRAMS Age Group	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	86.7	84.2 - 88.8	200	13.3	11.2 - 15.8	1,500
20-24	3,400	81.2	79.5 - 82.8	800	18.8	17.2 - 20.5	4,100
25-34	5,900	71.5	70.1 - 73.0	2,300	28.5	27.0 - 29.9	8,200
35+	1,800	62.4	59.7 - 65.0	1,100	37.6	35.0 - 40.3	2,900

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

Infant Gender	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,500	74.7	73.3 - 75.9	2,200	25.3	24.1 - 26.7	8,800
Female	5,900	72.8	71.3 - 74.2	2,200	27.2	25.8 - 28.7	8,000

Married	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,000	70.1	68.8 - 71.4	3,400	29.9	28.6 - 31.2	11,500
No	4,400	81.6	80.1 - 82.9	1,000	18.4	17.1 - 19.9	5,300

Poverty Level	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,400	79.9	78.5 - 81.2	1,400	20.1	18.8 - 21.5	6,700
131-185%	2,100	74.7	72.2 - 77.0	700	25.3	23.0 - 27.8	2,800
186+%	3,900	65.4	63.6 - 67.2	2,100	34.6	32.8 - 36.4	6,000

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

Birthweight	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	77.4	73.4 - 80.9	200	22.6	19.1 - 26.6	1,100
Normal birth weight	11,500	73.5	72.5 - 74.5	4,200	26.5	25.5 - 27.5	15,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	88.4	85.8 - 90.6	200	11.6	9.4 - 14.2	1,400
No	10,900	72.3	71.3 - 73.4	4,200	27.7	26.6 - 28.7	15,100

Smoke Before Pregnant	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	2,800	84.7	82.9 - 86.3	500	15.3	13.7 - 17.1	3,300
No	9,200	70.8	69.6 - 72.0	3,800	29.2	28.0 - 30.4	13,000

**PRAMS Health Indicator 5-Year Aggregate Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

WIC During Pregnancy	Yes			No			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,800	80.4	79.0 - 81.6	1,400	19.6	18.4 - 21.0	7,200
<b>No</b>	6,500	68.8	67.4 - 70.2	2,900	31.2	29.8 - 32.6	9,400