

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy-  
for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH  
Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before  
Pregnant;WIC During Pregnancy, for the Year(s)

2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted  
total responses for a dimension (row) is <30 that row will not be shown*

*The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. 5-Year Weighted Total  
Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the  
survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the  
population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered  
yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the  
counts for that particular dimension.*

*Note: Data collection for household income changed in 2004 and again in 2012. Because of this, 5-year aggregate data does not exist for all 5-year  
aggregate periods.*

*Below are the indicators included in this report and the long description for each indicator.*

PRAMS Health Indicator	Health Indicator Description
Number of cigarettes smoked per avg day before pregnancy	Number of cigarettes smoked on an average day 3 months before pregnancy among new mothers who reported smoking in the last 2 years

*Suggested citation:*

*Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and  
years), Report Created: 2/22/17*

*For documentation on other terms and coding in this report, please go to the following links:*

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact  
The Hawai'i Health Data Warehouse**

[profiles@hhdw.org](mailto:profiles@hhdw.org)

**(855) 946-5899 x15**

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy-  
for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

Years	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>2000-2004</b>	300	7.6	6.5 - 9.0	100	3.6	2.8 - 4.6	800	21.3	19.5 - 23.2	1,300	33.7	31.6 - 35.8	1,000	27.0	25.0 - 29.0	200	4.9	4.1 - 5.9
<b>2001-2005</b>	300	7.0	5.9 - 8.4	100	3.5	2.7 - 4.5	800	22.5	20.7 - 24.5	1,300	34.2	32.1 - 36.4	1,000	25.5	23.5 - 27.5	200	4.9	4.0 - 6.0
<b>2002-2006</b>	300	7.1	5.9 - 8.4	100	3.5	2.7 - 4.5	900	23.2	21.2 - 25.2	1,300	34.1	31.9 - 36.4	1,000	24.5	22.5 - 26.6	200	4.8	3.9 - 5.9
<b>2003-2007</b>	300	6.7	5.6 - 8.0	100	3.6	2.8 - 4.6	1,000	24.9	23.0 - 27.0	1,300	33.0	30.9 - 35.3	900	23.5	21.6 - 25.6	200	4.8	3.9 - 5.9
<b>2004-2008</b>	200	5.9	4.9 - 7.1	100	3.4	2.7 - 4.4	1,000	25.9	23.9 - 28.0	1,300	32.1	29.9 - 34.3	900	23.5	21.6 - 25.5	200	4.8	3.9 - 5.9
<b>2005-2009</b>	300	7.8	6.5 - 9.2	200	5.4	4.3 - 6.7	1,100	26.6	24.4 - 28.9	1,200	30.5	28.2 - 32.8	900	21.5	19.5 - 23.6	200	5.0	4.0 - 6.3
<b>2006-2010</b>	400	8.7	7.4 - 10.3	300	6.4	5.2 - 7.9	1,100	27.5	25.2 - 29.9	1,200	29.4	27.1 - 31.8	900	20.7	18.7 - 22.8	200	4.6	3.5 - 6.0
<b>2007-2011</b>	400	10.2	8.6 - 12.0	300	7.1	5.8 - 8.7	1,200	28.8	26.4 - 31.3	1,200	28.0	25.7 - 30.5	800	19.8	17.8 - 22.0	200	3.7	2.7 - 5.0
<b>2008-2012</b>	400	10.8	9.1 - 12.7	300	8.5	7.0 - 10.3	1,200	30.0	27.3 - 32.7	1,100	27.7	25.2 - 30.3	700	17.8	15.7 - 20.1	100	3.4	2.5 - 4.8
<b>2009-2013</b>	500	12.4	10.6 - 14.6	400	9.6	8.0 - 11.6	1,200	30.4	27.7 - 33.3	1,100	27.0	24.4 - 29.7	600	15.8	13.7 - 18.1	200	3.8	2.7 - 5.3

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy-  
for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**


*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

Years	41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
	#	%	C.I.(95%)	#
2000-2004	100	1.9	1.4 - 2.7	3,700
2001-2005	100	2.3	1.7 - 3.1	3,800
2002-2006	100	2.9	2.2 - 3.8	3,900
2003-2007	100	3.5	2.7 - 4.5	3,900
2004-2008	200	4.3	3.5 - 5.4	3,900
2005-2009	100	3.3	2.5 - 4.3	4,100
2006-2010	100	2.7	2.0 - 3.6	4,200
2007-2011	100	2.3	1.7 - 3.2	4,100
2008-2012	100	1.9	1.2 - 2.8	4,000
2009-2013	<50	1.0	0.5 - 1.8	4,000

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy-  
for the State of Hawaii, for the Year(s)**

**2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	500	12.4	10.6 - 14.6	400	9.6	8.0 - 11.6	1,200	30.4	27.7 - 33.3	1,100	27.0	24.4 - 29.7	600	15.8	13.7 - 18.1	200	3.8	2.7 - 5.3

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	100	9.1	6.9 - 12.0	100	9.4	7.1 - 12.2	200	30.4	26.4 - 34.7	200	29.0	25.1 - 33.2	100	18.4	15.2 - 22.2	<50	3.3	2.0 - 5.4
<b>Honolulu</b>	300	12.5	9.9 - 15.6	300	9.5	7.2 - 12.4	800	31.0	27.1 - 35.2	700	26.7	23.0 - 30.6	400	14.9	12.1 - 18.2	100	4.3	2.8 - 6.6
<b>Kauai</b>	<50	17.0	13.2 - 21.6	<50	9.5	6.7 - 13.3	100	27.6	22.9 - 32.9	100	27.3	22.6 - 32.6	<50	15.4	11.8 - 19.9	<50	2.7	1.4 - 5.4
<b>Maui</b>	100	15.0	12.0 - 18.5	100	10.9	8.4 - 14.0	100	28.3	24.3 - 32.6	100	25.8	22.0 - 29.9	100	17.3	14.1 - 21.0	<50	1.6	0.8 - 3.4

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.0	0.5 - 1.8	4,000

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	0.4	0.1 - 1.5	700
<50	1.1	0.5 - 2.4	2,700
<50	0.4	0.1 - 1.9	200
<50	1.2	0.5 - 2.9	500

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	200	19.9	15.1 - 25.7	100	14.3	10.6 - 19.1	200	24.1	19.0 - 30.0	200	22.3	17.4 - 28.1	100	16.4	12.2 - 21.7	<50	2.8	1.2 - 6.7
Native Hawaiian	100	7.4	5.4 - 10.1	100	7.4	5.2 - 10.5	500	32.2	27.9 - 36.8	500	30.3	26.3 - 34.7	300	16.8	13.7 - 20.5	100	4.6	2.9 - 7.1
Chinese	<50	8.9	2.6 - 26.4	<50	21.7	9.3 - 42.9	100	46.3	28.7 - 64.9	<50	17.1	8.0 - 33.1	<50	3.5	1.1 - 10.3	<50	1.8	0.5 - 6.1
Filipino	100	13.6	8.9 - 20.2	100	10.5	6.3 - 16.9	200	34.3	26.9 - 42.6	100	24.8	18.2 - 32.9	100	13.6	9.1 - 20.0	<50	3.0	1.0 - 8.7
Japanese	100	17.7	10.9 - 27.4	<50	5.5	2.7 - 10.9	100	28.3	19.7 - 39.0	100	30.4	21.4 - 41.2	100	14.9	8.7 - 24.3	<50	1.9	0.3 - 9.7
Other	100	10.7	6.6 - 16.8	<50	7.6	3.8 - 14.7	200	29.9	22.3 - 38.7	100	26.9	19.6 - 35.6	100	17.2	11.2 - 25.3	<50	5.5	2.3 - 12.4

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	5.3	3.1 - 8.9	<50	13.0	7.1 - 22.5	100	42.1	31.9 - 53.1	100	22.4	15.3 - 31.6	<50	14.4	8.6 - 23.0	<50	2.1	0.3 - 13.0
20-24	200	13.9	10.4 - 18.4	100	10.0	7.0 - 14.1	400	32.8	27.7 - 38.2	300	24.7	20.3 - 29.6	200	14.2	10.7 - 18.6	<50	3.9	2.1 - 6.9
25-34	200	11.8	9.4 - 14.7	200	8.8	6.8 - 11.5	600	29.6	25.7 - 33.7	600	28.8	25.0 - 32.9	300	15.8	13.0 - 19.1	100	3.8	2.4 - 6.0
35+	100	15.8	10.2 - 23.6	<50	10.0	5.7 - 16.9	100	19.8	14.0 - 27.2	100	28.4	21.0 - 37.2	100	21.2	14.6 - 29.6	<50	4.3	1.6 - 11.3

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	0.3	0.1 - 0.9	900
<50	1.2	0.5 - 2.9	1,600
<50	0.6	0.1 - 3.3	100
<50	0.2	0.0 - 1.0	600
<50	1.4	0.2 - 8.8	400
<50	2.3	0.7 - 7.5	500

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	0.7	0.2 - 2.5	300
<50	0.6	0.1 - 2.2	1,300
<50	1.4	0.6 - 3.0	2,000
<50	0.6	0.2 - 1.6	400



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	300	12.9	10.3 - 16.1	200	11.1	8.6 - 14.3	600	28.7	24.9 - 32.8	500	25.0	21.5 - 28.9	300	17.1	14.1 - 20.5	100	4.0	2.4 - 6.5
<b>Female</b>	300	12.0	9.5 - 15.0	200	8.2	6.2 - 10.8	700	32.0	28.2 - 36.2	600	28.8	25.1 - 32.8	300	14.6	11.8 - 17.8	100	3.5	2.2 - 5.6

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	300	15.9	12.7 - 19.7	200	11.3	8.6 - 14.7	500	30.0	25.7 - 34.7	500	27.6	23.4 - 32.1	200	13.7	10.8 - 17.2	<50	1.0	0.4 - 2.5
<b>No</b>	200	9.9	7.8 - 12.5	200	8.4	6.4 - 10.9	700	30.7	27.2 - 34.5	600	26.5	23.3 - 30.1	400	17.3	14.5 - 20.5	100	5.8	4.0 - 8.2

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	200	8.7	6.6 - 11.5	100	6.6	4.9 - 9.0	600	34.0	30.0 - 38.2	500	26.5	23.0 - 30.3	300	16.9	14.0 - 20.2	100	6.0	4.0 - 8.8
<b>131-185%</b>	100	13.4	8.7 - 20.0	100	9.0	5.0 - 15.4	200	29.1	22.5 - 36.7	200	28.7	22.0 - 36.3	100	17.9	12.5 - 24.9	<50	1.2	0.3 - 5.3
<b>186+%</b>	200	17.6	13.8 - 22.2	200	14.0	10.5 - 18.5	300	24.8	20.1 - 30.1	300	27.6	22.6 - 33.2	200	13.5	10.0 - 17.9	<50	1.7	0.7 - 4.1

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.1	0.5 - 2.7	1,900
<50	0.8	0.3 - 2.0	2,100

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	0.5	0.2 - 1.7	1,700
<50	1.3	0.7 - 2.7	2,300

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.3	0.6 - 2.9	1,900
<50	0.8	0.1 - 5.1	700
<50	0.7	0.2 - 2.4	1,200

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	12.8	10.0 - 16.3	<50	6.5	4.6 - 9.1	100	27.5	23.4 - 32.1	100	29.1	24.9 - 33.6	100	19.5	15.9 - 23.8	<50	3.5	2.2 - 5.6
Normal birth weight	500	12.4	10.4 - 14.8	400	9.9	8.1 - 12.1	1,100	30.8	27.8 - 33.9	1,000	26.7	23.9 - 29.7	600	15.4	13.2 - 17.9	100	3.8	2.6 - 5.4

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.6	0.1 - 3.2	<50	2.5	1.3 - 4.7	200	20.6	16.4 - 25.6	400	34.8	29.7 - 40.3	400	31.4	26.5 - 36.8	100	7.9	5.2 - 11.8
No	500	17.3	14.8 - 20.2	400	12.6	10.3 - 15.2	1,000	34.4	31.0 - 38.0	700	23.7	20.8 - 27.0	300	9.4	7.5 - 11.7	100	2.1	1.2 - 3.6

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	0	0.0	-	400	11.0	9.1 - 13.2	1,200	34.7	31.7 - 37.9	1,100	30.8	27.9 - 33.9	600	18.0	15.7 - 20.6	200	4.3	3.1 - 6.0
No	500	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.1	0.5 - 2.5	300
<50	1.0	0.5 - 1.9	3,700

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.1	1.0 - 4.4	1,200
<50	0.5	0.2 - 1.5	2,800

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.1	0.6 - 2.1	3,500
0	0.0	-	500

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009-2013**

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	10.9	8.5 - 13.8	200	7.2	5.4 - 9.5	700	31.8	28.2 - 35.6	600	26.4	23.1 - 30.0	400	17.9	15.1 - 21.1	100	4.7	3.2 - 7.0
<b>No</b>	300	14.5	11.6 - 17.9	200	12.7	9.9 - 16.3	500	28.5	24.3 - 33.1	500	27.8	23.7 - 32.3	200	13.3	10.4 - 16.9	<50	2.6	1.4 - 4.9

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.1	0.5 - 2.3	2,200
<50	0.5	0.1 - 1.8	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	400	10.8	9.1 - 12.7	300	8.5	7.0 - 10.3	1,200	30.0	27.3 - 32.7	1,100	27.7	25.2 - 30.3	700	17.8	15.7 - 20.1	100	3.4	2.5 - 4.8

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	100	9.3	6.8 - 12.5	100	8.5	6.3 - 11.3	200	29.7	25.5 - 34.2	200	30.8	26.6 - 35.4	100	18.4	15.0 - 22.5	<50	3.1	1.9 - 5.2
<b>Honolulu</b>	300	10.2	8.0 - 13.0	200	8.4	6.3 - 11.0	800	30.6	27.0 - 34.5	700	27.0	23.6 - 30.7	500	17.2	14.5 - 20.4	100	4.1	2.7 - 6.1
<b>Kauai</b>	<50	17.0	12.5 - 22.5	<50	8.3	5.6 - 12.1	100	28.5	22.8 - 35.1	<50	24.3	19.3 - 30.2	<50	19.0	13.9 - 25.5	<50	2.8	1.4 - 5.6
<b>Maui</b>	100	13.6	10.8 - 17.0	<50	9.2	6.9 - 12.0	100	27.3	23.1 - 32.0	100	28.3	24.0 - 33.1	100	19.6	15.8 - 23.9	<50	0.7	0.2 - 1.9

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	1.9	1.2 - 2.8	4,000

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	0.2	0.0 - 1.4	700
100	2.5	1.6 - 3.9	2,700
0	0.0	-	200
<50	1.3	0.5 - 3.5	500



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	100	15.7	11.8 - 20.8	100	13.6	10.1 - 18.0	200	26.5	21.4 - 32.2	200	19.9	15.5 - 25.1	200	20.3	15.9 - 25.5	<50	2.4	1.0 - 5.5
Native Hawaiian	100	6.7	5.0 - 9.0	100	6.8	4.6 - 10.0	500	30.4	26.4 - 34.7	500	32.2	28.2 - 36.4	300	18.4	15.1 - 22.1	100	4.0	2.5 - 6.3
Chinese	<50	10.8	3.5 - 28.8	<50	16.2	6.0 - 37.3	<50	45.4	27.9 - 64.1	<50	20.1	9.8 - 36.9	<50	4.8	2.0 - 11.1	<50	2.7	0.9 - 7.4
Filipino	100	11.7	7.4 - 17.9	<50	8.7	5.0 - 14.8	200	35.1	27.9 - 43.0	100	26.1	19.7 - 33.6	100	15.8	11.3 - 21.6	<50	2.5	0.9 - 6.5
Japanese	<50	12.7	7.3 - 21.3	<50	5.5	2.6 - 11.1	100	28.6	20.2 - 38.8	100	35.8	26.6 - 46.1	<50	13.4	8.0 - 21.6	<50	1.9	0.3 - 10.0
Other	100	12.1	7.8 - 18.3	<50	4.9	2.5 - 9.4	100	27.4	20.7 - 35.2	100	25.4	18.8 - 33.4	100	19.5	13.7 - 27.0	<50	5.8	2.5 - 12.6

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	5.0	2.8 - 8.6	<50	10.6	5.6 - 19.1	100	42.1	32.9 - 51.9	100	20.6	14.5 - 28.4	100	16.0	10.3 - 23.9	<50	3.8	1.3 - 10.5
20-24	100	11.6	8.6 - 15.6	100	7.8	5.3 - 11.4	400	32.9	28.0 - 38.1	300	26.7	22.4 - 31.5	200	17.2	13.6 - 21.5	<50	2.5	1.3 - 5.0
25-34	200	10.6	8.3 - 13.3	200	8.8	6.7 - 11.4	500	27.4	23.9 - 31.3	600	30.0	26.3 - 33.9	300	17.2	14.4 - 20.4	100	3.7	2.3 - 5.7
35+	100	13.9	8.9 - 21.1	<50	7.3	4.0 - 13.0	100	23.6	17.4 - 31.1	100	25.5	18.9 - 33.4	100	23.8	17.3 - 31.8	<50	4.9	1.9 - 11.8

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.7	0.8 - 3.4	900
<50	1.6	0.8 - 3.4	1,600
0	0.0	-	100
<50	0.2	0.0 - 1.0	500
<50	2.1	0.5 - 8.4	400
<50	5.0	2.5 - 9.7	500

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.0	0.8 - 5.5	300
<50	1.3	0.6 - 3.0	1,200
<50	2.4	1.4 - 4.1	2,000
<50	1.0	0.4 - 2.8	400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	200	10.9	8.7 - 13.7	200	9.3	7.1 - 12.1	600	29.0	25.4 - 32.9	500	27.0	23.6 - 30.8	400	18.5	15.7 - 21.7	100	3.6	2.3 - 5.7
<b>Female</b>	200	10.6	8.4 - 13.4	200	7.7	5.8 - 10.1	600	30.9	27.2 - 34.9	600	28.3	24.8 - 32.1	300	17.1	14.2 - 20.4	100	3.3	2.0 - 5.3

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	14.6	11.7 - 18.0	200	10.0	7.6 - 13.1	500	28.6	24.7 - 32.9	500	27.4	23.5 - 31.6	300	16.4	13.5 - 19.8	<50	1.5	0.7 - 3.2
<b>No</b>	200	8.0	6.2 - 10.2	200	7.3	5.5 - 9.7	700	31.0	27.5 - 34.6	600	27.9	24.7 - 31.3	400	18.9	16.1 - 22.0	100	4.8	3.3 - 7.0

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	200	8.3	6.3 - 10.9	100	5.6	4.0 - 7.8	600	31.3	27.6 - 35.2	500	27.9	24.5 - 31.6	400	19.3	16.4 - 22.6	100	5.6	3.8 - 8.2
<b>131-185%</b>	100	11.7	7.6 - 17.7	<50	6.8	3.6 - 12.4	200	29.7	23.3 - 36.9	200	29.5	23.3 - 36.7	100	18.6	13.4 - 25.2	<50	1.6	0.4 - 5.2
<b>186+%</b>	200	14.8	11.5 - 18.8	100	12.2	9.0 - 16.3	300	27.7	23.1 - 33.0	300	26.6	21.9 - 31.9	200	15.7	12.2 - 19.9	<50	1.3	0.6 - 2.9

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.6	0.8 - 3.1	2,000
<50	2.1	1.3 - 3.6	2,000

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.5	0.8 - 2.8	1,700
<50	2.1	1.3 - 3.6	2,300

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.0	1.1 - 3.6	1,900
<50	2.1	0.8 - 5.6	700
<50	1.7	0.8 - 3.4	1,200

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	12.7	9.5 - 16.6	<50	6.4	4.2 - 9.4	100	26.9	22.2 - 32.2	100	25.9	21.3 - 31.0	100	22.8	18.2 - 28.3	<50	3.8	2.1 - 6.8
Normal birth weight	400	10.6	8.9 - 12.7	300	8.7	7.1 - 10.7	1,100	30.3	27.5 - 33.2	1,000	27.7	25.0 - 30.5	600	17.4	15.2 - 19.8	100	3.4	2.4 - 4.9

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.6	0.1 - 2.9	<50	2.1	1.0 - 4.2	300	19.8	16.1 - 24.2	400	34.7	30.0 - 39.6	400	32.5	28.0 - 37.3	100	7.4	4.9 - 10.9
No	400	15.7	13.3 - 18.3	300	11.5	9.4 - 14.1	900	34.8	31.5 - 38.3	700	24.3	21.4 - 27.4	300	10.8	8.8 - 13.2	<50	1.6	0.9 - 2.8

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	0	0.0	-	300	9.5	7.8 - 11.5	1,200	33.6	30.7 - 36.6	1,100	31.0	28.3 - 33.9	700	20.0	17.7 - 22.5	100	3.9	2.8 - 5.4
No	400	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.5	0.6 - 3.9	300
100	1.9	1.2 - 2.9	3,700

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	3.0	1.7 - 5.2	1,300
<50	1.3	0.8 - 2.4	2,700

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.1	1.4 - 3.1	3,600
0	0.0	-	400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008-2012**

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	9.7	7.6 - 12.2	100	6.7	5.0 - 8.9	700	30.1	26.8 - 33.7	600	27.1	24.0 - 30.5	500	20.2	17.3 - 23.4	100	3.9	2.5 - 5.8
<b>No</b>	200	12.3	9.8 - 15.4	200	11.0	8.4 - 14.2	500	30.0	25.9 - 34.4	500	28.4	24.4 - 32.7	300	14.7	11.9 - 18.0	100	3.0	1.7 - 5.1

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.4	1.5 - 3.7	2,200
<50	0.8	0.3 - 2.1	1,700



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	400	10.2	8.6 - 12.0	300	7.1	5.8 - 8.7	1,200	28.8	26.4 - 31.3	1,200	28.0	25.7 - 30.5	800	19.8	17.8 - 22.0	200	3.7	2.7 - 5.0

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	100	7.8	5.6 - 10.9	100	7.5	5.4 - 10.3	200	27.6	23.4 - 32.2	200	32.5	28.1 - 37.3	200	22.0	18.1 - 26.5	<50	2.3	1.2 - 4.3
<b>Honolulu</b>	300	10.2	8.1 - 12.7	200	7.0	5.2 - 9.3	800	29.4	26.1 - 32.9	700	27.0	23.8 - 30.4	500	18.9	16.2 - 21.9	100	4.4	3.1 - 6.3
<b>Kauai</b>	<50	15.2	10.7 - 21.1	<50	5.5	3.4 - 8.8	100	26.5	20.2 - 33.9	<50	24.8	18.8 - 32.0	<50	23.5	16.8 - 31.8	<50	3.2	1.3 - 7.8
<b>Maui</b>	100	11.5	8.8 - 15.0	<50	7.9	5.7 - 10.9	100	27.7	23.2 - 32.8	100	29.2	24.6 - 34.4	100	20.9	16.8 - 25.7	<50	1.8	0.7 - 4.7

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.3	1.7 - 3.2	4,100

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	0.2	0.0 - 1.3	700
100	3.1	2.2 - 4.5	2,800
<50	1.3	0.2 - 8.2	200
<50	0.9	0.3 - 2.9	500

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	100	14.2	10.6 - 18.8	100	11.2	8.2 - 15.0	300	26.4	21.7 - 31.7	200	18.6	14.7 - 23.3	200	23.7	19.4 - 28.6	<50	3.5	1.9 - 6.5
Native Hawaiian	100	5.7	4.1 - 7.8	100	6.1	4.2 - 8.9	500	29.5	25.7 - 33.7	500	32.1	28.3 - 36.1	300	20.5	17.2 - 24.2	100	4.3	2.8 - 6.5
Chinese	<50	13.6	4.9 - 32.9	<50	16.9	5.7 - 40.6	<50	35.2	19.2 - 55.5	<50	24.4	12.4 - 42.6	<50	6.7	3.0 - 14.3	<50	3.1	1.1 - 8.6
Filipino	100	12.4	8.1 - 18.5	<50	6.2	3.3 - 11.4	200	35.0	28.1 - 42.5	200	29.9	23.5 - 37.1	100	14.1	10.2 - 19.1	<50	2.0	0.6 - 6.1
Japanese	<50	11.8	7.1 - 19.0	<50	5.3	2.6 - 10.4	100	25.2	18.1 - 34.0	200	37.5	29.0 - 46.9	100	16.6	11.0 - 24.3	<50	1.7	0.3 - 9.0
Other	100	11.7	7.5 - 17.6	<50	3.6	1.7 - 7.4	100	26.0	19.9 - 33.3	100	25.2	19.1 - 32.5	100	21.8	16.1 - 28.9	<50	5.9	2.7 - 12.3

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	3.6	2.0 - 6.6	<50	8.4	4.2 - 16.1	200	40.6	32.3 - 49.4	100	22.9	16.7 - 30.5	100	18.1	12.4 - 25.5	<50	4.9	2.1 - 11.0
20-24	200	10.9	8.1 - 14.4	100	6.7	4.5 - 9.7	400	30.7	26.4 - 35.4	400	26.3	22.4 - 30.7	300	20.1	16.6 - 24.2	<50	2.7	1.5 - 5.0
25-34	200	10.2	8.0 - 12.9	100	7.2	5.4 - 9.5	500	26.7	23.3 - 30.4	600	30.7	27.1 - 34.5	400	18.8	16.0 - 21.9	100	4.0	2.7 - 6.1
35+	100	13.5	8.9 - 20.1	<50	7.3	4.0 - 12.9	100	21.6	15.9 - 28.7	100	26.3	19.9 - 33.9	100	24.8	18.4 - 32.7	<50	4.4	1.6 - 11.2

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.4	1.3 - 4.4	1,000
<50	1.9	1.0 - 3.4	1,600
0	0.0	-	100
<50	0.5	0.1 - 2.0	600
<50	1.9	0.4 - 7.5	400
<50	5.9	3.2 - 10.6	600

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.6	0.5 - 4.7	400
<50	2.5	1.5 - 4.3	1,400
<50	2.4	1.4 - 4.0	1,900
<50	2.0	0.8 - 4.8	500

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	200	10.6	8.4 - 13.2	200	7.2	5.4 - 9.7	600	28.6	25.3 - 32.2	600	28.0	24.8 - 31.6	400	19.7	16.9 - 22.7	100	4.0	2.7 - 6.0
<b>Female</b>	200	9.8	7.7 - 12.4	100	7.0	5.3 - 9.2	600	29.0	25.5 - 32.7	600	28.1	24.7 - 31.6	400	20.0	17.1 - 23.3	100	3.4	2.2 - 5.3

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	13.9	11.2 - 17.0	200	9.2	7.0 - 12.0	500	28.3	24.6 - 32.2	500	27.3	23.7 - 31.2	300	17.6	14.7 - 20.8	<50	2.0	1.1 - 3.6
<b>No</b>	200	7.4	5.7 - 9.5	100	5.5	4.0 - 7.6	700	29.2	26.0 - 32.6	700	28.6	25.5 - 31.9	500	21.6	18.8 - 24.6	100	5.0	3.5 - 7.1

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	100	7.4	5.6 - 9.8	100	4.8	3.4 - 6.8	600	28.6	25.2 - 32.3	600	29.1	25.8 - 32.6	400	21.6	18.7 - 24.9	100	5.7	3.9 - 8.2
<b>131-185%</b>	100	12.8	8.6 - 18.7	<50	5.3	2.6 - 10.4	200	30.6	24.4 - 37.6	200	27.9	22.0 - 34.5	100	19.6	14.5 - 25.9	<50	1.8	0.6 - 5.2
<b>186+%</b>	200	13.7	10.7 - 17.4	100	9.6	7.1 - 13.0	300	28.3	24.0 - 33.1	300	26.1	21.8 - 30.9	200	18.2	14.8 - 22.3	<50	2.1	1.1 - 3.7

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.9	1.1 - 3.2	2,100
100	2.7	1.8 - 4.3	2,000

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.8	1.0 - 3.1	1,800
100	2.7	1.8 - 4.1	2,400

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.7	1.7 - 4.2	2,000
<50	2.1	0.8 - 5.4	700
<50	1.9	1.0 - 3.6	1,200

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Income (2004-2011)	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<\$10,000	100	7.3	4.8 - 11.0	<50	4.0	2.2 - 7.0	300	26.8	22.2 - 31.9	300	31.9	27.1 - 37.2	200	21.9	17.8 - 26.7	100	5.0	3.0 - 8.4
\$10,000-\$24,999	100	9.2	6.5 - 12.9	<50	4.7	2.9 - 7.3	300	28.0	23.6 - 33.0	300	26.5	22.4 - 31.0	300	24.6	20.3 - 29.5	100	5.7	3.4 - 9.5
\$25,000-\$49,999	100	10.8	7.7 - 14.9	100	7.1	4.7 - 10.8	400	34.7	29.5 - 40.3	300	26.1	21.5 - 31.4	200	16.0	12.6 - 20.0	<50	1.9	0.8 - 4.0
>=\$50,000	100	15.5	11.7 - 20.3	100	11.0	7.7 - 15.6	200	24.6	19.6 - 30.5	200	27.9	22.5 - 34.0	100	17.1	12.9 - 22.3	<50	2.6	1.3 - 5.1

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	10.8	7.9 - 14.5	<50	5.8	3.6 - 9.1	100	23.8	18.9 - 29.5	100	27.1	22.0 - 32.9	100	26.9	21.6 - 33.1	<50	4.8	2.7 - 8.5
Normal birth weight	400	10.1	8.5 - 12.1	300	7.3	5.8 - 9.0	1,100	29.3	26.7 - 32.0	1,100	28.0	25.5 - 30.7	700	19.2	17.1 - 21.6	100	3.6	2.6 - 5.0

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.7	0.2 - 2.5	<50	1.3	0.6 - 2.7	300	18.7	15.4 - 22.6	500	36.1	31.8 - 40.6	500	33.5	29.4 - 38.0	100	7.0	4.8 - 10.1
No	400	15.3	13.0 - 17.9	300	10.2	8.3 - 12.6	900	34.2	31.0 - 37.5	600	23.6	20.9 - 26.6	300	12.5	10.5 - 14.8	100	2.0	1.2 - 3.2

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	3.1	1.6 - 5.8	1,000
<50	1.3	0.6 - 2.9	1,100
<50	3.5	2.0 - 5.9	1,000
<50	1.2	0.5 - 3.0	800

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	0.9	0.2 - 3.5	400
100	2.4	1.7 - 3.4	3,800

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.6	1.4 - 4.6	1,400
100	2.2	1.5 - 3.2	2,700



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007-2011**

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	0	0.0	-	300	7.9	6.5 - 9.7	1,200	32.0	29.4 - 34.8	1,200	31.2	28.6 - 33.9	800	22.1	19.8 - 24.5	200	4.1	3.1 - 5.6
<b>No</b>	400	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	9.2	7.2 - 11.6	100	6.1	4.5 - 8.1	700	28.8	25.7 - 32.1	600	27.4	24.4 - 30.6	500	21.6	18.8 - 24.7	100	4.2	2.8 - 6.1
<b>No</b>	200	11.6	9.2 - 14.5	200	8.5	6.4 - 11.3	500	28.9	25.1 - 33.0	500	28.8	25.1 - 32.9	300	17.3	14.5 - 20.5	100	3.3	2.0 - 5.3

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.6	1.8 - 3.6	3,700
0	0.0	-	400

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.8	1.9 - 4.2	2,300
<50	1.6	0.9 - 3.0	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	400	8.7	7.4 - 10.3	300	6.4	5.2 - 7.9	1,100	27.5	25.2 - 29.9	1,200	29.4	27.1 - 31.8	900	20.7	18.7 - 22.8	200	4.6	3.5 - 6.0

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	100	7.4	5.2 - 10.5	100	7.3	5.1 - 10.3	200	24.5	20.3 - 29.3	200	34.0	29.2 - 39.1	200	23.8	19.5 - 28.6	<50	2.0	1.0 - 4.1
<b>Honolulu</b>	200	8.6	6.9 - 10.8	200	6.2	4.6 - 8.3	800	28.4	25.3 - 31.7	800	28.4	25.5 - 31.6	500	19.4	16.9 - 22.2	200	5.5	4.1 - 7.5
<b>Kauai</b>	<50	11.3	7.2 - 17.3	<50	4.4	2.6 - 7.4	100	29.4	22.2 - 37.9	100	26.5	19.4 - 35.1	<50	22.8	15.9 - 31.5	<50	3.0	1.2 - 7.4
<b>Maui</b>	<50	9.9	7.2 - 13.4	<50	6.9	4.7 - 9.9	100	26.0	21.1 - 31.7	100	29.2	24.1 - 34.8	100	23.1	18.4 - 28.6	<50	3.5	1.7 - 7.2

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.7	2.0 - 3.6	4,200

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	0.9	0.2 - 3.4	700
100	3.4	2.4 - 4.7	2,800
<50	2.5	0.7 - 9.1	200
<50	1.4	0.4 - 4.3	500

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	100	12.3	9.2 - 16.3	100	11.1	8.2 - 14.9	200	23.7	19.5 - 28.5	200	21.6	17.6 - 26.1	200	24.5	20.4 - 29.1	<50	4.4	2.6 - 7.4
Native Hawaiian	100	4.8	3.5 - 6.6	100	5.1	3.4 - 7.6	400	27.2	23.7 - 31.2	500	33.5	29.7 - 37.6	300	21.7	18.4 - 25.4	100	5.2	3.6 - 7.6
Chinese	<50	15.9	6.0 - 35.8	<50	17.4	5.6 - 43.0	<50	36.7	19.7 - 57.9	<50	20.3	9.9 - 37.1	<50	7.5	3.6 - 14.9	<50	2.2	0.6 - 7.4
Filipino	100	10.2	6.5 - 15.8	<50	4.9	2.5 - 9.3	200	37.3	30.5 - 44.8	200	28.4	22.5 - 35.1	100	15.6	11.4 - 21.0	<50	2.9	1.1 - 7.0
Japanese	<50	10.7	6.6 - 17.0	<50	4.3	2.1 - 8.6	100	23.2	16.6 - 31.3	100	37.8	29.7 - 46.7	100	18.9	13.3 - 26.3	<50	1.7	0.3 - 9.1
Other	100	9.3	5.9 - 14.4	<50	3.1	1.6 - 5.9	200	27.5	21.5 - 34.3	200	27.7	21.8 - 34.5	100	19.3	14.4 - 25.4	<50	7.0	3.7 - 13.1

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	3.0	1.6 - 5.6	<50	6.9	3.3 - 14.0	200	39.5	31.6 - 47.9	100	25.6	19.2 - 33.2	100	17.9	12.6 - 24.8	<50	5.1	2.3 - 10.6
20-24	100	8.3	6.2 - 10.9	100	6.5	4.5 - 9.4	400	29.2	25.1 - 33.7	400	27.8	23.9 - 32.0	300	21.2	17.8 - 25.1	100	3.9	2.4 - 6.4
25-34	200	9.0	7.0 - 11.5	100	6.4	4.8 - 8.5	500	24.7	21.5 - 28.1	600	32.6	29.1 - 36.2	400	19.7	17.0 - 22.8	100	5.0	3.5 - 7.1
35+	100	14.2	9.5 - 20.9	<50	5.4	2.8 - 10.2	100	23.2	17.6 - 29.8	100	24.1	18.3 - 31.1	100	25.9	19.6 - 33.5	<50	4.5	1.8 - 11.2

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.4	1.3 - 4.3	1,000
<50	2.3	1.4 - 3.9	1,600
0	0.0	-	100
<50	0.7	0.2 - 2.7	500
<50	3.2	1.2 - 8.6	400
<50	6.0	3.4 - 10.4	600

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.1	0.8 - 5.5	400
<50	3.0	1.9 - 4.9	1,300
<50	2.6	1.6 - 4.2	1,900
<50	2.7	1.2 - 5.8	500

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	200	9.6	7.7 - 12.0	100	6.8	5.0 - 9.0	600	27.0	23.9 - 30.3	600	29.1	26.0 - 32.5	400	20.2	17.6 - 23.1	100	5.0	3.5 - 7.1
<b>Female</b>	200	7.7	6.0 - 9.9	100	6.0	4.5 - 8.0	600	28.1	24.8 - 31.7	600	29.6	26.3 - 33.1	400	21.2	18.3 - 24.4	100	4.1	2.8 - 6.1

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	12.4	10.1 - 15.3	200	8.7	6.7 - 11.2	500	27.4	24.0 - 31.1	500	27.9	24.5 - 31.5	300	18.7	15.9 - 21.8	<50	2.7	1.7 - 4.3
<b>No</b>	100	5.8	4.5 - 7.5	100	4.6	3.3 - 6.5	600	27.6	24.6 - 30.9	700	30.5	27.5 - 33.8	500	22.3	19.6 - 25.3	100	6.1	4.4 - 8.2

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	100	5.6	4.3 - 7.4	100	3.8	2.6 - 5.4	600	27.4	24.1 - 30.9	600	30.8	27.5 - 34.2	500	22.6	19.7 - 25.8	100	6.4	4.6 - 8.9
<b>131-185%</b>	100	9.9	6.3 - 15.1	<50	4.3	2.1 - 8.6	200	30.0	24.1 - 36.7	200	31.3	25.4 - 37.9	100	19.7	15.0 - 25.4	<50	2.7	1.1 - 6.3
<b>186+%</b>	200	13.3	10.4 - 17.0	100	9.2	6.8 - 12.5	300	26.5	22.6 - 30.9	300	25.7	21.7 - 30.2	200	20.1	16.7 - 24.1	<50	3.5	2.2 - 5.4

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.2	1.4 - 3.5	2,200
100	3.2	2.1 - 4.8	2,000

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.2	1.4 - 3.6	1,800
100	3.0	2.1 - 4.4	2,300

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.5	2.4 - 5.1	2,000
<50	2.2	0.8 - 5.7	600
<50	1.5	0.8 - 2.9	1,200



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Income (2004-2011)	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<\$10,000	100	6.0	4.0 - 8.9	<50	3.0	1.5 - 5.7	300	25.6	21.1 - 30.5	400	33.6	28.9 - 38.8	200	22.3	18.3 - 26.9	100	5.0	3.0 - 8.2
\$10,000-\$24,999	100	6.3	4.4 - 8.7	<50	3.7	2.5 - 5.7	300	28.2	23.9 - 33.1	300	28.5	24.4 - 33.0	300	24.8	20.7 - 29.4	100	7.2	4.7 - 11.0
\$25,000-\$49,999	100	10.4	7.4 - 14.4	100	6.1	3.9 - 9.4	300	30.8	26.0 - 36.1	300	28.3	23.8 - 33.3	200	18.2	14.8 - 22.3	<50	3.0	1.6 - 5.4
>=\$50,000	100	14.0	10.4 - 18.5	100	10.7	7.4 - 15.2	200	24.4	19.8 - 29.8	200	26.8	21.8 - 32.4	100	18.9	14.7 - 23.8	<50	3.8	2.2 - 6.6

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	8.7	6.0 - 12.4	<50	4.2	2.4 - 7.4	100	24.8	19.3 - 31.3	100	28.7	22.8 - 35.4	100	29.0	22.9 - 35.9	<50	3.7	1.8 - 7.5
Normal birth weight	300	8.7	7.3 - 10.4	300	6.6	5.3 - 8.2	1,100	27.8	25.4 - 30.4	1,100	29.3	26.9 - 31.9	800	20.0	17.9 - 22.2	200	4.7	3.6 - 6.1

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.6	0.2 - 2.2	<50	1.4	0.7 - 2.7	300	16.6	13.6 - 20.2	600	36.6	32.7 - 40.8	500	32.7	28.8 - 36.7	100	8.4	6.2 - 11.4
No	400	13.8	11.7 - 16.2	200	9.6	7.7 - 11.8	900	34.4	31.3 - 37.6	600	24.7	22.1 - 27.6	300	13.2	11.2 - 15.4	100	2.2	1.4 - 3.5

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	4.6	2.8 - 7.4	1,000
<50	1.3	0.6 - 2.8	1,100
<50	3.1	1.8 - 5.4	1,000
<50	1.4	0.6 - 3.3	800

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	0.9	0.2 - 3.5	300
100	2.9	2.1 - 3.9	3,800

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.6	2.3 - 5.6	1,600
100	2.1	1.4 - 3.1	2,500

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006-2010**

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	0	0.0	-	300	7.0	5.7 - 8.6	1,100	30.2	27.7 - 32.7	1,200	32.2	29.7 - 34.7	900	22.7	20.5 - 25.0	200	5.0	3.9 - 6.5
<b>No</b>	400	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	7.0	5.4 - 8.9	100	5.3	3.9 - 7.2	600	27.8	24.7 - 31.0	700	30.1	27.1 - 33.3	500	21.6	18.9 - 24.6	100	5.2	3.7 - 7.1
<b>No</b>	200	11.1	8.8 - 13.8	100	7.9	5.9 - 10.4	500	27.4	23.9 - 31.1	500	28.5	25.0 - 32.3	300	19.1	16.3 - 22.2	100	4.0	2.6 - 6.1

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.9	2.2 - 4.0	3,800
0	0.0	-	400

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.1	2.2 - 4.5	2,300
<50	2.1	1.2 - 3.6	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	300	7.8	6.5 - 9.2	200	5.4	4.3 - 6.7	1,100	26.6	24.4 - 28.9	1,200	30.5	28.2 - 32.8	900	21.5	19.5 - 23.6	200	5.0	4.0 - 6.3

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	<50	6.3	4.2 - 9.5	<50	5.8	3.7 - 9.0	200	23.7	19.3 - 28.9	200	34.6	29.5 - 40.2	200	24.9	20.3 - 30.1	<50	2.6	1.4 - 5.0
<b>Honolulu</b>	200	7.9	6.3 - 9.9	200	5.5	4.1 - 7.3	700	26.9	24.1 - 29.8	800	29.8	27.1 - 32.8	600	20.4	18.1 - 23.0	200	5.7	4.3 - 7.4
<b>Kauai</b>	<50	9.5	5.3 - 16.4	<50	3.5	1.7 - 7.3	100	28.6	21.0 - 37.7	100	28.2	20.2 - 37.9	<50	22.9	15.7 - 32.2	<50	3.7	1.5 - 8.8
<b>Maui</b>	<50	8.4	5.7 - 12.1	<50	4.9	3.0 - 7.8	100	28.2	22.7 - 34.5	100	29.2	23.7 - 35.4	100	22.3	17.3 - 28.3	<50	5.3	2.9 - 9.5

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.3	2.5 - 4.3	4,100

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.0	0.8 - 4.6	700
100	3.8	2.8 - 5.1	2,700
<50	3.6	1.2 - 10.2	200
<50	1.6	0.5 - 4.8	500

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	100	11.0	8.2 - 14.6	100	9.2	6.7 - 12.5	200	23.8	19.9 - 28.3	200	22.7	18.8 - 27.1	200	25.4	21.4 - 29.9	<50	4.9	3.2 - 7.4
Native Hawaiian	100	4.5	3.1 - 6.4	100	3.7	2.4 - 5.7	400	26.3	22.8 - 30.1	500	35.3	31.4 - 39.3	300	21.8	18.6 - 25.4	100	5.4	3.8 - 7.6
Chinese	<50	20.1	7.8 - 42.9	<50	22.1	7.4 - 50.3	<50	30.0	14.6 - 51.7	<50	10.9	5.4 - 20.8	<50	11.8	6.0 - 21.8	<50	3.9	1.4 - 10.7
Filipino	<50	8.3	5.1 - 13.1	<50	4.9	2.4 - 9.8	200	33.3	27.3 - 40.0	100	30.0	24.4 - 36.3	100	18.7	14.2 - 24.3	<50	3.6	1.5 - 8.0
Japanese	<50	10.7	6.7 - 16.8	<50	3.8	1.8 - 8.0	100	23.2	17.1 - 30.8	100	34.7	27.6 - 42.5	100	21.0	15.6 - 27.6	<50	3.5	1.3 - 8.9
Other	<50	7.4	4.5 - 11.9	<50	3.0	1.5 - 5.9	200	28.5	22.7 - 35.2	200	30.2	24.5 - 36.6	100	17.6	13.2 - 23.0	<50	6.9	4.0 - 11.8

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	2.6	1.3 - 5.1	<50	5.0	2.5 - 9.9	200	34.0	26.8 - 41.9	100	31.4	24.7 - 38.9	100	17.9	12.7 - 24.7	<50	6.4	3.4 - 11.7
20-24	100	7.8	5.8 - 10.5	100	5.6	3.7 - 8.2	400	28.3	24.4 - 32.4	400	27.9	24.2 - 31.9	300	21.6	18.3 - 25.3	100	4.7	3.1 - 7.1
25-34	100	7.9	6.1 - 10.2	100	6.0	4.4 - 8.0	500	24.1	21.1 - 27.3	600	33.5	30.1 - 37.0	400	21.2	18.5 - 24.2	100	4.9	3.5 - 6.8
35+	100	12.6	8.3 - 18.7	<50	2.6	1.2 - 5.5	100	24.9	19.3 - 31.5	100	24.2	18.5 - 31.0	100	26.3	20.2 - 33.5	<50	5.2	2.5 - 10.3

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	3.1	1.8 - 5.1	1,000
<50	3.1	2.0 - 4.8	1,600
<50	1.2	0.2 - 6.2	100
<50	1.2	0.4 - 3.6	500
<50	3.1	1.1 - 8.2	400
<50	6.3	3.8 - 10.2	600

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.8	1.2 - 6.3	500
100	4.2	2.8 - 6.3	1,300
<50	2.5	1.6 - 4.0	1,900
<50	4.2	2.2 - 8.1	400



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	200	8.6	6.8 - 10.9	100	4.8	3.5 - 6.7	600	26.8	23.8 - 30.0	600	29.0	26.0 - 32.2	500	21.9	19.2 - 24.8	100	5.8	4.3 - 7.7
<b>Female</b>	100	6.9	5.3 - 8.8	100	5.9	4.4 - 7.9	500	26.4	23.3 - 29.7	600	32.1	28.8 - 35.5	400	21.0	18.3 - 24.1	100	4.2	3.0 - 6.0

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	11.9	9.6 - 14.6	100	7.8	5.9 - 10.2	500	27.7	24.4 - 31.2	500	27.2	24.1 - 30.6	300	18.9	16.3 - 21.9	100	3.3	2.3 - 4.8
<b>No</b>	100	4.7	3.5 - 6.3	100	3.5	2.5 - 5.1	600	25.8	23.0 - 28.8	800	32.9	29.8 - 36.0	500	23.4	20.7 - 26.3	100	6.3	4.8 - 8.3

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	100	4.9	3.6 - 6.5	100	3.2	2.2 - 4.6	500	25.5	22.5 - 28.8	700	33.1	29.9 - 36.5	500	23.0	20.1 - 26.1	100	6.3	4.7 - 8.4
<b>131-185%</b>	<50	7.9	4.7 - 13.0	<50	3.7	1.9 - 7.1	200	30.3	24.3 - 37.1	200	29.2	23.6 - 35.5	100	22.9	17.8 - 28.9	<50	3.1	1.3 - 7.3
<b>186+%</b>	100	12.8	9.9 - 16.2	100	7.2	5.1 - 10.1	300	27.1	23.4 - 31.2	300	25.3	21.6 - 29.4	200	20.7	17.4 - 24.4	100	4.6	3.2 - 6.6

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.1	2.1 - 4.5	2,100
100	3.5	2.4 - 5.0	2,000

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.1	2.1 - 4.7	1,700
100	3.4	2.4 - 4.8	2,300

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	4.1	2.9 - 5.7	2,100
<50	2.9	1.2 - 6.9	600
<50	2.2	1.3 - 3.8	1,200

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Income (2004-2011)	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<\$10,000	100	5.1	3.4 - 7.8	<50	2.2	1.1 - 4.2	200	23.0	19.0 - 27.6	400	34.7	30.0 - 39.6	200	22.7	18.7 - 27.2	100	7.0	4.7 - 10.3
\$10,000-\$24,999	100	5.1	3.5 - 7.4	<50	3.9	2.5 - 5.9	300	27.2	23.1 - 31.7	400	31.9	27.9 - 36.3	300	24.8	20.9 - 29.1	100	5.6	3.7 - 8.3
\$25,000-\$49,999	100	9.7	6.8 - 13.8	<50	3.7	2.3 - 5.7	300	29.5	25.0 - 34.4	200	27.1	22.8 - 31.8	200	22.0	18.2 - 26.4	<50	4.1	2.4 - 6.9
>=\$50,000	100	13.1	9.7 - 17.4	100	9.8	6.6 - 14.2	200	27.2	22.5 - 32.5	200	26.1	21.4 - 31.4	100	17.4	13.7 - 21.8	<50	4.4	2.7 - 7.0

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	6.3	3.8 - 10.3	<50	3.1	1.5 - 6.5	100	22.8	16.9 - 30.0	100	28.3	21.9 - 35.7	100	33.6	26.6 - 41.4	<50	4.6	2.3 - 9.1
Normal birth weight	300	7.9	6.6 - 9.5	200	5.6	4.4 - 7.0	1,000	27.0	24.7 - 29.4	1,100	30.5	28.2 - 33.0	800	20.5	18.5 - 22.6	200	5.1	4.0 - 6.4

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.8	0.2 - 2.3	<50	1.4	0.7 - 2.9	200	14.8	12.2 - 17.9	600	38.3	34.5 - 42.3	500	32.7	29.0 - 36.5	100	8.1	6.1 - 10.6
No	300	12.4	10.4 - 14.6	200	8.0	6.3 - 9.9	800	34.3	31.4 - 37.4	600	25.3	22.7 - 28.1	300	14.2	12.2 - 16.4	100	3.0	2.1 - 4.4

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	5.4	3.5 - 8.1	1,000
<50	1.6	0.8 - 3.2	1,100
<50	4.0	2.4 - 6.5	900
<50	2.1	1.0 - 4.2	700

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.3	0.3 - 4.7	300
100	3.5	2.6 - 4.5	3,700

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.9	2.6 - 5.8	1,600
100	2.8	2.0 - 4.0	2,500

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005-2009**

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	0	0.0	-	200	5.8	4.7 - 7.2	1,100	28.8	26.5 - 31.3	1,200	33.0	30.7 - 35.5	900	23.3	21.2 - 25.5	200	5.5	4.4 - 6.8
<b>No</b>	300	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	7.0	5.5 - 9.0	100	4.4	3.2 - 6.1	600	25.0	22.2 - 28.0	700	31.9	28.9 - 35.1	500	22.3	19.6 - 25.2	100	5.4	4.0 - 7.2
<b>No</b>	200	8.9	7.0 - 11.3	100	6.6	4.9 - 8.9	500	28.8	25.4 - 32.4	500	28.9	25.6 - 32.4	400	19.9	17.1 - 22.9	100	4.7	3.3 - 6.7

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.6	2.7 - 4.6	3,700
0	0.0	-	300

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	4.0	2.9 - 5.5	2,300
<50	2.3	1.5 - 3.7	1,800

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	200	5.9	4.9 - 7.1	100	3.4	2.7 - 4.4	1,000	25.9	23.9 - 28.0	1,300	32.1	29.9 - 34.3	900	23.5	21.6 - 25.5	200	4.8	3.9 - 5.9

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	<50	5.3	3.1 - 8.7	<50	3.7	1.9 - 6.9	100	24.1	19.3 - 29.6	200	36.1	30.5 - 42.0	200	25.3	20.4 - 30.8	<50	2.8	1.4 - 5.4
<b>Honolulu</b>	200	5.9	4.7 - 7.3	100	3.8	2.8 - 5.0	700	25.3	22.9 - 27.8	800	32.0	29.4 - 34.7	600	23.0	20.8 - 25.5	100	5.2	4.1 - 6.6
<b>Kauai</b>	<50	7.9	3.7 - 16.3	<50	1.1	0.2 - 6.8	100	26.6	18.3 - 36.9	100	31.3	22.2 - 42.1	<50	24.3	16.3 - 34.7	<50	3.6	1.2 - 10.2
<b>Maui</b>	<50	6.4	3.8 - 10.4	<50	1.9	0.7 - 5.0	100	32.3	26.1 - 39.1	100	27.0	21.2 - 33.7	100	23.5	18.1 - 29.9	<50	5.9	3.3 - 10.4

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
200	4.3	3.5 - 5.4	3,900

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.9	1.4 - 5.9	600
100	4.8	3.7 - 6.2	2,600
<50	5.1	2.0 - 12.3	200
<50	3.1	1.4 - 6.5	400



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	100	8.1	5.9 - 11.1	100	6.4	4.4 - 9.1	200	24.1	20.3 - 28.3	200	23.1	19.4 - 27.3	300	28.9	24.9 - 33.3	<50	5.2	3.4 - 7.7
Native Hawaiian	100	4.2	2.8 - 6.1	<50	2.4	1.4 - 4.0	400	25.7	22.3 - 29.3	500	36.8	33.0 - 40.8	300	22.0	18.8 - 25.5	100	5.4	3.8 - 7.6
Chinese	<50	9.6	4.2 - 20.5	<50	5.5	2.0 - 14.1	<50	25.9	16.8 - 37.7	<50	26.0	16.8 - 38.0	<50	27.0	17.7 - 38.8	<50	4.0	1.2 - 12.3
Filipino	<50	6.7	4.3 - 10.3	<50	1.8	0.7 - 4.5	200	32.4	27.1 - 38.3	200	34.9	29.4 - 40.8	100	18.8	14.6 - 23.9	<50	3.1	1.6 - 5.9
Japanese	<50	8.2	5.1 - 13.1	<50	3.4	1.7 - 6.8	100	23.4	17.9 - 30.0	100	34.7	28.1 - 41.9	100	23.7	18.2 - 30.2	<50	2.9	1.4 - 6.2
Other	<50	4.3	2.4 - 7.7	<50	2.6	1.2 - 5.6	100	25.8	20.7 - 31.7	200	30.3	24.9 - 36.2	100	22.4	17.7 - 27.9	<50	5.6	3.5 - 9.1

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	2.0	0.7 - 5.6	<50	2.7	1.1 - 6.3	100	33.1	26.6 - 40.4	100	32.8	26.3 - 40.0	100	19.2	14.1 - 25.5	<50	6.0	3.4 - 10.4
20-24	100	6.5	4.7 - 8.8	<50	3.7	2.5 - 5.6	300	26.3	22.9 - 30.0	400	30.9	27.2 - 34.7	300	22.8	19.6 - 26.4	100	4.5	3.1 - 6.6
25-34	100	5.6	4.3 - 7.4	100	3.7	2.6 - 5.2	400	23.9	21.1 - 27.0	600	33.7	30.5 - 37.1	400	24.6	21.7 - 27.7	100	5.1	3.8 - 6.9
35+	<50	9.9	6.4 - 14.9	<50	1.9	0.7 - 5.0	100	25.7	20.1 - 32.1	100	28.2	22.3 - 34.9	100	25.8	20.1 - 32.4	<50	3.2	1.4 - 7.0

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	4.3	2.8 - 6.6	900
100	3.6	2.3 - 5.5	1,500
<50	2.0	0.4 - 10.0	<50
<50	2.3	1.1 - 4.8	500
<50	3.6	1.7 - 7.7	400
<50	8.9	5.9 - 13.2	600

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	4.3	2.2 - 8.2	400
100	5.3	3.7 - 7.5	1,300
100	3.4	2.3 - 4.9	1,700
<50	5.5	3.0 - 9.7	400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	100	6.1	4.8 - 7.9	100	2.8	1.9 - 4.0	600	27.7	24.9 - 30.7	600	30.8	27.9 - 33.9	500	23.6	20.9 - 26.4	100	5.1	3.9 - 6.8
<b>Female</b>	100	5.7	4.3 - 7.4	100	4.1	3.0 - 5.7	400	24.0	21.2 - 27.0	600	33.4	30.3 - 36.7	400	23.5	20.7 - 26.5	100	4.5	3.3 - 6.1

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	9.0	7.2 - 11.1	100	4.7	3.5 - 6.4	500	26.1	23.3 - 29.2	500	30.1	27.0 - 33.3	400	22.5	19.8 - 25.4	100	3.8	2.7 - 5.4
<b>No</b>	100	3.4	2.4 - 4.8	<50	2.3	1.5 - 3.5	600	25.8	23.0 - 28.7	700	33.7	30.7 - 36.8	500	24.4	21.7 - 27.2	100	5.6	4.3 - 7.3

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	100	4.1	3.0 - 5.6	<50	2.4	1.5 - 3.6	500	23.8	21.1 - 26.8	700	34.6	31.4 - 37.8	500	23.8	21.1 - 26.8	100	5.5	4.1 - 7.3
<b>131-185%</b>	<50	4.4	2.4 - 7.9	<50	2.4	1.1 - 5.1	100	27.2	21.9 - 33.3	200	34.2	28.4 - 40.5	100	25.6	20.4 - 31.5	<50	3.0	1.5 - 6.2
<b>186+%</b>	100	9.2	7.0 - 11.8	100	4.6	3.2 - 6.7	300	27.8	24.2 - 31.6	300	27.0	23.4 - 30.9	300	23.5	20.2 - 27.2	100	5.3	3.8 - 7.5

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.9	2.8 - 5.3	2,000
100	4.8	3.5 - 6.6	1,900

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.8	2.7 - 5.3	1,800
100	4.8	3.6 - 6.4	2,100

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	5.8	4.4 - 7.6	2,000
<50	3.2	1.6 - 6.5	500
<50	2.7	1.6 - 4.3	1,100

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Income (2004-2011)	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<\$10,000	100	5.1	3.4 - 7.7	<50	0.7	0.2 - 2.0	200	21.7	18.0 - 25.9	400	36.0	31.5 - 40.7	200	22.7	18.9 - 27.0	100	6.3	4.3 - 9.0
\$10,000-\$24,999	<50	3.2	2.0 - 5.1	<50	3.8	2.4 - 5.8	300	26.2	22.5 - 30.2	400	33.1	29.1 - 37.3	300	25.5	21.9 - 29.5	100	5.2	3.5 - 7.6
\$25,000-\$49,999	<50	5.7	3.8 - 8.3	<50	3.3	2.0 - 5.5	200	26.7	22.6 - 31.2	300	30.0	25.7 - 34.6	200	26.8	22.7 - 31.3	<50	3.6	2.2 - 6.0
>=\$50,000	100	11.2	8.2 - 15.2	<50	4.9	3.1 - 7.8	200	28.1	23.6 - 33.2	200	28.8	24.1 - 34.1	100	18.9	15.1 - 23.5	<50	5.3	3.4 - 8.2

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	2.4	0.8 - 7.1	<50	2.0	0.6 - 6.1	100	25.9	19.2 - 33.9	100	29.1	21.9 - 37.4	100	33.1	25.6 - 41.7	<50	4.6	2.2 - 9.7
Normal birth weight	200	6.2	5.2 - 7.5	100	3.5	2.7 - 4.6	900	25.9	23.9 - 28.1	1,200	32.3	30.1 - 34.6	800	22.7	20.7 - 24.8	200	4.8	3.9 - 6.0

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.5	0.2 - 1.5	<50	0.9	0.4 - 1.9	200	15.7	13.2 - 18.7	600	37.5	33.9 - 41.2	500	33.5	30.0 - 37.2	100	6.9	5.2 - 9.1
No	200	9.5	7.9 - 11.3	100	5.1	3.9 - 6.6	800	32.6	29.9 - 35.5	700	28.5	25.9 - 31.3	400	17.0	14.9 - 19.3	100	3.5	2.6 - 4.7

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	7.6	5.3 - 10.7	1,000
<50	3.0	1.9 - 4.9	1,100
<50	4.0	2.5 - 6.3	900
<50	2.7	1.4 - 5.1	700

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.9	1.1 - 7.4	300
200	4.5	3.5 - 5.6	3,600

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	5.1	3.6 - 7.1	1,500
100	3.9	2.9 - 5.2	2,400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004-2008**

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	0	0.0	-	100	3.6	2.8 - 4.6	1,000	27.6	25.5 - 29.8	1,300	34.1	31.8 - 36.4	900	25.0	23.0 - 27.1	200	5.1	4.2 - 6.3
<b>No</b>	200	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	100	5.3	4.1 - 6.9	100	3.1	2.1 - 4.4	500	24.0	21.4 - 26.8	700	34.2	31.3 - 37.3	500	23.0	20.4 - 25.8	100	5.1	3.9 - 6.8
<b>No</b>	100	6.7	5.2 - 8.7	100	3.8	2.7 - 5.4	500	28.5	25.5 - 31.8	500	29.6	26.5 - 33.0	400	23.6	20.8 - 26.7	100	4.5	3.2 - 6.1

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
200	4.6	3.7 - 5.8	3,700
0	0.0	-	200

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	5.3	4.0 - 6.9	2,200
100	3.1	2.1 - 4.7	1,700



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	300	6.7	5.6 - 8.0	100	3.6	2.8 - 4.6	1,000	24.9	23.0 - 27.0	1,300	33.0	30.9 - 35.3	900	23.5	21.6 - 25.6	200	4.8	3.9 - 5.9

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	<50	4.8	3.0 - 7.5	<50	3.3	1.7 - 6.1	100	23.4	19.0 - 28.4	200	37.3	32.1 - 42.9	200	25.7	21.2 - 30.8	<50	2.6	1.4 - 5.0
<b>Honolulu</b>	200	7.0	5.6 - 8.7	100	3.9	2.9 - 5.2	700	24.6	22.2 - 27.1	900	32.8	30.2 - 35.6	600	23.1	20.7 - 25.7	100	5.0	3.9 - 6.4
<b>Kauai</b>	<50	7.2	3.5 - 14.3	<50	1.7	0.5 - 6.3	<50	23.4	16.1 - 32.6	100	33.9	25.2 - 43.8	<50	24.2	16.9 - 33.4	<50	4.8	2.1 - 10.8
<b>Maui</b>	<50	7.3	4.7 - 11.3	<50	2.7	1.2 - 5.7	100	30.1	24.3 - 36.6	100	27.6	22.1 - 33.8	100	22.6	17.6 - 28.6	<50	6.6	3.8 - 11.1

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.5	2.7 - 4.5	3,900

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.9	1.4 - 5.9	600
100	3.6	2.7 - 4.9	2,700
<50	4.8	1.9 - 11.6	200
<50	3.1	1.5 - 6.4	400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	100	9.9	7.2 - 13.3	100	6.3	4.3 - 9.2	200	21.6	18.0 - 25.8	200	26.1	22.0 - 30.7	300	28.0	23.8 - 32.6	<50	5.2	3.5 - 7.8
Native Hawaiian	100	4.6	3.4 - 6.4	<50	2.6	1.6 - 4.1	400	24.9	21.8 - 28.2	500	36.7	33.1 - 40.3	300	22.9	19.9 - 26.1	100	5.3	3.8 - 7.3
Chinese	<50	18.4	5.9 - 44.7	<50	5.7	2.0 - 15.3	<50	18.2	9.7 - 31.5	<50	19.8	10.8 - 33.5	<50	33.7	17.1 - 55.7	<50	2.1	0.4 - 10.7
Filipino	<50	7.0	4.5 - 10.6	<50	2.5	1.0 - 5.9	200	32.9	27.3 - 39.1	200	35.1	29.5 - 41.2	100	16.7	12.5 - 21.9	<50	2.9	1.3 - 6.1
Japanese	<50	8.1	5.0 - 12.9	<50	3.5	1.8 - 6.7	100	25.0	19.1 - 32.0	100	34.1	27.5 - 41.5	100	23.5	17.9 - 30.1	<50	3.0	1.5 - 6.0
Other	<50	4.5	2.4 - 8.5	<50	2.4	0.9 - 6.2	100	23.2	18.2 - 29.3	200	33.0	27.0 - 39.7	100	23.7	18.4 - 30.0	<50	6.1	3.8 - 9.7

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	2.3	1.0 - 5.1	<50	4.7	2.3 - 9.5	100	29.8	23.8 - 36.7	200	35.6	29.1 - 42.6	100	19.5	14.5 - 25.6	<50	5.1	2.9 - 9.0
20-24	100	5.9	4.3 - 8.0	<50	3.4	2.2 - 5.2	400	26.3	22.9 - 30.0	400	32.3	28.7 - 36.2	300	22.5	19.4 - 26.0	100	5.0	3.5 - 7.0
25-34	100	7.6	5.9 - 9.7	100	3.5	2.4 - 5.0	400	22.6	19.8 - 25.6	600	34.4	31.1 - 37.8	400	24.5	21.6 - 27.7	100	5.0	3.6 - 6.8
35+	<50	10.8	6.9 - 16.4	<50	2.9	1.1 - 7.6	100	24.8	19.1 - 31.5	100	26.5	20.8 - 33.2	100	27.4	21.3 - 34.5	<50	2.7	1.1 - 6.5

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.8	1.7 - 4.8	900
<50	3.1	2.0 - 4.9	1,500
<50	2.1	0.4 - 10.7	<50
<50	3.0	1.4 - 6.2	500
<50	2.8	1.2 - 6.5	400
<50	6.9	4.2 - 11.2	500

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	3.0	1.4 - 6.4	500
100	4.6	3.1 - 6.6	1,300
<50	2.5	1.6 - 3.9	1,700
<50	4.9	2.6 - 9.2	400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	100	6.6	5.2 - 8.4	100	2.8	1.9 - 4.2	500	25.8	23.1 - 28.7	700	33.1	30.1 - 36.2	500	23.2	20.5 - 26.0	100	5.1	3.8 - 6.8
<b>Female</b>	100	6.8	5.3 - 8.7	100	4.3	3.1 - 6.0	500	24.0	21.3 - 27.0	600	33.0	29.9 - 36.2	500	23.9	21.2 - 26.9	100	4.4	3.3 - 6.0

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	9.3	7.5 - 11.6	100	5.1	3.7 - 7.0	500	25.0	22.1 - 28.0	600	31.7	28.6 - 35.0	400	22.4	19.7 - 25.4	100	3.6	2.5 - 5.2
<b>No</b>	100	4.4	3.3 - 5.9	<50	2.2	1.5 - 3.4	500	24.9	22.2 - 27.8	700	34.1	31.2 - 37.2	500	24.5	21.8 - 27.3	100	5.7	4.4 - 7.4

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	100	5.2	3.9 - 6.8	100	2.9	2.0 - 4.3	500	22.6	20.0 - 25.5	700	35.9	32.8 - 39.1	500	23.1	20.4 - 26.0	100	5.2	3.9 - 6.9
<b>131-185%</b>	<50	7.3	4.3 - 12.1	<50	2.6	1.2 - 5.4	100	28.6	22.9 - 35.0	200	30.2	24.7 - 36.5	100	27.0	21.6 - 33.3	<50	2.8	1.3 - 5.8
<b>186+%</b>	100	9.1	7.0 - 11.9	<50	4.4	2.9 - 6.7	300	26.2	22.6 - 30.2	300	29.0	25.1 - 33.2	300	24.1	20.5 - 28.0	100	5.4	3.8 - 7.7

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.5	2.5 - 4.9	2,000
100	3.5	2.4 - 5.1	1,900

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.8	1.9 - 4.1	1,800
100	4.1	2.9 - 5.6	2,100

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	5.1	3.8 - 6.9	2,000
<50	1.5	0.6 - 3.8	500
<50	1.7	0.9 - 3.2	1,100

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	3.2	1.1 - 9.2	<50	3.0	0.9 - 9.1	100	25.8	19.1 - 33.9	100	29.5	22.3 - 37.8	100	31.5	24.1 - 39.9	<50	4.5	2.2 - 9.2
Normal birth weight	300	7.0	5.8 - 8.3	100	3.6	2.8 - 4.7	900	24.8	22.8 - 27.0	1,200	33.3	31.1 - 35.7	800	22.9	20.9 - 25.0	200	4.8	3.8 - 5.9

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.5	0.2 - 1.5	<50	1.4	0.7 - 2.9	200	15.5	12.9 - 18.5	600	37.5	33.9 - 41.3	500	33.6	30.1 - 37.3	100	7.0	5.3 - 9.3
No	300	10.7	9.0 - 12.7	100	5.0	3.8 - 6.5	700	31.0	28.4 - 33.8	700	30.0	27.3 - 32.8	400	17.1	14.9 - 19.4	100	3.4	2.5 - 4.5

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	0	0.0	-	100	3.8	3.0 - 4.9	1,000	26.7	24.6 - 28.9	1,300	35.4	33.1 - 37.8	900	25.2	23.2 - 27.4	200	5.1	4.1 - 6.3
No	300	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.6	0.9 - 6.8	300
100	3.6	2.8 - 4.6	3,600

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	4.5	3.1 - 6.4	1,500
100	2.9	2.0 - 4.1	2,400

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.7	2.9 - 4.8	3,700
0	0.0	-	300



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003-2007**

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	100	5.2	4.0 - 6.7	100	3.5	2.4 - 4.9	500	23.3	20.7 - 26.0	800	35.5	32.5 - 38.5	500	22.8	20.3 - 25.6	100	5.7	4.3 - 7.4
<b>No</b>	100	8.6	6.7 - 10.8	100	3.7	2.5 - 5.2	500	27.0	23.9 - 30.2	500	30.4	27.2 - 33.8	400	23.9	21.1 - 27.1	100	3.7	2.6 - 5.1

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	4.1	3.0 - 5.6	2,200
<50	2.8	1.9 - 4.2	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	300	7.1	5.9 - 8.4	100	3.5	2.7 - 4.5	900	23.2	21.2 - 25.2	1,300	34.1	31.9 - 36.4	1,000	24.5	22.5 - 26.6	200	4.8	3.9 - 5.9

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	<50	5.8	4.0 - 8.4	<50	2.6	1.3 - 5.2	100	23.6	19.5 - 28.3	200	35.7	30.9 - 40.8	200	26.4	22.2 - 31.0	<50	3.0	1.7 - 5.2
<b>Honolulu</b>	200	7.5	6.0 - 9.3	100	4.1	3.0 - 5.5	600	22.1	19.7 - 24.7	900	34.0	31.2 - 37.0	600	24.5	21.9 - 27.2	100	5.1	3.9 - 6.5
<b>Kauai</b>	<50	7.3	3.6 - 14.2	<50	1.8	0.5 - 6.6	<50	24.2	17.0 - 33.2	100	39.5	30.8 - 48.9	<50	19.5	13.5 - 27.4	<50	4.0	1.7 - 9.3
<b>Maui</b>	<50	6.0	3.7 - 9.6	<50	1.9	0.9 - 3.9	100	28.7	23.2 - 34.8	100	30.0	24.6 - 35.9	100	24.4	19.4 - 30.1	<50	5.9	3.4 - 10.1

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.9	2.2 - 3.8	3,900

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.9	1.4 - 5.9	600
100	2.8	1.9 - 3.9	2,700
<50	3.7	1.3 - 10.3	200
<50	3.2	1.5 - 6.6	400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	100	9.7	7.0 - 13.4	100	6.3	4.2 - 9.5	200	19.9	16.2 - 24.1	300	29.4	24.9 - 34.4	300	28.5	24.0 - 33.4	<50	4.3	2.6 - 6.9
Native Hawaiian	100	5.5	4.2 - 7.2	<50	2.3	1.5 - 3.7	300	22.7	19.9 - 25.7	600	37.5	34.2 - 40.9	400	24.0	21.3 - 27.0	100	5.5	4.1 - 7.4
Chinese	<50	17.4	4.5 - 48.5	<50	3.2	1.0 - 9.6	<50	22.9	8.0 - 50.2	<50	13.5	6.4 - 26.1	<50	38.3	18.7 - 62.7	<50	2.4	0.4 - 12.2
Filipino	<50	6.1	3.6 - 10.1	<50	2.1	0.7 - 5.6	200	30.2	24.5 - 36.6	200	35.0	29.1 - 41.3	100	20.3	15.3 - 26.4	<50	2.9	1.3 - 6.3
Japanese	<50	9.6	5.8 - 15.6	<50	3.3	1.7 - 6.3	100	26.1	19.7 - 33.7	100	33.8	26.9 - 41.5	100	21.4	16.0 - 28.1	<50	3.0	1.5 - 6.0
Other	<50	4.8	2.5 - 9.3	<50	3.5	1.5 - 8.2	100	20.6	15.3 - 27.1	200	33.2	26.7 - 40.4	100	24.9	19.1 - 31.8	<50	7.0	4.1 - 11.7

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	3.2	1.7 - 6.0	<50	4.6	2.2 - 9.5	100	27.9	22.1 - 34.5	200	36.7	30.3 - 43.5	100	20.3	15.4 - 26.1	<50	4.3	2.4 - 7.7
20-24	100	5.1	3.6 - 7.1	<50	2.8	1.8 - 4.4	300	24.1	20.8 - 27.7	500	35.2	31.4 - 39.1	300	24.1	20.8 - 27.7	100	5.3	3.7 - 7.5
25-34	200	9.3	7.3 - 11.8	100	4.0	2.8 - 5.8	400	21.4	18.6 - 24.5	600	34.1	30.8 - 37.6	400	24.3	21.4 - 27.5	100	4.7	3.4 - 6.4
35+	<50	8.4	5.0 - 13.7	<50	2.1	0.6 - 6.9	100	22.2	16.7 - 28.8	100	27.6	21.5 - 34.7	100	31.8	25.2 - 39.3	<50	3.8	1.8 - 7.6

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.9	1.0 - 3.5	900
<50	2.3	1.4 - 3.9	1,500
<50	2.4	0.4 - 12.2	<50
<50	3.4	1.6 - 7.2	500
<50	2.8	1.2 - 6.5	400
<50	5.9	3.4 - 10.0	500

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	3.1	1.4 - 6.5	500
<50	3.5	2.3 - 5.5	1,300
<50	2.1	1.2 - 3.4	1,700
<50	4.2	2.1 - 8.1	400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	100	7.0	5.5 - 9.0	100	3.1	2.1 - 4.5	500	23.0	20.4 - 25.9	700	33.6	30.5 - 36.8	500	25.1	22.3 - 28.0	100	5.2	3.9 - 7.0
<b>Female</b>	100	7.1	5.5 - 9.1	100	3.9	2.7 - 5.6	400	23.3	20.5 - 26.4	700	34.7	31.6 - 38.0	500	23.9	21.1 - 26.9	100	4.3	3.2 - 5.8

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	9.8	7.8 - 12.2	100	4.7	3.4 - 6.6	400	23.5	20.6 - 26.6	600	31.6	28.4 - 35.0	400	24.0	21.1 - 27.2	100	3.8	2.7 - 5.4
<b>No</b>	100	4.6	3.5 - 6.1	<50	2.4	1.6 - 3.6	500	22.9	20.3 - 25.7	700	36.4	33.4 - 39.5	500	24.9	22.2 - 27.8	100	5.6	4.3 - 7.3

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	100	5.6	4.3 - 7.3	100	2.5	1.7 - 3.8	400	21.3	18.7 - 24.1	700	36.9	33.8 - 40.1	500	24.3	21.6 - 27.2	100	5.4	4.0 - 7.1
<b>131-185%</b>	<50	8.0	4.7 - 13.2	<50	4.7	2.5 - 8.6	100	25.7	20.4 - 31.8	200	31.3	25.6 - 37.6	100	26.4	20.9 - 32.7	<50	2.6	1.3 - 5.3
<b>186+%</b>	100	9.9	7.4 - 13.1	<50	3.6	2.2 - 5.8	200	24.2	20.4 - 28.4	300	31.0	26.7 - 35.6	200	24.3	20.5 - 28.6	100	5.2	3.6 - 7.5

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.0	2.1 - 4.4	2,000
100	2.7	1.8 - 4.2	1,900

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.5	1.6 - 3.8	1,800
100	3.2	2.2 - 4.7	2,100

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	4.0	2.9 - 5.6	2,000
<50	1.4	0.5 - 3.5	600
<50	1.8	0.9 - 3.7	1,000



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	3.5	1.2 - 10.0	<50	2.7	0.8 - 9.0	100	25.6	18.8 - 33.8	100	29.3	21.9 - 38.0	100	32.2	24.2 - 41.3	<50	3.8	1.8 - 7.9
Normal birth weight	300	7.3	6.1 - 8.8	100	3.6	2.7 - 4.6	800	23.0	21.0 - 25.1	1,200	34.6	32.3 - 36.9	900	23.8	21.8 - 26.0	200	4.8	3.9 - 6.0

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.5	0.2 - 1.4	<50	2.1	1.2 - 3.9	200	14.2	11.7 - 17.1	600	37.0	33.4 - 40.8	500	34.1	30.6 - 37.9	100	7.6	5.8 - 10.0
No	300	11.3	9.5 - 13.5	100	4.2	3.2 - 5.6	700	28.9	26.3 - 31.8	800	32.2	29.5 - 35.1	400	18.3	16.1 - 20.7	100	3.0	2.2 - 4.2

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	0	0.0	-	100	3.8	2.9 - 4.9	900	24.9	22.8 - 27.1	1,300	36.7	34.4 - 39.1	1,000	26.4	24.3 - 28.6	200	5.1	4.1 - 6.3
No	300	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.8	1.0 - 7.4	300
100	2.9	2.2 - 3.9	3,600

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	4.4	3.0 - 6.4	1,500
<50	2.0	1.3 - 3.0	2,400

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.1	2.3 - 4.1	3,600
0	0.0	-	300

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002-2006**

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	100	5.5	4.2 - 7.0	100	3.1	2.1 - 4.5	500	21.9	19.3 - 24.6	800	36.4	33.4 - 39.5	500	24.2	21.6 - 27.0	100	5.5	4.2 - 7.2
<b>No</b>	100	8.7	6.7 - 11.1	100	4.1	2.9 - 5.9	400	24.8	21.9 - 28.1	500	31.6	28.3 - 35.1	400	24.8	21.8 - 28.1	100	3.8	2.8 - 5.3

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.5	2.5 - 5.0	2,100
<50	2.1	1.3 - 3.4	1,700

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	300	7.0	5.9 - 8.4	100	3.5	2.7 - 4.5	800	22.5	20.7 - 24.5	1,300	34.2	32.1 - 36.4	1,000	25.5	23.5 - 27.5	200	4.9	4.0 - 6.0

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	<50	6.5	4.6 - 9.0	<50	1.6	0.7 - 3.4	100	22.8	19.0 - 27.1	200	36.7	32.3 - 41.3	200	26.6	22.8 - 30.7	<50	3.8	2.4 - 5.9
<b>Honolulu</b>	200	7.3	5.8 - 9.1	100	4.2	3.2 - 5.7	600	22.0	19.6 - 24.6	900	33.5	30.8 - 36.4	700	25.7	23.1 - 28.4	100	5.0	3.9 - 6.4
<b>Kauai</b>	<50	6.9	3.5 - 13.5	<50	2.2	0.7 - 6.6	<50	21.9	15.8 - 29.6	100	38.3	30.5 - 46.7	<50	22.4	16.4 - 29.7	<50	5.9	3.1 - 11.1
<b>Maui</b>	<50	6.3	4.0 - 9.7	<50	2.1	1.1 - 3.9	100	26.1	21.2 - 31.7	100	33.3	28.2 - 38.9	100	24.1	19.7 - 29.1	<50	5.6	3.4 - 9.2

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.3	1.7 - 3.1	3,800

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.1	1.0 - 4.6	600
100	2.3	1.6 - 3.4	2,600
<50	2.4	0.6 - 8.3	200
<50	2.5	1.2 - 5.5	400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	100	9.3	6.5 - 13.0	<50	5.1	3.1 - 8.2	200	19.6	16.0 - 23.9	300	30.2	25.5 - 35.3	200	28.8	24.3 - 33.9	<50	4.7	2.9 - 7.3
Native Hawaiian	100	5.4	4.2 - 7.0	<50	2.9	2.0 - 4.1	300	22.4	19.9 - 25.1	600	38.2	35.2 - 41.2	400	24.1	21.7 - 26.8	100	5.4	4.2 - 6.9
Chinese	<50	18.1	5.5 - 45.6	<50	2.4	0.8 - 7.2	<50	26.7	10.9 - 52.0	<50	8.0	3.7 - 16.3	<50	41.3	21.7 - 64.1	<50	1.8	0.3 - 9.2
Filipino	<50	6.0	3.6 - 9.8	<50	2.0	0.8 - 5.1	100	26.7	21.4 - 32.9	200	37.3	31.5 - 43.5	100	22.3	17.3 - 28.3	<50	2.8	1.3 - 5.8
Japanese	<50	9.7	5.7 - 16.0	<50	4.6	2.4 - 8.7	100	27.3	20.7 - 35.2	100	32.2	25.3 - 40.0	100	20.6	15.2 - 27.3	<50	3.8	2.0 - 7.0
Other	<50	5.8	3.1 - 10.6	<50	3.3	1.3 - 8.4	100	19.0	13.8 - 25.7	100	29.8	23.4 - 37.1	100	30.1	23.6 - 37.5	<50	7.6	4.4 - 12.7

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	4.3	2.5 - 7.1	<50	5.0	2.5 - 9.7	100	25.4	20.1 - 31.4	200	37.0	31.1 - 43.4	100	20.5	16.0 - 26.0	<50	5.4	3.5 - 8.4
20-24	100	4.8	3.5 - 6.6	<50	2.1	1.3 - 3.3	300	23.3	20.3 - 26.7	500	37.5	33.9 - 41.3	300	24.6	21.5 - 28.1	100	4.9	3.5 - 6.8
25-34	200	9.6	7.5 - 12.2	100	4.3	3.0 - 6.2	400	21.7	18.8 - 24.8	500	32.1	28.8 - 35.5	400	26.1	23.1 - 29.4	100	4.6	3.4 - 6.2
35+	<50	7.3	4.0 - 13.0	<50	3.1	1.2 - 7.6	100	20.0	14.7 - 26.6	100	27.9	21.8 - 35.1	100	32.4	25.5 - 40.1	<50	5.8	3.1 - 10.7

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.3	1.3 - 4.1	900
<50	1.6	0.9 - 2.9	1,500
<50	1.8	0.3 - 9.2	<50
<50	2.9	1.3 - 6.6	500
<50	1.7	0.6 - 5.0	400
<50	4.4	2.4 - 8.0	500

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.4	1.0 - 5.4	500
<50	2.7	1.7 - 4.4	1,300
<50	1.7	1.0 - 3.0	1,600
<50	3.4	1.5 - 7.4	300



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	100	6.8	5.3 - 8.7	100	3.4	2.3 - 4.8	400	22.5	20.0 - 25.2	700	34.1	31.2 - 37.2	500	25.7	23.0 - 28.6	100	4.9	3.7 - 6.5
<b>Female</b>	100	7.3	5.6 - 9.3	100	3.6	2.5 - 5.2	400	22.6	19.9 - 25.6	600	34.3	31.3 - 37.5	500	25.2	22.4 - 28.2	100	4.9	3.8 - 6.3

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	9.3	7.4 - 11.7	100	4.6	3.3 - 6.4	400	22.9	20.1 - 25.9	600	31.3	28.2 - 34.6	500	25.4	22.5 - 28.6	100	4.4	3.2 - 5.9
<b>No</b>	100	4.9	3.8 - 6.4	<50	2.5	1.7 - 3.6	400	22.3	19.8 - 24.9	700	36.9	34.0 - 39.9	500	25.5	22.9 - 28.3	100	5.4	4.2 - 6.9

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	100	5.9	4.6 - 7.6	100	2.7	1.9 - 4.0	400	21.2	18.8 - 23.9	700	36.9	34.0 - 39.9	500	24.9	22.3 - 27.6	100	5.4	4.2 - 6.9
<b>131-185%</b>	<50	8.6	5.3 - 13.8	<50	4.8	2.7 - 8.6	100	23.5	18.5 - 29.3	200	31.2	25.8 - 37.1	200	27.6	22.3 - 33.6	<50	2.7	1.5 - 5.0
<b>186+%</b>	100	9.4	6.8 - 12.7	<50	3.6	2.1 - 5.9	200	24.2	20.3 - 28.6	300	30.7	26.3 - 35.5	200	25.4	21.3 - 29.9	<50	4.9	3.3 - 7.2

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.5	1.7 - 3.7	1,900
<50	2.1	1.3 - 3.4	1,800

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.1	1.3 - 3.3	1,800
<50	2.5	1.6 - 3.8	2,000

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	3.0	2.0 - 4.3	2,000
<50	1.6	0.7 - 3.7	600
<50	1.9	0.9 - 3.9	900

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	4.1	1.5 - 10.7	<50	2.6	0.7 - 9.8	100	22.5	16.5 - 29.9	100	30.4	23.1 - 38.9	100	31.1	23.4 - 40.0	<50	6.3	3.6 - 10.7
Normal birth weight	300	7.3	6.1 - 8.7	100	3.6	2.7 - 4.6	800	22.6	20.6 - 24.7	1,200	34.6	32.3 - 36.9	900	25.0	22.9 - 27.1	200	4.8	3.9 - 5.9

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.6	0.2 - 1.5	<50	2.7	1.6 - 4.5	200	15.0	12.5 - 17.9	500	36.3	32.8 - 39.9	500	34.9	31.4 - 38.5	100	7.6	5.9 - 9.8
No	300	11.2	9.4 - 13.3	100	3.8	2.8 - 5.1	600	27.4	24.8 - 30.1	800	33.0	30.3 - 35.8	400	19.4	17.2 - 21.9	100	3.3	2.4 - 4.4

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	0	0.0	-	100	3.8	2.9 - 4.8	800	24.3	22.3 - 26.4	1,300	36.8	34.6 - 39.1	1,000	27.4	25.3 - 29.6	200	5.3	4.3 - 6.4
No	300	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	3.0	1.1 - 8.0	200
100	2.3	1.6 - 3.1	3,500

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.9	1.9 - 4.6	1,400
<50	2.0	1.3 - 3.0	2,300

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.5	1.8 - 3.4	3,500
0	0.0	-	300

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001-2005**

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	100	6.1	4.8 - 7.7	100	3.0	2.0 - 4.3	400	20.4	18.0 - 23.0	800	36.8	33.9 - 39.7	500	25.8	23.3 - 28.6	100	5.1	3.9 - 6.6
<b>No</b>	100	7.7	5.9 - 10.1	100	4.2	2.9 - 5.9	400	25.3	22.3 - 28.5	500	31.3	28.1 - 34.7	400	25.2	22.1 - 28.4	100	4.7	3.5 - 6.2

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.8	1.9 - 4.1	2,100
<50	1.7	1.0 - 2.9	1,600

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

State	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Totals</b>	300	7.6	6.5 - 9.0	100	3.6	2.8 - 4.6	800	21.3	19.5 - 23.2	1,300	33.7	31.6 - 35.8	1,000	27.0	25.0 - 29.0	200	4.9	4.1 - 5.9

County	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Hawaii</b>	<50	7.6	5.6 - 10.2	<50	1.5	0.8 - 2.6	100	21.2	18.0 - 24.9	200	35.6	31.7 - 39.7	200	28.9	25.4 - 32.6	<50	3.9	2.5 - 5.9
<b>Honolulu</b>	200	7.9	6.3 - 9.8	100	4.4	3.3 - 5.9	500	20.7	18.3 - 23.2	800	33.1	30.4 - 36.0	700	27.0	24.3 - 29.7	100	4.9	3.8 - 6.2
<b>Kauai</b>	<50	5.6	2.9 - 10.6	<50	1.0	0.3 - 3.9	<50	24.9	19.0 - 31.9	100	36.3	29.6 - 43.6	<50	23.6	18.1 - 30.3	<50	7.3	4.3 - 12.3
<b>Maui</b>	<50	7.0	4.7 - 10.3	<50	2.8	1.6 - 5.0	100	23.4	19.2 - 28.2	100	33.2	28.6 - 38.2	100	26.1	21.9 - 30.7	<50	5.0	3.3 - 7.8

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	1.9	1.4 - 2.7	3,700

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.4	0.6 - 3.4	600
100	2.1	1.4 - 3.1	2,500
<50	1.2	0.2 - 7.1	200
<50	2.4	1.1 - 5.1	400



**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

DOH Race-Ethnicity	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Caucasian	100	9.3	6.6 - 13.1	<50	5.1	3.1 - 8.3	100	17.2	13.8 - 21.3	300	31.2	26.5 - 36.3	300	31.2	26.6 - 36.3	<50	4.2	2.6 - 6.6
Native Hawaiian	100	5.8	4.7 - 7.2	<50	2.6	1.9 - 3.5	300	21.6	19.5 - 24.0	500	37.1	34.6 - 39.8	400	25.8	23.6 - 28.1	100	5.9	4.8 - 7.2
Chinese	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r
Filipino	<50	6.7	4.1 - 10.7	<50	2.5	1.0 - 5.8	100	25.6	20.3 - 31.8	200	35.8	30.0 - 42.0	100	22.9	17.8 - 29.0	<50	4.0	2.1 - 7.4
Japanese	<50	10.2	6.1 - 16.5	<50	5.3	2.8 - 9.7	100	26.8	20.1 - 34.8	100	32.2	25.1 - 40.2	100	21.6	15.9 - 28.7	<50	2.1	1.0 - 4.6
Other	<50	8.1	4.7 - 13.5	<50	3.9	1.7 - 8.7	100	17.1	12.0 - 23.6	100	28.7	22.3 - 36.0	100	30.9	24.2 - 38.5	<50	6.8	3.9 - 11.8

PRAMS Age Group	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<20	<50	6.0	3.9 - 9.1	<50	3.8	1.9 - 7.4	100	25.7	20.9 - 31.2	200	35.5	30.2 - 41.1	100	22.2	17.9 - 27.2	<50	4.3	2.8 - 6.4
20-24	100	5.4	4.0 - 7.2	<50	2.6	1.7 - 3.9	300	21.8	19.0 - 25.0	500	37.7	34.2 - 41.3	300	25.9	22.7 - 29.3	100	4.8	3.5 - 6.5
25-34	200	10.2	8.0 - 13.0	100	4.5	3.1 - 6.5	300	20.2	17.4 - 23.4	500	30.8	27.6 - 34.2	400	27.0	23.9 - 30.3	100	5.4	4.0 - 7.1
35+	<50	7.1	4.0 - 12.4	<50	2.9	1.1 - 7.3	100	17.2	12.4 - 23.3	100	28.6	22.5 - 35.7	100	38.3	31.2 - 45.9	<50	3.9	2.0 - 7.5

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.7	0.9 - 3.2	800
<50	1.1	0.6 - 2.0	1,400
n/r	n/r	n/r	<50
<50	2.5	1.0 - 6.1	500
<50	1.8	0.6 - 5.1	400
<50	4.5	2.3 - 8.6	500

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.5	1.1 - 5.8	500
<50	1.8	1.0 - 3.2	1,300
<50	1.8	1.1 - 3.1	1,500
<50	2.0	0.8 - 5.0	400

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

Infant Gender	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Male</b>	100	7.5	5.9 - 9.4	100	4.2	3.1 - 5.8	400	21.0	18.6 - 23.5	700	33.8	31.0 - 36.8	500	27.0	24.4 - 29.9	100	4.6	3.5 - 6.1
<b>Female</b>	100	7.8	6.1 - 9.9	100	2.9	1.9 - 4.3	400	21.6	18.9 - 24.6	600	33.5	30.5 - 36.7	500	26.9	24.1 - 30.0	100	5.2	4.1 - 6.6

Married	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	200	9.3	7.4 - 11.6	100	4.8	3.5 - 6.6	400	21.5	18.8 - 24.4	600	30.9	27.9 - 34.1	500	27.7	24.8 - 30.8	100	4.3	3.2 - 5.8
<b>No</b>	100	6.0	4.7 - 7.6	<50	2.4	1.6 - 3.5	400	21.1	18.8 - 23.7	700	36.4	33.6 - 39.3	500	26.3	23.7 - 29.0	100	5.4	4.3 - 6.9

Poverty Level	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>0-130%</b>	100	6.6	5.2 - 8.3	<50	2.3	1.6 - 3.4	400	20.1	17.8 - 22.6	700	35.7	32.9 - 38.5	500	27.0	24.5 - 29.7	100	5.8	4.6 - 7.2
<b>131-185%</b>	100	9.1	5.8 - 14.0	<50	4.9	2.8 - 8.4	100	22.1	17.5 - 27.5	200	32.3	27.2 - 38.0	200	27.2	22.1 - 32.9	<50	2.6	1.4 - 4.7
<b>186+%</b>	100	9.0	6.5 - 12.3	<50	4.4	2.7 - 7.0	200	22.8	18.9 - 27.2	300	31.2	26.6 - 36.2	200	27.6	23.3 - 32.5	<50	3.8	2.5 - 5.9

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.8	1.2 - 2.8	2,000
<50	2.1	1.2 - 3.4	1,800

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	1.5	0.9 - 2.5	1,900
<50	2.4	1.6 - 3.6	1,900

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.5	1.7 - 3.8	2,000
<50	1.8	0.9 - 3.6	600
<50	1.1	0.4 - 3.2	800

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

Birthweight	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Low birth weight	<50	5.7	2.4 - 12.6	<50	2.7	0.7 - 10.0	100	25.0	18.8 - 32.5	100	29.9	23.0 - 37.8	100	29.0	21.9 - 37.2	<50	5.6	3.3 - 9.1
Normal birth weight	300	7.8	6.6 - 9.2	100	3.7	2.8 - 4.7	700	21.1	19.2 - 23.1	1,200	34.0	31.8 - 36.3	900	26.7	24.7 - 28.8	200	4.8	4.0 - 5.9

Smoke 3rd Trimester	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	<50	0.8	0.3 - 1.8	<50	2.9	1.7 - 4.7	200	14.7	12.3 - 17.4	500	33.9	30.6 - 37.4	500	37.4	34.0 - 41.0	100	7.9	6.2 - 10.0
No	300	12.1	10.2 - 14.3	100	3.8	2.8 - 5.1	600	25.5	23.0 - 28.1	800	33.6	30.9 - 36.4	500	20.3	18.0 - 22.8	100	3.1	2.3 - 4.0

Smoke Before Pregnant	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
Yes	0	0.0	-	100	3.9	3.0 - 5.0	800	23.0	21.1 - 25.1	1,300	36.5	34.3 - 38.7	1,000	29.2	27.1 - 31.4	200	5.3	4.4 - 6.3
No	300	100.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-	0	0.0	-

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by  
State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income  
(2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s)  
2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.2	0.7 - 7.1	200
100	1.9	1.4 - 2.7	3,500

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.5	1.5 - 3.9	1,400
<50	1.7	1.0 - 2.7	2,200

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
100	2.1	1.5 - 2.9	3,400
0	0.0	-	300

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000-2004**

WIC During Pregnancy	None (0 cigarettes)			Less than 1 cigarette			1 to 5 cigarettes			6 to 10 cigarettes			11 to 20 cigarettes			21 to 40 cigarettes		
	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)	#	%	C.I.(95%)
<b>Yes</b>	100	6.6	5.3 - 8.2	100	2.9	1.9 - 4.2	400	20.0	17.7 - 22.4	800	36.5	33.7 - 39.3	600	27.0	24.5 - 29.7	100	5.0	3.8 - 6.4
<b>No</b>	100	8.4	6.6 - 10.8	100	4.6	3.3 - 6.3	400	23.1	20.3 - 26.2	500	30.4	27.3 - 33.8	400	27.1	24.0 - 30.4	100	4.7	3.6 - 6.1

**PRAMS Health Indicator 5-Year Aggregate Report - Number of cigarettes smoked per avg day before pregnancy- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2004-2011);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000-2004;2001-2005;2002-2006;2003-2007;2004-2008;2005-2009;2006-2010;2007-2011;2008-2012;2009-2013**

*Note: Weighted counts (#) represent 5-yr averages, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

41 cigarettes or more			Weighted Total Responses (5-Yr Avg)
#	%	C.I.(95%)	#
<50	2.2	1.5 - 3.3	2,100
<50	1.7	0.9 - 2.9	1,600