

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

*The trend analysis section shows the trend for the selected health variable over selected years in a table and a graph. Weighted Total Responses: This number is a weighted representation of the mothers of babies delivered in a given year, who responded to a particular question in the survey. For most questions this will be the mothers of all babies delivered in a given year. For some secondary questions only a subset of the population was asked the question. For example, all survey participants were asked if they smoked during pregnancy, but only those who answered yes were asked how many cigarettes a day. If a dimension value is unknown for a particular survey record, that record will not be included in the counts for that particular dimension.*

*Note: Data collection for household income changed in 2004 and again in 2012, so there are three different income dimensions, depending on the year.*

*Below are the indicators included in this report and the long description for each indicator.*

PRAMS Health Indicator	Health Indicator Description
Discuss effects of smoking with HCW during PNC	Discuss effects of smoking with HCW during PNC

*Suggested citation:*

*Hawaii Health Data Warehouse, Hawaii State Department of Health, Pregnancy Risk Assessment Monitoring System, (report title and years), Report Created: 2/14/17*

*For documentation on other terms and coding in this report, please go to the following links:*

[Click here to access PRAMS documentation](#)

[Click here to access DOH-Race Ethnicity coding](#)

[Click here to access The Hawai'i Data Guide](#)

**For questions about this report, please contact  
The Hawai'i Health Data Warehouse**

[profiles@hhdw.org](mailto:profiles@hhdw.org)

**(855) 946-5899 x15**


**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

Year	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
2000	12,100	72.8	70.7 - 74.8	4,500	27.2	25.2 - 29.3	16,700
2001	12,000	73.3	71.2 - 75.3	4,400	26.7	24.7 - 28.8	16,400
2002	12,300	74.1	71.5 - 76.5	4,300	25.9	23.5 - 28.5	16,600
2003	13,000	76.1	73.6 - 78.3	4,100	23.9	21.7 - 26.4	17,100
2004	12,500	72.5	70.5 - 74.4	4,700	27.5	25.6 - 29.5	17,300
2005	12,300	73.0	70.9 - 75.0	4,600	27.0	25.0 - 29.1	16,900
2006	13,500	75.6	73.5 - 77.7	4,400	24.4	22.3 - 26.5	17,900
2007	12,800	71.6	69.4 - 73.7	5,100	28.4	26.3 - 30.6	17,900
2008	12,900	70.6	68.5 - 72.7	5,400	29.4	27.3 - 31.5	18,300
2009	12,900	71.9	69.0 - 74.7	5,000	28.1	25.3 - 31.0	17,900
2010	13,100	72.1	69.1 - 74.9	5,100	27.9	25.1 - 30.9	18,200
2011	13,400	73.6	70.6 - 76.3	4,800	26.4	23.7 - 29.4	18,200
2012	12,800	70.9	67.8 - 73.7	5,300	29.1	26.3 - 32.2	18,100
2013	12,400	68.9	65.6 - 72.1	5,600	31.1	27.9 - 34.4	18,100

## PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- for the State of Hawaii, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,400	68.9	65.6 - 72.1	5,600	31.1	27.9 - 34.4	18,100

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,700	72.3	66.9 - 77.0	600	27.7	23.0 - 33.1	2,300
<b>Honolulu</b>	8,900	68.1	63.6 - 72.3	4,100	31.9	27.7 - 36.4	13,000
<b>Kauai</b>	600	76.2	71.3 - 80.5	200	23.8	19.5 - 28.7	800
<b>Maui</b>	1,300	67.3	61.8 - 72.4	600	32.7	27.6 - 38.2	1,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,000	62.0	55.1 - 68.6	1,800	38.0	31.4 - 44.9	4,800
Native Hawaiian	4,300	79.9	74.7 - 84.2	1,100	20.1	15.8 - 25.3	5,400
Chinese	500	59.6	42.8 - 74.4	400	40.4	25.6 - 57.2	900
Filipino	2,100	70.9	62.4 - 78.1	900	29.1	21.9 - 37.6	3,000
Japanese	900	52.5	40.8 - 64.0	800	47.5	36.0 - 59.2	1,800
Other	1,600	71.2	61.0 - 79.6	600	28.8	20.4 - 39.0	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	700	83.9	69.3 - 92.4	100	16.1	7.6 - 30.7	900
20-24	3,100	77.3	70.0 - 83.2	900	22.7	16.8 - 30.0	4,100
25-34	6,700	68.4	63.8 - 72.7	3,100	31.6	27.3 - 36.2	9,800
35+	1,900	56.5	48.6 - 64.0	1,500	43.5	36.0 - 51.4	3,400

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	5,600	63.4	58.4 - 68.1	3,200	36.6	31.9 - 41.6	8,800
Female	6,900	74.2	69.7 - 78.2	2,400	25.8	21.8 - 30.3	9,300

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,200	63.6	59.1 - 67.8	4,200	36.4	32.2 - 40.9	11,400
No	5,200	78.1	73.2 - 82.3	1,500	21.9	17.7 - 26.8	6,700

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,100	77.0	71.9 - 81.4	1,500	23.0	18.6 - 28.1	6,600
131-185%	1,900	77.8	69.3 - 84.5	500	22.2	15.5 - 30.7	2,500
186+%	4,200	57.7	52.0 - 63.2	3,000	42.3	36.8 - 48.0	7,200

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

Income (2012+)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	3,300	79.5	73.1 - 84.7	800	20.5	15.3 - 26.9	4,100
\$17,001-\$26,000	1,900	78.8	70.3 - 85.3	500	21.2	14.7 - 29.7	2,400
\$26,001-\$51,000	2,700	69.9	62.5 - 76.4	1,200	30.1	23.6 - 37.5	3,900
>=\$51,001	3,500	56.8	50.6 - 62.7	2,700	43.2	37.3 - 49.4	6,100

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	77.2	72.9 - 80.9	300	22.8	19.1 - 27.1	1,300
Normal birth weight	11,400	68.2	64.6 - 71.6	5,300	31.8	28.4 - 35.4	16,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	84.8	71.4 - 92.6	200	15.2	7.4 - 28.6	1,000
No	11,300	67.8	64.3 - 71.2	5,300	32.2	28.8 - 35.7	16,600

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2013**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,600	81.1	73.7 - 86.9	600	18.9	13.1 - 26.3	3,200
<b>No</b>	9,500	66.1	62.2 - 69.7	4,900	33.9	30.3 - 37.8	14,400

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,000	75.9	71.3 - 80.0	1,900	24.1	20.0 - 28.7	7,900
<b>No</b>	6,400	63.4	58.7 - 67.9	3,700	36.6	32.1 - 41.3	10,100



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,800	70.9	67.8 - 73.7	5,300	29.1	26.3 - 32.2	18,100

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	71.6	66.6 - 76.1	600	28.4	23.9 - 33.4	2,300
<b>Honolulu</b>	9,200	70.6	66.5 - 74.3	3,800	29.4	25.7 - 33.5	13,100
<b>Kauai</b>	600	73.3	67.2 - 78.6	200	26.7	21.4 - 32.8	800
<b>Maui</b>	1,400	71.1	66.3 - 75.4	600	28.9	24.6 - 33.7	1,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	64.6	57.9 - 70.7	1,500	35.4	29.3 - 42.1	4,100
Native Hawaiian	4,000	79.6	74.5 - 83.9	1,000	20.4	16.1 - 25.5	5,100
Chinese	600	61.6	47.1 - 74.3	400	38.4	25.7 - 52.9	1,000
Filipino	2,100	65.1	57.4 - 72.2	1,100	34.9	27.8 - 42.6	3,300
Japanese	900	58.3	47.0 - 68.9	700	41.7	31.1 - 53.0	1,600
Other	2,400	80.6	73.2 - 86.3	600	19.4	13.7 - 26.8	3,000

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	800	80.7	66.4 - 89.9	200	19.3	10.1 - 33.6	1,000
20-24	2,900	78.1	71.4 - 83.5	800	21.9	16.5 - 28.6	3,700
25-34	7,200	70.7	66.6 - 74.4	3,000	29.3	25.6 - 33.4	10,200
35+	2,000	60.3	52.6 - 67.5	1,300	39.7	32.5 - 47.4	3,200

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,300	69.7	65.2 - 73.8	2,700	30.3	26.2 - 34.8	9,000
Female	6,600	72.0	67.8 - 75.9	2,500	28.0	24.1 - 32.2	9,100

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,500	67.1	63.1 - 70.9	3,700	32.9	29.1 - 36.9	11,200
No	5,300	77.0	72.3 - 81.2	1,600	23.0	18.8 - 27.7	6,900

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,000	78.2	73.5 - 82.3	1,400	21.8	17.7 - 26.5	6,400
131-185%	2,300	75.3	67.7 - 81.6	800	24.7	18.4 - 32.3	3,100
186+%	4,400	64.0	58.8 - 68.9	2,500	36.0	31.1 - 41.2	6,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

Income (2012+)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$17,000	3,300	82.2	76.3 - 86.8	700	17.8	13.2 - 23.7	4,000
\$17,001-\$26,000	1,500	69.4	60.3 - 77.2	700	30.6	22.8 - 39.7	2,200
\$26,001-\$51,000	3,400	76.1	70.2 - 81.2	1,100	23.9	18.8 - 29.8	4,400
>=\$51,001	3,700	62.4	56.8 - 67.7	2,200	37.6	32.3 - 43.2	6,000

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	69.6	64.3 - 74.5	400	30.4	25.5 - 35.7	1,200
Normal birth weight	11,900	71.0	67.8 - 74.1	4,900	29.0	25.9 - 32.2	16,800

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	700	92.3	80.8 - 97.2	100	7.7	2.8 - 19.2	700
No	11,700	69.8	66.6 - 72.8	5,100	30.2	27.2 - 33.4	16,700

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2012**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,200	80.1	72.8 - 85.8	600	19.9	14.2 - 27.2	2,800
<b>No</b>	10,100	69.0	65.5 - 72.2	4,600	31.0	27.8 - 34.5	14,700

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,400	77.7	73.4 - 81.5	1,800	22.3	18.5 - 26.6	8,200
<b>No</b>	6,400	65.2	60.9 - 69.3	3,400	34.8	30.7 - 39.1	9,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,400	73.6	70.6 - 76.3	4,800	26.4	23.7 - 29.4	18,200

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,900	77.7	73.2 - 81.6	500	22.3	18.4 - 26.8	2,400
<b>Honolulu</b>	9,500	72.7	68.7 - 76.4	3,600	27.3	23.6 - 31.3	13,000
<b>Kauai</b>	600	71.8	65.9 - 77.0	200	28.2	23.0 - 34.1	900
<b>Maui</b>	1,400	74.8	70.6 - 78.6	500	25.2	21.4 - 29.4	1,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	64.8	58.5 - 70.7	1,600	35.2	29.3 - 41.5	4,500
Native Hawaiian	4,200	80.4	75.4 - 84.6	1,000	19.6	15.4 - 24.6	5,300
Chinese	600	81.0	64.6 - 90.9	100	19.0	9.1 - 35.4	700
Filipino	2,700	81.4	74.9 - 86.6	600	18.6	13.4 - 25.1	3,300
Japanese	900	51.8	41.2 - 62.3	900	48.2	37.7 - 58.8	1,800
Other	2,100	77.5	69.0 - 84.3	600	22.5	15.7 - 31.0	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	900	92.2	83.5 - 96.5	100	7.8	3.5 - 16.5	1,000
20-24	3,700	86.0	80.9 - 90.0	600	14.0	10.0 - 19.1	4,300
25-34	7,000	70.9	66.6 - 74.7	2,900	29.1	25.3 - 33.4	9,800
35+	1,800	59.1	51.3 - 66.4	1,300	40.9	33.6 - 48.7	3,100

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,900	74.7	70.6 - 78.5	2,300	25.3	21.5 - 29.4	9,300
Female	6,500	72.4	68.0 - 76.3	2,500	27.6	23.7 - 32.0	8,900

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,700	67.4	63.3 - 71.1	3,700	32.6	28.9 - 36.7	11,400
No	5,700	83.9	79.8 - 87.3	1,100	16.1	12.7 - 20.2	6,800

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,400	84.3	80.1 - 87.7	1,000	15.7	12.3 - 19.9	6,400
131-185%	2,000	73.4	65.2 - 80.3	700	26.6	19.7 - 34.8	2,700
186+%	5,100	63.7	58.8 - 68.3	2,900	36.3	31.7 - 41.2	8,000



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,800	86.8	81.2 - 90.9	400	13.2	9.1 - 18.8	3,300
\$10,000-\$24,999	2,700	80.6	74.1 - 85.8	600	19.4	14.2 - 25.9	3,300
\$25,000-\$49,999	3,500	77.4	71.4 - 82.5	1,000	22.6	17.5 - 28.6	4,500
>=\$50,000	3,600	58.1	52.4 - 63.5	2,600	41.9	36.5 - 47.6	6,200

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	76.2	71.3 - 80.5	300	23.8	19.5 - 28.7	1,300
Normal birth weight	12,400	73.3	70.2 - 76.3	4,500	26.7	23.7 - 29.8	16,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	800	90.4	77.1 - 96.4	100	9.6	3.6 - 22.9	900
No	12,400	72.7	69.6 - 75.6	4,700	27.3	24.4 - 30.4	17,100

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2011**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,100	88.5	83.0 - 92.3	400	11.5	7.7 - 17.0	3,500
<b>No</b>	10,200	70.0	66.6 - 73.3	4,400	30.0	26.7 - 33.4	14,500

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,700	82.6	78.8 - 85.9	1,400	17.4	14.1 - 21.2	8,100
<b>No</b>	6,600	66.2	61.9 - 70.3	3,400	33.8	29.7 - 38.1	10,000

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,100	72.1	69.1 - 74.9	5,100	27.9	25.1 - 30.9	18,200

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,900	81.5	77.3 - 85.1	400	18.5	14.9 - 22.7	2,300
<b>Honolulu</b>	9,300	70.1	66.1 - 73.9	3,900	29.9	26.1 - 33.9	13,200
<b>Kauai</b>	600	73.6	68.1 - 78.5	200	26.4	21.5 - 31.9	800
<b>Maui</b>	1,400	73.3	68.8 - 77.4	500	26.7	22.6 - 31.2	1,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,200	70.5	64.3 - 76.0	1,300	29.5	24.0 - 35.7	4,500
Native Hawaiian	4,300	78.4	73.3 - 82.8	1,200	21.6	17.2 - 26.7	5,400
Chinese	400	56.6	40.3 - 71.6	300	43.4	28.4 - 59.7	800
Filipino	2,200	68.2	60.4 - 75.0	1,000	31.8	25.0 - 39.6	3,300
Japanese	800	48.1	37.7 - 58.8	900	51.9	41.2 - 62.3	1,700
Other	2,300	86.0	78.7 - 91.1	400	14.0	8.9 - 21.3	2,600

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,000	89.0	76.9 - 95.2	100	11.0	4.8 - 23.1	1,100
20-24	3,800	85.4	80.0 - 89.5	600	14.6	10.5 - 20.0	4,400
25-34	6,500	69.0	64.7 - 73.0	2,900	31.0	27.0 - 35.3	9,400
35+	1,900	57.4	49.8 - 64.7	1,400	42.6	35.3 - 50.2	3,300

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,500	71.2	66.9 - 75.1	2,600	28.8	24.9 - 33.1	9,100
Female	6,600	72.9	68.7 - 76.8	2,500	27.1	23.2 - 31.3	9,100

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,900	68.7	64.8 - 72.4	3,600	31.3	27.6 - 35.2	11,500
No	5,300	77.7	72.9 - 81.8	1,500	22.3	18.2 - 27.1	6,800

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,100	82.1	77.5 - 85.9	1,100	17.9	14.1 - 22.5	6,300
131-185%	2,200	74.0	66.1 - 80.6	800	26.0	19.4 - 33.9	3,000
186+%	4,800	61.9	56.9 - 66.6	3,000	38.1	33.4 - 43.1	7,800

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,800	84.2	78.0 - 88.9	500	15.8	11.1 - 22.0	3,300
\$10,000-\$24,999	2,600	83.1	76.4 - 88.1	500	16.9	11.9 - 23.6	3,200
\$25,000-\$49,999	3,200	70.6	64.2 - 76.2	1,400	29.4	23.8 - 35.8	4,600
>=\$50,000	3,700	59.2	53.6 - 64.5	2,500	40.8	35.5 - 46.4	6,200

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	73.6	68.6 - 78.0	300	26.4	22.0 - 31.4	1,300
Normal birth weight	12,200	71.9	68.7 - 74.9	4,800	28.1	25.1 - 31.3	16,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	92.2	82.9 - 96.7	100	7.8	3.3 - 17.1	1,400
No	11,700	70.4	67.2 - 73.4	4,900	29.6	26.6 - 32.8	16,600

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2010**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,100	82.2	75.9 - 87.1	700	17.8	12.9 - 24.1	3,800
<b>No</b>	9,900	69.4	66.0 - 72.7	4,400	30.6	27.3 - 34.0	14,200

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,300	79.1	74.9 - 82.8	1,700	20.9	17.2 - 25.1	8,000
<b>No</b>	6,700	66.2	62.0 - 70.3	3,400	33.8	29.7 - 38.0	10,100

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,900	71.9	69.0 - 74.7	5,000	28.1	25.3 - 31.0	17,900

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,700	73.0	68.3 - 77.2	600	27.0	22.8 - 31.7	2,400
<b>Honolulu</b>	9,000	71.4	67.3 - 75.1	3,600	28.6	24.9 - 32.7	12,600
<b>Kauai</b>	600	76.7	71.3 - 81.4	200	23.3	18.6 - 28.7	800
<b>Maui</b>	1,500	72.0	67.7 - 75.9	600	28.0	24.1 - 32.3	2,100



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	68.6	62.7 - 74.0	1,300	31.4	26.0 - 37.3	4,200
Native Hawaiian	4,600	79.5	74.7 - 83.6	1,200	20.5	16.4 - 25.3	5,800
Chinese	400	50.2	34.2 - 66.2	400	49.8	33.8 - 65.8	800
Filipino	2,300	73.0	65.8 - 79.3	900	27.0	20.7 - 34.2	3,200
Japanese	1,000	65.4	54.5 - 74.8	500	34.6	25.2 - 45.5	1,600
Other	1,600	69.0	59.9 - 76.8	700	31.0	23.2 - 40.1	2,400

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,700	93.5	88.8 - 96.4	100	6.5	3.6 - 11.2	1,800
20-24	3,400	82.7	77.0 - 87.2	700	17.3	12.8 - 23.0	4,200
25-34	6,000	67.0	62.7 - 71.0	3,000	33.0	29.0 - 37.3	9,000
35+	1,800	58.8	51.1 - 66.0	1,200	41.2	34.0 - 48.9	3,000

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,700	70.3	66.1 - 74.1	2,800	29.7	25.9 - 33.9	9,600
Female	6,200	73.8	69.6 - 77.6	2,200	26.2	22.4 - 30.4	8,400

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	6,700	63.4	59.3 - 67.2	3,900	36.6	32.8 - 40.7	10,600
No	6,200	84.3	80.3 - 87.5	1,200	15.7	12.5 - 19.7	7,300

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,100	81.8	77.4 - 85.6	1,100	18.2	14.4 - 22.6	6,300
131-185%	2,200	76.9	69.5 - 82.9	600	23.1	17.1 - 30.5	2,800
186+%	4,200	58.2	53.2 - 62.9	3,000	41.8	37.1 - 46.8	7,200

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,500	85.4	78.6 - 90.3	400	14.6	9.7 - 21.4	3,000
\$10,000-\$24,999	3,200	83.5	78.2 - 87.7	600	16.5	12.3 - 21.8	3,800
\$25,000-\$49,999	2,500	69.1	62.4 - 75.1	1,100	30.9	24.9 - 37.6	3,700
>=\$50,000	3,400	56.1	50.6 - 61.5	2,600	43.9	38.5 - 49.4	6,000

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	75.0	70.2 - 79.3	300	25.0	20.7 - 29.8	1,300
Normal birth weight	11,800	71.6	68.4 - 74.5	4,700	28.4	25.5 - 31.6	16,600

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,600	94.3	89.5 - 97.0	100	5.7	3.0 - 10.5	1,600
No	11,000	69.5	66.3 - 72.5	4,800	30.5	27.5 - 33.7	15,800

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2009**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,400	89.0	84.7 - 92.2	400	11.0	7.8 - 15.3	3,900
<b>No</b>	9,100	66.9	63.4 - 70.3	4,500	33.1	29.7 - 36.6	13,600

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,900	80.8	76.8 - 84.3	1,400	19.2	15.7 - 23.2	7,300
<b>No</b>	6,600	65.7	61.5 - 69.6	3,500	34.3	30.4 - 38.5	10,100

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,900	70.6	68.5 - 72.7	5,400	29.4	27.3 - 31.5	18,300

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,800	75.8	69.6 - 81.1	600	24.2	18.9 - 30.4	2,400
<b>Honolulu</b>	9,000	70.2	67.6 - 72.7	3,800	29.8	27.3 - 32.4	12,800
<b>Kauai</b>	600	65.3	54.4 - 74.7	300	34.7	25.3 - 45.6	900
<b>Maui</b>	1,500	69.5	62.6 - 75.6	700	30.5	24.4 - 37.4	2,100

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	66.1	61.4 - 70.6	1,500	33.9	29.4 - 38.6	4,300
Native Hawaiian	3,700	72.3	67.8 - 76.5	1,400	27.7	23.5 - 32.2	5,200
Chinese	400	60.1	53.4 - 66.5	200	39.9	33.5 - 46.6	600
Filipino	2,600	77.3	72.6 - 81.4	800	22.7	18.6 - 27.4	3,400
Japanese	1,200	57.8	50.8 - 64.5	900	42.2	35.5 - 49.2	2,000
Other	2,200	78.2	72.9 - 82.7	600	21.8	17.3 - 27.1	2,800

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,200	87.1	79.8 - 92.0	200	12.9	8.0 - 20.2	1,400
20-24	3,200	79.1	74.5 - 83.1	800	20.9	16.9 - 25.5	4,000
25-34	6,400	67.1	64.0 - 70.1	3,100	32.9	29.9 - 36.0	9,500
35+	2,100	63.6	58.4 - 68.5	1,200	36.4	31.5 - 41.6	3,300

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,500	68.0	64.9 - 71.0	3,100	32.0	29.0 - 35.1	9,600
Female	6,400	73.5	70.4 - 76.4	2,300	26.5	23.6 - 29.6	8,700

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,700	66.4	63.7 - 69.1	3,900	33.6	30.9 - 36.3	11,500
No	5,200	77.9	74.3 - 81.0	1,500	22.1	19.0 - 25.7	6,700

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,100	76.4	72.8 - 79.7	1,600	23.6	20.3 - 27.2	6,700
131-185%	1,900	72.7	66.6 - 78.0	700	27.3	22.0 - 33.4	2,600
186+%	4,900	63.5	60.1 - 66.8	2,800	36.5	33.2 - 39.9	7,800

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,300	78.8	73.3 - 83.4	600	21.2	16.6 - 26.7	2,900
\$10,000-\$24,999	3,000	75.8	71.0 - 80.1	1,000	24.2	19.9 - 29.0	4,000
\$25,000-\$49,999	3,000	71.7	67.1 - 75.9	1,200	28.3	24.1 - 32.9	4,200
>=\$50,000	3,800	61.1	57.2 - 64.8	2,400	38.9	35.2 - 42.8	6,200

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	69.1	60.0 - 77.0	400	30.9	23.0 - 40.0	1,200
Normal birth weight	12,000	70.8	68.5 - 73.0	5,000	29.2	27.0 - 31.5	17,000

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	88.4	81.7 - 92.8	200	11.6	7.2 - 18.3	1,500
No	11,300	68.8	66.5 - 71.1	5,100	31.2	28.9 - 33.5	16,400



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2008**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,800	84.4	79.9 - 88.0	500	15.6	12.0 - 20.1	3,400
<b>No</b>	9,700	67.2	64.7 - 69.6	4,800	32.8	30.4 - 35.3	14,500

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,200	78.9	75.6 - 81.8	1,700	21.1	18.2 - 24.4	7,800
<b>No</b>	6,700	64.5	61.5 - 67.4	3,700	35.5	32.6 - 38.5	10,400

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,800	71.6	69.4 - 73.7	5,100	28.4	26.3 - 30.6	17,900

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	64.2	57.5 - 70.4	900	35.8	29.6 - 42.5	2,400
<b>Honolulu</b>	9,100	73.4	70.8 - 75.8	3,300	26.6	24.2 - 29.2	12,400
<b>Kauai</b>	800	75.3	65.9 - 82.9	300	24.7	17.1 - 34.1	1,100
<b>Maui</b>	1,400	67.7	60.6 - 74.0	700	32.3	26.0 - 39.4	2,100

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,900	70.1	65.3 - 74.4	1,200	29.9	25.6 - 34.7	4,100
Native Hawaiian	3,900	79.1	74.8 - 82.8	1,000	20.9	17.2 - 25.2	4,900
Chinese	400	59.4	52.6 - 65.9	300	40.6	34.1 - 47.4	600
Filipino	2,300	68.8	63.8 - 73.4	1,000	31.2	26.6 - 36.2	3,300
Japanese	1,200	57.7	50.7 - 64.3	900	42.3	35.7 - 49.3	2,100
Other	2,200	77.0	71.5 - 81.8	700	23.0	18.2 - 28.5	2,900

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,500	87.5	80.4 - 92.3	200	12.5	7.7 - 19.6	1,700
20-24	3,400	76.9	72.4 - 80.8	1,000	23.1	19.2 - 27.6	4,400
25-34	6,100	69.3	66.2 - 72.3	2,700	30.7	27.7 - 33.8	8,800
35+	1,900	61.7	56.3 - 66.9	1,200	38.3	33.1 - 43.7	3,100

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	6,500	70.5	67.5 - 73.5	2,700	29.5	26.5 - 32.5	9,200
<b>Female</b>	6,300	72.7	69.5 - 75.6	2,400	27.3	24.4 - 30.5	8,700

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	7,500	67.0	64.2 - 69.7	3,700	33.0	30.3 - 35.8	11,200
<b>No</b>	5,300	79.3	75.8 - 82.4	1,400	20.7	17.6 - 24.2	6,700

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	5,100	76.9	73.3 - 80.1	1,500	23.1	19.9 - 26.7	6,700
<b>131-185%</b>	1,800	69.4	63.5 - 74.8	800	30.6	25.2 - 36.5	2,600
<b>186+%</b>	4,900	66.8	63.4 - 70.1	2,400	33.2	29.9 - 36.6	7,300

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,400	80.5	75.2 - 85.0	600	19.5	15.0 - 24.8	3,000
\$10,000-\$24,999	2,900	77.6	72.9 - 81.7	800	22.4	18.3 - 27.1	3,800
\$25,000-\$49,999	2,800	70.2	65.4 - 74.6	1,200	29.8	25.4 - 34.6	4,000
>=\$50,000	3,800	63.6	59.7 - 67.3	2,200	36.4	32.7 - 40.3	5,900

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,100	73.6	66.1 - 80.0	400	26.4	20.0 - 33.9	1,500
Normal birth weight	11,700	71.4	69.1 - 73.6	4,700	28.6	26.4 - 30.9	16,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	89.4	82.8 - 93.6	200	10.6	6.4 - 17.2	1,500
No	11,400	70.2	67.9 - 72.5	4,800	29.8	27.5 - 32.1	16,200

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2007**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,900	83.6	79.0 - 87.3	600	16.4	12.7 - 21.0	3,500
<b>No</b>	9,800	68.9	66.4 - 71.3	4,400	31.1	28.7 - 33.6	14,200

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,800	77.2	73.9 - 80.3	1,700	22.8	19.7 - 26.1	7,600
<b>No</b>	6,900	67.2	64.3 - 70.0	3,400	32.8	30.0 - 35.7	10,200

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,500	75.6	73.5 - 77.7	4,400	24.4	22.3 - 26.5	17,900

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,600	75.8	69.1 - 81.4	500	24.2	18.6 - 30.9	2,100
<b>Honolulu</b>	9,800	75.4	72.9 - 77.7	3,200	24.6	22.3 - 27.1	13,000
<b>Kauai</b>	700	79.0	68.2 - 86.9	200	21.0	13.1 - 31.8	900
<b>Maui</b>	1,400	75.7	68.6 - 81.6	500	24.3	18.4 - 31.4	1,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	3,000	71.2	66.5 - 75.5	1,200	28.8	24.5 - 33.5	4,300
Native Hawaiian	3,900	80.1	75.8 - 83.9	1,000	19.9	16.1 - 24.2	4,900
Chinese	400	62.4	55.2 - 69.1	200	37.6	30.9 - 44.8	600
Filipino	2,700	81.3	76.7 - 85.2	600	18.7	14.8 - 23.3	3,300
Japanese	1,300	63.2	56.2 - 69.6	800	36.8	30.4 - 43.8	2,100
Other	2,200	80.0	74.5 - 84.6	500	20.0	15.4 - 25.5	2,700

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	87.7	80.2 - 92.6	200	12.3	7.4 - 19.8	1,500
20-24	3,400	78.0	73.4 - 82.0	1,000	22.0	18.0 - 26.6	4,400
25-34	6,700	76.7	73.7 - 79.5	2,000	23.3	20.5 - 26.3	8,700
35+	2,100	63.9	58.5 - 68.9	1,200	36.1	31.1 - 41.5	3,200



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	6,900	76.1	73.1 - 78.8	2,200	23.9	21.2 - 26.9	9,000
<b>Female</b>	6,600	75.2	72.1 - 78.1	2,200	24.8	21.9 - 27.9	8,800

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	8,100	71.1	68.4 - 73.7	3,300	28.9	26.3 - 31.6	11,400
<b>No</b>	5,400	83.6	80.1 - 86.6	1,100	16.4	13.4 - 19.9	6,500

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	5,100	83.0	79.5 - 86.0	1,100	17.0	14.0 - 20.5	6,200
<b>131-185%</b>	1,900	78.9	72.6 - 84.0	500	21.1	16.0 - 27.4	2,400
<b>186+%</b>	5,500	68.3	65.0 - 71.5	2,500	31.7	28.5 - 35.0	8,000

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,400	81.5	76.1 - 86.0	500	18.5	14.0 - 23.9	2,900
\$10,000-\$24,999	3,100	83.2	78.8 - 86.9	600	16.8	13.1 - 21.2	3,700
\$25,000-\$49,999	3,000	72.2	67.4 - 76.5	1,200	27.8	23.5 - 32.6	4,200
>=\$50,000	4,000	69.2	65.3 - 72.8	1,800	30.8	27.2 - 34.7	5,800

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,100	79.5	72.0 - 85.4	300	20.5	14.6 - 28.0	1,400
Normal birth weight	12,400	75.3	73.1 - 77.4	4,100	24.7	22.6 - 26.9	16,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,500	93.7	88.0 - 96.7	100	6.3	3.3 - 12.0	1,600
No	11,800	73.9	71.6 - 76.1	4,200	26.1	23.9 - 28.4	15,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2006**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,200	87.3	83.0 - 90.6	500	12.7	9.4 - 17.0	3,700
<b>No</b>	10,100	72.7	70.2 - 75.0	3,800	27.3	25.0 - 29.8	13,900

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,100	80.8	77.5 - 83.7	1,500	19.2	16.3 - 22.5	7,600
<b>No</b>	7,300	71.8	68.9 - 74.5	2,900	28.2	25.5 - 31.1	10,200

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,300	73.0	70.9 - 75.0	4,600	27.0	25.0 - 29.1	16,900

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,700	78.6	72.5 - 83.7	500	21.4	16.3 - 27.5	2,100
<b>Honolulu</b>	8,900	71.6	69.0 - 74.0	3,500	28.4	26.0 - 31.0	12,400
<b>Kauai</b>	600	79.7	69.0 - 87.3	100	20.3	12.7 - 31.0	700
<b>Maui</b>	1,200	73.4	66.2 - 79.6	400	26.6	20.4 - 33.8	1,700

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,400	65.6	60.5 - 70.3	1,300	34.4	29.7 - 39.5	3,700
Native Hawaiian	4,000	83.2	79.2 - 86.5	800	16.8	13.5 - 20.8	4,800
Chinese	400	68.1	61.7 - 73.8	200	31.9	26.2 - 38.3	600
Filipino	2,400	75.8	71.1 - 80.0	800	24.2	20.0 - 28.9	3,100
Japanese	1,300	56.6	50.2 - 62.8	1,000	43.4	37.2 - 49.8	2,200
Other	1,900	76.8	71.3 - 81.6	600	23.2	18.4 - 28.7	2,500

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	87.1	80.7 - 91.6	200	12.9	8.4 - 19.3	1,500
20-24	3,400	81.5	77.4 - 84.9	800	18.5	15.1 - 22.6	4,200
25-34	5,800	70.2	67.0 - 73.2	2,500	29.8	26.8 - 33.0	8,300
35+	1,800	61.3	55.8 - 66.5	1,100	38.7	33.5 - 44.2	2,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,400	74.4	71.5 - 77.2	2,200	25.6	22.8 - 28.5	8,600
Female	5,900	71.5	68.4 - 74.4	2,400	28.5	25.6 - 31.6	8,300

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,300	67.8	65.0 - 70.4	3,500	32.2	29.6 - 35.0	10,800
No	5,000	82.2	78.9 - 85.1	1,100	17.8	14.9 - 21.1	6,100

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	4,800	78.8	75.4 - 81.9	1,300	21.2	18.1 - 24.6	6,100
131-185%	1,300	74.3	67.3 - 80.3	500	25.7	19.7 - 32.7	1,800
186+%	5,100	66.4	63.0 - 69.5	2,600	33.6	30.5 - 37.0	7,600

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,400	82.8	78.0 - 86.8	500	17.2	13.2 - 22.0	2,900
\$10,000-\$24,999	2,700	76.6	71.9 - 80.7	800	23.4	19.3 - 28.1	3,500
\$25,000-\$49,999	3,000	74.8	70.2 - 78.9	1,000	25.2	21.1 - 29.8	4,000
>=\$50,000	3,300	61.9	57.9 - 65.8	2,000	38.1	34.2 - 42.1	5,300

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	76.5	68.0 - 83.2	300	23.5	16.8 - 32.0	1,200
Normal birth weight	11,400	72.7	70.5 - 74.8	4,300	27.3	25.2 - 29.5	15,700

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,300	96.1	91.3 - 98.3	100	3.9	1.7 - 8.7	1,300
No	10,800	70.9	68.6 - 73.1	4,400	29.1	26.9 - 31.4	15,200

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2005**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,000	85.7	81.5 - 89.0	500	14.3	11.0 - 18.5	3,500
<b>No</b>	9,100	69.5	67.0 - 71.9	4,000	30.5	28.1 - 33.0	13,100

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,900	80.1	77.0 - 82.9	1,500	19.9	17.1 - 23.0	7,300
<b>No</b>	6,300	67.4	64.4 - 70.3	3,100	32.6	29.7 - 35.6	9,400



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,500	72.5	70.5 - 74.4	4,700	27.5	25.6 - 29.5	17,300

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	71.6	65.5 - 77.0	600	28.4	23.0 - 34.5	2,100
<b>Honolulu</b>	9,100	72.6	70.3 - 74.8	3,400	27.4	25.2 - 29.7	12,500
<b>Kauai</b>	600	74.3	64.4 - 82.2	200	25.7	17.8 - 35.6	800
<b>Maui</b>	1,300	71.7	65.3 - 77.3	500	28.3	22.7 - 34.7	1,800

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	68.3	63.9 - 72.4	1,200	31.7	27.6 - 36.1	3,900
Native Hawaiian	3,800	78.7	74.7 - 82.2	1,000	21.3	17.8 - 25.3	4,800
Chinese	400	60.9	56.2 - 65.3	200	39.1	34.7 - 43.8	600
Filipino	2,400	76.1	71.9 - 79.8	800	23.9	20.2 - 28.1	3,200
Japanese	1,300	56.3	50.0 - 62.4	1,000	43.7	37.6 - 50.0	2,200
Other	2,000	79.8	74.7 - 84.0	500	20.2	16.0 - 25.3	2,500

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,100	85.7	78.7 - 90.7	200	14.3	9.3 - 21.3	1,300
20-24	3,500	79.8	75.9 - 83.2	900	20.2	16.8 - 24.1	4,400
25-34	5,900	70.3	67.4 - 73.0	2,500	29.7	27.0 - 32.6	8,400
35+	2,000	63.1	58.3 - 67.7	1,200	36.9	32.3 - 41.7	3,200

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,900	72.8	70.1 - 75.3	2,600	27.2	24.7 - 29.9	9,500
Female	5,600	72.1	69.1 - 74.9	2,200	27.9	25.1 - 30.9	7,700

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,400	69.9	67.5 - 72.2	3,600	30.1	27.8 - 32.5	12,100
No	4,100	78.6	74.9 - 81.8	1,100	21.4	18.2 - 25.1	5,200

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,100	79.4	76.2 - 82.2	1,300	20.6	17.8 - 23.8	6,400
131-185%	1,500	73.8	67.9 - 79.0	500	26.2	21.0 - 32.1	2,000
186+%	4,900	65.7	62.6 - 68.7	2,600	34.3	31.3 - 37.4	7,500

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

Income (2004-2011)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<\$10,000	2,200	82.0	77.1 - 86.1	500	18.0	13.9 - 22.9	2,600
\$10,000-\$24,999	3,300	78.5	74.5 - 82.0	900	21.5	18.0 - 25.5	4,300
\$25,000-\$49,999	3,100	72.5	68.4 - 76.2	1,200	27.5	23.8 - 31.6	4,300
>=\$50,000	2,900	61.2	57.2 - 64.9	1,900	38.8	35.1 - 42.8	4,800

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	74.9	66.7 - 81.6	300	25.1	18.4 - 33.3	1,100
Normal birth weight	11,700	72.3	70.3 - 74.3	4,500	27.7	25.7 - 29.7	16,200

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,200	88.7	82.5 - 92.9	200	11.3	7.1 - 17.5	1,400
No	11,100	70.8	68.7 - 72.9	4,600	29.2	27.1 - 31.3	15,600

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2004**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,900	83.9	79.8 - 87.2	600	16.1	12.8 - 20.2	3,500
<b>No</b>	9,300	69.2	66.9 - 71.5	4,100	30.8	28.5 - 33.1	13,500

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,100	78.8	76.0 - 81.4	1,600	21.2	18.6 - 24.0	7,800
<b>No</b>	6,300	67.5	64.8 - 70.2	3,000	32.5	29.8 - 35.2	9,300

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	13,000	76.1	73.6 - 78.3	4,100	23.9	21.7 - 26.4	17,100

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	77.4	73.2 - 81.1	400	22.6	18.9 - 26.8	1,900
<b>Honolulu</b>	9,400	75.2	71.9 - 78.1	3,100	24.8	21.9 - 28.1	12,500
<b>Kauai</b>	700	84.3	78.3 - 88.9	100	15.7	11.1 - 21.7	800
<b>Maui</b>	1,400	77.1	72.7 - 81.0	400	22.9	19.0 - 27.3	1,800

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,800	75.1	69.3 - 80.1	900	24.9	19.9 - 30.7	3,700
Native Hawaiian	3,800	80.5	78.0 - 82.8	900	19.5	17.2 - 22.0	4,700
Chinese	500	71.1	54.7 - 83.4	200	28.9	16.6 - 45.3	700
Filipino	2,900	76.2	70.3 - 81.3	900	23.8	18.7 - 29.7	3,800
Japanese	1,200	61.9	52.7 - 70.3	800	38.1	29.7 - 47.3	2,000
Other	1,800	82.2	74.5 - 87.9	400	17.8	12.1 - 25.5	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,400	89.8	84.3 - 93.6	200	10.2	6.4 - 15.7	1,600
20-24	3,400	85.5	81.3 - 88.8	600	14.5	11.2 - 18.7	4,000
25-34	6,300	72.4	68.7 - 75.8	2,400	27.6	24.2 - 31.3	8,600
35+	1,900	66.5	59.6 - 72.7	1,000	33.5	27.3 - 40.4	2,900

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	7,000	77.4	74.1 - 80.4	2,000	22.6	19.6 - 25.9	9,000
Female	6,000	74.5	70.8 - 78.0	2,100	25.5	22.0 - 29.2	8,100

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,300	73.1	69.8 - 76.1	3,000	26.9	23.9 - 30.2	11,300
No	4,800	81.8	78.3 - 84.9	1,100	18.2	15.1 - 21.7	5,800

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	6,100	80.0	76.5 - 83.1	1,500	20.0	16.9 - 23.5	7,600
131-185%	1,800	74.3	67.4 - 80.2	600	25.7	19.8 - 32.6	2,500
186+%	3,800	69.3	64.4 - 73.8	1,700	30.7	26.2 - 35.6	5,400



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

Income (2000-2003)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	3,400	83.2	78.5 - 87.0	700	16.8	13.0 - 21.5	4,100
\$15,001-\$25,000	1,700	77.6	70.8 - 83.2	500	22.4	16.8 - 29.2	2,200
\$25,001-\$40,000	2,400	75.4	69.4 - 80.5	800	24.6	19.5 - 30.6	3,200
>=\$40,001	4,500	69.9	65.5 - 74.0	1,900	30.1	26.0 - 34.5	6,500

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	1,000	78.7	68.9 - 86.0	300	21.3	14.0 - 31.1	1,200
Normal birth weight	12,100	75.9	73.3 - 78.2	3,800	24.1	21.8 - 26.7	15,900

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	88.2	81.0 - 92.9	200	11.8	7.1 - 19.0	1,500
No	11,400	74.8	72.1 - 77.2	3,900	25.2	22.8 - 27.9	15,300

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2003**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	3,000	86.8	82.3 - 90.3	500	13.2	9.7 - 17.7	3,500
<b>No</b>	9,700	73.5	70.6 - 76.2	3,500	26.5	23.8 - 29.4	13,200

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	6,000	81.6	78.1 - 84.5	1,400	18.4	15.5 - 21.9	7,300
<b>No</b>	6,900	71.9	68.4 - 75.2	2,700	28.1	24.8 - 31.6	9,700

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,300	74.1	71.5 - 76.5	4,300	25.9	23.5 - 28.5	16,600

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,500	78.3	74.3 - 81.8	400	21.7	18.2 - 25.7	1,900
<b>Honolulu</b>	9,000	73.5	70.1 - 76.6	3,200	26.5	23.4 - 29.9	12,200
<b>Kauai</b>	500	73.2	65.6 - 79.7	200	26.8	20.3 - 34.4	700
<b>Maui</b>	1,300	74.0	69.4 - 78.2	400	26.0	21.8 - 30.6	1,700

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,400	67.5	61.1 - 73.3	1,200	32.5	26.7 - 38.9	3,500
Native Hawaiian	3,600	79.2	76.6 - 81.6	900	20.8	18.4 - 23.4	4,600
Chinese	400	62.8	45.8 - 77.1	200	37.2	22.9 - 54.2	600
Filipino	2,700	81.0	75.1 - 85.8	600	19.0	14.2 - 24.9	3,400
Japanese	1,300	55.0	46.5 - 63.2	1,000	45.0	36.8 - 53.5	2,300
Other	1,900	87.1	80.2 - 91.8	300	12.9	8.2 - 19.8	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,300	84.3	76.1 - 90.0	200	15.7	10.0 - 23.9	1,500
20-24	3,400	83.1	78.9 - 86.6	700	16.9	13.4 - 21.1	4,100
25-34	5,900	72.1	68.3 - 75.6	2,300	27.9	24.4 - 31.7	8,200
35+	1,700	61.0	53.7 - 67.8	1,100	39.0	32.2 - 46.3	2,700

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Male</b>	6,200	76.0	72.5 - 79.2	2,000	24.0	20.8 - 27.5	8,200
<b>Female</b>	6,000	72.2	68.5 - 75.7	2,300	27.8	24.3 - 31.5	8,400

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	7,900	69.1	65.8 - 72.3	3,500	30.9	27.7 - 34.2	11,400
<b>No</b>	4,400	84.9	81.5 - 87.8	800	15.1	12.2 - 18.5	5,200

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>0-130%</b>	5,200	81.3	77.7 - 84.4	1,200	18.7	15.6 - 22.3	6,400
<b>131-185%</b>	2,500	74.1	68.1 - 79.3	900	25.9	20.7 - 31.9	3,400
<b>186+%</b>	3,400	63.7	58.6 - 68.5	1,900	36.3	31.5 - 41.4	5,300

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

Income (2000-2003)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,600	81.0	76.2 - 85.1	600	19.0	14.9 - 23.8	3,200
\$15,001-\$25,000	2,000	84.5	78.7 - 88.9	400	15.5	11.1 - 21.3	2,400
\$25,001-\$40,000	2,700	74.5	68.5 - 79.7	900	25.5	20.3 - 31.5	3,600
>=\$40,001	4,100	64.7	60.1 - 69.1	2,200	35.3	30.9 - 39.9	6,400

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	73.2	61.5 - 82.4	300	26.8	17.6 - 38.5	1,100
Normal birth weight	11,500	74.2	71.6 - 76.7	4,000	25.8	23.3 - 28.4	15,400

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	90.2	84.0 - 94.1	100	9.8	5.9 - 16.0	1,300
No	10,900	72.7	70.0 - 75.3	4,100	27.3	24.7 - 30.0	15,000

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2002**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,700	86.1	81.3 - 89.9	400	13.9	10.1 - 18.7	3,100
<b>No</b>	9,200	70.8	67.8 - 73.7	3,800	29.2	26.3 - 32.2	13,000

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,800	83.2	79.9 - 86.1	1,200	16.8	13.9 - 20.1	7,000
<b>No</b>	6,400	67.2	63.5 - 70.7	3,100	32.8	29.3 - 36.5	9,500

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,000	73.3	71.2 - 75.3	4,400	26.7	24.7 - 28.8	16,400

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,400	76.0	71.7 - 79.8	400	24.0	20.2 - 28.3	1,800
<b>Honolulu</b>	8,800	72.8	70.2 - 75.3	3,300	27.2	24.7 - 29.8	12,100
<b>Kauai</b>	600	75.7	68.7 - 81.5	200	24.3	18.5 - 31.3	800
<b>Maui</b>	1,200	73.1	68.2 - 77.5	400	26.9	22.5 - 31.8	1,700



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,300	67.2	62.4 - 71.7	1,100	32.8	28.3 - 37.6	3,500
Native Hawaiian	3,600	80.7	78.7 - 82.5	900	19.3	17.5 - 21.3	4,400
Chinese	500	66.0	52.8 - 77.1	300	34.0	22.9 - 47.2	700
Filipino	2,700	77.9	72.9 - 82.3	800	22.1	17.7 - 27.1	3,500
Japanese	1,100	55.0	47.5 - 62.4	900	45.0	37.6 - 52.5	2,000
Other	1,800	80.4	74.0 - 85.5	400	19.6	14.5 - 26.0	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,400	88.0	83.9 - 91.2	200	12.0	8.8 - 16.1	1,600
20-24	3,400	79.7	76.0 - 83.0	900	20.3	17.0 - 24.0	4,200
25-34	5,800	72.9	69.9 - 75.8	2,200	27.1	24.2 - 30.1	7,900
35+	1,500	55.8	49.7 - 61.7	1,200	44.2	38.3 - 50.3	2,600

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,100	73.1	70.1 - 75.9	2,200	26.9	24.1 - 29.9	8,300
Female	5,900	73.6	70.6 - 76.4	2,100	26.4	23.6 - 29.4	8,100

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	7,600	69.4	66.6 - 72.0	3,400	30.6	28.0 - 33.4	11,000
No	4,400	81.5	78.4 - 84.2	1,000	18.5	15.8 - 21.6	5,400

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,000	81.4	78.6 - 84.0	1,100	18.6	16.0 - 21.4	6,100
131-185%	2,600	73.7	69.3 - 77.7	900	26.3	22.3 - 30.7	3,600
186+%	3,400	62.8	58.6 - 66.7	2,000	37.2	33.3 - 41.4	5,500

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

Income (2000-2003)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	2,700	84.1	80.6 - 87.1	500	15.9	12.9 - 19.4	3,300
\$15,001-\$25,000	1,800	79.8	74.8 - 84.0	500	20.2	16.0 - 25.2	2,200
\$25,001-\$40,000	2,700	74.7	70.3 - 78.6	900	25.3	21.4 - 29.7	3,500
>=\$40,001	4,100	63.7	59.9 - 67.3	2,300	36.3	32.7 - 40.1	6,500

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	900	79.9	72.1 - 85.9	200	20.1	14.1 - 27.9	1,100
Normal birth weight	11,100	72.9	70.7 - 75.0	4,100	27.1	25.0 - 29.3	15,300

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,400	89.4	83.5 - 93.3	200	10.6	6.7 - 16.5	1,500
No	10,400	71.7	69.4 - 73.9	4,100	28.3	26.1 - 30.6	14,500

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2001**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,700	82.0	77.9 - 85.5	600	18.0	14.5 - 22.1	3,300
<b>No</b>	8,800	70.7	68.2 - 73.1	3,700	29.3	26.9 - 31.8	12,500

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,500	80.3	77.5 - 82.8	1,400	19.7	17.2 - 22.5	6,900
<b>No</b>	6,400	68.4	65.4 - 71.3	3,000	31.6	28.7 - 34.6	9,400

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

State	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Totals</b>	12,100	72.8	70.7 - 74.8	4,500	27.2	25.2 - 29.3	16,700

County	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Hawaii</b>	1,400	72.2	67.8 - 76.2	500	27.8	23.8 - 32.2	1,900
<b>Honolulu</b>	8,900	71.8	69.2 - 74.3	3,500	28.2	25.7 - 30.8	12,400
<b>Kauai</b>	600	83.1	76.1 - 88.4	100	16.9	11.6 - 23.9	700
<b>Maui</b>	1,300	76.7	72.1 - 80.8	400	23.3	19.2 - 27.9	1,700

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

DOH Race-Ethnicity	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Caucasian	2,700	67.5	62.9 - 71.7	1,300	32.5	28.3 - 37.1	4,000
Native Hawaiian	3,600	79.2	77.1 - 81.1	900	20.8	18.9 - 22.9	4,500
Chinese	500	71.0	57.2 - 81.8	200	29.0	18.2 - 42.8	600
Filipino	2,500	74.6	69.4 - 79.2	800	25.4	20.8 - 30.6	3,300
Japanese	1,200	61.2	53.7 - 68.2	800	38.8	31.8 - 46.3	2,000
Other	1,700	77.8	71.1 - 83.3	500	22.2	16.7 - 28.9	2,200

PRAMS Age Group	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<20	1,600	85.5	80.2 - 89.6	300	14.5	10.4 - 19.8	1,800
20-24	3,100	78.3	74.5 - 81.6	900	21.7	18.4 - 25.5	4,000
25-34	5,600	70.0	66.9 - 73.0	2,400	30.0	27.0 - 33.1	8,000
35+	1,800	64.8	59.2 - 70.1	1,000	35.2	29.9 - 40.8	2,800

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

Infant Gender	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Male	6,400	74.0	71.2 - 76.6	2,300	26.0	23.4 - 28.8	8,700
Female	5,700	71.5	68.4 - 74.4	2,300	28.5	25.6 - 31.6	8,000

Married	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	8,000	69.2	66.5 - 71.7	3,600	30.8	28.3 - 33.5	11,500
No	4,200	80.9	77.9 - 83.6	1,000	19.1	16.4 - 22.1	5,100

Poverty Level	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
0-130%	5,500	77.6	74.6 - 80.3	1,600	22.4	19.7 - 25.4	7,000
131-185%	1,800	78.5	73.5 - 82.7	500	21.5	17.3 - 26.5	2,300
186+%	4,000	65.3	61.5 - 68.9	2,100	34.7	31.1 - 38.5	6,100

**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

Income (2000-2003)	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<=\$15,000	3,100	80.8	76.9 - 84.2	700	19.2	15.8 - 23.1	3,800
\$15,001-\$25,000	1,900	72.0	66.5 - 77.0	700	28.0	23.0 - 33.5	2,600
\$25,001-\$40,000	2,500	76.4	71.9 - 80.4	800	23.6	19.6 - 28.1	3,200
>=\$40,001	4,100	66.2	62.5 - 69.7	2,100	33.8	30.3 - 37.5	6,200

Birthweight	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Low birth weight	800	80.1	71.9 - 86.4	200	19.9	13.6 - 28.1	1,100
Normal birth weight	11,200	72.1	70.0 - 74.2	4,300	27.9	25.8 - 30.0	15,500

Smoke 3rd Trimester	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
Yes	1,100	85.7	79.5 - 90.2	200	14.3	9.8 - 20.5	1,300
No	10,600	71.7	69.4 - 73.8	4,200	28.3	26.2 - 30.6	14,800



**PRAMS Health Indicator Report - Discuss effects of smoking with HCW during PNC- by State;County;DOH Race-Ethnicity;PRAMS Age Group;Infant Gender;Married;Poverty Level;Income (2000-2003);Income (2004-2011);Income (2012+);Birthweight;Smoke 3rd Trimester;Smoke Before Pregnant;WIC During Pregnancy, for the Year(s) 2000;2001;2002;2003;2004;2005;2006;2007;2008;2009;2010;2011;2012;2013**

*Note: Weighted responses, rounded to the nearest hundred; n/r (Not Reportable) - If the unweighted total responses for a dimension (row) is <30 that row will not be shown*

**Year: 2000**

Smoke Before Pregnant	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	2,700	84.8	81.0 - 87.9	500	15.2	12.1 - 19.0	3,200
<b>No</b>	9,000	69.8	67.3 - 72.1	3,900	30.2	27.9 - 32.7	12,800

WIC During Pregnancy	Yes			No			Weighted Total Responses
	#	%	C.I.(95%)	#	%	C.I.(95%)	#
<b>Yes</b>	5,600	78.1	75.2 - 80.7	1,600	21.9	19.3 - 24.8	7,200
<b>No</b>	6,400	69.0	66.0 - 71.8	2,900	31.0	28.2 - 34.0	9,200